Condition	**Continuous high rainfall in a short span leading to water logging**			
	Suggested contingency measure			
1	**2**	**3**	**4**	**5**
Outbreak of pests and diseases due to unseasonal rains				
Crop1 Soybean	• Early planting to minimize the incidence of girdle beetle and green semilooper • Foliar spray with 5% NSKE or dimethoate 30EC 1 ml/l to protect against semilooper	• Monitor adult moth activity of Spodoptera through pheromone traps (10 traps/ha) • Apply Quinalphos 25 EC 2ml/l or Emamectin benzoate 5 SG 4g/10 lit to control spodoptera		
Crop2 Jowar	Whorl application of phorate 10G or carbofuran 3 G @ 8-10 kg/ha to control shoot borer attack -	Spray of mancozeb @ 0.25-0.4% at 8-10 days interval to control leaf blight	Trichoderma mixed with FYM @10g/kg at 10 days prior to its use in the field can be applied to control stalk rot incidence which is likely	

Condition	**Continuous high rainfall in a short span leading to water logging**			
	Suggested contingency measure			
1	**2**	**3**	**4**	**5**
			during post flowering	
Crop3 Maize	Whorl application of phorate 10G or carbofuran 3 G @ 8-10 kg/ha to control shoot borer attack	Spray of mancozeb @ 0.25-0.4% at 8-10 days interval to control *Turcicum* leaf blight	Trichoderma mixed with FYM @10g/kg at 10 days prior to its use in the field can be applied to control stalk rot incidence which is likely during post flowering	
Crop4 Arhar	Soil application of *Trichoderma harzianum* along with FYM as side dressing to prevent *Fusarium* wilt	• Drenching of carbendazim 0.1% at plant base to control wilt • Foliar application of acephate 1.5 gm / lt or Miticide to prevent sterility mosaic virus	Drench with carbendazim 0.1% at plant base to control wilt	Quick drying to prevent molds
Crop5 Cotton	Protect against incidence of wilt and root rot. Drenching of Copper oxy chloride (COC) 0.3% or	• When marginal yellowing of leaves due to jassid injury is seen, spray neem oil 0.3% with sticker or imidacloprid 0.6 ml / lit or acetamiprid 0.1-0.2 ml /lit • Protect against Bacterial	To control grey mildew and boll rot, apply carbendazim 1 gm/ lit or mancozeb 3 gm / lit	Proper storage of seed cotton to prevent wetting and incidence of molds

Condition	**Continuous high rainfall in a short span leading to water logging**			
	Suggested contingency measure			
1	**2**	**3**	**4**	**5**
	carbendazim 0.1%	leaf blight (BLB) with foliar application of streptocycline sulphate 6 gm + 30 gm COC for every 10 lt of water		
Horticulture				
Crop1 (**Tomato,**	Spraying of contact insecticide for control of caterpillar Stacking for protecting fungal diseases	Spraying of contact insecticide for control of caterpillar/ fruit borer Stacking for protecting fungal diseases		
Crop2 **Cabbage,**				
Crop3 **chilly**	Drench the nursery beds with coc3 g/l to prevent damping off	Spray coc 30g+1g streptocycline in 10 lit. of water, 2-3 times against bacterial leaf spot and blight	Spray carbondizim 0.1% to control fruit rot	Quick drying of fruit to prevent fruit rot
Crop4				
Crop5				