Condition	Suggested contingency measure			
	Vegetative stage	**Flowering stage**	**Crop maturity stage**	**Post harvest**
Outbreak of pests and diseases due to unseasonable rains				
Soybean	Early planting to minimize the incidence of gridle beetle and green semilooper Foliar spray with 5% NSKE or dimethoate 30EC 1 ml/l to protect against semi looper	Monitor moth activity of spodoptera through pheromone traps (10 traps/ha) Apply Quinalphos 25EC 2ml/l or Emameetin benzoate 5 SG 4 g/10 lit to control spodoptera		-
Maize		Foliar application of Mancozeb .25-.4% at 8-10 days interval to control turcicum leaf blight	Trichoderma mixed with FYM 10 gm/kg at 10 days prior to its use in the field can be applied to control stalk rot which is likely during post flowering	Proper storage of seed cotton to prevent wetting and incidence of molds
Sorghum	Early sowing of sorgum to control Shootfly. Use of carbofuran granules 3G 8-10kg/ha to control stem	Use of carbofuran granules to control midge	Use of systematic insecticide as dusting with carbrabryl	Quick drying to prevent molds

	borer		powder(25kg/ha) to control Ear head bug	
Wheat	Spray 0.2% mancozeb 76% WP against wheat rust.	Spray 0.2% mancozeb 76% WP against wheat rust.	Spray 0.2% mancozeb 76% WP against wheat rust.	
Chickpea	Spray triazophos 40 % EC @ 1-1.5 l/ha in chickpea against pest incidence. · "T" shaped pegs placed in late sown chickpea field for biological control of pod borer. Chemical control spraying of Quinolphos 25 EC or Chlorpyriphos 20 EC C or Methyl Parathion 50 EC @ 600 ml dissolve in 500 L of water should be used. Dusting of Felvunerate 0.4% or Endosulphan 4% 15-20 kg or Quinolphos 1.5 WP 20-25 per hectare with duster	Spray triazophos 40 % EC @ 1-1.5 l/ha in chickpea against pest incidence. "T" shaped pegs placed in late sown chickpea field for biological control of pod borer Chemical control spraying of Quinolphos 25 EC or Chlorpyriphos 20 EC C or Methyl Parathion 50 EC @ 600 ml dissolve in 500 L of water should be used. Dusting of Felvunerate 0.4% or Endosulphan 4% 15-20 kg or Quinolphos 1.5 WP 20-25 per hectare with duster	Spray triazophos 40 % EC @ 1-1.5 l/ha in chickpea against pest incidence. Carry out critical survey of fields for insect and disease attack in crops	