Condition	Suggested contingency measure			
	Vegetative stage[k]	**Flowering stage[l]**	**Crop maturity stage[m]**	**Post harvest[n]**
1	2	3	4	5
Outbreak of pests and diseases due to unseasonal rains				
Crop1 (specify) Soybean	• Early planting to minimize the incidence of girdle beetle and green semilooper • Foliar spray with 5% NSKE or dimethoate 30EC 1 ml/l to protect against semilooper	• Monitor adult moth activity of Spodoptera through pheromone traps (10 traps/ha) • Apply Quinalphos 25 EC 2ml/l or Emamectin benzoate 5 SG 4g/10 lit to control spodoptera	apply spray of insecticides & fungicide for protecting from fungus	
Crop2 Maize	Spray imidachloprit 0.3 ml/l or Dimethoate 1.0 ml/l to control leaf hopper	Foliar application of Mancozeb @0.25 - 0.4% at 8-10 days interval to control *Turcicum* leaf blight	Trichoderma mixed with FYM @ 10 g/kg at 10 days prior to its use in the field can be applied to control stalk rot incidence which is likely during post	

			flowering	
Chickpea	Spray triazophos 40 % EC @ 1-1.5 l/ha in chickpea against pest incidence. · "T" shaped pegs placed in late sown chickpea field for biological control of pod borer and for chemical control spraying of Quinolphas 25 EC or Chlorpyriphos 20 EC C or Methyle Parathiyan 50 EC @ 600 ml dissolve in 500 L of water should be used. Dusting of Felvunerate 0.4% or Endosulphan 4% 15-20 kg or Quinolphas 1.5 WP 20-25 per hectare with duster			
Wheat	Spray 0.2 % mancozeb	Spray 0.2 % mancozeb	Trichoderma mixed	

	76% WP against wheat rust.	76% WP against wheat rust.	with FYM @ 10 g/kg at 10 days prior to its use in the field can be applied to control stalk rot incidence which is likely during post flowering	
Horticulture	NA			