Un-timely (unseasonal) rains

Condition	Suggested contingency measure			
	Vegetative stage	**Flowering stage**	**Crop maturity stage**	**Post harvest**
Outbreak of pests and diseases due to unseasonal rains				
Cotton	Spray Larwin@250g Or Ekalux 800ml/acre to check Mealy bug		**1.Insect/Pests:** Spray Imidachloprid 40 ml/ Pride20ml/acre for Jassid; Hostathion 600 ml/acre against white fly; Larwin@250gOr Ekalux 800ml/acre to check Mealy bug; synthetic pyrethroids/Carbamate insecticides against Pink/ spotted /American(small size) boll worm ; Organophosphate/Naturalite/oxadiazine against American(big size) boll worm and Carbamate/ Organochlorinate/ Organophosphates against Tobacco boll worm. **2.Diseases:** Grow LH 144/LH 2076 against Leaf curl; Cobalt chloride($COCl_2$) to check para wilt disease, Spray Blitox+streptocycline against Bacterial Blight and Blitox/Captan for control of Anthracnose,leaf blight and leaf	Storage of produce in dry place

		spot .	
Rice	Spray Nuvacron/Monocil@ 560 ml/acre against leaf folder and stem borer.	**1. Insect/Pests**: Spray Nuvacron /Monocil@ 560 ml/acre against leaf folder and stem borer; Confidor @40 ml/acre/ Ekalux @ 800 ml/acre against Plant hoppers/ Rice ear cutting caterpillar. **2. Diseases**: Grow PR 120, PAU 201, PR 111 against Bacterial leaf blight (BLB); spray Blitox(500ml)/Tilt (200ml) per acre to control False smut; Spray Tilt @ 200m l/acre against sheath blight ,Sheath rot and Bunt diseases.	-do-
Wheat	Spray pesticide to control Pink boll worm especially in rice fields.	Spray Nuvacron @150ml/acre to control sucking pest (Aphid), Ekalux for Army worm (@400 ml); Boll worm(800 ml) per acre and Tilt @200ml/acre to check Kernel bunt & rusts.	Treat the produce meant for seed with 250gm Malathion dust(5%)and disinfest 10 gunny bags with 5 ml cymbush/10 litres water ,Godowns with 100 ml Malathion/10 litres water.

Rapeseed/Mustard	Spray the pesticides when the insects (Aphids,jassids) have completely covered the flowers and pods	Aphids : Spray Actara 25 WG@ 40 g or Rogar 30 EC in 80-125 litres of water. Diseases**:** Two **s**prays of Blitox or Indofil M-45 in 100 litres of water at interval of 15 days to check white rust and Alternaria blight		Storage of produce in dry place
Horticulture				
Citrus	Chemical control of Phytophthora / foot rot with Ridomil-MZ/ Alliette as per recommendation (Decorticate and disinfect the wound on trunk either with disinfect solution and cover the wounds with Bordeaux paste folled by Bordeaux mixtureand Ridomil MZ as paint(2g/100ml of linseed oil) to the infected tree Control of sucking pests with systemic pesticides **(1250 ml Roger (dimethoate)30 EC)**	Chemical control of Phytophthora / Foot rot with Ridomil-MZ/ Alliette as per recommendation**(** Decorticate and disinfect the wound on trunk either with disinfect solution and cover the wounds with Bordeaux paste folled by Bordeaux mixtureand Ridomil MZ as paint(2g/100ml	Chemical control of Phytophthora / Foot rot with Ridomil-MZ/ Alliette as per recommendation **(**Decorticate and disinfect the wound on trunk either with disinfect solution and cover the wounds with Bordeaux paste folled by Bordeaux mixture and Ridomil MZ as paint(2g/100ml of linseed oil) to the infected tree), Control of sucking pests with systemic	Application of fungicides/ nutrients (GA 3 @ 30 ppm and wrapping in polythene of 100 gauze**)** to check post harvest losses

		of linseed oil) to the infected tree **)**, Control of sucking pests with systemic pesticides	pesticides	
Grapes	Chemical control **(**Prune the shoots in in Jan and Feb, Spray Bordeaux mixture in last week of March, Spray Bavistan 50 WP @ 500g in last week of May in 500 L of water , Spray Bavistan 50 WP @ 500g in mid July in 500 L of water**)**of sucking pests and diseases like Powdery mildew/ Anthracnose	Chemical control of sucking pests and diseases like powdery mildew/ Anthracnose	Chemical control of sucking pests and diseases like powdery mildew/ anthracnose/ hen and chicken disease/shot berry etc.	Timely harvesting of grapes, storage in proper CFB boxes.
Guava	Chemical control**(**Anthracnose/wilt with @300g and insects like fruit fly with Fenvelrate @1250 mi**)** of sucking pests	Chemical control **(**Anthracnose/wilt with @300g and insects like fruit fly with Fenvelrate	Chemical control **(**Anthracnose/wilt with @300g and insects like fruit fly with Fenvelrate @1250 mi**)** of fruit fly and	

		@1250 mi**)** of sucking pests and diseases like anthracnose.	anthracnose of guava. Harvesting at proper maturity level.	
Ber	Chemical Control of powdery mildew (spray karathene /Bayleton@0.5g/liter or sulfur @ 2.5 g/liter**)** of Leaf eating caterpillar and diseases like powdery mildew.**)** of fruit fly Control of leaf eating catterpillar	Spray sevin (Hexavin)@3g/liter Pumping out of excess rain water to check wilt	Sray Endosulfan@ 2.5g/liter (fruit fly) Spray sevin (Hexavin)@3g/liter (leaf eating caterpillar) Spray Blitox @ 5 g/ l water to check rottening of fruit	- Keep in dry place
Chilli	--			
Cucurbits	-	Spray Indofil M 45 @ 3 g/l water against downy mildew	Spray Blitox @ 5 g/ l water to check rottening of fruit. Also destroy the infested fruits and spray the crop with Endosulfan @ 8 ml/l or Sevin @ 5 g/l water to control fruit fly	-