

---

# PH3205-Computational Physics

## Spring 2022

[Bipradeep Saha \(19MS135\)](#)

---

### Assignment - 4


**Problem:** Write a program to solve eigenvectors and eigenvalues of three finite square wells of equal width separated by two equal widths of infinite square wells.

**Solution:** We solve the problem in Matlab

We first define the potential

```
L = 5; % Interval Length.
N = 1000; % No of points.
x = linspace(-L, L, N).'; % Coordinate vector.
dx = x(2) - x(1); % Coordinate step.
a = L/20; % width
b = L/10; % separation
D = 200; % depth
U = -D*(heaviside(x+ 1.5*a + b) - heaviside(x + 0.5*a +b) + heaviside(x+0.5*a) ...
- heaviside(x-0.5*a) + heaviside(x - (0.5*a + b)) -heaviside(x-((a*3/2)+b)));
```

And this is what we get:



Then we set up the hamiltonian

```
%% Setting up the hamiltonian
hbar = 1;
m = 1;
e = ones(N,1);
Lap = spdiags([e -2*e e],[-1 0 1],N,N) / dx^2;
H = -(1/2)*(hbar^2/m)*Lap + spdiags(U,0,N,N);
```

And we then obtain the first 3 eigenvalues and eigenstates and then we plot it:

```
%% Getting the 3 lowest eigen values
nmodes = 3;
[V,E] = eigs(H,nmodes,'smallestreal');
[E,ind] = sort(diag(E)); % Convert E to vector and sort low to high.
V = V(:,ind);

usc = D*U*max(abs(V))/max(abs(U));

figure(1)
ax = plot(x,U/500, Color='black', LineStyle='--', DisplayName='Scaled Potential');
hold on
plot(x,V(:,1), DisplayName='Energy='+string(E(1)))
plot(x,V(:,2), DisplayName='Energy='+string(E(2)))
plot(x,V(:,3), DisplayName='Energy='+string(E(3)))
ylim([-0.5 0.25])
legend(Location="best")
xlabel("x (in m)")
ylabel("Un-normalized wave funcitons")
```

The eigenstates along with their energy and the rescaled potential is:

