## Problem Set 3

Instructions: Answer the following questions in their entirety in a separate document. Submit your completed problem set as a PDF document via email to sean.corcoran@vanderbilt.edu. Use your last name and problem set number as the filename (e.g., Ginsburg Problem Set 3.pdf). Working together is encouraged, but it is expected that all submitted work be that of the individual student.

1. (16 points—2 each) In a population of students, the number of absences during the school year ranges from 0 to 8. The probabilities of a randomly drawn student from this population having 0, 1, 2, ..., 8 absences are shown in the table below. Define the event A as the student being absent fewer than 4 days, and the event B as the student being absent more than 3 days.

# of Days	0	1	2	3	4	5	6	7	8
Probability	0.2	0.14	0.25	0.11	0.1	0.09	0.05	0.03	0.03

- (a) What is the probability of event A?
- (b) What is the probability of event B?
- (c) What is the probability of  $\sim A$ ?
- (d) Are events A and B mutually exclusive? Explain why or why not.
- (e) What is the probability of  $A \cap B$ ?
- (f) What is the probability of  $A \cup B$ ?
- (g) Show using values from the table that  $P((A \cap B) \cup (\sim A \cap B)) = P(B)$ .
- (h) Show using values from the table that  $P(A \cup (\sim A \cap B)) = P(A \cup B)$ .
- 2. (6 points—3 each) Using the probability distribution in Question 1, find the following (and show your work):
  - (a) E(# of absences)
  - (b) Var(# of absences) and SD(# of absences)

3. (8 points—2 each) Shown below is a 2 x 2 table that reports the fraction of the population in each cell:

		Education level		
		HS	<hs< th=""><th>Totals</th></hs<>	Totals
Current smoker:	NO YES	0.614 0.194	0.130 0.062	0.744 0.256
	Totals	0.808	0.192	1.000

- (a) For a randomly drawn person, what is P(smoker)?
- (b) For a randomly drawn person, what is P(smoker | <HS diploma)?
- (c) For a randomly drawn person, what is P(smoker | HS diploma+)?
- (d) Are education and smoking status independent? Why or why not?
- 4. (5 points) Shown below is a 2 x 2 table. In Period 1, events A or B can happen. In Period 2, outcome C or D will result. If P(C|B) = 0.150 and P(D|A) = 0.7, then fill in the missing boxes below:

		Period 1		
		Event A	Event B	
Period 2	Event C Event D		0.030	

- 5. (5 points) After the attacks of September 11, 2001, the TSA implemented a program called SPOT (Screening of Passengers by Observation Techniques) in which passengers were flagged for suspicious behavior and given additional searching or screening. Suppose that:
  - There are 2 billion + 100 passenger trips per year (2,000,000,100).
  - 100 of these passengers are terrorists.
  - Nearly all (99%) terrorists exhibit the kinds of behaviors that were flagged.
  - Some non-terrorists exhibit these suspicious behaviors, but it is rare (1%).

The SPOT test has low false negative and false positive rates, suggesting it is an effective way to catch would-be terrorists. Use Bayes' Theorem to calculate the probability that a flagged passenger is, in fact, a terrorist.

- 6. (6 points—3 each) Paul and Natasha live in Los Angeles. Paul hates cold weather but Natasha has been transferred to a cold Northeastern city. Paul notes that he cannot move go to a city where more than 30% of the days have an average daily high below freezing. Suppose the average daily high temperatures (X) in a city can be described by a uniform distribution where the minimum and maximum average daily highs are -2 and 105, respectively.
  - (a) What is the PDF for X, and what is  $P(x \le 32)$ ? Should Natasha look for a one or a two bedroom apartment? (Hint: you do not need calculus to find the requested probability).
  - (b) What are E(X) and Var(X)?
- 7. (4 points) Assume the random variable z has a standard normal distribution. Use Stata, an online calculator, or a textbook table to answer the following:
  - (a) The probability is 0.75 that z is less than what number?
  - (b) The probability is 0.30 that z is less than what number?
  - (c) The probability is 0.10 that z is greater than what number?
  - (d) The probability is 0.60 that z is greater than what number?
- 8. (6 points) Applicants to Local U. have SAT composite scores that follow a normal distribution with a mean of 1100 and a standard deviation of 160.
  - (a) What is the probability that a Local U. applicant will have a composite SAT score of 1300 or higher?
  - (b) What is the probability that a Local U. applicant will have a composite score above 900 but below 1300?
  - (c) Showing strikingly bad judgment, Local U. wishes to offer a tuition-free scholar-ship to applicants who score in the top 2% of applicants on the SAT. Above what score will they offer this scholarship?

In each case, show or explain how you obtained your result.

- 9. (6 points—3 each) Suppose the probability that a teenage driver gets into an accident during a one-year period is 0.12, and assume the probability of getting into an accident is independent across drivers.
  - (a) A particular family has 5 teenage drivers. What is the probability that at least one driver in this family will have an accident over the coming year? Show or explain how you obtained your answer.

(b) Now consider the population of families with 5 teenage drivers, and define X as the number of accidents that occurred among their 5 drivers. For these families in a typical year, what is E(X) and sd(X)? Show or explain how you obtained your answer.