

# Assignment-5

Name: Sai Pravallika Danda, Roll Number: CS20BTECH11013

Download all latex-tikz codes from

<https://github.com/spdanda/AI1103/blob/main/Assignment5/Assignment5.tex>

## CSIR-UGC-NET June-2016 Q50 :

Let  $X$  and  $Y$  be independent and identically distributed random variables such that  $\Pr(X = 0) = \Pr(X = 1) = \frac{1}{2}$ . Let  $Z = X + Y$  and  $W = |X - Y|$ . Then which statement is not correct?

- 1)  $X$  and  $W$  are independent.
- 2)  $Y$  and  $W$  are independent.
- 3)  $Z$  and  $W$  are uncorrelated.
- 4)  $Z$  and  $W$  are independent.

### Solution:

$X, Y \in \{0, 1\} \implies Z \in \{0, 1, 2\}$  and  $W \in \{0, 1\}$ .

Also,

$$\Pr(X = x) = \begin{cases} \frac{1}{2} & x \in \{0, 1\} \\ 0 & \text{otherwise} \end{cases} \quad (0.0.1)$$

$$\Pr(Y = y) = \begin{cases} \frac{1}{2} & y \in \{0, 1\} \\ 0 & \text{otherwise} \end{cases} \quad (0.0.2)$$

So Probability Generating functions(PGF's) for  $X$  and  $Y$  are

$$\mathcal{G}_X(z) = E[z^X] \quad (0.0.3)$$

$$= \sum_{i=0}^1 p_i z^i \quad (0.0.4)$$

$$= \frac{1+z}{2} \quad (0.0.5)$$

Similarly,

$$\mathcal{G}_Y(z) = \frac{1+z}{2} \quad (0.0.6)$$

a)

$$\Pr(W = 0) = \Pr(X = 0, Y = 0) + \Pr(X = 1, Y = 1) \quad (0.0.7)$$

$$= \frac{1}{2} \times \frac{1}{2} + \frac{1}{2} \times \frac{1}{2} \quad (0.0.8)$$

$$= \frac{1}{2} \quad (0.0.9)$$

$$\Pr(W = 1) = 1 - \Pr(W = 0) \quad (0.0.10)$$

$$= \frac{1}{2} \quad (0.0.11)$$

$$\therefore \Pr(W = w) = \begin{cases} \frac{1}{2} & w \in \{0, 1\} \\ 0 & \text{otherwise} \end{cases} \quad (0.0.12)$$

$$\text{Also } E[W] = 0 \times \frac{1}{2} + 1 \times \frac{1}{2} \quad (0.0.13)$$

$$= \frac{1}{2} \quad (0.0.14)$$

b) PGF of  $Z$ :

$$\mathcal{G}_Z(z) = E[z^{X+Y}] \quad (0.0.15)$$

$$= E[z^X \times z^Y] \quad (0.0.16)$$

$$= E[z^X] \times E[z^Y] \quad (0.0.17)$$

( $\because X$  and  $Y$  are independent)

$$\therefore \mathcal{G}_Z(z) = \left(\frac{1+z}{2}\right)^2 \quad (0.0.18)$$

$$= \frac{1+2z+z^2}{4} \quad (0.0.19)$$

$$= \frac{1}{4} + \frac{1}{2}z + \frac{1}{4}z^2 \quad (0.0.20)$$

∴ From the PGF of  $Z$ ; pmf of  $Z$  is

$$\Pr(Z = z) = \begin{cases} \frac{1}{4} & z = 0 \\ \frac{1}{2} & z = 1 \\ \frac{1}{4} & z = 2 \end{cases} \quad (0.0.21)$$

$$\text{Also, } E[Z] = 0 \times \frac{1}{4} + 1 \times \frac{1}{2} + 2 \times \frac{1}{4} \quad (0.0.22)$$

$$= 1 \quad (0.0.23)$$

Now, for checking each option,

1) Checking if  $X$  and  $W$  are independent

$$p_1 = \Pr(X = x, W = w) \quad (0.0.24)$$

$$= \Pr(X = x, Y = x \pm w) \quad (0.0.25)$$

$$= \Pr(X = x) \times \Pr(Y = x \pm w) \quad (0.0.26)$$

$$= \frac{1}{2} \times \frac{1}{2} \quad (0.0.27)$$

$$= \begin{cases} \frac{1}{4} & (x \pm w) \in \{1, 2, 3, 4, 5, 6\} \\ 0 & \text{otherwise} \end{cases} \quad (0.0.28)$$

(only one value for  $Y$  is obtained for each case when  $x$  and  $w$  are substituted)

$$\Pr(X = x) \times \Pr(W = w) = \frac{1}{2} \times \frac{1}{2} \quad (0.0.29)$$

$$= \frac{1}{4} \quad (0.0.30)$$

$$\Pr(X = x) \Pr(W = w) = \Pr(X = x, W = w) \quad (0.0.31)$$

$\implies X$  and  $W$  are independent and hence Option 1 is true.

2) Checking if  $Y$  and  $W$  are independent

Solving of this case is identical to the first option except the variable  $X$  is replaced by  $Y$ . Hence on solving, you get  $Y$  and  $W$  are independent.

∴ Option 2 is also true.

3) Checking if  $W$  and  $Z$  are uncorrelated  
**Uncorrelated random variables:** Two variables are said to be uncorrelated if the expected value of their joint distribution is equal to product of the expected values of their respective marginal distributions.

Also,  $WZ \in \{0, 1, 2\}$

$$\Pr(WZ = 2) = \Pr(W = 1, Z = 2) = 0 \quad (0.0.32)$$

(∵ If  $Z = 2, X = Y = 1 \implies W = 0$  i.e.,  $\neq$  to 1)

$$\Pr(WZ = 0) = \frac{1}{2} \quad (0.0.33)$$

(∵ either  $W = 0$  or  $Z = 0$ )

$$\Pr(WZ = 1) = 1 - \Pr(WZ = 0) - \Pr(WZ = 2) \quad (0.0.34)$$

$$= 1 - \frac{1}{2} \quad (0.0.35)$$

$$= \frac{1}{2} \quad (0.0.36)$$

∴ Expected value of  $WZ, E[WZ]$

$$= \sum_{k=0,1,2} k \Pr(WZ = k) \quad (0.0.37)$$

$$= 1 \times \frac{1}{2} + 0 \quad (0.0.38)$$

$$= \frac{1}{2} \quad (0.0.39)$$

$$\text{Also } E[W] \times E[Z] = \frac{1}{2} \times 1 = \frac{1}{2} \quad (0.0.40)$$

$$= E[WZ] \quad (0.0.41)$$

Hence from the above equation  $W$  and  $Z$  are uncorrelated random variables.

∴ Option 3 is also true.

4) Let's check for one particular case.

$$\Pr(W = 0 | Z = 0) = 1 \quad (0.0.42)$$

$$(\because Z = 0 \implies X = Y = 0 \implies W = 0)$$

$$\therefore \frac{\Pr(W = 0, Z = 0)}{\Pr(Z = 0)} = 1 \quad (0.0.43)$$

$$\begin{aligned} \implies \Pr(W = 0, Z = 0) &= \Pr(Z = 0) \\ &\neq \Pr(W = 0) \times \Pr(Z = 0) \quad (0.0.44) \end{aligned}$$

$\therefore W$  and  $Z$  are not independent random variables

Hence Option 4 is incorrect.

$\therefore$  Answer is Option4.