# NEET UG BIOLOGY

## MASTER TEXTBOOK

Complete Master Theory, Formulas & Experimental Compendium strictly aligned with NEET UG 2027 NMC Syllabus

Authoritative First Edition · Global Academic Series

# NEET UG BIOLOGY

## MASTER TEXTBOOK

### Complete Master Theory, Formulas & Experimental Compendium

**Strictly Aligned with NEET UG 2027 NMC Syllabus**

**Authored by**

**Satya Prakash**

**Comprehensive Coverage of Botany & Zoology Domains**

Cell Biology, Plant Morphology & Families, Plant Physiology, Human Physiology, Reproduction, Genetics & Genomics, Biotechnology, Human Health, Evolution & Ecology

Published for Independent Medical Aspirants, Educators & Academic Study Circles

Comprehensive Academic Edition · Global Print-on-Demand

**NEET UG BIOLOGY MASTER TEXTBOOK**

*Complete Master Theory, Formulas & Experimental Compendium strictly aligned with NEET UG 2027 NMC Syllabus*

**Author:** Satya Prakash
**Editorial Correspondence & Academic Feedback:** spdedsec@hotmail.com
**Source Repository & Official Errata:**
`https://github.com/spdedsec/neet-study-materials`

© 2026–2027 Satya Prakash. All rights reserved.

**First Edition:** October 2026
**Publishing Imprint:** Independent Academic Publishing / Satya Prakash Editorial Imprint
**Target Cycle:** NEET UG 2027 (Undergraduate Medical Admissions Entrance Examination)
**ISBN-13 (Paperback):** Pending Official Assignment (Volume 3)
**ISBN-13 (Hardcover):** Pending Official Assignment (Volume 3)

**All Rights Reserved.** No part of this publication may be reproduced, stored in a retrieval system, or transmitted in any form or by any means—electronic, mechanical, photocopying, recording, scanning, or otherwise—without the prior written permission of the author, except for brief quotations embodied in critical reviews or scholarly analysis as permitted under copyright law.

**Academic & Legal Disclaimer for Medical Entrance Preparation:** This textbook is an independent preparation manual authored by Satya Prakash to assist candidates preparing for the National Eligibility cum Entrance Test (NEET UG). While every effort has been made to verify biological accuracy and strict concordance with the National Medical Commission (NMC) curriculum and rationalized NCERT textbooks, the author and publisher make no official claims of affiliation with or endorsement by the National Testing Agency (NTA), the National Medical Commission (NMC), or the Ministry of Education, Government of India. The author and publisher disclaim any direct or indirect liability for errors, omissions, or examination outcomes.

**Typesetting & Typography:** Typeset in LaTeX 2$\varepsilon$ with the Tectonic XeTeX engine using Latin Modern typefaces. Biological taxonomic nomenclature follows the International Code of Nomenclature for algae, fungi, and plants (ICN) and the International Code of Zoological Nomenclature (ICZN). Produced and bound under rigorous academic publishing standards.

Printed in the United States of America / United Kingdom / India.

*To my parents, whose selfless dedication, faith, and patience illuminated every step of my journey;*

*To the biologists, naturalists, and physicians whose tireless investigations into the living world revealed the exquisite molecular and cellular unity of life;*

*And to you, future physician—*
*may the biological principles detailed in these pages inspire not only your success in NEET UG,*
*but a lifelong reverence for the living human body and the physiological systems you will soon heal.*

# Acknowledgements

Writing an exhaustive, 360-mark reference textbook for NEET UG Biology requires balancing two vital pedagogical imperatives: absolute fidelity to rationalized NCERT curriculum benchmarks, and clinical conceptual depth that renders a student immune to complex Assertion-Reasoning and multi-statement evaluations.

I express my deepest gratitude to my family for their unending understanding, encouragement, and sacrifices throughout the compilation of this manuscript.

I thank the academic biology faculty, medical researchers, and botanists whose critical discussions helped ensure that every floral formula, enzyme kinetic relation, metabolic pathway, and endocrine regulatory loop was represented with anatomical and physiological precision.

I am especially grateful to the hundreds of medical aspirants whose insightful questions, mock examination error logs, and diagnostic doubts directly shaped the high-yield comparative matrices and "NEET Trap" callouts throughout this book. Your relentless pursuit of medical excellence was the primary catalyst for this work.

Finally, I acknowledge the developers of LaTeX $2_\varepsilon$ and the Tectonic typesetting ecosystem, whose modern publishing toolchain enabled us to integrate comprehensive taxonomic matrices, biochemical pathways, and color-coded diagnostic summaries into a publication-ready authoritative manuscript.

**Satya Prakash**
*October 2026*

# Author's Preface

The National Eligibility cum Entrance Test (NEET UG) allocates exactly **360 marks out of 720**—precisely half the total examination score—to Biology. Consequently, medical aspirants frequently encounter the ubiquitous pedagogical advice: *"Simply read NCERT multiple times and complete mastery is assured."*

However, empirical examination trends over recent testing cycles demonstrate that passive textual memorization without deep mechanistic comprehension leaves candidates acutely vulnerable. Modern NEET papers increasingly feature multi-statement evaluative questions, experimental deduction problems, Assertion-Reason pairings with subtle causal disconnects, and quantitative genetics calculations. Memorizing isolated sentences without understanding underlying biochemical cascades, comparative anatomical distinctions, or evolutionary dynamics creates an illusion of competence that falters under examination pressure.

Furthermore, the National Medical Commission (NMC) has significantly rationalized the curriculum:

- **Curriculum Exclusions:** Entire legacy topics have been formally removed, including *Digestion and Absorption*, *Transport in Plants*, *Mineral Nutrition*, *Strategies for Enhancement in Food Production*, *Environmental Issues*, and legacy secondary growth sections. Focusing finite preparation hours on deleted topics is counter-productive.
- **Explicit Additions & Emphases:** The NMC has explicitly mandated comprehensive study of **seven distinct angiosperm families** (Malvaceae, Cruciferae/Brassicaceae, Fabaceae/Leguminosae, Asteraceae/Compositae, Poaceae/Gramineae, Solanaceae, and Liliaceae) and reinstated detailed animal morphology for both **Frog** (*Rana tigrina*) and **Cockroach** (*Periplaneta americana*).

This master textbook is constructed around an uncompromising, seven-part architectural framework strictly aligned with the revised NMC syllabus:

1. **Part I: Cell Biology, Biomolecules & Cell Division:** Rigorous coverage of prokaryotic and eukaryotic ultrastructure, endomembrane dynamics, full Michaelis-Menten enzyme kinetics ($K_m$, Lineweaver-Burk dynamics, inhibition modalities), and meiotic chromosomal mechanics.
2. **Part II: Structural Organisation & Diversity:** Complete plant morphology, floral organography, full comparative floral formulas for all seven prescribed angiosperm families, animal taxonomy across all phyla, and exhaustive anatomical systems of Frog and Cockroach.

3. **Part III: Plant Physiology:** Detailed bioenergetics of light reactions, chemiosmotic coupling, Calvin $C_3$, Hatch-Slack $C_4$, Crassulacean Acid Metabolism (CAM), photorespiration ($C_2$), glycolysis, TCA cycle, mitochondrial ETS, and phytohormonal regulatory cascades.
4. **Part IV: Human Physiology:** Clinical and mechanical integration of respiration, cardiovascular dynamics, nephron counter-current multiplication, sliding filament musculoskeletal mechanics, and neuro-endocrine coordination.
5. **Part V: Reproduction & Reproductive Health:** Angiosperm micro/megasporogenesis, double fertilization, apomixis, human gametogenesis, menstrual endocrinology, contraceptive mechanics, and assisted reproductive technologies (ART).
6. **Part VI: Genetics & Molecular Biology:** Mendelian combinatorics, non-Mendelian extensions, Morgan's linkage and chromosome mapping, semi-conservative replication, transcription, translation, the Lac operon, Genomics (Human Genome Project), and DNA fingerprinting.
7. **Part VII: Biotechnology, Human Health & Ecology:** Recombinant DNA tools and processes, transgenics, infectious pathogens, innate/adaptive immunology, evolutionary mechanisms, population growth kinetics, ecosystem bioenergetics, and biodiversity conservation.
8. **Appendix: Quantitative Biology & Master Diagnostic Matrices:** Unified mathematical formulations for genetic probability, recombination frequency, Hardy-Weinberg population equilibria, pulmonary/cardiac constants, and side-by-side diagnostic matrices for all seven plant families.

To achieve true mastery, active retrieval must replace passive rereading. When analyzing physiological feedback loops or biochemical cycles, sketch the regulatory cascades from memory. When studying genetics, solve the combinatorial probability trees by hand. Treat this volume as your analytical companion, analyze your error logs with intellectual honesty, and maintain unwavering dedication toward your goal of medical service.

**Satya Prakash**
Author & Curriculum Researcher
spdedsec@hotmail.com
*October 2026*

# Syllabus Blueprint & NMC Rationalization Guide

This section details the official NEET UG 2027 Biology curriculum prescribed by the National Medical Commission (NMC), covering both Botany and Zoology domains, examination weightage, and critical syllabus exclusions.

## Official Biology Examination Architecture

| Subject Domain | Compulsory Qs | Marks | Pace Budget | Question Typology |
|---|---|---|---|---|
| **Botany** | 45 (All Compulsory) | 180 | 40–45 min | Single Choice MCQs, Statement I/II, |
| **Zoology** | 45 (All Compulsory) | 180 | 40–45 min | Assertion-Reason, Diagram/Matrix Match |
| **Master Biology Total** | **90 Compulsory** | **360** | **80–90 min** | High-Speed Conceptual Retrieval |

**Table 1:** Reformed NEET UG Biology Examination Scheme (+4 for correct response, −1 for incorrect response, 0 for unattempted. Optional Section B discontinued by NTA).

## Official 10-Unit Biology Syllabus Structure

1. **Unit 1: Diversity in Living World** (What is living? Biodiversity, taxonomic hierarchy, binomial nomenclature, five-kingdom classification: Monera, Protista, Fungi, Plantae, Animalia; Salient features of non-chordates up to phyla level and chordates up to class level).
2. **Unit 2: Structural Organisation in Animals and Plants** (Morphology, anatomy and functions of different parts of flowering plants: Root, stem, leaf, inflorescence, flower, fruit and seed; Description of families: Malvaceae, Cruciferae/Brassicaceae, Fabaceae/Leguminosae, Asteraceae/Compositae, Poaceae/Gramineae, Solanaceae, Liliaceae; Animal tissues; Morphology, anatomy and functions of digestive, circulatory, respiratory, nervous and reproductive systems of Frog and Cockroach).

3. **Unit 3: Cell Structure and Function** (Cell theory, prokaryotic and eukaryotic cells, plant and animal cells, cell envelope, organelles: nucleus, mitochondria, chloroplast, endomembrane system; Biomolecules: carbohydrates, proteins, nucleic acids, enzymes; Cell cycle: interphase, mitosis and meiosis).
4. **Unit 4: Plant Physiology** (Photosynthesis in higher plants: light reactions, photolysis, cyclic and non-cyclic photophosphorylation, Calvin cycle, $C_4$ pathway, photorespiration; Respiration in plants: glycolysis, TCA cycle, electron transport system, oxidative phosphorylation; Plant growth and development: auxins, gibberellins, cytokinins, ethylene, abscisic acid).
5. **Unit 5: Human Physiology** (Breathing and respiration: pulmonary volumes, alveolar gas exchange, oxygen dissociation; Body fluids and circulation: blood composition, ABO/Rh grouping, cardiac cycle, ECG; Excretory products and elimination: nephron function, counter-current mechanism, RAAS; Locomotion and movement: sliding filament theory, skeletal joints; Neural control: action potentials, synapses; Chemical coordination: endocrine hormones and mechanisms).
6. **Unit 6: Reproduction** (Sexual reproduction in flowering plants: microsporogenesis, megasporogenesis, double fertilization, apomixis; Human reproduction: male and female reproductive systems, gametogenesis, menstrual cycle, fertilization, embryogeny; Reproductive health: contraception, sexually transmitted infections, assisted reproductive technologies).
7. **Unit 7: Genetics and Evolution** (Heredity and variation: Mendelian ratios, incomplete dominance, codominance, multiple alleles, linkage and crossing over, sex determination, sex-linked disorders; Molecular basis of inheritance: DNA structure, replication, transcription, genetic code, translation, lac operon, DNA fingerprinting; Evolution: origin of life, Darwinian selection, evidence for evolution, Hardy-Weinberg equilibrium, human evolution).
8. **Unit 8: Biology and Human Welfare** (Health and disease: pathogens causing typhoid, pneumonia, malaria, amoebiasis, ringworm, filariasis; Basic principles of immunology: innate and acquired immunity, vaccines; AIDS and cancer: etiology, diagnosis, treatment; Microbes in human welfare: household products, industrial processing, sewage treatment, biofertilizers).
9. **Unit 9: Biotechnology and Its Applications** (Principles and processes: recombinant DNA technology, restriction enzymes, cloning vectors, PCR; Applications in agriculture and medicine: genetically modified crops, Bt cotton, RNA interference, human insulin, gene therapy, transgenic animals).
10. **Unit 10: Ecology and Environment** (Organisms and populations: population growth models, species interactions; Ecosystem: structure, productivity, decomposition, energy flow, ecological pyramids; Biodiversity and conservation: patterns of biodiversity, red data book, in-situ and ex-situ conservation).

**Zero-Tolerance NMC Exclusions: Topics Strictly Deleted from Biology**

Under revised NMC regulations, the following topics and chapters have been **completely eliminated**:

- **Digestion and Absorption** (Entire human digestion chapter eliminated).
- **Transport in Plants** (Xylem/phloem transport mechanisms, water potential, transpiration pull eliminated from Plant Physiology).
- **Mineral Nutrition** (Macro/micronutrients, nitrogen metabolism, deficiency symptoms eliminated).
- **Strategies for Enhancement in Food Production** (Animal husbandry and plant breeding eliminated).
- **Environmental Issues** (Pollution, solid waste, global warming eliminated from Ecology).
- **Anatomy of Flowering Plants (legacy sections)** (Secondary growth, anomalous structures eliminated).

**Important Prescribed Additions to Note:**

- **Prescribed 7 Plant Families:** Malvaceae, Cruciferae/Brassicaceae, Fabaceae/Leguminosae, Asteraceae/Compositae, Poaceae/Gramineae, Solanaceae, and Liliaceae must be studied thoroughly.
- **Frog & Cockroach Morphology:** Both are explicitly in the syllabus.

## Empirical Question Weightage Matrix

| Biology Domain | Typical Qs | Marks Weightage |
|---|---|---|
| Genetics & Molecular Basis of Inheritance (Units 7A, 7B) | 15–18 | 60–72 Marks |
| Human Physiology (Respiration, Circulation, Excretion, Locomotion, Neuro-Endocrine) | 12–15 | 48–60 Marks |
| Plant Physiology (Photosynthesis, Respiration, Plant Growth Regulators) | 6–8 | 24–32 Marks |
| Cell Biology & Cell Division (Cell Structure, Biomolecules, Mitosis, Meiosis) | 7–9 | 28–36 Marks |
| Reproduction & Biotechnology (Plant/Human Reproduction, Health, rDNA, Biotech) | 14–16 | 56–64 Marks |
| Ecology & Environment (Populations, Ecosystems, Biodiversity) | 8–10 | 32–40 Marks |
| Diversity in Living World & 7 Plant Families (Taxonomy, Frog/Cockroach) | 10–12 | 40–48 Marks |
| Immunology, Human Health & Evolutionary Biology | 6–8 | 24–32 Marks |
| **Total Biology Paper** | **90 Qs (Compulsory)** | **360 Marks** |

**Table 2:** Empirical Question Distribution in Recent NEET UG Papers.

# Contents

**Acknowledgements** **ix**

**Author's Preface** **xi**

**Syllabus Blueprint & NMC Rationalization Guide** **xiii**

**I Cell Biology, Biomolecules & Cell Division** **1**

**1 Cell Structure, Organelles & Biomolecules** **3**

1.1 Cell Theory & Prokaryotic vs Eukaryotic Architecture . . . . . 3

1.2 The Endomembrane System & Organelle Specialization . . . . . 4

1.3 Biomolecules, Structural Conformation & Enzyme Kinetics . . . 6

1.3.1 Enzyme Catalytic Action & Michaelis-Menten Saturation Kinetics . . . . . 7

1.3.2 Competitive vs. Non-Competitive Enzyme Inhibition . . . 9

1.3.3 IUBMB Enzyme Classification & Cofactor Architecture . 9

**2 Cell Cycle, Mitosis & Meiosis** **11**

2.1 Cell Cycle Progression & DNA/Chromosome Quantitation . . . . 11

2.2 Mitosis: Spindle Dynamics & Cytokinesis . . . . . 12

2.3 Meiosis: Prophase I Molecular Substages & Segregation . . . . . 13

**II Structural Organisation & Diversity** **15**

**3 Plant Morphology & Systematic Angiosperm Families** **17**

3.1 Vegetative Morphology: Root, Stem, Leaf & Adaptations . . . . 17

3.2 Floral Morphology, Inflorescence, Aestivation & Placentation . . 19

3.3 Systematic Flora: Diagnostic Profiles & Floral Formulas of the 7 NMC Families . . . . . 20

3.3.1 1. Family Malvaceae (Mallow / Cotton Family) . . . . . 21

3.3.2 2. Family Cruciferae / Brassicaceae (Mustard Family) . . 21

3.3.3 3. Family Fabaceae / Leguminosae (Legume / Pea Family) 21
3.3.4 4. Family Asteraceae / Compositae (Sunflower Family) . 22
3.3.5 5. Family Poaceae / Gramineae (Grass / Cereal Family) . 22
3.3.6 6. Family Solanaceae (Potato / Nightshade Family) . . . 23
3.3.7 7. Family Liliaceae (Lily Family) . . . . . 23

**4 Animal Diversity & Comparative Anatomy: Frog and Cockroach 27**
4.1 Diagnostic Hallmarks of Major Animal Phyla . . . . . 27
4.2 Anatomy of the Cockroach (*Periplaneta americana*) . . . . . 27
4.3 Anatomy of the Frog (*Rana tigrina*) . . . . . 30

**III Plant Physiology 33**

**5 Photosynthesis in Higher Plants 35**
5.1 Chloroplast Ultrastructure, Pigments & Action Spectra . . . . . 35
5.2 Photochemical Phase: Z-Scheme, Water Photolysis & Photophosphorylation . . . . . 36
5.3 Chemiosmotic ATP Synthesis & Proton Motive Force . . . . . 36
5.4 Biosynthetic Phase: Calvin $C_3$ Cycle Stoichiometry & Energetics 37
5.5 Hatch-Slack $C_4$ Pathway & Kranz Anatomy . . . . . 37
5.6 Crassulacean Acid Metabolism (CAM) & Photorespiration . . . . 38
5.7 Factors Regulating Photosynthesis & Blackman's Law . . . . . 39

**6 Respiration in Plants & Phytohormonal Coordination 41**
6.1 Glycolytic Pathway (EMP) & Mitochondrial Link Reaction . . . 41
6.2 Tricarboxylic Acid (TCA) Cycle & Amphibolic Intermediates . . 42
6.3 Mitochondrial Electron Transport System (ETS) & ATP Synthase 42
6.4 Respiratory Quotient ($RQ$) Formulations . . . . . 43
6.5 Phytohormones: Plant Growth Regulators & Morphogenesis . . . 43

**IV Human Physiology 45**

**7 Human Respiration, Circulation & Excretion 47**
7.1 Breathing & Alveolar Gas Exchange . . . . . 47

7.1.1 Alveolar Diffusion Gradients & Respiratory Membrane Dynamics . . . . . 47
7.1.2 Oxygen Transport & The Oxygen-Hemoglobin Dissociation Curve . . . . . 48
7.1.3 Carbon Dioxide Transport Modalities & The Chloride Shift 49
7.1.4 Neural Regulation of Respiration & Clinical Pathologies . 50
7.2 Body Fluids, Circulation & Coagulation Cascade . . . . . 50
7.3 Excretory Products & Renal Physiology . . . . . 51

**8 Locomotion, Nervous Integration & Chemical Coordination 53**
8.1 Musculoskeletal Mechanics & Sliding Filament Theory . . . . . 53
8.2 Neural Control & Action Potential Propagation . . . . . 55
8.3 Endocrine Integration & Hormone Action Mechanisms . . . . . 55

**V Reproduction & Reproductive Health 57**

**9 Sexual Reproduction in Flowering Plants (Angiosperms) 59**
9.1 Microsporogenesis & Male Gametophyte . . . . . 59
9.2 Megasporogenesis & The 7-Celled 8-Nucleate Embryo Sac . . . . 60
9.3 Pollination Ecology, Outbreeding Devices & Pollen-Pistil Interaction . . . . . 60
9.4 Double Fertilization, Endosperm Energetics, Embryogenesis & Apomixis . . . . . 61

**10 Human Reproduction & Reproductive Health 63**
10.1 Male & Female Reproductive Architectures . . . . . 63
10.2 Gametogenesis: Spermatogenesis vs Oogenesis . . . . . 64
10.3 Menstrual Cycle Hormonal Regulation & Implantation . . . . . 64
10.4 Reproductive Health, Contraceptive Modalities & Assisted Reproductive Technologies (ART) . . . . . 65

**VI Genetics & Molecular Biology 67**

**11 Principles of Inheritance & Variation 69**
11.1 Mendelian Principles & Cross Combinatorics . . . . . 69
11.2 Non-Mendelian Extensions: Dominance, Allelism & Pleiotropy . 70

11.3 Chromosomal Theory, Sex Determination & Morgan's Linkage . 71
11.4 Genetic Disorders & Pedigree Analysis . . . . . 72

**12 Molecular Basis of Inheritance & Genomics 73**
12.1 Watson-Crick B-DNA Structural Dimensions & Chromatin Packaging . . . . . 73
12.2 Historic Proof of DNA as the Genetic Material . . . . . 74
12.3 Semi-Conservative DNA Replication Machinery & Enzymology . 74
12.4 Transcription in Prokaryotes vs Eukaryotes & RNA Processing . 75
12.5 The Genetic Code, Wobble Phenomenon & Translation . . . . . 76
12.6 Operon Model: The Lac Operon . . . . . 77
12.7 Genomics & The Human Genome Project (HGP) . . . . . 77
12.8 DNA Fingerprinting & Satellite DNA Polymorphism . . . . . 78

**VII Biotechnology, Human Health & Ecology 81**

**13 Biotechnology: Principles, Processes & Modern Applications 83**
13.1 Tools of Recombinant DNA Technology & Restriction Endonucleases . . . . . 83
13.2 Cloning Vectors, pBR322 Architecture & Host Transformation . 84
13.3 PCR Amplification, Bioreactor Kinetics & Downstream Processing 84
13.4 Transgenic Applications in Agriculture, Medicine & Biopiracy . . 85

**14 Immunology, Human Pathogens & Evolutionary Biology 87**
14.1 Innate & Adaptive Immunology, Antibody Architecture ($H_2L_2$) . 87
14.2 Major Infectious Diseases & *Plasmodium* Digenetic Life Cycle . . 88
14.3 Pathologies: HIV/AIDS Replication, Cancer Biology & Addiction Pharmacology . . . . . 90
14.4 Evolutionary Biology: Chemical Evolution, Natural Selection & Human Lineage . . . . . 91

**15 Ecology, Ecosystems & Biodiversity Conservation 93**
15.1 Organisms & Abiotic Adaptations . . . . . 93
15.2 Population Growth Formulations & Interspecific Interactions . . 94
15.3 Ecosystem Bioenergetics, Trophic Pyramids & Ecological Succession . . . . . 95
15.4 Global Biodiversity Metrics, Species-Area Law & Conservation . 96

**A High-Yield Quantitative Biology & Master Diagnostic Matrices 99**

A.1 Quantitative Genetics & Probability Calculus . . . . . 99

A.2 Recombination Frequency & Chromosome Mapping Calculus . . 100

A.3 Population Genetics: Hardy-Weinberg Mathematical Model . . . 101

A.4 Quantitative Human Physiology Matrices . . . . . 101

A.5 Master Systematic Comparison of All 7 Prescribed Plant Families 101

**About the Author & Errata Submissions 105**

PART I

# Cell Biology, Biomolecules & Cell Division

CHAPTER 1

# Cell Structure, Organelles & Biomolecules

The cell represents the fundamental structural and functional unit of all living organisms. In this chapter, we explore prokaryotic and eukaryotic cellular organization, the coordinated dynamics of the endomembrane system, autonomous and semi-autonomous organelles, the molecular architecture of biomacromolecules (carbohydrates, lipids, proteins, and nucleic acids), and the quantitative principles of enzyme catalysis, including Michaelis-Menten kinetics, $K_m$ constants, and competitive/non-competitive inhibition.

## 1.1 Cell Theory & Prokaryotic vs Eukaryotic Architecture

### Foundational Cell Theory & Prokaryotic Envelope Architecture

**1. Historical Milestones in Cell Biology:**

- **Anton van Leeuwenhoek (1674):** First observed and described live microscopic cells (bacteria, protozoa, spermatozoa).
- **Robert Brown (1831):** Discovered and named the **nucleus** in orchid root cells.
- **Cell Theory Formulation:**
  - **Matthias Schleiden (1838):** German botanist; examined large varieties of plants and concluded that all plants are composed of different kinds of cells that form plant tissues.
  - **Theodore Schwann (1839):** British zoologist; studied diverse animal cells and reported that animal cells have a thin outer layer (plasma membrane). He also concluded that the presence of a **cell wall is a unique diagnostic feature of plant cells**.
- **Rudolf Virchow (1855):** Resolved the origin of new cells with his famous aphorism:

  *Omnis cellula-e cellula*

  (All living cells arise from pre-existing cells via division).

**2. Prokaryotic Diagnostic Architecture ($1 - 2\,\mu$m):** Prokaryotes (Bacteria, Blue-green algae, Mycoplasma, PPLO) lack membrane-bound nuclei and organelles:

- **Three-Tiered Cell Envelope:**
  1. **Glycocalyx:** Outermost layer. Present either as a loose, gelatinous sheath called a **slime layer**, or as a thick, rigid, tough protective **capsule**.
  2. **Cell Wall:** Middle rigid layer composed of **peptidoglycan (murein)**; provides structural shape and prevents osmotic bursting or collapsing.
  3. **Plasma Membrane:** Innermost selectively permeable lipid bilayer; interacts with the external environment.
- **Mesosomes:** Characteristic membranous invaginations of the plasma membrane into the cytoplasm forming vesicles, tubules, and lamellae. Essential for: (i) cell wall synthesis, (ii) DNA replication and distribution to daughter cells, (iii) cellular respiration, and (iv) increasing enzymatic surface area.
- **Plasmids:** Small, circular, self-replicating extrachromosomal double-stranded DNA molecules; confer specific phenotypic traits such as antibiotic resistance (e.g., ampicillin resistance).
- **Inclusion Bodies:** Non-membrane-bound reserve storage granules lying free in the prokaryotic cytoplasm (e.g., phosphate granules, cyanophycean granules, glycogen granules, and gas vacuoles in photosynthetic prokaryotes).
- **Ribosomes ($70S$):** Composed of two subunits: $50S$ (large) and $30S$ (small). Several ribosomes can attach to a single mRNA strand forming a **polyribosome (polysome)**.

## 1.2 The Endomembrane System & Organelle Specialization

### The Eukaryotic Endomembrane System

The endomembrane system includes functionally coordinated membrane-bound organelles whose activities are tightly interlinked: **Endoplasmic Reticulum (ER), Golgi Apparatus, Lysosomes, and Vacuoles**. Mitochondria, Chloroplasts, and Peroxisomes are **excluded** because their functions are not coordinated with these components.

- **Fluid Mosaic Model of Plasma Membrane (Singer & Nicolson, 1972):** A quasi-fluid bilayer of phospholipids with embedded peripheral and integral proteins. Lipids provide fluidity ($L_\alpha$ phase), allowing lateral diffusion of proteins measured as membrane viscosity. Phospholipids have hydrophilic polar heads oriented outwards and hydrophobic non-polar hydrocarbon tails tucked inwards.
- **Endoplasmic Reticulum (ER):** Divides intracellular space into two compartments: luminal (inside ER) and extra-luminal (cytoplasm).

  - **Rough ER (RER):** Outer surface studded with $80S$ ribosomes; extensive and continuous with the outer nuclear membrane; actively involved in **protein synthesis and secretory trafficking**.
  - **Smooth ER (SER):** Lacks ribosomes; major site for the synthesis of **lipids and steroid hormones** (e.g., testosterone, estrogen); plays a vital role in drug and xenobiotic detoxification in hepatocytes.
- **Golgi Apparatus (Camillo Golgi, 1898):** Consists of flat, disc-shaped membranous sacs or cisternae ($0.5 - 1.0\,\mu$m) stacked concentrically near the nucleus. Exhibits distinct morphological polarity:
  - **Cis-Face (Convex / Forming Face):** Receives transitional transport vesicles budding from the ER.
  - **Trans-Face (Concave / Maturing Face):** Releases packaged secretory vesicles into the cytoplasm.
  - **Primary Function:** Crucial site for post-translational modification, specifically **glycosylation of proteins (forming glycoproteins)** and **lipids (forming glycolipids)**.
- **Lysosomes:** Membrane-bound vesicular structures formed by Golgi packaging. Rich in all types of **acid hydrolases** (lipases, proteases, carbohydrases, nucleases) optimally active at **acidic pH** ($\approx 4.5 - 5.0$) maintained by an active proton-pumping $H^+$-ATPase in the lysosomal membrane.
- **Vacuoles:** Bound by a single semi-permeable membrane called the **Tonoplast**. In plant cells, vacuoles occupy up to 90% of total cell volume. The tonoplast actively transports ions against concentration gradients into the vacuole; hence, luminal ion concentration is significantly higher than in cytoplasm. Specialized vacuoles include **contractile vacuoles** for osmoregulation in *Amoeba* and **food vacuoles** in *Paramecium*.

### Semi-Autonomous Organelles, Cytoskeleton & Nucleus

- **Mitochondria (Powerhouses of the Cell):** Double-membrane bound. Outer membrane is smooth; inner membrane forms deep inward foldings called **Cristae** (increase surface area for oxidative phosphorylation). Matrix possesses a single circular dsDNA molecule, a few RNA molecules, $70S$ ribosomes, and enzymes for the Krebs cycle. Inner membrane houses complexes of the Electron Transport System (ETS) and $F_0 - F_1$ ATP Synthase. Divide by **fission**.
- **Plastids (Chloroplasts, Chromoplasts, Leucoplasts):**
  - **Chloroplasts:** Double-membrane semi-autonomous organelles. Stroma contains circular dsDNA, $70S$ ribosomes, and soluble Calvin cycle enzymes (e.g., RuBisCO). Membrane thylakoids are stacked into **Grana**, interconnected by unstacked **Stroma Lamellae**.
  - **Chromoplasts:** Contain fat-soluble carotenoid pigments (carotene, xanthophylls) imparting yellow, orange, or red colors to flowers and fruits.

  - **Leucoplasts:** Colorless storage plastids: **Amyloplasts** store starch (potato); **Elaioplasts** store oils and fats; **Aleuroplasts** store proteins.
- **Cilia & Flagella (Axoneme Architecture):** Core axoneme possesses $9+2$ **microtubular doublet arrangement** (9 peripheral doublets and 2 central singlets) composed of the protein **tubulin**, enclosed by the plasma membrane. Cilia and flagella emerge from centriole-like basal bodies possessing a $9+0$ **cartwheel array**.
- **Nucleus & Chromatin (Robert Brown, 1831; Flemming):** Enclosed by a double nuclear envelope perforated by nuclear pores ($10-20\,\text{nm}$) formed by the fusion of inner and outer membranes. The nucleolus is a non-membrane bound site for **active ribosomal RNA (rRNA) synthesis**. Chromatin condenses during cell division into chromosomes characterized by a primary constriction called the **Centromere** bearing protein discs called **Kinetochores**.

## 1.3 Biomolecules, Structural Conformation & Enzyme Kinetics

### Macromolecular Architecture: Carbohydrates, Lipids & Proteins

**1. Carbohydrates:**

- Monosaccharides (e.g., glucose, ribose) are building blocks joined by **glycosidic bonds** formed by dehydration synthesis.
- Polysaccharides: **Starch** (energy reserve in plants, forms helical secondary structure that traps $I_2$ molecules turning blue); **Cellulose** (linear polymer of $\beta$-D-glucose with $\beta(1\rightarrow4)$ linkages; unbranched, no helices, cannot hold $I_2$); **Glycogen** (animal storage, branched polymer with $\alpha(1\rightarrow4)$ and $\alpha(1\rightarrow6)$ linkages; right end is reducing, left end non-reducing); **Inulin** (homopolymer of fructose).

**2. Lipids:** Water-insoluble organic molecules. Simple lipids are esters of fatty acids with glycerol (triacylglycerols). Complex phospholipids (e.g., **Lecithin**) contain a phosphorylated choline head and two fatty acid tails, forming fundamental cell membrane bilayers.

**3. Protein Structural Hierarchy:** Proteins are heteropolymers of $\alpha$-amino acids linked by **peptide bonds** ($-\text{CO}-\text{NH}-$):

- **Primary Structure ($1^\circ$):** Linear positional amino acid sequence; determines genetic identity.
- **Secondary Structure ($2^\circ$):** Local folding stabilized by intramolecular hydrogen bonds: right-handed $\alpha$-helix and $\beta$-pleated sheets.

- **Tertiary Structure ($3^\circ$):** Overall 3D spatial conformation of a single polypeptide folded upon itself; stabilized by disulfide bonds ($-\mathrm{S}-\mathrm{S}-$), hydrogen bonds, ionic interactions, and hydrophobic interactions. Absolutely **essential for biological/catalytic activity**.
- **Quaternary Structure ($4^\circ$):** Spatial arrangement of multiple folded polypeptide subunits (e.g., adult human hemoglobin consists of 4 subunits: $2\alpha$ and $2\beta$ chains).

### 1.3.1 Enzyme Catalytic Action & Michaelis-Menten Saturation Kinetics

Enzymes are specialized biological catalysts. While the vast majority are globular proteins, catalytic RNA molecules known as **Ribozymes** (such as the $23S$ rRNA peptidyl transferase in prokaryotic ribosomes) also carry out enzymatic cleavage and ligation.

Enzymes accelerate biochemical reactions by factors of $10^6$ to $10^{12}$ strictly through **lowering the activation energy barrier ($E_a$)** required to transition substrates into the unstable transition-state complex. An enzyme does not alter the overall free energy change ($\Delta G^\circ$) of a reaction, nor does it shift the equilibrium constant ($K_{\text{eq}}$). It merely accelerates the rate at which equilibrium is reached.

**Hyperbolic Saturation Kinetics:** Consider an enzyme $E$ binding reversibly to substrate $S$ to form the non-covalent enzyme-substrate intermediate complex $ES$, which subsequently dissociates to yield regenerated enzyme and product $P$:

$$E + S \underset{k_{-1}}{\overset{k_1}{\rightleftharpoons}} ES \xrightarrow{k_2} E + P \tag{1.1}$$

Under steady-state conditions ($d[ES]/dt = 0$), the initial reaction velocity $v$ as a function of substrate concentration $[S]$ follows the Michaelis-Menten hyperbolic relation:

$$\mathbf{v} = \frac{\mathbf{V}_{\max}[\mathbf{S}]}{\mathbf{K_m} + [\mathbf{S}]} \tag{1.2}$$

where $V_{\max}$ is the theoretical maximum velocity when all catalytic active sites are saturated with substrate, and $K_m$ is the **Michaelis constant**, defined mathematically as the substrate concentration at which reaction velocity reaches half of its maximal rate ($v = \frac{1}{2}V_{\max}$).

**Physiological Significance of $K_m$:** The Michaelis constant $K_m$ is an intrinsic physical constant that measures enzyme-substrate binding affinity:

$$K_m \propto \frac{1}{\text{Substrate Affinity}} \tag{1.3}$$

An enzyme with a low $K_m$ achieves half-maximal saturation at low substrate concentrations, indicating high affinity. Conversely, an enzyme with a high $K_m$ requires elevated substrate levels to achieve catalysis, indicating lower affinity.





**Figure 1.1:** Michaelis-Menten Enzyme Kinetics: Hyperbolic Saturation Curve and the Modulating Effect of a Competitive Inhibitor.

### Michaelis-Menten Equation & Lineweaver-Burk Double-Reciprocal Formulation

Taking the algebraic reciprocal of the Michaelis-Menten equation yields a linear formulation ($y = mx + c$):

$$\frac{\mathbf{1}}{\mathbf{v}} = \left(\frac{\mathbf{K_m}}{\mathbf{V}_{\max}}\right)\frac{\mathbf{1}}{[\mathbf{S}]} + \frac{\mathbf{1}}{\mathbf{V}_{\max}} \tag{1.4}$$

- Y-intercept $= \frac{1}{V_{\max}}$ X-intercept $= -\frac{1}{K_m}$ Slope $= \frac{K_m}{V_{\max}}$

### 1.3.2 Competitive vs. Non-Competitive Enzyme Inhibition

| Inhibition | $K_m$ | $V_{\max}$ | Mechanism & Canonical NCERT Examples |
|---|---|---|---|
| **Competitive** | **Increases**($K'_m > K_m$) | **Unchanged** | Inhibitor competes directly for active catalytic site. Excess $[S]$ overcomes inhibition. *Example:* **succinate dehydrogenase** inhibited by **malonate**. |
| **Non-Competitive** | **Unchanged** | **Decreases**($V'_{\max} < V_{\max}$) | Inhibitor binds allosteric site, altering enzyme conformation. Excess $[S]$ cannot overcome. *Example:* Cyanide inhibiting **cytochrome** $c$ **oxidase**. |

**High-Frequency Enzyme Inhibition Traps in NEET**

- **Competitive Reversibility:** Why does $V_{\max}$ remain unchanged in competitive inhibition? Because at infinitely high $[S]$, substrate molecules outcompete inhibitor molecules for every active site, achieving full theoretical $V_{\max}$.
- **Malonate vs. Succinate:** Malonate has two carboxyl groups ($-OOC - CH_2 - COO^-$) mimicking succinate ($-OOC - CH_2 - CH_2 - COO^-$), but lacks the two-carbon backbone required for dehydrogenation to fumarate.

### 1.3.3 IUBMB Enzyme Classification & Cofactor Architecture

The International Union of Biochemistry and Molecular Biology (IUBMB) classifies enzymes into six systematic classes:

1. **Oxidoreductases / Dehydrogenases:** Catalyze oxidation-reduction reactions between two substrates ($S_{\text{red}} + S'_{\text{ox}} \rightleftharpoons S_{\text{ox}} + S'_{\text{red}}$).
2. **Transferases:** Catalyze group transfer (other than hydrogen) between a pair of substrates ($S - G + S' \rightleftharpoons S' - G + S$).
3. **Hydrolases:** Catalyze cleavage of ester, ether, peptide, glycosidic, or C − C bonds via hydrolytic addition of water.

4. **Lyases:** Catalyze non-hydrolytic cleavage of bonds, resulting in the formation of double bonds without adding water.
5. **Isomerases:** Catalyze interconversion of geometric, positional, or optical isomers.
6. **Ligases:** Catalyze covalent coupling of two molecules coupled to ATP hydrolysis ($C-O, C-S, C-N, P-O$ bonds).

**Enzyme Cofactors (Non-Protein Components):** Many enzymes require non-protein chemical components termed **cofactors** to become catalytically active. The inactive protein portion alone is termed an **apoenzyme**; the catalytically active enzyme-cofactor complex is termed a **holoenzyme** (Holoenzyme = Apoenzyme + Cofactor).

- **Prosthetic Groups:** Organic compounds distinguished by being tightly and permanently bound to the apoenzyme (e.g., **Haem** in catalase and peroxidase).
- **Coenzymes:** Organic compounds that are only transiently associated with the apoenzyme during catalytic turnover; almost all coenzymes are derived from dietary vitamins (e.g., **NAD** and **NADP** contain niacin).
- **Metal Ions:** Form coordinate bonds with active site side-chains and substrate molecules (e.g., $Zn^{2+}$ in carbonic anhydrase and carboxypeptidase).

CHAPTER 2

# Cell Cycle, Mitosis & Meiosis

The continuity of life depends fundamentally upon cell division. This chapter analyzes the eukaryotic cell cycle, the biochemical checkpoints governing phase transitions, the quantitative dynamics of chromosome number ($2n$) and DNA content ($C$), the mechanical stages of mitosis and cytokinesis, and the specialized molecular stages of meiotic Prophase I that drive genetic recombination.

## 2.1 Cell Cycle Progression & DNA/Chromosome Quantitation

### The Eukaryotic Cell Cycle: Phase Duration & Quantitative Dynamics

A typical human cell in culture divides once every $\approx 24$ hours (yeast completes division in only 90 minutes). The cell cycle is partitioned into two core phases:

$$\begin{aligned}\text{Cell Cycle} = &\ \text{Interphase } (> 95\% \text{ duration}) \\ &+ \text{M-Phase / Mitosis } (< 5\% \text{ duration})\end{aligned}$$

**1. Interphase Sub-Phases:**

- $G_1$ **Phase (Gap 1 / Pre-DNA Synthesis):** Interval between mitosis and initiation of DNA replication. Cell is metabolically active and continuously grows, synthesizing RNA and proteins without replicating DNA.
- **S Phase (Synthesis Phase): DNA replication occurs**. The amount of DNA per cell doubles from $2C \rightarrow 4C$. However, **chromosome number remains strictly unchanged** ($2n \rightarrow 2n$). Centriole duplications occur in the cytoplasm of animal cells.
- $G_2$ **Phase (Gap 2 / Post-DNA Synthesis):** Proteins (notably tubulin for mitotic spindle) are synthesized while cell growth continues in preparation for mitosis.
- $G_0$ **Phase (Quiescent Stage):** Cells that do not divide exit $G_1$ to enter an inactive metabolic resting state (e.g., adult heart muscle cells, neurons). Cells remain metabolically active but divide only when required to replace damaged tissue.

**2. Mathematical Matrix: Chromosome Number ($2n$) vs DNA Content ($C$):**

| Cell Stage | $G_1$ | S | $G_2$ | Meta | Ana (Mitosis) |
|---|---|---|---|---|---|
| **Chromosomes** | $2n$ | $2n$ | $2n$ | $2n$ | **4n** (Split) |
| **DNA Content** | $2C$ | **4C** ($\times 2$) | $4C$ | $4C$ | $4C$ |

## 2.2 Mitosis: Spindle Dynamics & Cytokinesis

### Mitotic Stages: Equational Nuclear Division

Mitosis produces genetically identical diploid daughter cells ($2n \rightarrow 2n$), partitioned into four sequential karyokinetic stages:

1. **Prophase:** Chromatin condenses into distinct chromosomes consisting of two sister chromatids joined at the centromere. Centrosomes migrate to opposite cell poles and radiate microtubule arrays called **asters** (2 asters + spindle = mitotic apparatus). Nucleolus, Golgi complexes, ER, and nuclear envelope break down completely.
2. **Metaphase:** Nuclear envelope is completely absent. Chromosomes lie aligned along the equatorial plane (**Metaphase Plate**). Spindle fibers from opposite poles attach to proteinaceous discs called **Kinetochores** located on centromeres. Metaphase is the **optimal stage to study chromosome morphology and size**.
3. **Anaphase:** Centromere of each chromosome splits synchronously, separating sister chromatids into individual daughter chromosomes. Kinetochore microtubules shorten, pulling chromosomes toward opposite poles. Centromeres lead toward the pole with chromatid arms trailing behind, displaying characteristic shapes:
   - **V-shape:** Metacentric chromosome (median centromere).
   - **L-shape:** Sub-metacentric chromosome (sub-median centromere).
   - **J-shape:** Acrocentric chromosome (sub-terminal centromere).
   - **I-shape:** Telocentric chromosome (strictly terminal centromere).
4. **Telophase:** Chromosomes reach opposite poles, decondense, and lose their morphological individuality. Nuclear envelope reassembles around each chromosome cluster. Nucleolus, Golgi complex, and ER reform.

**Cytokinesis (Division of Cytoplasm):**

- **Animal Cells:** Accomplished by formation of a peripheral contractile **cleavage furrow** in plasma membrane that deepens centripetally.
- **Plant Cells:** Rigid cell walls prevent furrowing. Cytokinesis proceeds centrifugally via formation of a precursor **Cell Plate** (phragmoplast) that develops into the middle lamella (calcium and magnesium pectate).

- **Syncytium Formation:** When karyokinesis is not followed by cytokinesis, a multinucleate condition arises (e.g., liquid endosperm in coconut).

## 2.3 Meiosis: Prophase I Molecular Substages & Segregation

### Meiotic Sub-Stages: Reductional Division & Genetic Recombination

Meiosis reduces diploid chromosome number to haploid ($2n \rightarrow n$), ensuring conservation of species-specific chromosome counts across sexual generations while generating genetic diversity.

**1. Five Sequential Substages of Prophase I:**

1. **Leptotene:** Chromatin compaction proceeds continuously; chromosomes become progressively visible under light microscopy as slender threads.
2. **Zygotene:** Homologous chromosomes begin pairing lengthwise via an active alignment process called **Synapsis**. Pairing is mediated by a complex tripartite protein scaffold called the **Synaptonemal Complex**. The paired homologous structure is termed a **Bivalent** or a **Tetrad** (four chromatids).
3. **Pachytene:** Characterized by appearance of **Recombination Nodules**. Non-sister chromatids of homologous chromosomes exchange reciprocal genetic segments (**Crossing Over**), mediated by the multienzyme complex **Recombinase**. Pachytene generates novel allelic combinations.
4. **Diplotene:** Marked by the dissolution of the synaptonemal complex. Homologous chromosomes separate from each other except at crossover points, creating distinct X-shaped configurations called **Chiasmata**. In vertebrate oocytes, diplotene can remain arrested for months or years (**Dictyotene stage**).
5. **Diakinesis:** Final substage marked by the complete **Terminalization of chiasmata**. Chromosomes reach maximum condensation; nucleolus disappears and nuclear envelope breaks down.

**2. Anaphase I vs Anaphase II Segregation Mechanics:**

- **Anaphase I (Reductional Disjunction):** Homologous chromosomes separate to opposite poles, but **sister chromatids remain connected at their centromeres**. Consequently, chromosome number is halved ($2n \rightarrow n$).
- **Anaphase II (Equational Disjunction): Centromeres split synchronously**, allowing sister chromatids to separate and move toward opposite poles, identical to mitotic anaphase.

PART II

# Structural Organisation & Diversity

CHAPTER 3

# Plant Morphology & Systematic Angiosperm Families

Angiosperm morphology encompasses the structural diversification of vegetative and reproductive organs enabling survival across varied habitats. This chapter details root and stem adaptations, leaf venation and phyllotaxy, inflorescence architectures, floral organography, aestivation patterns, placentation modalities, and the complete diagnostic profiles and floral formulas of all seven plant families explicitly prescribed by the National Medical Commission (NMC).

## 3.1 Vegetative Morphology: Root, Stem, Leaf & Adaptations

### Root System Architecture & Functional Modifications

The root originates from the embryonic radicle during seed germination.

- **Root System Types:**
  - **Taproot System:** Direct elongation of radicle forming primary root with lateral branches of secondary and tertiary order (e.g., Mustard, Gram, Dicotyledonous plants).
  - **Fibrous Root System:** Primary root is short-lived in monocots and replaced by large numbers of slender roots originating from base of stem (e.g., Wheat).
  - **Adventitious Root System:** Roots arising from any plant part other than embryonic radicle (e.g., Grass, *Monstera*, Banyan tree).
- **Four Morphological Regions of Root (from apex upwards):**
  1. **Root Cap:** Thimble-like parenchymatous structure protecting tender apical meristem.
  2. **Region of Meristematic Activity:** Small, thin-walled cells with dense protoplasm dividing repeatedly.
  3. **Region of Elongation:** Rapid cellular elongation and enlargement; responsible for growth in root length.
  4. **Region of Maturation:** Cells differentiate into mature tissues. Bears delicate, thread-like **Root Hairs** (tubular epidermal outgrowths) maximizing water and mineral absorption.
- **High-Yield Physiological Modifications of Roots:**

  - **Food Storage:** Swollen taproots (Carrot, Turnip) and swollen adventitious roots (Sweet Potato / *Ipomoea batatas*).
  - **Mechanical Support:** Heavy pillar-like hanging **Prop Roots** of Banyan tree (*Ficus benghalensis*); **Stilt Roots** arising from lower stem nodes in Maize and Sugarcane.
  - **Pneumatophores (Respiratory Roots):** In mangrove halophytes (e.g., *Rhizophora*) growing in waterlogged, saline, oxygen-deficient swamps; roots grow vertically upwards (negatively geotropic) and possess microscopic lenticels (**pneumathodes**) to obtain atmospheric oxygen.

### Stem & Leaf Morphology, Phyllotaxy & Evolutionary Modifications

**1. Stem Modifications:** Stems originate from embryonic plumule and possess nodes, internodes, and axillary buds:

- **Underground Storage Stems:** Perennating organs storing reserve food: Stem Tuber (**Potato**), Rhizome (**Ginger, Turmeric**), Corm (**Zaminkand, *Colocasia***).
- **Stem Tendrils:** Slender, spirally coiled climbing organs modified from axillary buds (Gourds: Cucumber, Pumpkin, Watermelon; Grapevines).
- **Thorns:** Hard, woody, straight, pointed protective organs modified from axillary buds (e.g., *Citrus*, *Bougainvillea*).
- **Phylloclades:** Green, photosynthetic flattened stems (e.g., *Opuntia*) or cylindrical fleshy stems (e.g., *Euphorbia*) storing mucilaginous water in arid xerophytic habitats (leaves modified into spines).
- **Vegetative Propagation Stems: Runners** (Grass, Strawberry creeping horizontally), **Stolons** (Mint, Jasmine arched branches), **Offsets** (short lateral branch with rosette of leaves: *Pistia*, *Eichhornia*), **Suckers** (underground horizontal stem growing obliquely upwards: Banana, Pineapple, *Chrysanthemum*).

**2. Leaf Morphology & Venation:** Comprises leaf base (with pulvinus in legumes), petiole, and expanded lamina:

- **Venation: Reticulate venation** (veinlets form network, characteristic of Dicots); **Parallel venation** (veins run parallel to each other, characteristic of Monocots).
- **Phyllotaxy (Pattern of Leaf Arrangement on Stem/Branch):**
  - **Alternate:** Single leaf arises at each node alternately (China rose, Mustard, Sunflower).
  - **Opposite:** Pair of leaves arises at each node opposite to each other (*Calotropis*, Guava).
  - **Whorled:** More than two leaves arise at a node forming a whorl (*Alstonia*).

- **Leaf Modifications:** Tendrils for climbing (Pea); Spines for defense (Cacti); Fleshy leaves for food storage (Onion, Garlic); **Phyllode** (petiole expands and becomes green and photosynthetic: Australian *Acacia*); Insect traps (Pitcher plant, Venus flytrap).

## 3.2 Floral Morphology, Inflorescence, Aestivation & Placentation

### Inflorescence, Flower Symmetry & Floral Organ Insertion

**1. Inflorescence Architectures:** The arrangement of flowers on the floral axis (peduncle):

- **Racemose Inflorescence:** Main axis continues to grow indefinitely; flowers are borne laterally in an **acropetal succession** (oldest flowers at base, youngest at apex).
- **Cymose Inflorescence:** Main axis terminates in a flower, limiting indefinite growth; flowers are borne in **basipetal succession** (oldest flower at apex, youngest at base).

**2. Flower Symmetry Types:**

- **Actinomorphic (Radial Symmetry, $\oplus$):** Flower can be divided into two equal radial halves in any radial plane passing through center (e.g., Mustard, *Datura*, Chilli).
- **Zygomorphic (Bilateral Symmetry, %):** Flower can be divided into two similar halves only in one particular vertical plane (e.g., Pea, Gulmohur, Bean, *Cassia*).
- **Asymmetric / Irregular:** Flower cannot be divided into two equal halves by any vertical plane (e.g., *Canna*).

**3. Insertion of Floral Whorls Relative to Ovary:**

- **Hypogynous Flower (Superior Ovary, $\underline{\mathrm{G}}$):** Calyx, corolla, and stamens situated below ovary on thalamus (e.g., Mustard, China rose, Brinjal).
- **Perigynous Flower (Half-Inferior Ovary, G$-$):** Gynoecium is situated in center, and other parts lie on rim of thalamus at same level (e.g., Plum, Rose, Peach).
- **Epigynous Flower (Inferior Ovary, $\overline{\mathrm{G}}$):** Margin of thalamus grows upward enclosing ovary completely and getting fused with it; other parts arise above ovary (e.g., Guava, Cucumber, ray florets of Sunflower).

### Aestivation Patterns & Placentation Modalities

### 1. Aestivation (Arrangement of Sepals/Petals in Floral Bud):

| Aestivation Type | Structural Pattern | NCERT Examples |
|---|---|---|
| **Valvate** | Margins meet without overlapping. | *Calotropis* |
| **Twisted** | One margin overlaps that of next sequentially. | China rose, Lady's finger, Cotton |
| **Imbricate** | Margins overlap irregularly without direction. | *Cassia*, Gulmohur |
| **Vexillary** | Five petals: posterior **Standard** overlaps 2 **Wings**, which overlap 2 anterior **Keel**. | Pea, Bean, Fabaceae family |

### 2. Placentation (Arrangement of Ovules within the Ovary):

- **Marginal:** Placenta forms a ridge along ventral suture of ovary; ovules borne in two rows (e.g., **Pea**).
- **Axile:** Placenta is axial and ovules attached to it in multilocular ovary (e.g., **China rose, Tomato, Lemon**).
- **Parietal:** Ovules develop on inner wall of ovary or peripheral part; ovary unilocular but becomes two-celled due to formation of a false septum called **Replum** (e.g., **Mustard, *Argemone***).
- **Free Central:** Ovules borne on central axis and septa are completely absent (e.g., ***Dianthus*, Primrose**).
- **Basal:** Placenta develops at base of ovary and a **single ovule** is attached to it (e.g., **Sunflower, Marigold**).

### 3. Fruit & Seed Typology:

- **Drupe Fruit:** Developed from monocarpellary superior ovaries and are one-seeded: **Mango** (fleshy edible mesocarp, thin epicarp, stony hard endocarp) and **Coconut** (mesocarp is fibrous, enclosing liquid endosperm).
- **Seed Architecture:** Dicotyledonous seeds (Gram, Pea) are typically non-endospermic (endosperm consumed during embryogeny); Monocotyledonous seeds (Maize, Wheat) are typically endospermic, featuring an outer proteinaceous **aleurone layer**, a single shield-shaped cotyledon called **scutellum**, a plumule sheath (**coleoptile**), and a radicle sheath (**coleorhiza**). Castor is an endospermic dicot; Orchids are non-endospermic monocots.

## 3.3 Systematic Flora: Diagnostic Profiles & Floral Formulas of the 7 NMC Families

The National Medical Commission (NMC) explicitly mandates comprehensive comparative mastery of seven angiosperm families for NEET UG. These encompass five dicotyledonous families (Malvaceae, Cruciferae, Fabaceae, Asteraceae, Solanaceae) and two monocotyledonous families (Poaceae and Liliaceae).

### 3.3.1 1. Family Malvaceae (Mallow / Cotton Family)

- **Floral Formula:** $\oplus\, \mathrm{Epi}_{\mathbf{3-8}}\, \mathrm{K}_{(\mathbf{5})}\, \mathrm{C}_{\mathbf{5}}\, \mathrm{A}_{(\infty)}\, \underline{\mathrm{G}}_{(\mathbf{5-\infty})}$
- **Vegetative Characters:** Mucilaginous sap throughout tissues, stellate hairs on epidermis, alternate stipulate leaves with reticulate venation.
- **Floral Architecture:** Solitary axillary or terminal flowers; **Epicalyx** (involucel of 3–8 bracteoles) situated immediately below the calyx; corolla of 5 free petals with characteristically **twisted aestivation**.
- **Androecium & Gynoecium:** Stamens indefinite ($\infty$), **monadelphous** (filaments fused into a hollow cylindrical staminal tube surrounding the style); anthers are **monothecous and reniform (kidney-shaped)**. Gynoecium multicarpellary, syncarpous, superior, with axile placentation.
- **Economic Significance:** Textile fiber from epidermal seed hairs of Cotton (*Gossypium hirsutum*); edible pods of Lady's finger / Okra (*Abelmoschus esculentus*); ornamental shrubs like China rose (*Hibiscus rosa-sinensis*).

### 3.3.2 2. Family Cruciferae / Brassicaceae (Mustard Family)

- **Floral Formula:** $\oplus\, \mathrm{K}_{\mathbf{2+2}}\, \mathrm{C}_{\mathbf{4}}\, \mathrm{A}_{\mathbf{2+4}}\, \underline{\mathrm{G}}_{(\mathbf{2})}$
- **Vegetative Characters:** Herbaceous plants with pungent watery juice containing sulfur-rich glucosinolates (mustard oil odor).
- **Floral Architecture:** Corymbose raceme inflorescence; cruciform corolla consisting of 4 clawed petals arranged diagonally in the shape of a Maltese cross.
- **Androecium & Gynoecium:** Stamens 6, **tetradynamous** (2 short outer lateral stamens, 4 long inner median stamens). Gynoecium bicarpellary syncarpous; ovary unilocular at inception, becoming bilocular due to the development of a false membranous longitudinal septum called the **Replum**; parietal placentation; fruit is a specialized dehiscent capsule termed **siliqua** or silicula.
- **Economic Significance:** Edible oil from Mustard (*Brassica campestris*); edible taproots of Radish (*Raphanus sativus*) and Turnip (*Brassica rapa*); vegetables like Cabbage and Cauliflower.

### 3.3.3 3. Family Fabaceae / Leguminosae (Legume / Pea Family)

- **Floral Formula:** $\%\, \mathrm{K}_{(\mathbf{5})}\, \mathrm{C}_{\mathbf{1+2+(2)}}\, \mathrm{A}_{\mathbf{(9)+1}}\, \underline{\mathrm{G}}_{\mathbf{1}}$
- **Vegetative Characters:** Roots possess characteristic nodules hosting symbiotic nitrogen-fixing *Rhizobium* bacteria; pinnately compound leaves with swollen, sensitive **pulvinate leaf bases**.
- **Floral Architecture:** Zygomorphic pentamerous flower; papilionaceous corolla displaying **vexillary aestivation** consisting of 1 large posterior stan-

dard (vexillum), 2 lateral wings (alae), and 2 anterior petals fused to form a boat-shaped keel (carina) enclosing the essential organs.

- **Androecium & Gynoecium:** Stamens 10, **diadelphous** [(9)+1] (nine stamens fused by filaments into a gutter-shaped sheath, tenth posterior stamen free); dithecous anthers. Gynoecium monocarpellary, unilocular, superior ovary with ovules arranged along the ventral suture on **marginal placentation**; fruit is a **legume** dehiscing along both sutures.
- **Economic Significance:** Primary protein source through pulses (Gram, Arhar, Sem, Moong); edible oils (Soybean, Groundnut); blue dye (*Indigofera*); bast fibers (Sunnhemp); fodder (*Sesbania*, *Trifolium*); medicine (*Muliathi*).

### 3.3.4 4. Family Asteraceae / Compositae (Sunflower Family)

- **Floral Formula:**

$$\text{Ray Florets: } \% \, \mathrm{K}_{\text{pappus}} \, \mathrm{C}_{(\mathbf{5})} \, \mathrm{A}_{\mathbf{0}} \, \overline{\mathrm{G}}_{(\mathbf{2})}$$

$$\text{Disc Florets: } \oplus \, \mathrm{K}_{\text{pappus}} \, \overbrace{\mathrm{C}_{(5)} \, \mathrm{A}_{(5)}} \, \overline{\mathrm{G}}_{(\mathbf{2})}$$

- **Inflorescence Architecture:** Dense pseudanthium inflorescence known as a **Capitulum (Head)**, surrounded at its base by a protective involucre of green bracts. Central actinomorphic, bisexual tubular disc florets are flanked peripherally by zygomorphic, pistillate or sterile ligulate ray florets.
- **Diagnostic Hallmarks:** Calyx is modified into a persistent hairy or bristly tuft called the **Pappus**, which serves as a parachute for wind dispersal (anemochory). Androecium is **syngenesious** (anthers are fused into a cylinder around the style, while filaments remain free); epipetalous.
- **Gynoecium & Fruit:** Bicarpellary, syncarpous, **inferior unilocular ovary with a single basal ovule**; fruit is a dry indehiscent **cypsela** crowned with pappus.
- **Economic Significance:** Edible and commercial oil from Sunflower (*Helianthus annuus*) and Safflower; ornamentals like Marigold (*Tagetes*) and *Chrysanthemum*; natural botanical insecticide pyrethrum extracted from *Chrysanthemum cinerariifolium*.

### 3.3.5 5. Family Poaceae / Gramineae (Grass / Cereal Family)

- **Floral Formula:** $\% \, \mathrm{P}_{\mathbf{2}} \, \mathrm{A}_{\mathbf{3 \text{ or } 6}} \, \underline{\mathrm{G}}_{\mathbf{1}}$
- **Vegetative Characters:** Herbaceous monocots with cylindrical, hollow-jointed stems called **culms**; alternate, two-ranked distichous leaves with parallel venation, open sheathing leaf base, and a small membranous outgrowth called a **ligule** at the blade-sheath junction.

- **Inflorescence & Perianth:** Basic unit is a **Spikelet** bearing sterile bracts (empty glumes) at the base; each floret is enclosed between an outer bract (**lemma**, often bearing an awn) and an inner bracteole (**palea**). The perianth is reduced to 2 (rarely 3) minute, scale-like, hygroscopic fleshy structures termed **Lodicules**.
- **Androecium & Gynoecium:** Stamens typically 3 (or 6 in Rice and Bamboo) with long, slender filaments bearing versatile, pendulous anthers adapted for wind pollination (anemophily). Gynoecium monocarpellary, unilocular, superior ovary containing a single basal ovule, bearing 2 long **feathery bifid stigmas**. Fruit is a **caryopsis** (grain) in which the pericarp is completely fused with the seed coat.
- **Economic Significance:** Humanity's chief caloric cereals: Wheat (*Triticum aestivum*), Rice (*Oryza sativa*), Maize (*Zea mays*); commercial sugar from Sugarcane (*Saccharum officinarum*); structural culms from Bamboo (*Bambusa*).

### 3.3.6 6. Family Solanaceae (Potato / Nightshade Family)

- **Floral Formula:** $\oplus\, \mathrm{K}_{(\mathbf{5})}\ \overbrace{\mathrm{C}_{(5)}\ \mathrm{A}_{5}}\ \underline{\mathrm{G}}_{(\mathbf{2})}$
- **Diagnostic Hallmarks:** Persistent accrescent calyx that remains attached to the fruit; epipetalous stamens alternating with corolla lobes.
- **Gynoecium Architecture:** Bicarpellary, syncarpous, superior ovary positioned **obliquely at an angle of** $45^\circ$ **to the posterior-anterior axis**; bilocular with a **swollen axile placenta** bearing numerous amphitropous ovules. Fruit is a succulent berry or a septicidal capsule.
- **Economic Significance:** Staple food crops: Potato (*Solanum tuberosum*), Tomato (*Solanum lycopersicum*), Brinjal (*Solanum melongena*); spices: Chilli (*Capsicum annuum*); medicinal alkaloids: Atropine from *Atropa belladonna*, withanolides from *Withania somnifera* (Ashwagandha); fumigatory: Tobacco (*Nicotiana tabacum*); ornamental: *Petunia*.

### 3.3.7 7. Family Liliaceae (Lily Family)

- **Floral Formula:** $\oplus\ \overbrace{\mathrm{P}_{(3+3)}\ \mathrm{A}_{3+3}}\ \underline{\mathrm{G}}_{(\mathbf{3})}$
- **Diagnostic Hallmarks:** Perennial monocot herbs with underground bulbs, corms, or rhizomes; scapigerous inflorescence with umbellate clusters subtended by spathaceous bracts.
- **Perianth & Androecium:** Non-differentiated perianth composed of 6 petaloid tepals arranged in two trimerous whorls $(3+3)$. Stamens 6, **epiphyllous** $(3+3)$ inserted on tepals; dithecous anthers.
- **Gynoecium & Fruit:** Tricarpellary, syncarpous, trilocular superior ovary with numerous ovules on **axile placentation**. Fruit is a loculicidal capsule

(rarely berry); seeds are endospermic.

- **Economic Significance:** Vegetables and sulfur condiments: Onion (*Allium cepa*), Garlic (*Allium sativum*), *Asparagus*; skin pharmaceuticals: *Aloe vera*; horticultural ornamentals: Tulip (*Tulipa*), *Gloriosa superba*; mitotic spindle inhibitor: **Colchicine** extracted from the corms of *Colchicum autumnale*.

| Family | Floral Formula | Inflorescence & Perianth | Androecium Details | Gynoecium & Placentation | Fruit & Seed | Key Economic & Medicinal Species |
|---|---|---|---|---|---|---|
| **Malvaceae**<br><br>epicalyx (3 − 8);<br>twisted corolla<br>**monothecous reniform anthers**<br>**axile placentation**<br>hairy seeds | $\oplus\ \mathrm{Epi}_{3-8}\ \mathrm{K}_{(5)}\ \mathrm{C}_5\ \mathrm{A}_{(\infty)}\ \underline{\mathrm{G}}_{(5-\infty)}$ | Solitary or axillary; | **Monadelphous** staminal tube; | Multicarpellary syncarpous; | Capsule or schizocarp; | Cotton (*Gossypium*), Lady finger (*Abelmoschus*), China rose (*Hibiscus*). |
| **Cruciferae**<br><br>cruciform corolla (clawed petals)<br>(2 outer short + 4 inner long)<br>**replum (false septum)**;<br>parietal placentation<br>non-endospermic | $\oplus\ \mathrm{K}_{2+2}\ \mathrm{C}_4\ \mathrm{A}_{2+4}\ \underline{\mathrm{G}}_{(2)}$ | Racemose raceme; | **Tetradynamous** | Bicarpellary syncarpous; | **Siliqua** or silicula; | Mustard (*Brassica campestris*), Radish (*Raphanus*), Turnip, Cabbage. |
| **Fabaceae**<br><br>(1 standard + 2 wings + 2 keel)<br>dithecous anthers<br>**marginal placentation**<br>non-endospermic; root nodules | $\%\ \mathrm{K}_{(5)}\ \mathrm{C}_{1+2+(2)}\ \mathrm{A}_{(9)+1}\ \underline{\mathrm{G}}_1$ | Racemose; **vexillary aestivation** | **Diadelphous** [(9) + 1], | Monocarpellary unilocular; | **Legume** (pod); | Gram, Arhar, Moong, Soybean, Groundnut, *Indigofera*, Sunnhemp, *Sesbania*, *Muliathi*. |
| **Asteraceae**<br><br>calyx modified to **Pappus**<br>filaments free), epipetalous<br>**inferior ovary**, **basal placentation**<br>non-endospermic | Ray: $\%\ \mathrm{K}_{\mathrm{pappus}}\ \mathrm{C}_{(5)}\ \mathrm{A}_0\ \overline{\mathrm{G}}_{(2)}$<br>Disc: $\oplus\ \mathrm{K}_{\mathrm{pappus}}\ \overbrace{\mathrm{C}_{(5)}\ \mathrm{A}_{(5)}}\ \overline{\mathrm{G}}_{(2)}$ | **Capitulum (Head)** with involucre; | **Syngenesious** (anthers fused, | Bicarpellary syncarpous; | **Cypsela** with pappus; | Sunflower (*Helianthus*), Marigold (*Tagetes*), *Chrysanthemum*, Pyrethrum. |

**Table 3.1:** Definitive Diagnostic Matrix for NMC Angiosperm Families (Part I: Malvaceae, Cruciferae, Fabaceae, Asteraceae).

| Family | Floral Formula | Inflorescence & Perianth | Androecium Details | Gynoecium & Placentation | Fruit & Seed | Key Economic & Medicinal Species |
|---|---|---|---|---|---|---|
| **Poaceae**<br><br>lemma, palea; 2 **Lodicules** (3 or 6)<br>**feathery bifid stigmas**; basal placentation with testa); endospermic | $\% \, P_2 \, A_{3 \text{ or } 6} \, \underline{G}_1$<br><br>Versatile pendulous stamens<br>Monocarpellary unilocular;<br><br>**Caryopsis** (pericarp fused<br>Wheat (*Triticum*), Rice (*Oryza*), Maize (*Zea mays*), Sugarcane, Bamboo. | **Spikelet** with sterile glumes, | | | | |
| **Solanaceae**<br><br>persistent accrescent calyx<br>alternipetalous<br>**obliquely placed ovary**; **swollen axile placenta**<br>numerous endospermic seeds | $\oplus \, K_{(5)} \, \overbrace{C_{(5)} \, A_5} \, \underline{G}_{(2)}$<br><br>Epipetalous stamens (5),<br>Bicarpellary syncarpous;<br><br>Berry or capsule;<br>Potato, Tomato, Brinjal, Chilli, *Belladonna*, *Ashwagandha*, Tobacco, *Petunia*. | Solitary, axillary or cymose; | | | | |
| **Liliaceae**<br><br>perianth $3+3$ tepals in two whorls<br>trilocular, **axile placentation**<br>endospermic seeds | $\oplus \, \overbrace{P_{(3+3)} \, A_{3+3}} \, \underline{G}_{(3)}$<br><br>**Epiphyllous** stamens $(3+3)$<br>Tricarpellary syncarpous;<br>Capsule or berry;<br>Onion, Garlic (*Allium*), *Aloe*, *Asparagus*, Tulip, *Gloriosa*, Colchicine (*Colchicum*). | Scapigerous umbellate clusters; | | | | |

**Table 3.2:** Definitive Diagnostic Matrix for NMC Angiosperm Families (Part II: Poaceae, Solanaceae, Liliaceae).

Authored by Satya Prakash · spdedsec@hotmail.com

CHAPTER 4

# Animal Diversity & Comparative Anatomy: Frog and Cockroach

The animal kingdom displays remarkable organizational diversity across twelve major phyla. Furthermore, the revised NEET syllabus mandates detailed comparative anatomical mastery of two key model organisms: the amphibian *Rana tigrina* (Frog) and the arthropod *Periplaneta americana* (Cockroach). This chapter provides an integrated anatomical compendium encompassing their digestive, circulatory, respiratory, excretory, nervous, and reproductive systems.

## 4.1 Diagnostic Hallmarks of Major Animal Phyla

The Animal Kingdom comprises multicellular, heterotrophic eukaryotes lacking cell walls. Phylogenetic classification is established upon grades of organization (cellular, tissue, organ, organ-system), body symmetry (asymmetry, radial, bilateral), germ layers (diploblastic, triploblastic), coelomic cavity (acoelomate, pseudocoelomate, coelomate), and metameric segmentation.

## 4.2 Anatomy of the Cockroach (*Periplaneta americana*)

### Cockroach Morphological Systems & Internal Organography

**1. Exoskeleton & Segmentation:** Covered by a hard, brown, chitinous exoskeleton consisting of hardened plates called **Sclerites**: dorsal **Tergites** and ventral **Sternites**, joined laterally by a delicate, flexible articular membrane (**arthrodial membrane**).

**2. Cephalic Architecture & Mouthparts:**

- Head is triangular, hypognathous (directed downward at 90°), formed by fusion of **6 embryonic segments**.
- Bears a pair of compound eyes, each composed of approximately **2000 hexagonal ommatidia**, generating mosaic vision characterized by high sensitivity but low resolution.
- **Biting and Chewing Mouthparts:** Labrum (upper lip), paired Mandibles with chitinous grinding and incising teeth, paired Maxillae, Labium (lower lip), and median flexible lobe **Hypopharynx** (tongue) into which salivary duct opens.

| Phylum | Level / Symmetry | Coelom / Germ | Diagnostic Hallmarks & NCERT Examples |
|---|---|---|---|
| **Porifera** | Cellular; Asymmetric | Acoelomate; Diploblastic | Spongocoel lined by flagellated **Choanocytes (Collar cells)**; Water canal system; Spicules/spongin skeleton. *Sycon, Spongilla, Euspongia.* |
| **Cnidaria** | Tissue; Radial | Acoelomate; Diploblastic | **Cnidoblasts/Nematocysts** for anchorage/defense; Polyp (sessile) and Medusa (free-swimming); **Metagenesis** in *Obelia. Hydra, Physalia, Adamsia.* |
| **Ctenophora** | Tissue; Radial | Acoelomate; Diploblastic | Eight external ciliated **comb plates** for locomotion; **Bioluminescence**; Strictly marine. *Pleurobrachia, Ctenoplana.* |
| **Platyhelminthes** | Organ/System; Bilateral | **Acoelomate**; Triploblastic | Dorso-ventrally flattened; **Flame cells (Protonephridia)** for osmoregulation; Parasitic hooks/suckers. *Taenia, Fasciola, Planaria* (high regeneration). |
| **Aschelminthes** | Organ-system; Bilateral | **Pseudocoelomate**; Triploblastic | Syncytial epidermis, muscular pharynx, sexual dimorphism (females longer). *Ascaris, Wuchereria, Ancylostoma.* |
| **Annelida** | Organ-system; Bilateral | Coelomate; Triploblastic | **Metameric segmentation**; Longitudinal/circular muscles; **Nephridia** for excretion; Closed circulation. *Nereis, Pheretima, Hirudinaria.* |
| **Arthropoda** | Organ-system; Bilateral | Coelomate; Triploblastic | **Largest phylum** ($>$ 2/3 animal species); Chitinous exoskeleton; **Malpighian tubules**; Open circulation; Statocysts. *Apis, Bombyx, Periplaneta, Limulus.* |
| **Mollusca** | Organ-system; Bilateral | Coelomate; Triploblastic | **Second largest phylum**; Calcareous shell; Visceral hump with mantle cavity; Rasping **Radula**. *Pila, Pinctada, Sepia, Octopus.* |
| **Echinodermata** | Organ-system; Radial (adult), Bilateral (larva) | Coelomate; Triploblastic | Endoskeleton of calcareous ossicles; **Water vascular system** (ambulacral system); Excretory system absent. *Asterias, Echinus, Antedon.* |
| **Hemichordata** | Organ-system; Bilateral | Coelomate; Triploblastic | Worm-like marine; Proboscis, collar (**stomochord**), and trunk; Proboscis gland for excretion. *Balanoglossus, Saccoglossus.* |
| **Chordata** | Organ-system; Bilateral | Coelomate; Triploblastic | (1) **Notochord**, (2) **Dorsal hollow nerve cord**, (3) **Pharyngeal gill slits**, (4) **Post-anal tail**. Subphyla: Urochordata, Cephalochordata, Vertebrata. |

**Table 4.1:** Systematic Comparative Diagnostic Matrix for the Major Animal Phyla.

**3. Thoracic & Abdominal Anatomy:**

- Thorax consists of Prothorax, Mesothorax, and Metathorax.
- **Forewings (Tegmina):** Mesothoracic origin; opaque, leathery, dark, protective, covering hindwings at rest.
- **Hindwings:** Metathoracic origin; transparent, membranous, utilized in flight.
- Abdomen consists of **10 segments** in both sexes.
- **Sexual Dimorphism:** Males bear a pair of short, unjointed thread-like **Anal Styles** on the $9^{th}$ sternite (strictly absent in females). Both sexes bear jointed filamentous **Anal Cerci** on the $10^{th}$ abdominal segment.

**4. Visceral Organ Systems:**

- **Alimentary Canal:** Mouth → Pharynx → Oesophagus → **Crop** (large thin-walled sac for food storage) → **Gizzard / Proventriculus** (outer thick circular muscle layer and inner cuticular lining forming **6 hard chitinous teeth** for grinding food).
  - **Hepatic / Gastric Caeca:** Ring of **6–8 blind tubules** at the junction of foregut and midgut; secrete digestive enzymes.
  - **Malpighian Tubules:** Ring of **100–150 thin yellow filamentous tubules** at junction of midgut and hindgut; absorb nitrogenous wastes from hemolymph and excrete them as potassium urate → uric acid (**Uricotelic**).
- **Circulatory System:** Open type; hemolymph flows in hemocoel bathing visceral organs. Heart consists of an elongated muscular tube with **13 funnel-shaped contractile chambers** with ostia.
- **Respiratory System:** Network of tracheae branching into tracheoles; opens externally via **10 pairs of spiracles** (2 pairs thoracic, 8 pairs abdominal).
- **Nervous System:** Fused ganglia linked by paired longitudinal connectives: **3 ganglia in thorax and 6 ganglia in abdomen**. If head is decapitated, cockroach survives for up to a week because head contains only a small portion of nervous system.
- **Reproductive Systems:** Male has paired testes in $4^{th} - 6^{th}$ segments, mushroom-shaped gland in $6^{th} - 7^{th}$ segments, and chitinous asymmetric **phallomeres (gonapophyses)**; sperms glued into bundles called **spermatophores**. Female has paired ovaries in $2^{nd} - 6^{th}$ segments (each ovary composed of 8 ovarian tubules/ovarioles), paired spermathecae in $6^{th}$ segment, and collateral glands. Eggs enclosed in dark reddish-brown capsules called **Oothecae** (8 mm long, containing 14–16 eggs). Development is **paurometabolous** (13 nymphal instars).

## 4.3 Anatomy of the Frog (*Rana tigrina*)

### Frog Morphological Systems & Internal Organography

**1. Morphology & External Characters:** Amphibian, cold-blooded (**Poikilotherm**); exhibits protective color change (**Mimicry / Camouflage**). Skin is smooth, moist, and slimy due to mucus secretion, functioning as a major respiratory organ (**Cutaneous Respiration**). Body divided into **Head and Trunk** (neck and tail absent). Eyes are protected by a transparent **Nictitating Membrane**.

- **Sexual Dimorphism:** Male frogs possess sound-amplifying **Vocal Sacs** under throat and a specialized **Copulatory / Nuptial Pad** on the first digit of the forelimb (both absent in female frogs).

**2. Internal Anatomical Systems:**

- **Digestive System:** Alimentary canal is short (carnivorous diet). Mouth leads into buccal cavity with bilobed muscular tongue attached anteriorly. Stomach secretes HCl and gastric juice. Partially digested food (**Chyme**) passes to duodenum. Duodenum receives bile and pancreatic juice through common bile duct. Undigested waste enters rectum and exits via **Cloaca** (common chamber receiving digestive, excretory, and reproductive products).
- **Circulatory System:** Closed vascular system with a **3-chambered heart (2 Atria and 1 Ventricle)**.
  - Dorsal triangular **Sinus Venosus** receives deoxygenated blood from vena cava and joins right atrium.
  - Ventral muscular **Conus Arteriosus** arises from ventricle and distributes blood to aortic arches.
  - **RBCs:** Oval, biconvex, and **nucleated** with hemoglobin.
  - Special portal systems: **Hepatic Portal System** (intestine to liver) and **Renal Portal System** (lower body to kidneys).
- **Respiratory System:**
  - **Cutaneous Respiration:** Skin functions in water and during aestivation (summer sleep) and hibernation (winter sleep).
  - **Buccopharyngeal & Pulmonary Respiration:** Lungs operate on land as paired pink sacs.
- **Excretory System:** Compact dark-red paired **Mesonephric Kidneys**. Functional unit is uriniferous tubule. Excretes urea (**Ureotelic**). In males, ureters act as **urogenital ducts** conveying both urine and spermatozoa into the cloaca.
- **Male Reproductive System:** Paired yellowish ovoid testes adhered to upper kidneys by peritoneal fold called **Mesorchium**. $10-12$ **Vasa Efferentia** emerge from testes, enter kidneys, and open into **Bidder's Canal**, communicating with the urogenital duct.

- **Female Reproductive System:** Paired ovaries near kidneys (no structural connection with kidneys). Oviducts open independently into the cloaca. A mature female lays $2500 - 3000$ **eggs** at a time. Fertilization is **external in water**.
- **Development:** Indirect development via free-swimming aquatic larva called **Tadpole**, which undergoes metamorphosis under control of **thyroxine** hormone.

PART III

# Plant Physiology

CHAPTER 5

# Photosynthesis in Higher Plants

Photosynthesis is the fundamental anabolic process through which green plants transduce solar radiant energy into stable chemical bonds of carbohydrates. This chapter investigates chloroplast ultrastructure, photosynthetic pigment spectra, photochemical light reactions and photophosphorylation, Mitchell's chemiosmotic ATP synthesis, Calvin $C_3$ stoichiometry, Hatch-Slack $C_4$ dicarboxylic acid compartmentalization, Crassulacean Acid Metabolism (CAM), photorespiratory $C_2$ energetics, and the physiological factors governing photosynthetic rate.

## 5.1 Chloroplast Ultrastructure, Pigments & Action Spectra

### Chloroplast Architecture & Photosynthetic Pigment Spectra

Photosynthesis occurs within the chloroplasts of chlorenchyma (mesophyll) cells:

- **Membrane Compartmentalization:**
  - **Thylakoid Membrane System (Grana & Stroma Lamellae):** Traps light energy and synthesizes ATP and NADPH (**Light Reactions / Photochemical Phase**).
  - **Fluid Stroma Matrix:** Contains soluble enzymes for enzymatic reduction of $CO_2$ into carbohydrates (**Dark Reactions / Biosynthetic Phase**).
- **Paper Chromatographic Separation of Leaf Pigments:**
  - **Chlorophyll** $a$**:** Bright / blue-green; primary reaction center pigment.
  - **Chlorophyll** $b$**:** Yellow-green; accessory light-harvesting pigment.
  - **Xanthophylls:** Yellow; accessory pigment.
  - **Carotenoids:** Yellow to yellow-orange; accessory pigments; protect chlorophyll $a$ against lethal **photo-oxidation**.
- **Absorption vs Action Spectrum (T.W. Engelmann, 1843–1909):** Engelmann split light into spectral components using a prism and illuminated filamentous green alga *Cladophora* in suspension of aerobic bacteria. Bacteria accumulated predominantly in regions of **blue** ($400-450\,\text{nm}$) and **red** ($650-700\,\text{nm}$) light, demonstrating that maximum rate of photosynthesis coincides with chlorophyll $a$ absorption peaks.

## 5.2 Photochemical Phase: Z-Scheme, Water Photolysis & Photophosphorylation

### Photochemical Reactions, Z-Scheme & Photophosphorylation Dynamics

Photosynthetic units comprise antenna complexes bound to reaction centers in the thylakoid membrane:

- **Photosystem I (PS I):** Reaction center Chl *a* absorbs maximally at **700** nm ($P_{700}$).
- **Photosystem II (PS II):** Reaction center Chl *a* absorbs maximally at **680** nm ($P_{680}$).
- **Photolysis of Water (OEC on inner lumenal thylakoid surface):**

$$2H_2O \xrightarrow[Mn^{2+},Cl^-]{OEC} 4H^+ + O_2 \uparrow + 4e^-$$

**Comparative Matrix: Non-Cyclic vs Cyclic Photophosphorylation:**

| Parameter | Non-Cyclic (Z-Scheme) | Cyclic Photophosphorylation |
|---|---|---|
| Photosystems | **PS II and PS I** operate in series | **PS I alone** operates |
| Electron Pathway | Non-cyclic ($e^-$ reduce $NADP^+$) | Cyclic ($e^-$ cycle back to $P_{700}$) |
| Products | **ATP, NADPH and** $O_2$ evolved | **ATP alone** (no NADPH, no $O_2$) |
| Location | Grana thylakoids (appressed) | **Stroma lamellae** (lack PS II) |
| Wavelength | Full sunlight ($\lambda \leq 680$ nm) | Beyond 680 nm ($\lambda > 680$ nm) |

## 5.3 Chemiosmotic ATP Synthesis & Proton Motive Force

### Chemiosmotic Hypothesis of ATP Synthesis (Peter Mitchell, 1961)

Photophosphorylation is coupled directly to the generation of a **transmembrane proton gradient ($\Delta$pH) across the thylakoid membrane**. Protons accumulate in high concentration within the **thylakoid lumen**, dropping lumenal pH to $\approx 4.5 - 5.0$ relative to stroma ($\approx 8.0$):

1. **Water Photolysis:** The splitting of $H_2O$ on the lumenal face of the thylakoid membrane releases $H^+$ directly into the lumen.
2. **Plastoquinone (PQ) Proton Shuttle:** Plastoquinone acts as an electron and proton carrier. As it accepts electrons from PS II, it picks up protons from the stroma; upon passing electrons to Cytochrome $b_6f$, it releases protons into the thylakoid lumen.

3. **NADP Reductase Localization:** Located on the stroma-facing surface of the thylakoid membrane. Removes protons from the stroma to reduce $NADP^+$ into $NADPH + H^+$.

**Proton Motive Force & ATP Synthase ($CF_0 - CF_1$ Complex):** Protons exit the thylakoid lumen along the concentration gradient through the transmembrane channel of the $CF_0$ **subunit**. The flow of protons induces rotational conformational changes in the catalytic $CF_1$ **headpiece** projecting into the stroma, synthesizing ATP:

$$ADP + P_i \xrightarrow{CF_0-CF_1 \text{ ATP Synthase}} ATP \tag{5.1}$$

## 5.4 Biosynthetic Phase: Calvin $C_3$ Cycle Stoichiometry & Energetics

### The Calvin $C_3$ Cycle (Melvin Calvin, 1954)

Occurs within the stroma matrix. Deciphered using radioactive $^{14}C$ in algal photosynthesis (*Chlorella*):

1. **Carboxylation:**

$$\text{RuBP (5C)} + CO_2 + H_2O \xrightarrow{\textbf{RuBisCO}} 2 \times \text{3-PGA (3C)}$$

2. **Reduction (per 2 PGA):**

$$2 \times \text{3-PGA} + 2ATP + 2NADPH \longrightarrow 2 \times \text{Triose-P} + 2ADP + 2NADP^+$$

3. **Regeneration:**

$$\text{Triose Phosphate} + 1\,ATP \longrightarrow \text{RuBP (5C)} + 1\,ADP + P_i$$

- **Per Single $CO_2$ Fixed: 3** ATP + **2** NADPH
- **Per Net Hexose Glucose ($6\,CO_2$): 18** ATP + **12** NADPH

## 5.5 Hatch-Slack $C_4$ Pathway & Kranz Anatomy

### The $C_4$ Dicarboxylic Acid Pathway & Kranz Leaf Anatomy

$C_4$ plants (Maize, Sugarcane, Sorghum) have evolved spatial compartmentalization to eliminate photorespiration:

- **Kranz Anatomy (Wreath-like Architecture):**
  - **Bundle Sheath Cells:** Concentric multi-layered cylinders surrounding vascular bundles; thick suberized walls impermeable to gases; lack intercellular spaces; possess large **agranal chloroplasts rich in RuBisCO** (lack PEPcase).
  - **Mesophyll Cells:** Thin-walled, granal chloroplasts; **rich in PEPcase** (strictly lack RuBisCO).
- **Biochemical Reaction Sequence:**
  1. **Primary Carboxylation (Mesophyll):** PEP (3C) + $HCO_3^-$ $\xrightarrow{\textbf{PEPcase}}$ **OAA** (4C).
  2. **Reduction & Transport:** OAA → Malic acid / Aspartic acid, transported through plasmodesmata to bundle sheath cells.
  3. **Decarboxylation (Bundle Sheath):** Malate decarboxylated to release $CO_2$ and Pyruvate (3C). The released $CO_2$ saturates RuBisCO in bundle sheath cells, completely preventing oxygenase activity!
  4. **Regeneration (Mesophyll):** Pyruvate returns to mesophyll and converts to PEP at the cost of 2 ATP.
- **Energetic Cost of $C_4$ Pathway:** Requires 30 ATP+12 NADPH per glucose molecule (12 ATP extra compared to $C_3$).

## 5.6 Crassulacean Acid Metabolism (CAM) & Photorespiration

### Crassulacean Acid Metabolism (CAM) & The Wasteful $C_2$ Photorespiratory Cycle

**1. Crassulacean Acid Metabolism (CAM Pathway):** Evolved in succulent xerophytic plants (e.g., family *Crassulaceae*, *Opuntia, Kalanchoe, Agave, Ananas* [Pineapple]) to achieve extreme water conservation in arid environments. CAM plants utilize **temporal (day-night) separation** of carbon fixation within the *same mesophyll cell*:

- **Night (Scotoactive Stomata Open):** Stomata open during the cool, humid night, minimizing transpirational water loss. Atmospheric $CO_2$ enters and is fixed by cytoplasmic **PEP carboxylase** into Oxaloacetate (4C), reduced to **Malic acid (4C)**. Malic acid is actively pumped and stored in the large central vacuole, causing a sharp drop in nocturnal vacuolar pH (**Nocturnal Acidification**).
- **Day (Stomata Tightly Closed):** Stomata remain closed under intense solar heat, preventing transpirational water loss. Malic acid leaves the vacuole and is decarboxylated in the cytoplasm/chloroplast by malic enzyme to release $CO_2$ and Pyruvate (3C). The released $CO_2$ enters the chloroplast

stroma, where **RuBisCO** fixes it via the standard Calvin $C_3$ cycle powered by ATP and NADPH generated from the light reactions.

- **Water-Use Efficiency (WUE):** CAM plants boast the highest water-use efficiency among all terrestrial plants, consuming only $50 - 100\,\text{g}\,H_2O$ per gram of $CO_2$ fixed (compared to $250 - 350\,\text{g}$ in $C_4$ and $400 - 500\,\text{g}$ in $C_3$).

**2. Photorespiration ($C_2$ Glycolate Cycle):** Occurs in $C_3$ plants under conditions of high $O_2$, low $CO_2$, and elevated temperatures when RuBisCO acts as an **oxygenase**:

$$\text{RuBP (5C)} + O_2 \xrightarrow{\textbf{RuBisCO}} \text{3-PGA (3C)} + \text{2-Phosphoglycolate (}\mathbf{2}\text{C)} \quad (5.2)$$

- Operates sequentially across three intracellular organelles: **Chloroplast $\longrightarrow$ Peroxisome $\longrightarrow$ Mitochondria**.
- **Metabolic Inefficiency:** Yields **neither ATP nor NADPH nor sugar**, but consumes ATP and releases previously fixed $CO_2$. Reduces photosynthetic output by up to $25 - 30\%$ in $C_3$ crops.

## 5.7 Factors Regulating Photosynthesis & Blackman's Law

**NEET Trap: Blackman's Law & Limiting Factors**

- **Blackman's Law of Limiting Factors (F.F. Blackman, 1905):** If a chemical process is affected by more than one factor, its rate is determined by the factor nearest to its minimal value; it is the factor which directly affects the process if its quantity is changed.
- $CO_2$ **Saturation Levels:** $C_4$ plants saturate at **360** $\mu$L/L (ppm); $C_3$ plants saturate only beyond **450** $\mu$L/L. Current ambient $CO_2$ is $\approx 400$ ppm; thus, $CO_2$ **is the major limiting factor for $C_3$ plants** (greenhouse crops like tomatoes and bell peppers show marked yield increases in $CO_2$-enriched atmospheres).
- **Light Saturation:** Occurs at **10**% of full sunlight. Except for plants in dense forest understories or deep shade, light is rarely a limiting factor in nature.

CHAPTER 6

# Respiration in Plants & Phytohormonal Coordination

Cellular respiration is the catabolic release of metabolic energy stored in the chemical bonds of respiratory substrates. In this chapter, we explore cytosolic glycolysis, pyruvate oxidation in the link reaction, the mitochondrial Krebs cycle, complexes I–V of the electron transport system, respiratory quotients, and the physiological roles of the five classical phytohormone classes.

## 6.1 Glycolytic Pathway (EMP) & Mitochondrial Link Reaction

### Glycolysis (EMP Pathway) Stoichiometry & Link Reaction

Discovered by Gustav Embden, Otto Meyerhof, and J. Parnas. Occurs in the **cytoplasm** of all living cells; independent of $O_2$.

- **Preparatory Phase (Invests** $2\,\text{ATP}$**):**

$$\text{Glucose} \xrightarrow{\text{HK, 1 ATP}} \text{G-6-P} \longrightarrow \text{F-6-P} \xrightarrow{\text{PFK, 1 ATP}} \text{F-1,6-BP}$$

- **Cleavage Phase:**

$$\text{F-1,6-BP (6C)} \xrightarrow{\text{Aldolase}} \text{PGAL (3C)} + \text{DHAP (3C)}$$

- **Payoff Phase (**$\times 2$ **per glucose):**

$$2\,\text{PGAL} + 2\,\text{NAD}^+ \longrightarrow 2 \times 1,3\text{-BPGA} + \mathbf{2}\,\text{NADH} + \mathbf{2}\,\text{H}^+$$

$$2 \times 1,3\text{-BPGA} + 2\,\text{ADP} \xrightarrow{\text{PGK}} 2 \times 3\text{-PGA} + \mathbf{2}\,\text{ATP (SLP)}$$

$$2\,\text{PEP} + 2\,\text{ADP} \xrightarrow{\text{PK}} 2\,\text{Pyruvate} + \mathbf{2}\,\text{ATP (SLP)}$$

- **Net Glycolytic Yield:** $\mathbf{2}\,\text{Pyruvate} + \mathbf{2}\,\text{ATP} + \mathbf{2}\,\text{NADH} + \mathbf{2}\,\text{H}^+$ (zero $CO_2$ or $O_2$).

**The Link Reaction (Oxidative Decarboxylation of Pyruvate):** Occurs in the **mitochondrial matrix**, catalyzed by the *pyruvate dehydrogenase* multienzyme complex (requires $Mg^{2+}$, TPP, Lipoamide, $NAD^+$, CoA):

$$2\,\text{Pyruvate (3C)} + 2\,\text{CoA} + 2\,\text{NAD}^+ \longrightarrow \mathbf{2}\,\text{Acetyl-CoA (2C)} + \mathbf{2}\,\text{CO}_2 \uparrow + \mathbf{2}\,\text{NADH} + \mathbf{2}\,\text{H}^+$$

## 6.2 Tricarboxylic Acid (TCA) Cycle & Amphibolic Intermediates

### The Tricarboxylic Acid (TCA / Krebs) Cycle (Hans Krebs, 1937)

Occurs within the **mitochondrial matrix**:

- **Condensation & Decarboxylations (per Acetyl-CoA):**

$$\begin{aligned} \text{Acetyl-CoA} + \text{OAA} + \text{H}_2\text{O} &\xrightarrow{\text{Citrate Synthase}} \text{Citrate (6C)} + \text{CoA} \\ \text{Isocitrate} &\longrightarrow \alpha\text{-ketoglutarate (5C)} + \text{CO}_2 + \text{NADH} \\ \alpha\text{-ketoglutarate} &\longrightarrow \text{Succinyl-CoA (4C)} + \text{CO}_2 + \text{NADH} \end{aligned}$$

- **Substrate-Level Phosphorylation & Oxidations:**

$$\begin{aligned} \text{Succinyl-CoA} + \text{GDP} + \text{P}_\text{i} &\longrightarrow \text{Succinate} + \mathbf{1}\,\text{GTP/ATP} + \text{CoA} \\ \text{Succinate} &\longrightarrow \text{Fumarate} + \mathbf{1}\,\text{FADH}_2 \\ \text{Malate} + \text{NAD}^+ &\longrightarrow \text{OAA} + \mathbf{1}\,\text{NADH} + \text{H}^+ \end{aligned}$$

- **Yield per Glucose (2 Cycles):**

$$\mathbf{4}\,\text{CO}_2 + \mathbf{6}\,\text{NADH} + \mathbf{2}\,\text{FADH}_2 + \mathbf{2}\,\text{ATP/GTP}$$

## 6.3 Mitochondrial Electron Transport System (ETS) & ATP Synthase

### ETS Complex Architecture in the Inner Mitochondrial Membrane

- **Complex I (NADH Dehydrogenase):** Contains FMN and Fe-S centers; transfers electrons from NADH to Ubiquinone (CoQ), pumping $4\text{H}^+$ into intermembrane space.
- **Complex II (Succinate Dehydrogenase):** Contains FAD and Fe-S centers; transfers electrons from succinate to Ubiquinone (does not pump protons).
- **Complex III (Cytochrome $bc_1$ Complex):** Contains Cytochrome $b$, Fe-S, and Cytochrome $c_1$; transfers electrons from ubiquinol to mobile carrier Cytochrome $c$, pumping $4\text{H}^+$.
- **Cytochrome $c$:** Small soluble protein attached to outer surface of inner membrane; acts as mobile electron shuttle from Complex III to Complex IV.
- **Complex IV (Cytochrome $c$ Oxidase):** Contains Cytochrome $a, a_3$ and two copper centers ($\text{Cu}_\text{A}, \text{Cu}_\text{B}$); reduces the terminal electron acceptor $\text{O}_2$:

$$\frac{1}{2}\text{O}_2 + 2\text{H}^+ + 2e^- \longrightarrow \text{H}_2\text{O} \quad \text{(Metabolic Water)} \tag{6.1}$$

- **Complex V (ATP Synthase / $F_0 - F_1$ Complex):** Protons flow down concentration gradient from intermembrane space to matrix through $F_0$ transmembrane channel, driving $F_1$ ATP synthesis.

## 6.4 Respiratory Quotient ($RQ$) Formulations

### Respiratory Quotient ($RQ$) Mathematical Formulations

$$\text{RQ} = \frac{\text{Volume of } CO_2 \text{ evolved}}{\text{Volume of } O_2 \text{ consumed}} \tag{6.2}$$

- **Carbohydrates (Glucose):** $\text{RQ} = \frac{6\,CO_2}{6\,O_2} = \mathbf{1.0}$
- **Fats (Tripalmitin):**

$$2C_{51}H_{98}O_6 + 145O_2 \longrightarrow 102CO_2 + 98H_2O \implies \text{RQ} = \frac{102}{145} \approx \mathbf{0.7}$$

- **Proteins:** $\text{RQ} \approx \mathbf{0.9}$
- **Organic Acids (Malic Acid):**

$$C_4H_6O_5 + 3O_2 \longrightarrow 4CO_2 + 3H_2O \implies \text{RQ} = \frac{4}{3} = \mathbf{1.33} > 1.0$$

- **Anaerobic Respiration:** $\text{RQ} = \frac{2\,CO_2}{0\,O_2} = \infty$

## 6.5 Phytohormones: Plant Growth Regulators & Morphogenesis

### Five Classical Phytohormone Classes: Functions & Agricultural Roles

| Hormone | Discovery / Chemistry | Key Physiological Functions & Applications |
|---|---|---|
| **Auxins** | Darwin; isolated by F.W. Went from *Avena* tip. Natural: IAA, IBA; Synth: NAA, 2,4-D. | • Promotes **apical dominance**; decapitation induces bushy hedge.<br>• Initiates root formation in stem cuttings; promotes flowering in pineapples.<br>• **2,4-D** is selective weedicide for dicots.<br>• Induces parthenocarpy in tomatoes. |
| **Gibberellins** | E. Kurosawa from *Gibberella fujikuroi* (*bakanae* disease). > 100 known ($GA_3$ common). Terpenes. | • Internode elongation; grape stalks and apples.<br>• **Bolting** in rosette plants (cabbage, beet).<br>• Lengthens sugarcane stem (+20 tons/acre).<br>• Accelerates malting in brewing. |
| **Cytokinins** | Skoog discovered kinetin from herring sperm; Letham isolated **Zeatin** from corn. Aminopurines. | • Rapid cell division; high cytokinin:auxin ratio induces shoots.<br>• Overcomes apical dominance; delays leaf senescence (**Richmond-Lang effect**). |
| **Ethylene** | H.H. Cousins from oranges. Only **gaseous hormone**; source: **Ethephon**. | • Fruit ripening with **Respiratory Climacteric**.<br>• Promotes senescence and abscission of organs.<br>• Promotes root growth and root hair formation.<br>• Breaks seed/bud dormancy; female flowers in cucumbers. |
| **ABA** | Inhibitor-$\beta$, Abscission II, Dormin. Sesquiterpene; growth inhibitor. | • **Stress Hormone**: rapid stomatal closure in drought.<br>• Induces/maintains **seed dormancy**; anti-GA antagonist. |

**Photoperiodism & Vernalization Highlights**

- **Photoperiodism (Garner & Allard):** Differential flowering response to relative day/night durations. **Short-Day Plants (SDP)** require day lengths below critical photoperiod (Tobacco, Soyabean); **Long-Day Plants (LDP)** require photoperiod exceeding critical threshold (Wheat, Radish, Spinach); **Day-Neutral Plants (DNP)** flower independently of photoperiod (Tomato, Sunflower). The site of light/dark perception is the **leaves**, which synthesize the hypothetical floral hormone **Florigen**.
- **Vernalization:** Qualitative or quantitative promotion of flowering by low temperature exposure ($1 - 6^{\circ}$C); perceived by the **shoot apical meristem**.

PART IV

# Human Physiology

CHAPTER 7

# Human Respiration, Circulation & Excretion

Human physiology represents the essential clinical foundation for medical training. In this chapter, we explore pulmonary mechanics, alveolar gas exchange, and hemoglobin oxygen dissociation; cardiovascular hemodynamics, cardiac cycle phases, ECG deflections, and the blood coagulation cascade; and renal nephron microanatomy, glomerular filtration dynamics, counter-current concentration mechanics, and the hormonal regulation of osmoregulation.

## 7.1 Breathing & Alveolar Gas Exchange

#### Respiratory Volumes & Pulmonary Capacities Matrix

All values represent healthy young adult averages measured by spirometry (residual volume cannot be measured by spirometry):

- **Tidal Volume (TV):** 500 mL ($\approx 6000 - 8000$ mL/min total ventilation).
- **Inspiratory Reserve Volume (IRV):** $2500 - 3000$ mL (extra forced inspiration).
- **Expiratory Reserve Volume (ERV):** $1000 - 1100$ mL (extra forced expiration).
- **Residual Volume (RV):** $1100 - 1200$ mL (cannot be measured by spirometry).
- **Inspiratory Capacity (IC):** TV + IRV $= 3000 - 3500$ mL.
- **Expiratory Capacity (EC):** TV + ERV $= 1500 - 1600$ mL.
- **Functional Residual Capacity (FRC):** ERV + RV $= 2100 - 2300$ mL.
- **Vital Capacity (VC):** ERV + TV + IRV $= 3500 - 4500$ mL.
- **Total Lung Capacity (TLC):** VC + RV $= 5000 - 6000$ mL.

### 7.1.1 Alveolar Diffusion Gradients & Respiratory Membrane Dynamics

Gas exchange across the pulmonary alveoli and systemic capillary beds occurs strictly via simple physical diffusion driven by partial pressure gradients ($\Delta P$). The respiratory diffusion membrane is extremely thin ($< 0.2\,\mu$m), consisting of three layers: (1) the single-layered squamous epithelium of alveoli, (2) the

endothelium of alveolar capillaries, and (3) the thin intervening basement substance.

| Gas | Atm. Air | Alveoli | Deox. Blood | Ox. Blood | Tissues |
|---|---|---|---|---|---|
| $O_2$ | 159 mmHg | **104** mmHg | 40 mmHg | **95** mmHg | 40 mmHg |
| $CO_2$ | 0.3 mmHg | **40** mmHg | 45 mmHg | **40** mmHg | 45 mmHg |

**Table 7.1:** Partial Pressures (mmHg) of Oxygen and Carbon Dioxide across Physiological Compartments.

**Solubility Differential:** Although the partial pressure gradient for oxygen ($\Delta P_{O_2} = 104 - 40 = 64$ mmHg) is vastly greater than that for carbon dioxide ($\Delta P_{CO_2} = 45 - 40 = 5$ mmHg), the solubility of $CO_2$ in blood plasma is **20 to 25 times higher** than that of $O_2$. Consequently, $CO_2$ diffuses across the respiratory membrane with comparable overall volume throughput.

### 7.1.2 Oxygen Transport & The Oxygen-Hemoglobin Dissociation Curve

Nearly 97% of oxygen is transported bound to hemoglobin in erythrocytes as **oxyhemoglobin** ($Hb_4(O_2)_4$), while the remaining 3% travels dissolved in aqueous blood plasma. Each tetrameric hemoglobin molecule can cooperatively bind up to four molecules of $O_2$.

A plot of the percentage saturation of hemoglobin against the partial pressure of oxygen ($pO_2$) yields a characteristic **Sigmoidal (S-shaped) curve** (Figure 7.1). This sigmoidal shape reflects positive allosteric cooperativity: the binding of the first $O_2$ molecule structurally alters the globin chains, progressively increasing the binding affinity for subsequent oxygen molecules.

**Bohr Effect (Shift to the Right):** In actively respiring metabolizing tissues, elevated metabolic byproducts decrease hemoglobin's affinity for oxygen, shifting the dissociation curve to the right:

$$\uparrow pCO_2, \quad \uparrow [H^+] \ (\downarrow \text{pH / acidity}), \quad \uparrow \text{Temperature}, \quad \uparrow 2,3\text{-BPG} \tag{7.1}$$

A right shift increases $P_{50}$ (the partial pressure required for half-saturation, rising from $\approx 27$ mmHg to $\approx 38$ mmHg). This physiological adaptation unloads oxygen precisely where cellular respiration is most active.

**Shift to the Left (Alveolar Loading):** In the pulmonary capillaries of the alveoli, where oxygen partial pressure is high, temperature is lower, and $CO_2$ and $H^+$ concentrations are low, hemoglobin exhibits heightened oxygen affinity, shifting the curve to the left and facilitating rapid saturation.





**Figure 7.1:** Oxygen-Hemoglobin Dissociation Curve: Normal Sigmoid Curve, Right Shift (Bohr Effect), and Left Shift.

### Gas Delivery Metrics & Physiological Constants

**1. Oxygen Delivery Rate:** Under basal resting physiological conditions, every 100 mL of oxygenated blood delivers approximately 5 mL **of** $O_2$ to the systemic tissues. **2. Carbon Dioxide Clearance Rate:** Every 100 mL of deoxygenated blood returning to the lungs delivers approximately 4 mL **of** $CO_2$ into the alveoli for exhalation.

## 7.1.3 Carbon Dioxide Transport Modalities & The Chloride Shift

Carbon dioxide is transported from respiring tissues to the lungs through three distinct biochemical routes:

1. **As Bicarbonate Ions ($HCO_3^-$) (70%):** RBCs possess a very high concentration of the zinc-containing enzyme **carbonic anhydrase** (with minute quantities in plasma). In tissue capillaries, high $pCO_2$ drives:

$$CO_2 + H_2O \underset{\text{anhydrase}}{\overset{\text{carbonic}}{\rightleftharpoons}} H_2CO_3 \underset{\text{anhydrase}}{\overset{\text{carbonic}}{\rightleftharpoons}} HCO_3^- + H^+ \tag{7.2}$$

Bicarbonate diffuses out into the plasma; to preserve electrical neutrality, chloride ions ($Cl^-$) diffuse from plasma into the RBC. This phenomenon

is known as the **Chloride Shift (Hamburger phenomenon)**. At the alveoli, the entire reaction reverses, releasing gaseous $CO_2$.

2. **As Carbamino-Hemoglobin ($20 - 25\%$):** $CO_2$ binds directly and reversibly to the free amino groups of hemoglobin globin chains, forming carbamino-hemoglobin.
3. **Dissolved in Plasma ($7\%$):** Transported as physically dissolved $CO_2$ molecules in aqueous blood plasma.

### 7.1.4 Neural Regulation of Respiration & Clinical Pathologies

The human respiratory rate ($\approx 12 - 16$ breaths/min) is maintained by neural feedback circuits:

- **Respiratory Rhythm Center:** Located in the **medulla oblongata**; generates the basal inspiratory rhythm.
- **Pneumotaxic Center:** Situated in the dorsal **pons varolii**; acts as a "switch-off" point for inspiration, regulating tidal volume and respiratory rate.
- **Chemosensitive Area:** Located adjacent to the rhythm center; highly sensitive to arterial $\mathbf{p}CO_2$ and $[H^+]$. **Critical Trap:** Arterial oxygen tension ($pO_2$) plays an entirely **insignificant** role in modulating the central respiratory rhythm.
- **Clinical Disorders:**
  - **Asthma:** Allergic bronchial inflammation characterized by wheezing, bronchospasm, and mucosal edema.
  - **Emphysema:** Chronic progressive disorder where alveolar septa are destroyed, drastically reducing the respiratory surface area (primary cause: cigarette smoking; genetically deficient patients are treated with $\alpha$-1-antitrypsin).
  - **Occupational Lung Diseases:** Chronic exposure to silica dust (silicosis) or asbestos fibers (asbestosis) in stone-crushing and grinding industries causes chronic inflammation, pulmonary fibrosis, and irreversible lung damage.

## 7.2 Body Fluids, Circulation & Coagulation Cascade

#### Blood Formed Elements, Hemostasis & Cardiac Cycle Dynamics

**1. Blood Components:** Plasma ($55\%$, contains fibrinogen, globulins, albumins) and formed elements ($45\%$):

- **Erythrocytes (RBCs):** Biconcave, enucleated in adult mammals, $5.0 - 5.5 \times 10^6/mm^3$; lifespan $= 120$ days; destroyed in spleen; hemoglobin $=$

$12 - 16\,\text{g}/100\,\text{mL}$.

- **Leukocytes (WBCs):** $6000-8000/\text{mm}^3$. Granulocytes: Neutrophils ($60-65\%$, phagocytic), Eosinophils ($2-3\%$, allergic response), Basophils ($0.5-1\%$, secrete histamine, serotonin, heparin). Agranulocytes: Monocytes ($6-8\%$), Lymphocytes ($20-25\%$, B and T cells).
- **Platelets:** Cell fragments from bone marrow **megakaryocytes** ($1.5-3.5\times 10^5/\text{mm}^3$).

**2. Blood Coagulation Cascade (Enzyme Cascade Theory):**

1. Injured Tissue/Platelets $\longrightarrow$ Thromboplastin $\xrightarrow{Ca^{2+}}$ Thrombokinase
2. Prothrombin (Inactive) $\xrightarrow[Ca^{2+}]{\text{Thrombokinase}}$ Thrombin (Active Protease)
3. Fibrinogen (Soluble) $\xrightarrow{\text{Thrombin}}$ Fibrin (Mesh) $\xrightarrow{\text{Traps Cells}}$ **Clot**

**3. Cardiac Cycle ($0.8\,\text{sec}$ at $72\,\text{bpm}$) & Hemodynamics:**

- **Atrial Systole:** $0.1\,\text{sec}$ (increases ventricular filling by $30\%$).
- **Ventricular Systole:** $0.3\,\text{sec}$ (isovolumetric contraction and ejection).
- **Joint Diastole:** $0.4\,\text{sec}$ (passive filling of atria and ventricles).
- **Stroke Volume (SV):** $\text{EDV} - \text{ESV} = 120\,\text{mL} - 50\,\text{mL} = \mathbf{70}\,\text{mL}$.
- **Cardiac Output (CO):** $\text{SV} \times \text{HR} = 70\,\text{mL} \times 72/\text{min} \approx \mathbf{5040}\,\text{mL/min}$ ($\approx \mathbf{5}\,\text{L/min}$).

**Heart Sounds:** First sound (**LUBB**) produced by closure of atrioventricular (bicuspid/tricuspid) valves; second sound (**DUBB**) by closure of semilunar valves.

**4. Electrocardiogram (ECG) Deflections: P-wave:** Depolarization of atria; **QRS-complex:** Depolarization of ventricles (marks onset of ventricular contraction); **T-wave:** Repolarization of ventricles.

## 7.3 Excretory Products & Renal Physiology

### Nephron Architecture, GFR & Counter-Current Osmoregulation

**1. Nitrogenous Excretion:**

- **Ammonotelism:** Highly toxic $NH_3$ requiring vast water volumes (bony fishes, aquatic amphibians, aquatic insects).
- **Ureotelism:** Urea synthesized in liver via Ornithine cycle (mammals, terrestrial amphibians, marine cartilaginous fishes).
- **Uricotelism:** Uric acid excreted as insoluble pellet/paste with minimum water loss (reptiles, birds, land snails, insects).

**2. Nephron Typology & Glomerular Filtration:**

- **Cortical Nephrons (85%):** Short Loop of Henle extending only slightly into medulla; vasa recta reduced or absent.
- **Juxtamedullary Nephrons (15%):** Long Loop of Henle running deep into inner medulla; well-developed vasa recta; **essential for producing concentrated urine**.
- **Glomerular Filtration Rate (GFR): 125** mL/min = **180** L/day. Daily urine output is $\approx 1.5$ L, meaning $> 99\%$ of filtrate is obligatorily and facultatively reabsorbed.
- **Net Filtration Pressure (NFP):**

$$\begin{aligned} \text{NFP} &= \text{GBHP} - (\text{BCOP} + \text{CHP}) \\ &= 60\,\text{mmHg} - (32 + 18)\,\text{mmHg} = \mathbf{10}\,\text{mmHg} \end{aligned}$$

**3. Counter-Current Multiplier (Henle's Loop) & Exchanger (Vasa Recta):**

- **Descending Limb of Henle:** Permeable to water, impermeable to electrolytes $\Longrightarrow$ filtrate concentrates (300 $\rightarrow$ 1200 mOsmol/L).
- **Ascending Limb of Henle:** Impermeable to water, actively transports NaCl into medullary interstitium $\Longrightarrow$ filtrate dilutes (1200 $\rightarrow$ 200 mOsmol/L).
- Establishes a steep medullary interstitial osmotic gradient (300 mOsmol/L in cortex $\rightarrow$ 1200 mOsmol/L in inner medulla), enabling human kidneys to excrete urine **4 times more concentrated** than initial filtrate.

**4. Hormonal Regulation of Renal Function:**

- **RAAS Complex:** Fall in GFR stimulates Juxtaglomerular (JG) cells to release **Renin** $\rightarrow$ converts Angiotensinogen into Angiotensin I $\xrightarrow{\text{ACE}}$ **Angiotensin II** (potent vasoconstrictor) $\rightarrow$ stimulates adrenal cortex to release **Aldosterone** (promotes active $Na^+$ and $H_2O$ reabsorption from DCT).
- **ADH / Vasopressin:** Released from posterior pituitary upon hypothalamic osmoreceptor stimulation; inserts aquaporin channels into DCT and collecting ducts, preventing diuresis.
- **ANF (Atrial Natriuretic Factor):** Released by atrial myocardium upon elevated blood pressure; causes vasodilation and natriuresis, acting as an antagonist to RAAS.

CHAPTER 8

# Locomotion, Nervous Integration & Chemical Coordination

Vertebrate locomotion and physiological homeostasis require tight mechanical and neuro-endocrine coordination. In this chapter, we investigate the sliding filament theory of skeletal muscle contraction, human skeletal architecture, resting membrane potentials and action potential generation, chemical and electrical synaptic transmission, and the mechanisms of action of peptide versus steroid hormones.

## 8.1 Musculoskeletal Mechanics & Sliding Filament Theory

### Sarcomere Ultrastructure & Cross-Bridge Cycling Mechanics

The **Sarcomere** is the fundamental contractile unit of striated muscle, bounded between two successive **Z-lines**:

- **Sarcomere Banding Pattern:**
  - **A-Band (Anisotropic):** Contains both thick (myosin) and overlapping thin (actin) filaments; length **remains completely constant** during muscle contraction.
  - **I-Band (Isotropic):** Contains only thin actin filaments; bisected by elastic Z-line; **shortens** during contraction.
  - **H-Zone:** Central region of A-band not overlapped by thin filaments; bisected by fibrous M-line; **disappears** during maximal contraction.
- **Myofilament Biochemistry:**
  - **Thin Filament:** Composed of two helically wound polymeric **F-actin** filaments, two filamentous **Tropomyosin** ribbons, and complex **Troponin** triplets (TnT, TnI, TnC). **Troponin-C (TnC)** binds $Ca^{2+}$, unmasking active myosin-binding sites on actin.
  - **Thick Filament:** Composed of polymerized **Myosin** monomers (**Meromyosins**):
    - **Heavy Meromyosin (HMM):** Globular head and short arm; projects outward as cross-bridge; contains actin-binding site and active **ATPase catalytic site**.
    - **Light Meromyosin (LMM):** Tail portion.

- **Sliding Filament Cycle:** Motor neuron impulse → acetylcholine release at neuromuscular junction → action potential spreads along sarcolemma and T-tubules → sarcoplasmic reticulum releases $Ca^{2+}$ → $Ca^{2+}$ binds Troponin-C → tropomyosin shifts → myosin head hydrolyzes ATP (ADP + $P_i$) and binds actin forming cross-bridge → **Power Stroke** pulls thin filaments towards center of A-band → new ATP binds myosin head causing cross-bridge detachment.

### Human Skeletal Framework (206 Bones) & Articulations

- **Axial Skeleton (80 Bones):**
  - **Skull (22 bones):** Cranial (8) + Facial (14). Associated: Hyoid (1) + Ear Ossicles ($2 \times 3 = 6$: Malleus, Incus, Stapes).
  - **Vertebral Column (26 bones in adult):** Cervical (7, $C_1$ = Atlas, $C_2$ = Axis), Thoracic (12), Lumbar (5), Sacral (1 fused), Coccygeal (1 fused).
  - **Thoracic Cage:** Sternum (1) + Ribs (12 pairs = 24). True ribs ($1^{st} - 7^{th}$ pair, vertebrosternal), False ribs ($8^{th}, 9^{th}, 10^{th}$ pair, vertebrochondral), Floating ribs ($11^{th}, 12^{th}$ pair, vertebral).
- **Appendicular Skeleton (126 Bones):**
  - **Forelimb ($30 \times 2 = 60$):** Humerus (1), Radius (1), Ulna (1), Carpals (8 wrist), Metacarpals (5 palm), Phalanges (14 digits, formula 2,3,3,3,3).
  - **Pectoral Girdle ($2 \times 2 = 4$):** Clavicle (collar bone) + Scapula (flat triangular bone with acromion process, spine, and glenoid cavity articulating with head of humerus).
  - **Hindlimb ($30 \times 2 = 60$):** Femur (thigh, longest and strongest bone), Patella (sesamoid kneecap), Tibia (1), Fibula (1), Tarsals (7 ankle), Metatarsals (5), Phalanges (14).
  - **Pelvic Girdle (2 Coxal bones):** Each formed by fusion of Ilium, Ischium, and Pubis; meet ventrally at pubic symphysis (fibrous cartilage); cavity = **Acetabulum** (articulates with femur head).
- **Joint Classifications:**
  - **Fibrous (Synarthroses):** Immovable (sutures between flat cranial bones).
  - **Cartilaginous (Amphiarthroses):** Limited movement (pubic symphysis, intervertebral discs).
  - **Synovial (Diarthroses):** Freely movable fluid-filled synovial joints: Ball and socket (shoulder, hip); Hinge (knee, elbow); Pivot (Atlas and Axis, $C_1 - C_2$); Gliding (between carpals); Saddle (between carpal and metacarpal of thumb).
- **Clinical Disorders: Myasthenia gravis:** Autoimmune disorder affecting neuromuscular junction leading to fatigue and paralysis; **Muscular dystro-**

**phy:** Progressive degeneration of skeletal muscle (genetic); **Tetany:** Rapid spasms due to low $Ca^{2+}$ in body fluid; **Arthritis:** Inflammation of joints; **Gout:** Inflammation due to accumulation of uric acid crystals; **Osteoporosis:** Age-related bone mass reduction (decreased estrogen).

## 8.2 Neural Control & Action Potential Propagation

### Neurophysiology: Membrane Potentials & Synaptic Transmission

- **Resting Membrane Potential (RMP):** $-\mathbf{70}\,\text{mV}$ (inside negative; permeable to $K^+$, impermeable to $Na^+$).
- $Na^+/K^+$ **ATPase Pump:** Actively pumps $\mathbf{3}\,Na^+$ out for every $\mathbf{2}\,K^+$ in (1 ATP).
- **Depolarization (Action Potential):** $+\mathbf{30}\,\text{mV}$ ($Na^+$ influx via voltage-gated channels).
- **Repolarization Phase:** $Na^+$ channels close, voltage-gated $K^+$ channels open $\implies$ $K^+$ efflux.
- **Chemical Synaptic Transmission:**

$$\text{AP at terminal} \longrightarrow Ca^{2+} \text{ influx} \longrightarrow \text{ACh exocytosis} \longrightarrow \text{Receptors}$$

## 8.3 Endocrine Integration & Hormone Action Mechanisms

### Endocrine Glands, Hormones & Dual Action Mechanisms

| Gland | Hormone | Type | Action & Pathology |
|---|---|---|---|
| Ant. Pituitary | GH | Peptide | High in child: **Gigantism**; adult: **Acromegaly**; low: **Dwarfism**. |
| Post. Pituitary | Oxytocin, ADH | Peptide | Oxytocin: Milk ejection/labor. ADH: Water reabsorption; low: **Diabetes Insipidus**. |
| Thyroid | Thyroxine ($T_4, T_3$) | Iodinated | BMR. Child low: **Cretinism**; adult: **Myxoedema**; high: **Graves'**. |
| Thyroid C-cells | TCT | Peptide | **Hypocalcemic**; lowers blood $Ca^{2+}$ via bone mineralization. |
| Parathyroid | PTH | Peptide | **Hypercalcemic**; bone resorption; antagonist to TCT. |
| Adrenal Cortex | Aldosterone, Cortisol | Steroid | $Na^+$ retention. Low: **Addison's**; high: **Cushing's**. |
| Adrenal Medulla | Adrenaline, Norad. | Catechol. | Emergency fight-or-flight; elevates HR, BP, glycogenolysis. |
| Pancreas ($\beta$) | Insulin | Peptide | Hypoglycemic; promotes glucose uptake; low: **Diabetes Mellitus**. |
| Pancreas ($\alpha$) | Glucagon | Peptide | Hyperglycemic; stimulates glycogenolysis; antagonist to insulin. |

**Dual Mechanisms of Hormone Action:**

- **Peptide / Protein Hormones (Water-Soluble):** Insulin, glucagon, pituitary hormones. Cannot cross lipid bilayer; bind to **extracellular membrane-bound receptors** $\implies$ generate second messengers (**cyclic AMP,** $IP_3, Ca^{2+}$) $\implies$ biochemical amplification cascade regulates cellular metabolism.
- **Steroid & Iodothyronine Hormones (Lipid-Soluble):** Estrogen, progesterone, testosterone, cortisol, thyroxine ($T_3/T_4$). Cross plasma membrane freely; bind to **intracellular / nuclear receptors** $\implies$ hormone-receptor complex interacts with chromosome to regulate gene transcription and protein synthesis directly.

PART V

# Reproduction & Reproductive Health

CHAPTER 9

# Sexual Reproduction in Flowering Plants (Angiosperms)

Sexual reproduction in angiosperms achieves genetic diversification through specialized floral organs, highly orchestrated micro- and megasporogenesis, and the unique phenomenon of double fertilization. This chapter explores anther microsporangial architectures, pollen wall chemistry, the 7-celled 8-nucleate embryo sac, pollination ecology, outbreeding adaptations, endosperm bioenergetics, and apomictic seed formation.

## 9.1 Microsporogenesis & Male Gametophyte

### Anther Histology, Microsporogenesis & Pollen Wall Chemistry

**1. Anther Architecture:** Typically bilobed, dithecous (two thecae per lobe), and tetrasporangiate (four microsporangia):

- **Four Anther Wall Layers (from outside to inside):**
  1. **Epidermis:** Outermost single protective layer.
  2. **Endothecium:** Sub-epidermal layer with fibrous hygroscopic $\alpha$-cellulose bands; assists in anther dehiscence.
  3. **Middle Layers:** $1-3$ transient ephemeral layers that degenerate to provide nutrition to developing microspores.
  4. **Tapetum:** Innermost nutritive layer; cells possess dense cytoplasm and are characteristically **multinucleate / polyploid**; synthesizes and secretes **sporopollenin**, **callase**, and pollenkitt.
- **Microsporogenesis:** Sporogenous tissue in microsporangia undergoes meiosis:

$$1 \text{ MMC } (2n) \xrightarrow{\text{Meiosis}} \text{Microspore Tetrad (4 Haploid Microspores, } n)$$

- **Pollen Grain Wall (Sporoderm):**
  - **Exine:** Hard outer layer composed of **Sporopollenin** (the most chemically resistant biological polymer known; withstands high temperatures, concentrated acids, and strong alkalis; no known enzyme degrades sporopollenin). Possesses prominent apertures called **Germ Pores** where sporopollenin is absent.

  - **Intine:** Thin, delicate, continuous inner wall composed of **cellulose and pectin**.
- **Pollen Shedding Stage:** In $> 60\%$ of angiosperms, pollen is shed at the **2-celled stage** (large vegetative cell with irregular nucleus + small spindle-shaped generative cell floating in vegetative cytoplasm). In $< 40\%$, the generative cell divides mitotically to form 2 male gametes prior to shedding (**3-celled stage**).

## 9.2 Megasporogenesis & The 7-Celled 8-Nucleate Embryo Sac

### Ovule Anatomy & Monosporic Embryo Sac Development

**1. Ovule (Integumented Megasporangium):** Attached to placenta by **funicle**; junction between funicle and ovule body is the **hilum**. Central parenchymatous tissue is the **nucellus**, enclosed by 1–2 protective integuments except at the apical **micropyle**. The basal region opposite the micropyle is the **chalaza**.

**2. Megasporogenesis & Polygonum-Type Development:**

- Megaspore Mother Cell (MMC, $2n$) near micropyle undergoes meiosis forming a linear tetrad of 4 haploid megaspores.
- Three micropylar megaspores degenerate; only **one chalazal megaspore remains functional** (**Monosporic Development**).
- The functional megaspore undergoes **three sequential free-nuclear mitotic divisions** producing an 8-nucleate stage, followed by cytokinesis into a **7-celled, 8-nucleate mature embryo sac**:
  - **Micropylar Egg Apparatus (3 cells):** 1 central **Egg cell** ($n$) + 2 flanking **Synergids** bearing finger-like **Filiform Apparatus** that guides the entering pollen tube.
  - **Chalazal Antipodal Cells (3 cells):** Nutritive vegetative cells; degenerate following fertilization.
  - **Large Central Cell (1 cell):** Contains **2 Polar Nuclei** that fuse to form a diploid **Secondary Nucleus** ($2n$).

## 9.3 Pollination Ecology, Outbreeding Devices & Pollen-Pistil Interaction

### Pollination Modes & Contrivances for Cross-Pollination

**1. Pollination Modes:**

- **Autogamy:** Transfer of pollen from anther to stigma of the same flower. Requires chasmogamous or **cleistogamous flowers** (never open, ensuring complete seed set even without pollinators, e.g., *Viola*, *Oxalis*, *Commelina*).
- **Geitonogamy:** Transfer of pollen from anther to stigma of another flower on the same plant. *Functionally cross-pollination involving a pollinator, but genetically identical to autogamy.*
- **Xenogamy (Cross-Pollination):** Transfer of pollen to stigma of a different plant; introduces genuine genetic variation.

**2. Outbreeding Devices (Mechanisms Preventing Inbreeding Depression):**

- **Dichogamy:** Non-synchronous pollen release and stigma receptivity: **Protandry** (anthers mature first, e.g., Sunflower) vs **Protogyny** (stigma matures first, e.g., *Aristolochia*).
- **Heterostyly / Anther-Stigma Placement:** Anther and stigma placed at different physical levels preventing contact.
- **Self-Incompatibility:** Genetically controlled mechanism that prevents self-pollen from fertilizing ovules by inhibiting pollen germination or pollen tube growth in the pistil.
- **Dioecy (Unisexuality):** Male and female flowers borne on separate individual plants (e.g., Papaya, Date palm); completely prevents both autogamy and geitonogamy.

## 9.4 Double Fertilization, Endosperm Energetics, Embryogenesis & Apomixis

### Double Fertilization Dynamics (S.G. Nawaschin, 1898)

Unique to angiosperms; involves two distinct syngamic fusion events:

- **Syngamy (Generative Fertilization):**

$$\text{Male Gamete } (n) + \text{Egg Cell } (n) \longrightarrow \text{Zygote } (\mathbf{2n}) \longrightarrow \text{Embryo}$$

- **Triple Fusion (Vegetative Fertilization):**

$$\text{Male Gamete } (n) + \text{Secondary Nucleus } (2n) \longrightarrow \text{PEN } (\mathbf{3n}) \longrightarrow \text{Endosperm}$$

- **Total Nuclei Involved:** 1 (egg) + 2 (sperm) + 2 (polar nuclei) = **5** Nuclei.

**Endosperm Bioenergetics:** Primary Endosperm Nucleus (PEN, $3n$) divides repeatedly prior to zygotic cleavage to establish nutrient reserves:

- **Free-Nuclear Endosperm:** PEN undergoes repeated karyokineses without cell wall synthesis (e.g., coconut water is free-nuclear endosperm containing thousands of free nuclei; surrounding white meat is cellular endosperm).

**Apomixis & Polyembryony:**

- **Apomixis:** Form of asexual reproduction that **mimics sexual reproduction**, producing viable seeds without fertilization (common in *Asteraceae* and grasses; nucellar cells protrude into embryo sac forming diploid embryos). Eliminates segregation of hybrid traits in agriculture.
- **Polyembryony:** Occurrence of multiple embryos within a single seed (e.g., *Citrus*, Mango).

CHAPTER 10

# Human Reproduction & Reproductive Health

Human reproduction involves internal fertilization, viviparous embryonic development, and endocrine feedback loops. In this chapter, we analyze male and female reproductive anatomy, gametogenesis, the 28-day menstrual cycle, fertilization, blastocyst implantation, placental endocrinology, contraceptive modalities, and clinical Assisted Reproductive Technologies (ART).

## 10.1 Male & Female Reproductive Architectures

### Male & Female Reproductive Histology & Duct Systems

**1. Male Reproductive System:**

- **Scrotum:** Cutaneous pouch maintaining testes at $2 - 2.5^{\circ}\text{C}$ **below core body temperature**, essential for normal spermatogenesis.
- **Testes Histology:** Each testis contains $\approx 250$ testicular lobules, each housing $1 - 3$ highly coiled **Seminiferous Tubules**:
  - **Spermatogonia (Male Germ Cells):** Undergo meiotic divisions to produce spermatozoa.
  - **Sertoli Cells (Nurse Cells):** Provide structural and metabolic nourishment to differentiating sperms; secrete **Inhibin** (inhibits FSH) and **Androgen Binding Protein (ABP)**.
  - **Leydig Cells (Interstitial Cells):** Located in connective tissue spaces outside tubules; synthesize and secrete androgens (**Testosterone**) under LH stimulation.
- **Accessory Glands:** Paired **Seminal Vesicles** ($60 - 70\%$ of semen; rich in fructose, prostaglandins, and calcium), single **Prostate Gland** (milky alkaline secretion), and paired **Bulbourethral (Cowper's) Glands** (lubricating mucus).

**2. Female Reproductive System:**

- **Ovaries:** Primary female sex organs producing oocytes and steroid hormones (Estrogen, Progesterone).
- **Fallopian Tubes (Oviducts,** $10 - 12\,\text{cm}$**):**
  - **Infundibulum:** Funnel-shaped part with finger-like **fimbriae** collecting ovum after ovulation.

  - **Ampulla:** Wider central segment; **the physiological site of fertilization**.
  - **Isthmus:** Narrow terminal segment joining the uterus.
- **Uterine Wall (3 Layers):** Outer thin serous **Perimetrium**; middle thick smooth-muscle **Myometrium** (undergoes powerful oxytocin-induced contractions during parturition); inner glandular vascular **Endometrium** (undergoes cyclic cyclical shedding during menstruation).

## 10.2 Gametogenesis: Spermatogenesis vs Oogenesis

### Comparative Gametogenic Sequences & Nuclear Timing

**1. Spermatogenesis (Continuous from Puberty through Old Age):**

$$\begin{aligned}\text{Spermatogonia } (2n) &\xrightarrow{\text{Mitosis}} \text{Primary Spermatocyte } (2n)\\ &\xrightarrow{\text{Meiosis I}} 2 \times \text{Secondary Spermatocytes } (n)\\ &\xrightarrow{\text{Meiosis II}} 4 \times \text{Spermatids } (n)\\ &\xrightarrow{\text{Spermiogenesis}} 4 \times \text{Spermatozoa } (\mathbf{n})\end{aligned}$$

**2. Oogenesis (Initiated during Fetal Life, Suspended until Puberty):**

$$\begin{aligned}\text{Oogonia } (2n) &\xrightarrow{\text{Mitosis}} \text{Primary Oocyte } (2n) \quad \text{[Prophase I]}\\ \text{At Puberty: } &\longrightarrow \text{Secondary Oocyte } (\mathbf{n}) + \mathbf{1}^{\text{st}} \text{ Polar Body } (\mathbf{n})\\ &\quad \text{[Arrested at Metaphase II; released at ovulation]}\\ \text{At Fertilization: } &\longrightarrow \text{Ootid / Ovum } (\mathbf{n}) + \mathbf{2}^{\text{nd}} \text{ Polar Body } (\mathbf{n})\end{aligned}$$

## 10.3 Menstrual Cycle Hormonal Regulation & Implantation

### The 28-Day Menstrual Cycle & Pregnancy Endocrinology

1. **Menstrual Phase (Days 1–5):** Endometrial sloughing and blood loss ($50 - 100\,\text{mL}$) triggered by the sudden **withdrawal of progesterone** following luteal degeneration.
2. **Follicular / Proliferative Phase (Days 6–13):** Pituitary **FSH** stimulates primary follicle maturation into mature **Graafian Follicle**. Follicular granulosa cells secrete **Estrogen**, regenerating the endometrium.

3. **Ovulatory Phase (Day 14):** Peak levels of LH and FSH; rapid LH surge (**LH Surge**) induces rupture of Graafian follicle and release of the secondary oocyte (**Ovulation**).
4. **Luteal / Secretory Phase (Days 15–28):** Ruptured follicle transforms into endocrine **Corpus Luteum**, which secretes massive amounts of **Progesterone** (maintains endometrium for implantation). In the absence of fertilization, corpus luteum degenerates into **Corpus Albicans**, triggering progesterone drop and menses.

**Fertilization & Implantation Mechanics:**

- Occurs at the **ampulla** of the fallopian tube. Sperm contact with **zona pellucida** triggers the acrosome reaction, depolarizing the membrane and blocking polyspermy.
- Cleavage divisions form a 16-celled **morula** → fluid-filled **blastocyst**. Trophoblast attaches to endometrium ≈ 7 days post-fertilization (**Implantation**).
- **Placental Hormones exclusive to pregnancy:** Human Chorionic Gonadotropin (**hCG**), Human Placental Lactogen (**hPL**), and **Relaxin** (secreted by ovary in late pregnancy).

## 10.4 Reproductive Health, Contraceptive Modalities & Assisted Reproductive Technologies (ART)

### Contraceptive Classifications & Assisted Reproductive Protocols

**1. Contraceptive Methodologies:**

- **Natural Methods:** Periodic abstinence (avoiding coitus during days 10 – 17 of menstrual cycle), coitus interruptus, and **lactational amenorrhea** (intense suckling suppresses gonadotropins for up to 6 months postpartum).
- **Barrier Methods:** Condoms (protect against STIs/AIDS), Diaphragms, cervical caps, vaults.
- **Intrauterine Devices (IUDs):**
  - **Non-medicated IUDs:** *Lippes loop* (promotes phagocytosis of sperms).
  - **Copper-releasing IUDs: CuT, Cu7, Multiload 375** ($Cu^{2+}$ ions suppress sperm motility and fertilizing ability).
  - **Hormone-releasing IUDs: Progestasert, LNG-20** (render uterus unsuitable for implantation and cervix hostile to sperms).
- **Oral Contraceptive Pills:** Combined daily estrogen-progestogen pills; or **Saheli** (developed by CDRI Lucknow; non-steroidal, once-a-week pill with negligible side effects).

- **Surgical / Sterilization Methods: Vasectomy** (vas deferens ligated in males) and **Tubectomy** (fallopian tubes ligated in females); highly effective, irreversible.

**2. Assisted Reproductive Technologies (ART Protocols):**

- **IVF-ET (In Vitro Fertilization - Embryo Transfer):**
  - **ZIFT (Zygote Intra-Fallopian Transfer):** Zygote or early embryo **up to 8 blastomeres** transferred into Fallopian tube.
  - **IUT (Intra-Uterine Transfer):** Embryos with **more than 8 blastomeres** transferred directly into the Uterus.
- **GIFT (Gamete Intra-Fallopian Transfer):** Transfer of ovum collected from donor into fallopian tube of female recipient.
- **ICSI (Intra-Cytoplasmic Sperm Injection):** Single spermatozoon injected directly into ovum cytoplasm in laboratory.
- **IUI (Intra-Uterine Insemination):** Semen collected from husband/donor introduced artificially into uterus.

PART VI

# Genetics & Molecular Biology

CHAPTER 11

# Principles of Inheritance & Variation

Genetics is the highest-weightage conceptual domain in the NEET examination. In this chapter, we investigate classical Mendelian principles of segregation and independent assortment, non-Mendelian extensions (incomplete dominance, codominance, multiple allelism, pleiotropy, and polygenic inheritance), the chromosomal theory of inheritance, Morgan's linkage and crossing-over experiments on *Drosophila*, sex determination mechanisms, and human pedigree genetics.

## 11.1 Mendelian Principles & Cross Combinatorics

### Mendel's Experimental Framework & Contrasting Traits

Gregor Johann Mendel conducted hybridization experiments on garden peas (*Pisum sativum*) for 7 years (1856–1863), analyzing **7 pairs of contrasting traits** located across 4 chromosomes (Chromosomes 1, 4, 5, and 7):

| Character | Dominant | Recessive | Chr. |
|---|---|---|---|
| Stem Height | Tall ($T$) | Dwarf ($t$) | 4 |
| Flower Color | Violet ($W$) | White ($w$) | 1 |
| Flower Position | Axial ($A$) | Terminal ($a$) | 4 |
| Pod Shape | Inflated ($I$) | Constricted ($i$) | 4 |
| Pod Color | Green ($G$) | Yellow ($g$) | 5 |
| Seed Shape | Round ($R$) | Wrinkled ($r$) | 7 |
| Seed Color | Yellow ($Y$) | Green ($y$) | 1 |

**Mendel's Laws of Inheritance:**

1. **Law of Dominance:** Characters are controlled by discrete unit factors (genes) occurring in pairs. In a dissimilar pair of factors, one member dominates the other.
2. **Law of Segregation (Purity of Gametes):** Alleles do not show blending; both traits are recovered in $F_2$. During gamete formation, alleles of a pair segregate such that a gamete receives only one of the two factors. This law is **universal without exception**.
3. **Law of Independent Assortment:** When two pairs of traits are combined in a hybrid, segregation of one pair of characters is independent of the other pair (valid strictly for unlinked genes).

### Mendelian Ratios & Combinatorial Cross Calculus

For parents heterozygous at $n$ independently assorting gene loci:

- **Monohybrid Cross ($Aa \times Aa$):** Phenotypic 3 : 1; Genotypic 1 : 2 : 1.
- **Dihybrid Cross ($AaBb \times AaBb$):** Phenotypic 9 : 3 : 3 : 1; Genotypic $(1:2:1)^2$.
- **Trihybrid Phenotypic Ratio:** $(3+1)^3 = 27:9:9:9:3:3:3:1$.
- **Combinatorics for $n$ Heterozygous Loci:**
  - Unique gametes $= 2^n$ Phenotypic classes $= 2^n$
  - Genotypic classes $= 3^n$ Punnett squares $= 4^n$
- **Test Cross Ratios:** Monohybrid = 1 : 1; Dihybrid = 1 : 1 : 1 : 1.

## 11.2 Non-Mendelian Extensions: Dominance, Allelism & Pleiotropy

### Deviations from Standard Mendelian Ratios

1. **Incomplete Dominance:** Heterozygote displays an intermediate phenotype between homozygous extremes (e.g., flower color in *Mirabilis jalapa* [4 o'clock plant] and *Antirrhinum majus* [Snapdragon / Dog flower]). Crossing Red ($RR$) with White ($rr$) gives Pink ($Rr$). **Phenotypic ratio equals genotypic ratio:** 1 : 2 : 1.
2. **Codominance:** Both alleles express themselves fully in the heterozygote without blending (e.g., ABO blood groups in humans; $I^AI^B$ individuals express both A and B surface antigens simultaneously).
3. **Multiple Allelism:** More than two alternative allelic forms govern a single gene locus in a population:
   - Human ABO blood locus ($I$) on Chromosome 9 has 3 alleles: $I^A, I^B, i$.
   - Total number of genotypes for $n$ alleles:

$$\text{Genotypes} = \frac{n(n+1)}{2} = \frac{3(3+1)}{2} = \mathbf{6} \text{ Genotypes}$$

   Yielding **4** Phenotypes (Blood groups A, B, AB, and O).
4. **Pleiotropy:** A single gene exerts multiple, seemingly unrelated phenotypic effects:
   - **Phenylketonuria (PKU):** Mutation in gene coding for *phenylalanine hydroxylase* causes mental retardation, reduced hair, and cutaneous hypopigmentation.
   - **Starch Synthesis in Pea Seeds:** Gene $B$ controls starch grain size (incomplete dominance: $BB$ large, $Bb$ intermediate, $bb$ small) and seed shape (complete dominance: $BB, Bb$ round, $bb$ wrinkled).

5. **Polygenic (Quantitative) Inheritance:** Continuous traits governed by multiple additive genes interacting with environment (e.g., human skin color, height). Governed by 3 genes $(A, B, C)$; $F_2$ distribution follows a Gaussian bell curve $(1:6:15:20:15:6:1)$.

## 11.3 Chromosomal Theory, Sex Determination & Morgan's Linkage

### Chromosomal Linkage & Sturtevant Gene Mapping

**1. Chromosomal Theory of Inheritance (Sutton & Boveri, 1902):** Chromosomes, like Mendelian factors, occur in pairs. Homologous chromosomes segregate during Anaphase I and assort independently during Metaphase I.

**2. Morgan's Linkage Experiments on *Drosophila melanogaster* (T.H. Morgan, 1910):** Morgan chose the fruit fly because of: (i) simple synthetic culture medium, (ii) short 2-week life cycle, (iii) large progeny per mating, (iv) distinct sexual dimorphism, and (v) low chromosome count $(2n = 8)$:

- **Cross A (Tightly Linked Genes):** Yellow body $(y)$ and white eye $(w)$ crossed with wild brown body $(y^+)$ and red eye $(w^+)$. Recombination Frequency = **1.3%** (Parental types = **98.7%**).
- **Cross B (Loosely Linked Genes):** White eye $(w)$ and miniature wing $(m)$ crossed with wild red eye $(w^+)$ and normal wing $(m^+)$. Recombination Frequency = **37.2%** (Parental types = **62.8%**).

**3. Chromosome Mapping Formulation (Alfred Sturtevant):** Recombination frequency is directly proportional to physical distance between genes on a chromosome:

$$\text{Distance (cM)} = \left(\frac{\text{Total Recombinant Progeny}}{\text{Total Progeny}}\right) \times 100\% \tag{11.1}$$

$$1\,\text{cM} = 1\,\text{Map Unit (m.u.)} = 1\%\ \text{Recombination Frequency} \tag{11.2}$$

*Upper Limit:* Maximum observable recombination frequency for two linked genes is strictly **50%** (at 50%, genes behave as unlinked).

**4. Sex Determination Systems:**

- **XX–XY (Male Heterogamety):** Humans and *Drosophila*. Male produces two sperm types (50% X, 50% Y); female is homogametic (XX).
- **XX–XO (Male Heterogamety):** Grasshopper. Male has $2A + XO$; female has $2A + XX$.
- **ZZ–ZW (Female Heterogamety):** Birds. Female produces two egg types $(2A + ZW)$; male is homogametic $(2A + ZZ)$.

- **Haplodiploidy:** Honeybees. Queens/workers develop from fertilized diploid eggs ($2n = 32$); drones develop from unfertilized eggs by **parthenogenesis** ($n = 16$). Drones have no father and cannot have sons, but have grandfathers and grandsons.

## 11.4 Genetic Disorders & Pedigree Analysis

**NEET Trap: Mendelian vs Chromosomal Disorders**

**1. Mendelian Single-Gene Disorders:**

- **Sickle-Cell Anemia (Autosomal Recessive):** Point mutation (transversion) in the $\beta$-globin gene on Chromosome 11. GAG codon mutates to GUG, causing substitution of **Glutamic acid (Glu) by Valine (Val) at position 6** of the $\beta$-chain. Under hypoxia, $Hb^S$ polymerizes into rigid sickle shapes. Heterozygotes ($Hb^A Hb^S$) exhibit malaria resistance.
- **Thalassemia (Autosomal Recessive):** Quantitative defect (reduced synthesis of normal globin chains), unlike sickle cell which is qualitative (abnormal globin synthesized). $\alpha$-Thalassemia is governed by $HBA1$ and $HBA2$ on Chromosome 16; $\beta$-Thalassemia is governed by $HBB$ on Chromosome 11.
- **Hemophilia (Sex-Linked Recessive):** Criss-cross inheritance from carrier mother to son. Hemophilia A (Factor VIII deficiency); Hemophilia B / Christmas disease (Factor IX deficiency).
- **Phenylketonuria (Autosomal Recessive):** Deficiency of liver enzyme *phenylalanine hydroxylase*; phenylalanine accumulates and converts to phenylpyruvic acid.

**2. Chromosomal Aneuploidies:**

- **Down's Syndrome (Trisomy 21):** Karyotype: $47, XX + 21$ or $47, XY + 21$. Round face, furrowed tongue, partially open mouth, broad palm with simian crease, congenital heart disease, mental retardation.
- **Klinefelter's Syndrome (Sex Chromosome Trisomy):** Karyotype: $47, XXY$. Overall masculine development with feminine features (e.g., **Gynaecomastia** / breast development); sterile male.
- **Turner's Syndrome (Sex Chromosome Monosomy):** Karyotype: $45, XO$. Phenotypic sterile female with rudimentary ovaries, webbed neck, short stature, lack of secondary sexual characteristics.

CHAPTER 12

# Molecular Basis of Inheritance & Genomics

Molecular genetics deciphers the chemical basis of heredity and cellular information transfer. In this chapter, we explore Watson-Crick B-DNA geometry, histone nucleosome packaging, experimental validations of DNA as genetic material, replication enzymology, prokaryotic and eukaryotic transcription, the universal genetic code, translation mechanics, the Lac operon, the Human Genome Project (HGP), and forensic DNA fingerprinting.

## 12.1 Watson-Crick B-DNA Structural Dimensions & Chromatin Packaging

### Watson-Crick B-DNA Double Helix Dimensions (1953)

Constructed from X-ray diffraction data of Rosalind Franklin and Maurice Wilkins:

- **Helical Symmetry:** Right-handed antiparallel double helix ($5' \to 3'$ paired with $3' \to 5'$).
- **Pitch ($P$):** $\mathbf{3.4}\,\text{nm} = \mathbf{34}\,\text{Å}$ (one complete 360° turn, 10 bp/turn).
- **Distance between Base Pairs:** $\mathbf{0.34}\,\text{nm} = \mathbf{3.4}\,\text{Å}$.
- **Diameter of Double Helix:** $\mathbf{2.0}\,\text{nm} = \mathbf{20}\,\text{Å}$.
- **Hydrogen Bonding:** $\text{A} = \text{T}$ (2 H-bonds), $\quad \text{G} \equiv \text{C}$ (3 H-bonds).
- **Chargaff's Rules:** $[\text{A}] = [\text{T}], \quad [\text{G}] = [\text{C}], \quad \frac{[\text{A}]+[\text{G}]}{[\text{T}]+[\text{C}]} = 1.0.$
- **Length of Human Diploid DNA:**

$$6.6 \times 10^{9}\,\text{bp} \times 0.34 \times 10^{-9}\,\text{m/bp} \approx \mathbf{2.2}\,\text{meters}$$

**Nucleosome Packaging Architecture:** To package a 2.2 m DNA strand inside a $\sim 10\,\mu$m nucleus, DNA wraps around basic proteins called **Histones**:

- Histones are rich in positively charged basic amino acids: **Lysine and Arginine**.
- **Histone Octamer:** Core complex composed of two molecules each of $H_2A, H_2B, H_3$, and $H_4$.
- **Nucleosome Core:** Exactly 200 bp **of negatively charged DNA** is wrapped around the octamer core.
- **Linker Histone ($H_1$):** Seals DNA at entry/exit points.

- **Euchromatin:** Loosely packed, stains light, transcriptionally active.
- **Heterochromatin:** Densely packed, stains dark, transcriptionally inactive.

## 12.2 Historic Proof of DNA as the Genetic Material

### Classic Experiments Demonstrating DNA as Genetic Material

1. **Griffith's Transformation Experiment (Frederick Griffith, 1928):** Used *Streptococcus pneumoniae* (virulent smooth encapsulated S-strain vs non-virulent rough non-capsulated R-strain):
   - Heat-killed S-strain + Living R-strain injected into mice $\implies$ Mice died of pneumonia, and living virulent S-strain bacteria were recovered!
   - Proved that a **"Transforming Principle"** transferred from heat-killed S to R strain, conferring capsule synthesis.
2. **Biochemical Characterization (Avery, MacLeod & McCarty, 1944):** Purified biomolecules from heat-killed S cells. Proteases and RNases did not abolish transformation; only **Deoxyribonuclease (DNase) completely abolished transformation**, proving DNA is the transforming principle.
3. **Unequivocal Proof: Hershey-Chase Experiment (1952):** Used Bacteriophage T2 and *E. coli*:
   - Phages grown on radioactive phosphorus ($^{32}$P) labeled DNA; phages grown on radioactive sulfur ($^{35}$S) labeled protein coats (cysteine and methionine).
   - Infection $\rightarrow$ Blending (detached coats) $\rightarrow$ Centrifugation.
   - Bacteria infected with $^{32}$P phages were radioactive (pellet had radioactivity); $^{35}$S remained in supernatant viral ghosts. Conclusively proved that **DNA is the hereditary material**.

## 12.3 Semi-Conservative DNA Replication Machinery & Enzymology

### Replication Fork Enzymology & Experimental Proof

**1. Experimental Proof of Semi-Conservative Replication:**

- **Meselson & Stahl (1958):** Grew *E. coli* in $^{15}NH_4Cl$ (heavy isotope) $\rightarrow$ transferred to $^{14}NH_4Cl$. Density gradient centrifugation in CsCl: Generation I (20 min) yielded **100% hybrid** $^{15}N - ^{14}N$; Generation II (40 min) yielded **50% hybrid** +50% **light** $^{14}N - ^{14}N$.

- **Taylor (1958):** Used tritiated thymidine ($^3$H-thymidine) in root cells of *Vicia faba* proving semi-conservative replication in chromosomes.

**2. Replication Fork Enzymes & Functional Roles:** Replication is bidirectional, proceeding strictly in the $5' \to 3'$ **direction**:

| Enzyme | Specific Biological Function |
|---|---|
| **DNA Helicase** | Unwinds double helix by breaking H-bonds, creating replication fork. |
| **SSB Proteins** | Prevent premature re-annealing of single DNA strands. |
| **Topoisomerase** | Relieves torsional strain ahead of replication fork. |
| **RNA Primase** | Synthesizes short complementary RNA primers ($5-10$ nt) with $3'$-OH. |
| **DNA Pol III** | Main enzyme; polymerizes $5' \to 3'$ ($\approx 2000$ bp/s); proofreads $3' \to 5'$. |
| **Leading Strand** | Continuous synthesis proceeding toward replication fork ($5' \to 3'$). |
| **Lagging Strand** | Discontinuous synthesis generating short **Okazaki fragments**. |
| **DNA Pol I** | Removes RNA primers ($5' \to 3'$ exonuclease) and fills gaps (Kornberg enzyme). |
| **DNA Ligase** | Seals nicks by forming phosphodiester bonds between Okazaki fragments. |

## 12.4 Transcription in Prokaryotes vs Eukaryotes & RNA Processing

### Transcription Units, RNA Polymerases & Eukaryotic RNA Processing

A transcription unit comprises: (1) **Promoter** ($5'$ upstream of coding strand), (2) **Structural gene**, and (3) **Terminator** ($3'$ downstream of coding strand):

$$\text{Template Strand: } 3' \longrightarrow 5' \qquad \text{Coding Strand (Non-template): } 5' \longrightarrow 3' \tag{12.1}$$

**1. Prokaryotic Transcription:** Single RNA polymerase catalyzes all phases: binds promoter with transient assistance of **Sigma ($\sigma$) initiation factor**; elongates; terminates via **Rho ($\rho$) factor**. Translation can occur **coupled simultaneously with transcription** due to lack of nuclear envelope.

**2. Eukaryotic Transcription (Three Distinct RNA Polymerases):**

- **RNA Polymerase I:** Transcribes rRNAs ($28S, 18S$, and $5.8S$).
- **RNA Polymerase II:** Transcribes precursor of mRNA called **heterogeneous nuclear RNA (hnRNA)**.
- **RNA Polymerase III:** Transcribes **tRNA**, $5S$ **rRNA**, and **snRNAs** (small nuclear RNAs).

**3. Post-Transcriptional Processing of hnRNA $\to$ Mature mRNA:**

1. **Capping:** Unusual nucleotide **7-methylguanosine triphosphate ($\mathbf{m^7Gppp}$)** added to 5′-end.
2. **Tailing (Polyadenylation):** Adenylate residues ($200 - 300$) added to 3′-end in template-independent manner.
3. **Splicing:** Non-coding intervening sequences (**Introns**) excised, and coding sequences (**Exons**) ligated by spliceosomes (complex of snRNPs).

## 12.5 The Genetic Code, Wobble Phenomenon & Translation

### The Universal Genetic Code & Translation Mechanisms

Deciphered by George Gamow, Har Gobind Khorana, Marshall Nirenberg, and Severo Ochoa:

- **Triplet Nature:** 64 codons ($4^3$): 61 sense codons, 3 stop codons.
- **Stop Codons: UAA** (Ochre), **UAG** (Amber), **UGA** (Opal).
- **Dual Function: AUG** codes for Met AND acts as initiator codon.
- **Degeneracy:** Amino acids coded by $>1$ codon (Leu, Ser, Arg have 6 codons).
- **Exceptions:** AUG (Met) and UGG (Trp) have only 1 codon each.
- **Unambiguous:** One specific codon codes for strictly one specific amino acid.
- **Universal:** From bacteria to humans, UUU codes for Phenylalanine.

**Wobble Hypothesis (Francis Crick, 1966):** Base-pairing between the 3′ base of the mRNA codon and the 5′ base of the tRNA anticodon is flexible (non-standard Watson-Crick pairing). Allows a single tRNA species to recognize multiple synonymous codons.

**Translation Steps:**

1. **Aminoacylation (Charging) of tRNA:** Amino Acid + ATP $\rightarrow$ Aminoacyl-AMP + $PP_i$ $\xrightarrow{\text{tRNA}}$ Aminoacyl-tRNA + AMP.
2. **Initiation:** Small ribosomal subunit ($30S/40S$) binds mRNA 5′-end; initiator tRNA binds P-site.
3. **Elongation:** Incoming aminoacyl-tRNA enters A-site. Peptide bond formation is catalyzed by the ribozyme **Peptidyl Transferase**:

$$\text{Peptidyl Transferase} = \begin{cases} \textbf{23S} \text{ rRNA} & \text{(Prokaryotes)} \\ \textbf{28S} \text{ rRNA} & \text{(Eukaryotes)} \end{cases}$$

4. **Translocation & Termination:** Ribosome translocates $5' \rightarrow 3'$ along mRNA. Release factors bind stop codon in A-site, releasing the completed polypeptide.

## 12.6 Operon Model: The Lac Operon

### The Lac Operon of *Escherichia coli* (Jacob & Monod, 1961)

An inducible genetic unit subject to negative transcriptional regulation:

- **Gene $i$ (Regulatory Gene):** Synthesizes the active Lac Repressor protein constitutively.
- **Promoter ($p$):** Binding site for RNA polymerase.
- **Operator ($o$):** Sequence adjacent to structural genes where repressor binds.
- **Structural Genes:**
  - **Gene $z$:** Codes for **$\beta$-galactosidase** (hydrolyzes lactose into glucose and galactose).
  - **Gene $y$:** Codes for **Permease** (increases cellular membrane permeability to lactose).
  - **Gene $a$:** Codes for **Transacetylase** (transfers acetyl group from acetyl-CoA to $\beta$-galactosides).
- **Regulatory Dynamics:**
  - **In Absence of Inducer (Lactose):** Repressor protein binds operator $\implies$ blocks RNA polymerase $\implies$ Operon switched **OFF**.
  - **In Presence of Inducer (Lactose / Allolactose):** Inducer binds repressor, inducing conformational inactivation $\implies$ repressor detaches from operator $\implies$ RNA polymerase transcribes $z, y, a$ $\implies$ Operon switched **ON**.

## 12.7 Genomics & The Human Genome Project (HGP)

**Genomics & Human Genome Project (HGP) High-Yield Blueprint**

The Human Genome Project (1990–2003) was an international mega-project coordinated by the US Department of Energy and NIH, joined by the Wellcome Trust (UK):

- **Two Major Methodological Approaches:**
  1. **Expressed Sequence Tags (ESTs):** Identifying and sequencing all genes that are actively expressed as RNA.
  2. **Sequence Annotation:** Blindly sequencing the entire genome (coding and

non-coding) followed by computational assignment of functional regions.

- **Methodological Workflow:** Total genomic DNA isolated → sheared into random fragments → cloned into specialized host systems using vectors: **BAC (Bacterial Artificial Chromosomes)** and **YAC (Yeast Artificial Chromosomes)** → sequenced using automated Sanger dideoxy chain-termination methods → assembled by overlapping contigs using computer algorithms.
- **Chromosome 1 Sequence Completion:** Chromosome 1, the largest human chromosome, was the last to be sequenced, completed in **May 2006**.
- **Salient Metrics of the Human Genome:**
  - Total base pairs: $\mathbf{3.1647 \times 10^9}$ bp (3164.7 million bp).
  - Average gene size: **3000 bases**; largest known human gene is **Dystrophin** on X-chromosome ($2.4 \times 10^6$ bp).
  - Total estimated protein-coding genes: $\approx 30,000$.
  - Less than **2% of the human genome** codes for proteins.
  - Chromosome 1 has the most genes (**2968**); the Y-chromosome has the fewest (**231**).
  - Scientists have mapped $\approx$ **1.4 million single nucleotide polymorphisms** (SNPs).

## 12.8 DNA Fingerprinting & Satellite DNA Polymorphism

### DNA Fingerprinting Protocol (Alec Jeffreys, 1984)

Explores **DNA polymorphism** in non-coding repetitive DNA (**Satellite DNA**):

- **Density Gradient Centrifugation:** Genomic DNA separates into major bulk genomic peak and minor satellite peaks (micro- and mini-satellites).
- **VNTRs (Variable Number of Tandem Repeats):** Belong to **minisatellite DNA** ($0.1 - 20$ kb); show high degree of copy number variation among individuals, inherited from parents.
- **Sequential Experimental Protocol:**
  1. **Isolation:** Extraction of high-molecular-weight DNA from blood, semen, hair roots.
  2. **Digestion:** Cleavage of DNA by specific restriction endonucleases.
  3. **Electrophoresis:** Separation of fragments according to molecular size on agarose gel.
  4. **Southern Blotting:** Transfer (blotting) of separated DNA fragments onto synthetic nylon/nitrocellulose membranes.
  5. **Hybridization:** Incubation with radioactive labeled VNTR probe DNA.

6. **Autoradiography:** Detection of hybridized VNTR bands on X-ray film, producing individual-specific band patterns.

PART VII

# Biotechnology, Human Health & Ecology

CHAPTER 13

# Biotechnology: Principles, Processes & Modern Applications

Biotechnology harnesses cellular and biomolecular processes to develop technologies and products that improve human health and agriculture. This chapter details recombinant DNA tools (restriction endonucleases, cloning vectors, pBR322 architecture), host transformation methodologies, polymerase chain reaction (PCR) amplification, industrial bioreactor kinetics, downstream bioprocessing, and modern transgenic applications including Bt crops, RNA interference, recombinant human insulin, gene therapy, and biopiracy ethics.

## 13.1 Tools of Recombinant DNA Technology & Restriction Endonucleases

### Restriction Enzymes: Molecular Scissors & Palindromic Recognition

Discovered by Werner Arber, Hamilton Smith, and Daniel Nathans (1970):

- **First Discovered Restriction Endonuclease:** *Hind II* (isolated from *Haemophilus influenzae*; cleaves DNA at specific 6-bp recognition sites).
- **Nomenclature (*Eco*RI):** $E$ = genus *Escherichia*, *co* = species *coli*, R = strain RY13, I = first enzyme isolated from this strain.
- **Palindromic Recognition Sequence:** A sequence of base pairs that reads identical on both strands in the $5' \to 3'$ direction:

$$5' - \text{G} \downarrow \text{A A T T C} - 3'$$
$$3' - \text{C T T A A} \uparrow \text{G} - 5'$$

- **Sticky vs Blunt Ends:** *Eco*RI cleaves between G and A, leaving single-stranded staggered overhangs called **Sticky Ends** that facilitate hydrogen-bonding with complementary vector overhangs.
- **Enzymatic Ligation: DNA Ligase** catalyzes formation of phosphodiester bonds, joining insert and vector.

## 13.2 Cloning Vectors, pBR322 Architecture & Host Transformation

### Cloning Vectors & Competent Host Transformation Protocols

**1. Salient Features of Ideal Plasmid Vector pBR322:**

- **Origin of Replication (*Ori*):** DNA sequence where replication initiates; controls copy number of linked foreign DNA.
- **Selectable Markers:** Identify and select transformants and eliminate non-transformants. Example: genes conferring antibiotic resistance ($amp^R$ **and** $tet^R$).
- **Insertional Inactivation:** Ligation of foreign DNA at the *BamHI* restriction site in the $tet^R$ gene inactivates tetracycline resistance $\implies$ recombinant clones grow on ampicillin but perish on tetracycline.
- **Blue-White Screening:** Foreign insert ligated into gene coding for enzyme **$\beta$-galactosidase** ($\alpha$-complementation). Non-recombinant colonies metabolize X-gal producing **blue colonies**; recombinants with insertional inactivation yield **white colonies**.
- **Cloning Sites:** Unique recognition sites for restriction enzymes (e.g., *EcoRI, BamHI, SalI, PvuI, PstI, ClaI, HindIII*).

**2. Host Transformation Techniques:**

- **Chemical Competence & Heat Shock:** Bacterial cells treated with divalent cation ($Ca^{2+}$) to increase membrane pore permeability $\rightarrow$ incubated on ice $\rightarrow$ **Heat Shock at** 42°C **for 2 minutes** $\rightarrow$ returned to ice.
- **Microinjection:** Recombinant DNA injected directly into the nucleus of an animal zygote/cell using a glass micropipette.
- **Biolistics / Gene Gun:** Plant cells bombarded with high-velocity microprojectiles of **gold or tungsten** coated with recombinant DNA.
- **Disarmed Pathogen Vectors:** Modified Ti plasmid of soil bacterium ***Agrobacterium tumefaciens*** delivers genes to dicots; disarmed retroviruses used in animals.

## 13.3 PCR Amplification, Bioreactor Kinetics & Downstream Processing

### Polymerase Chain Reaction (PCR) & Bioreactor Operations

Invented by Kary Mullis (1983). Generates over a billion-fold ($2^n$) amplification of DNA in vitro:

1. **Denaturation (94−96°C):** Thermal separation of dsDNA strands by breaking hydrogen bonds.

2. **Annealing** ($50 - 60^{\circ}$C)**:** Hybridization of two oligonucleotide primers to flanking $3'$-ends.
3. **Extension** ($72^{\circ}$C)**:** Polymerization by thermostable Taq Polymerase from *Thermus aquaticus*.

**Bioreactors & Downstream Processing:**

- **Bioreactors** ($100 - 1000$ L)**:** Vessels providing optimal temperature, pH, substrate, oxygen, and salts:
  - **Simple Stirred-Tank Bioreactor:** Motor-driven agitator ensures continuous mixing and oxygen availability.
  - **Sparged Stirred-Tank Bioreactor:** Sterile air bubbles sparged through medium, drastically increasing oxygen transfer area.
- **Downstream Processing:** Separation and purification of biosynthesized products, followed by preservation, clinical trials, and strict quality control.

## 13.4 Transgenic Applications in Agriculture, Medicine & Biopiracy

**High-Yield Transgenic Applications & Bioethics**

1. **Bt Cotton (Pest-Resistant Transgenic Crop):** Soil bacterium *Bacillus thuringiensis* produces crystal endotoxin proteins (**Cry proteins**). The inactive protoxin is solubilized into active toxin in the **alkaline pH of the insect midgut**, binding to midgut epithelial cells, creating pores, causing cell lysis and insect death:
   - **CryIAc** and **CryIIAb** control Cotton Bollworms.
   - **CryIAb** controls Corn Borer.
2. **RNA Interference (RNAi):** Cellular defense against mobile genetic elements. Root-knot nematode *Meloidogyne incognita* infects roots of tobacco plants. Using *Agrobacterium* vectors, nematode-specific genes are introduced producing both sense and anti-sense RNA $\implies$ form **double-stranded RNA (dsRNA)** $\implies$ silences target mRNA of nematode.
3. **Genetically Engineered Insulin (Humulin):** Human insulin consists of chain A (21 amino acids) and chain B (30 amino acids) linked by **disulfide bonds**. Synthesized as a prohormone containing an extra **C-peptide** (33 amino acids). In 1983, American company **Eli Lilly** produced two separate DNA sequences for chains A and B in *E. coli*, purified them, and combined them via disulfide bonds.
4. **Gene Therapy for ADA Deficiency:** First clinical gene therapy performed in 1990 on a 4-year-old girl with **Adenosine Deaminase (ADA) deficiency** (causes Severe Combined Immunodeficiency - SCID). Lymphocytes from patient's blood transduced with functional ADA cDNA using a retroviral vector and returned to patient. (Cured permanently only if introduced into embryonic stem cells).

5. **Transgenic Animals & Biopiracy:**
   - **Rosie Cow (1997):** Produced human protein-enriched milk (2.4 g/L) containing **human alpha-lactalbumin**, nutritionally balanced for infants.
   - **Biopiracy Case:** In 1997, US company RiceTec was granted a patent on Basmati rice cross-bred with semi-dwarf varieties, misappropriating traditional Indian farmers' strains.

CHAPTER 14

# Immunology, Human Pathogens & Evolutionary Biology

Human health reflects complex interactions between host immune defense systems, pathogenic virulence, and evolutionary adaptations. This chapter investigates innate and adaptive immunity, antibody molecular architecture, infectious pathogens, the digenetic life cycle of *Plasmodium*, retroviral HIV pathogenesis, cancer biology, abiogenetic chemical evolution, homologous versus analogous evidence, and the geological timeline of human evolution.

## 14.1 Innate & Adaptive Immunology, Antibody Architecture ($H_2L_2$)

### Innate Defense Barriers & Adaptive Lymphocyte Differentiation

**1. Innate Immunity (Non-Specific First & Second Lines of Defense):**

1. **Physical Barriers:** Keratinized skin stratum corneum; mucous coating of respiratory, gastrointestinal, and urogenital tracts.
2. **Physiological Barriers:** Gastric hydrochloric acid (pH $\approx 1.5 - 2.0$); Lysozyme in tears and saliva.
3. **Cellular Barriers:** Phagocytes: Polymorphonuclear leukocytes (**PMNL** / neutrophils, $60 - 65\%$), tissue macrophages, and **Natural Killer (NK) cells**.
4. **Cytokine Barriers:** Virus-infected cells secrete glycoproteins called **Interferons** ($\alpha, \beta, \gamma$), protecting uninfected cells from viral infection.

**2. Adaptive (Acquired) Immunity: Humoral (AMI) vs Cell-Mediated (CMI):**

- **Humoral / Antibody-Mediated Immunity (AMI):** Mediated by **B-lymphocytes** differentiating into plasma cells that secrete soluble antibodies ($H_2L_2$) into blood and lymph.
- **Cell-Mediated Immunity (CMI):** Mediated by **T-lymphocytes** (Helper $CD4^+T_H$ and Cytotoxic $CD8^+T_C$ cells). **Primary mediator of graft/allograft rejection!** (Immune suppression achieved using **Cyclosporin A** from *Trichoderma polysporum*).

**3. Antibody Molecular Architecture ($H_2L_2$) & Classes:** Each antibody comprises two identical Heavy (H) chains ($\approx 50\,\text{kDa}$) and two identical Light (L) chains ($\approx 25\,\text{kDa}$), cross-linked by **interchain disulfide bonds** ($-S-S-$):

$$\text{Antibody Formula: } H_2L_2 \tag{14.1}$$

- IgG**:** Monomer; most abundant ($75-80\%$); **only class that crosses the human placenta**.
- IgA**:** Dimer with J-chain; secretory antibody abundant in **Colostrum** (first milk) and saliva.
- IgM**:** Pentamer; largest molecular weight; first antibody synthesized in primary immune response.
- IgE**:** Monomer; mediates **Type I Hypersensitivity (Allergies)**; triggers mast cell histamine release.
- IgD**:** Monomer; B-cell antigen receptor.

**4. Lymphoid Organs:**

- **Primary (Generative): Bone marrow** (origin of all blood cells; B-cell maturation) and **Thymus** (T-cell maturation).
- **Secondary (Peripheral): Spleen** (blood filter, major erythrocyte reservoir), **Lymph nodes**, and **MALT** (Mucosa-Associated Lymphoid Tissue, constitutes $\approx 50\%$ **of all lymphoid tissue in humans**).

## 14.2 Major Infectious Diseases & *Plasmodium* Digenetic Life Cycle

### Human Pathogenic Spectrum & Malarial Cycle Mechanics

**1. Major Infectious Diseases:**

| Disease | Pathogen | Transmission | Diagnostic Pathology & Hallmarks |
|---|---|---|---|
| **Typhoid Mary Mallon.** | *S. typhi* High sustained fever ($39-40^{\circ}$C), intestinal perforation. Test: **Widal Test**. | Food / water; | |
| **Pneumonia** ***H. influenzae*** **aerosols.** | *S. pneumoniae*, Droplets / **Infects pulmonary alveoli**; fluid accumulation, gray-blue fingernails. | | |
| **Common Cold** **fomites.** | **Rhino viruses** **Infects nose/nasal passage, NOT lungs!** Lasts $3-7$ days. | Droplets / | |
| **Amoebiasis** **vector.** | *E. histolytica* Large intestine; **excess mucous and blood clots in stools**. | Houseflies | |
| **Ascariasis** **food / water.** | *A. lumbricoides* Internal bleeding, fever, mechanical **intestinal obstruction**. | Fecal-oral; | |
| **Filariasis** ***W. malayi*** ***Culex.*** | *W. bancrofti*, Female Chronic inflammation of **lymphatic vessels of lower limbs** (Elephantiasis). | | |
| **Ringworm** ***Trichophyton*** | *Microsporum*, Direct contact. | Dry, scaly, intensely itchy lesions on skin, scalp, groin. | |

## 2. Digenetic Life Cycle of *Plasmodium* (Malarial Parasite):

- **Human Host (Asexual Cycle / Intermediate Host):** Female *Anopheles* mosquito inoculates infective **Sporozoites** with saliva → invade hepatocytes (liver schizogony) → burst hepatocytes releasing merozoites → invade RBCs (erythrocytic schizogony) → burst RBCs synchronously every $48-72$ hours, releasing pyrogenic **Haemozoin** causing chills and recurring high fever → develop into male/female gametocytes.
- **Mosquito Host (Sexual Cycle / Definitive Host):** Female *Anopheles* ingests gametocytes during blood meal → fertilization occurs in the

**mosquito gut** → motile **ookinete** penetrates gut wall forming **oocyst** → sporogony generates thousands of sporozoites which lodge in **salivary glands**.

## 14.3 Pathologies: HIV/AIDS Replication, Cancer Biology & Addiction Pharmacology

### HIV Retrovirus Pathogenesis, Oncogenesis & Drug Pharmacology

**1. HIV/AIDS Pathogenesis:** Caused by Human Immunodeficiency Virus, an enveloped retrovirus harboring **two identical +ssRNA molecules** and Reverse Transcriptase:

1. gp120 envelope protein binds CD4 **receptors** on macrophages and Helper T ($T_H$) lymphocytes.
2. Reverse Transcriptase synthesizes viral cDNA → Integrase integrates viral DNA into host genome.
3. Macrophages act as **HIV factories**. Virus infects and destroys $CD4^+T_H$ cells; cell count drops below $200/\mu L$, allowing fatal opportunistic infections (*M. tuberculosis*, *Toxoplasma*).
4. Screening: **ELISA**; Confirmatory: **Western Blot**; Treatment: ART (e.g., Zidovudine/AZT).

**2. Cancer Biology:** Loss of **contact inhibition** leads to uncontrolled proliferation into neoplasms:

- **Malignant Tumors:** Exhibit **Metastasis** (tumor cells slough off into circulation and colonize distant organs; the most dreaded characteristic of cancer).
- **Carcinogens:** Physical (ionizing X/$\gamma$-rays, non-ionizing UV-B thymine dimers); Chemical (tobacco nitrosamines); Biological (oncogenic viruses, mutated proto-oncogenes).
- **Therapies:** Surgery, Radiotherapy, Chemotherapy (Vincristine, Vinblastine), and Immunotherapy ($\alpha$**-Interferon**).

**3. Addiction Pharmacology:**

- **Opioids:** Extracted from latex of *Papaver somniferum* (Opium poppy). Heroin (**Smack** / diacetylmorphine) synthesized by acetylation of morphine; CNS and gastrointestinal depressant.
- **Cannabinoids:** Extracted from *Cannabis sativa*; bind cannabinoid receptors in brain, impacting cardiovascular system.
- **Cocaine:** Extracted from *Erythroxylum coca*; inhibits dopamine reuptake, producing euphoria; induces hallucinations in excess.

- **Nicotine:** Tobacco alkaloid; stimulates adrenal medulla to release adrenaline/noradrenaline.

## 14.4 Evolutionary Biology: Chemical Evolution, Natural Selection & Human Lineage

### Abiogenesis, Morphological Evidences & Human Evolution Timeline

**1. Origin of Life (Oparin-Haldane & Miller-Urey Experiment, 1953):** Prebiotic Earth had a reducing atmosphere lacking free $O_2$, rich in $CH_4, NH_3, H_2O$ vapor, $H_2$. Stanley Miller discharged electrical sparks ($\approx 75,000\,V$) into a gas chamber containing $CH_4 : NH_3 : H_2$ in ratio **2 : 1 : 2** at **800**°C. Effluent contained amino acids: **Glycine, Alanine, and Aspartic acid**, validating chemical abiogenesis.

**2. Homology vs Analogy:**

- **Homologous Organs (Divergent Evolution):** Same anatomical origin and basic structure, modified for different functions (e.g., forelimb bones of Whale, Bat, Cheetah, Human; Thorns of *Bougainvillea* vs Tendrils of *Cucurbita*).
- **Analogous Organs (Convergent Evolution):** Different anatomical origin, modified for similar functions due to similar selective pressures (e.g., wings of Butterfly vs Bird; eye of Octopus vs Mammal; flippers of Penguin vs Dolphin; Sweet potato [root] vs Potato [stem]).

**3. Chronological Timeline of Human Evolution:**

| Hominid | Era | Origin | Cranial Key Anatomical Traits |
|---|---|---|---|
| *Dryopithecus* | ≈ 15 mya | Africa / | |
| Eurasia | 400 cc | Arboreal, hairy, walked like apes. | |
| *Ramapithecus* | ≈ 15 mya | Shivalik | |
| hills | 450 cc | More man-like; upright tendency. | |
| *Australopithecus* | 2 – 3 mya | East | |
| Africa | 500 cc | **Bipedal**; stone weapons; **ate fruit**. | |
| *Homo habilis* | 1.7 – 2 mya | Olduvai | |
| Gorge | **700** cc | **First tool maker**; **no meat**. | |
| *Homo erectus* | ≈ 1.5 mya | Java | |
| fossils | **900** cc | Erect posture; **fire**; **ate meat**. | |
| *Neanderthal* | 100k – 40k ya | Neander | |
| Valley | **1400** cc | Used hides; **buried dead with rituals**. | |
| *Homo sapiens* | 75k – 10k ya | Africa | **1400** cc Modern man; cave art; agriculture. |

CHAPTER 15

# Ecology, Ecosystems & Biodiversity Conservation

Ecology examines the interactions that determine the distribution and abundance of living organisms. This chapter covers organismal abiotic adaptations, mathematical population growth models, interspecific symbiotic interactions, ecosystem bioenergetics and trophic pyramids, decomposition kinetics, and global biodiversity conservation strategies under revised NMC guidelines.

## 15.1 Organisms & Abiotic Adaptations

### Abiotic Factors & Evolutionary Homeostatic Strategies

1. **Key Abiotic Factors:**

- **Temperature:** Most ecologically relevant factor: **Eurythermal** (tolerate wide temperature range) vs **Stenothermal** (restricted to narrow range).
- **Water:** Salt tolerance determines distribution: **Euryhaline** (wide salinity tolerance) vs **Stenohaline** (narrow salinity tolerance).
- **Light:** Governs photosynthesis, photoperiodism, and circadian foraging rhythms.

2. **Organismal Responses to Environmental Stress:**

- **Regulators:** Maintain physiological homeostasis (constant body temperature and osmolarity) through thermoregulation and osmoregulation (all birds, mammals, very few lower vertebrates).
- **Conformers:** Cannot maintain constant internal environment; internal body temperature/osmolarity changes with ambient conditions (99% of animals and nearly all plants).
- **Migrate:** Temporary movement to hospitable areas (e.g., migratory birds from Siberia to Keoladeo National Park, Bharatpur, Rajasthan).
- **Suspend:** Escape in time: **Hibernation** (winter sleep in bears), **Aestivation** (summer sleep in snails/fish), **Diapause** (stage of suspended development in zooplankton), and bacterial **endospores**.

3. **Morphological Adaptations:**

- **Kangaroo Rat:** Meets 100% of water requirement through metabolic oxidation of fats; excretes solid urine.

- **Allen's Rule:** Mammals from colder climates have **shorter ears and shorter limbs** to minimize heat loss.
- **Blubber:** Thick subcutaneous fat layer in polar seals acting as thermal insulator.

## 15.2 Population Growth Formulations & Interspecific Interactions

### Population Growth Formulations: Exponential vs Logistic

**1. Exponential Growth (Unlimited Resources / Geometric Progression):**

$$\frac{\mathbf{dN}}{\mathbf{dt}} = (\mathbf{b} - \mathbf{d})\mathbf{N} = \mathbf{rN} \implies \mathbf{N_t = N_0 e^{rt}} \quad \textbf{(J-shaped curve)} \tag{15.1}$$

where $r$ is the **intrinsic rate of natural increase** (Norway rat: 0.015, Flour beetle: 0.12, Human population India 1981: 0.0205).

**2. Logistic Growth / Verhulst-Pearl Model (Limited Resources / Realistic):**

$$\frac{\mathbf{dN}}{\mathbf{dt}} = \mathbf{rN}\left(\frac{\mathbf{K - N}}{\mathbf{K}}\right) \quad \textbf{(Sigmoid / S-shaped curve)} \tag{15.2}$$

where $K$ is the **Carrying Capacity** (maximum population density the habitat can sustain indefinitely). When $N \to K$, environmental resistance $\left(\frac{K-N}{K}\right) \to 0$, and population growth rate ceases.

### Interspecific Population Interactions Matrix

| Interaction | Sp. A | Sp. B | NCERT Canonical Ecological Examples |
|---|---|---|---|
| **Mutualism** | + | + | Lichens (Algae + Fungi); Mycorrhizae (Fungi + Roots of higher plants); Fig tree and pollinator Fig Wasp; Mediterranean orchid *Ophrys* (**sexual deceit** mimicking female bee *Colpa*). |
| **Competition** | − | − | Abingdon tortoise in Galapagos became extinct within a decade after goats were introduced (browsing efficiency); **Gause's Competitive Exclusion Principle**; **MacArthur's Warblers** (5 species coexisted via **resource partitioning**). |
| **Predation** | + | − | Biological control: *Cactoblastis* moth controlled invasive prickly pear cactus in Australia (1920); Pisaster starfish removal caused extinction of $>$ 10 invertebrate species; Monarch butterfly accumulates unpalatable cardiac glycosides in caterpillar stage. |
| **Parasitism** | + | − | Ectoparasites: Lice on humans, ticks on dogs, marine copepods on fish, *Cuscuta* on hedge plants; Endoparasites: Liver fluke, *Plasmodium*; **Brood Parasitism** (Cuckoo/Koel lays eggs in Crow's nest). |
| **Commensalism** | + | 0 | Epiphytic orchid on mango tree; Barnacles on whale; Cattle egret foraging alongside grazing cattle; Clownfish living among stinging tentacles of Sea anemone. |
| **Amensalism** | − | 0 | *Penicillium* secreting penicillin antibiotic inhibiting bacterial growth; Black walnut (*Juglans nigra*) secreting juglone inhibiting understory vegetation. |

## 15.3 Ecosystem Bioenergetics, Trophic Pyramids & Ecological Succession

### Productivity Equations, Decomposition Stages & Trophic Pyramids

**1. Primary Productivity Formulations:**

- **Gross Primary Productivity (GPP):** Total rate of organic matter synthesis during photosynthesis.
- **Net Primary Productivity (NPP):** NPP = GPP − $R$ (where $R$ is respiratory loss).
- **Global Annual Biosphere NPP:** $\approx$ **170** billion tons (dry organic matter).
- **Oceans Productivity:** **55** billion tons (oceans cover 70% area but yield only 32% NPP).

**2. Five Stages of Decomposition:** Detritus undergoes simultaneous physical and biochemical processing:

1. **Fragmentation:** Breakdown of detritus into smaller particles by detritivores (e.g., Earthworms).
2. **Leaching:** Water-soluble inorganic nutrients percolate into soil horizons and precipitate as unavailable salts.

3. **Catabolism:** Bacterial and fungal enzymes degrade detritus into simpler inorganic substances.
4. **Humification:** Accumulation of dark, amorphous, colloidal **Humus** (highly resistant to microbes; acts as nutrient reservoir).
5. **Mineralization:** Slow microbial release of inorganic nutrients from humus degradation.

**3. Energy Flow & Ecological Pyramids:**

- **10% Law of Lindeman:** Only **10**% of energy entering a trophic level is transferred to the next higher level; 90% is dissipated as metabolic heat.
- **Photosynthetically Active Radiation (PAR):** Out of total incident solar radiation, less than **50**% is PAR. Plants capture only $\mathbf{2-10}\%$ of PAR ($1-5\%$ of total incident radiation).
- **Pyramid of Numbers:** Mostly upright (grassland); **inverted** in tree ecosystem supporting thousands of parasites.
- **Pyramid of Biomass:** Upright on land; **Inverted in Aquatic / Marine Ecosystems** (phytoplankton biomass is significantly smaller than zooplankton/fishes).
- **Pyramid of Energy: Always and strictly upright**; can never be inverted (Second Law of Thermodynamics).
- **Succession:** Both **Hydrarch** (water) and **Xerarch** (bare rock) successions converge toward identical moderate water conditions called **Mesic conditions**.

## 15.4 Global Biodiversity Metrics, Species-Area Law & Conservation

### Species-Area Relationship (Alexander von Humboldt)

Within a region, species richness increases with increasing explored area, up to a limit:

$$S = CA^{Z} \implies \log \mathbf{S} = \log \mathbf{C} + \mathbf{Z} \log \mathbf{A} \tag{15.3}$$

where $S$ = Species richness, $A$ = Area, $C$ = Y-intercept, $Z$ = **Slope of regression line (extinction sensitivity)**:

- For regional areas (plants in Britain, birds in California): $Z$ is strictly 0.1 **to** 0.2.
- For entire continents (frugivorous birds and mammals in tropical forests): $Z$ is much steeper, 0.6 **to** 1.2 (typically 1.15).

**Global Biodiversity Metrics & The "Evil Quartet" Causes of Loss**

**1. Global Biodiversity (Robert May's Estimate):** Total species on Earth $\approx$ **7** million. Animals constitute $> 70\%$ of recorded species, plants $\leq 22\%$. Among animals, **insects constitute** $> 70\%$ (7 out of 10 animals are insects). India possesses **2.4% of global land area** but accounts for **8.1% of global species diversity** (one of 12 mega-diversity nations).

**2. The "Evil Quartet" (Major Drivers of Biodiversity Extinction):**

1. **Habitat Loss & Fragmentation:** The single most catastrophic cause. Tropical rainforests once covered 14% of Earth's land, now reduced to $< 6\%$. Amazon rainforest is being cleared for soybean cultivation and cattle pasture.
2. **Over-Exploitation:** Excessive hunting led to extinction of *Steller's sea cow* and *Passenger pigeon*.
3. **Alien Species Invasions:**
   - Nile perch introduced into Lake Victoria caused extinction of $> 200$ endemic *Cichlid fish* species.
   - Invasive weeds: *Parthenium* (Carrot grass), *Lantana*, and *Eichhornia crassipes* (Water Hyacinth / "Terror of Bengal").
   - African catfish *Clarias gariepinus* introduced illegally for aquaculture threatens indigenous catfishes.
4. **Co-Extinctions:** Obligate co-evolved relationships (e.g., plant-pollinator mutualism, host fish and parasite).

**3. In-Situ vs Ex-Situ Conservation:**

- **In-Situ (On-Site):** 14 Biosphere Reserves, 90 National Parks, 448 Wildlife Sanctuaries, Sacred Groves (Khasi and Jaintia Hills in Meghalaya, Aravalli Hills in Rajasthan, Western Ghats).
- **Ex-Situ (Off-Site):** Zoological parks, Botanical gardens, Wildlife safari parks, Cryopreservation of gametes at $-196^{\circ}$C in liquid nitrogen, Seed banks, Tissue culture.

APPENDIX A

# High-Yield Quantitative Biology & Master Diagnostic Matrices

To achieve a perfect score of 360 in NEET Biology, students must master both qualitative concepts and quantitative formulations. This appendix provides a unified analytical compendium of genetic probability calculations, recombination frequencies, population genetics models, physiological capacity values, floral formula matrices for all seven prescribed plant families, and high-frequency Assertion-Reasoning anchors.

## A.1 Quantitative Genetics & Probability Calculus

### Foundational Mendelian Probability Principles

Mendelian inheritance patterns obey the fundamental laws of probability:

1. **Rule of Multiplication (Product Rule):** The probability of two independent events occurring simultaneously is the product of their individual probabilities:
$$P(A \cap B) = P(A) \times P(B) \tag{A.1}$$
*Application:* Probability of obtaining a homozygous recessive genotype ($aabb$) from a dihybrid cross ($AaBb \times AaBb$) is $P(aa) \times P(bb) = \frac{1}{4} \times \frac{1}{4} = \frac{1}{16}$.
2. **Rule of Addition (Sum Rule):** The probability of either of two mutually exclusive events occurring is the sum of their individual probabilities:
$$P(A \cup B) = P(A) + P(B) \tag{A.2}$$
*Application:* In a monohybrid cross ($Aa \times Aa$), the probability of obtaining a heterozygote ($Aa$) via either maternal or paternal allele donation is $\frac{1}{4} + \frac{1}{4} = \frac{1}{2}$.

### Master Combinatorial Formulas for Multi-Gene Crosses

For a cross involving $n$ independently assorting heterozygous gene pairs ($AaBbCc\ldots$):

- **Genetically Distinct Gametes:** $N_{\text{gametes}} = 2^n$
- **Total Offspring Genotypes:** $N_{\text{genotypes}} = 3^n$

- **Total Offspring Phenotypes:** $N_{\text{phenotypes}} = 2^n$ (assuming complete dominance)
- **Squares in Punnett Grid:** $N_{\text{grid}} = 2^n \times 2^n = 4^n$
- **Fraction Completely Recessive:** $F_{\text{recessive}} = (1/4)^n$
- **Fraction Completely Dominant:** $F_{\text{dominant}} = (1/4)^n$
- **Fraction Completely Heterozygous:** $F_{\text{hetero}} = (1/2)^n$

### Chi-Square ($\chi^2$) Goodness-of-Fit Test in Genetic Analysis

To determine whether observed offspring numbers deviate significantly from theoretical Mendelian expectations:

$$\chi^2 = \sum \frac{(O - E)^2}{E} \tag{A.3}$$

where $O$ is the observed frequency, $E$ is the expected frequency based on Mendelian ratio, and degrees of freedom $df = k - 1$ (where $k$ is the number of phenotypic classes). If $\chi^2_{\text{calc}} < \chi^2_{\text{critical}}$ at $p = 0.05$, the observed deviation is attributed strictly to random chance and the Mendelian hypothesis is accepted.

## A.2 Recombination Frequency & Chromosome Mapping Calculus

### Morgan-Sturtevant Gene Mapping Formulations

The recombination frequency ($RF$) between two linked genes on a chromosome is directly proportional to the physical distance between them:

$$RF = \left( \frac{\text{Total Number of Recombinant Offspring}}{\text{Total Number of Offspring Progeny}} \right) \times 100\% \tag{A.4}$$

**Units & Metric Standards:**

- 1% Recombination Frequency = 1 Map Unit (m.u.) = 1 centimorgan (cM).
- Maximum observable recombination frequency for two genes on the same chromosome is **50**% (at which point they assort independently, indistinguishable from unlinked genes on different chromosomes).

$$C = \frac{\text{Observed Double Crossovers}}{\text{Expected Double Crossovers}} \tag{A.5}$$

$$I = 1 - C \quad \text{(Interference)} \tag{A.6}$$

*Interpretation:* If $I > 0$ (positive interference), a crossover in one interval suppresses crossover formation in an adjacent chromosomal region. If $I = 1$, double crossovers are completely suppressed.

## A.3 Population Genetics: Hardy-Weinberg Mathematical Model

### Hardy-Weinberg Equilibrium Equations

For a diploid population with two alleles ($A$ and $a$) at a single locus with frequencies $p$ and $q$:

$$p + q = 1 \quad \text{(Allelic Frequencies)} \tag{A.7}$$

$$p^2 + 2pq + q^2 = 1 \quad \text{(Genotypic Frequencies)} \tag{A.8}$$

where:

- $p^2$: Frequency of homozygous dominant individuals ($AA$).
- $2pq$: Frequency of heterozygous carrier individuals ($Aa$).
- $q^2$: Frequency of homozygous recessive affected individuals ($aa$).

**NEET Trap: Allelic Frequency vs Genotypic Frequency**

Pay meticulous attention to the wording in Hardy-Weinberg exam questions:

- If given "frequency of recessive **allele**": that is $q$.
- If given "frequency of recessive **disease / phenotype / individuals**": that is $q^2$. You must compute $\sqrt{q^2} = q$ first, then find $p = 1 - q$, and finally compute carrier frequency $2pq$.
- **Equilibrium Preconditions:** Strictly valid only for: (1) large population size, (2) random mating, (3) no gene flow / migration, (4) no mutation, and (5) no natural selection.

## A.4 Quantitative Human Physiology Matrices

## A.5 Master Systematic Comparison of All 7 Prescribed Plant Families

| Physiological Parameter | Standard Value | Mathematical Definition / Formula |
|---|---|---|
| Tidal Volume (TV) | 500 mL | Volume inspired or expired during normal breathing |
| Inspiratory Reserve Volume (IRV) | 2500–3000 mL | Extra volume inspired forcibly above TV |
| Expiratory Reserve Volume (ERV) | 1000–1100 mL | Extra volume expired forcibly after normal expiration |
| Residual Volume (RV) | 1100–1200 mL | Gas remaining in lungs after maximal expiration |
| Vital Capacity (VC) | 3500–4500 mL | $VC = TV + IRV + ERV$ |
| Total Lung Capacity (TLC) | 5000–6000 mL | $TLC = VC + RV = TV + IRV + ERV + RV$ |
| Cardiac Output (CO) | $\approx 5000$ mL/min | $CO =$ Stroke Volume (70 mL) $\times$ Heart Rate (72 bpm) |
| Glomerular Filtration Rate (GFR) | 125 mL/min | Equivalent to 180 L/day across both kidneys |
| Net Filtration Pressure (NFP) | 10 mmHg | $NFP = GHP\,(60) - [BCOP\,(32) + CHP\,(18)]$ |
| Urine Excretion Rate | 1.0–1.5 L/day | $>$ 99% of filtrate is reabsorbed by renal tubules |

**Table A.1:** Master Diagnostic Matrix: Quantitative Human Physiological Constants.

| Family | Floral Formula | Androecium | Gynoecium | Key Diagnostic Examples |
|---|---|---|---|---|
| **Malvaceae** | $\oplus\ \mathrm{Epi}_{3-8}\ \mathrm{K}_{(5)}\ \mathrm{C}_5\ \mathrm{A}_{(\infty)}\ \underline{\mathrm{G}}_{(5-\infty)}$ | Monadelphous stamens, monothecous reniform anthers | Syncarpous, superior, axile placentation | *Hibiscus*, Cotton (*Gossypium*), Lady's finger (*Abelmoschus*) |
| **Cruciferae** | $\oplus\ \mathrm{K}_{2+2}\ \mathrm{C}_4\ \mathrm{A}_{2+4}\ \underline{\mathrm{G}}_{(2)}$ | Tetradynamous stamens (2 short outer + 4 long inner) | Bicarpellary, unilocular becoming bilocular via false septum (replum) | Mustard (*Brassica*), Radish (*Raphanus*), Turnip |
| **Fabaceae** | $\%\ \mathrm{K}_{(5)}\ \mathrm{C}_{1+2+(2)}\ \mathrm{A}_{(9)+1}\ \underline{\mathrm{G}}_1$ | Diadelphous stamens $[(9)+1]$, dithecous | Monocarpellary, superior, marginal placentation | Gram, Pea, Soybean, Sunnhemp, Lupin, *Sesbania* |
| **Asteraceae** | $\oplus$ or $\%\ \mathrm{K}_{\mathrm{pappus}}\ \mathrm{C}_{(5)}\ \mathrm{A}_{(5)}\ \overline{\mathrm{G}}_{(2)}$ | Syngenesious (anthers fused, filaments free) | Bicarpellary, inferior, basal placentation, single ovule | Sunflower (*Helianthus*), Marigold, *Taraxacum* |
| **Poaceae** | $\%\ \mathrm{P}_2\ \mathrm{A}_{3\ \text{or}\ 6}\ \underline{\mathrm{G}}_1$ | Stamens versatile, pendulous | Monocarpellary, superior, basal placentation, feathery stigma | Wheat (*Triticum*), Rice, Maize, Bamboo, Sugarcane |
| **Solanaceae** | $\oplus\ \mathrm{K}_{(5)}\ \overbrace{\mathrm{C}_{(5)}\ \mathrm{A}_5}\ \underline{\mathrm{G}}_{(2)}$ | Epipetalous stamens, alternipetalous | Bicarpellary, syncarpous, superior, obliquely placed ovary, swollen placenta | Potato, Tomato, Brinjal, Chilli, Belladonna, Ashwagandha, Tobacco |
| **Liliaceae** | $\oplus\ \overbrace{\mathrm{P}_{(3+3)}\ \mathrm{A}_{3+3}}\ \underline{\mathrm{G}}_{(3)}$ | Epiphyllous stamens in two whorls | Tricarpellary, syncarpous, superior, axile placentation | *Allium* (Onion, Garlic), *Aloe*, *Asparagus*, *Colchicum autumnale* |

**Table A.2:** Systematic Quick-Diagnostic Matrix for the 7 NMC-Prescribed Angiosperm Families.

**Direct NCERT High-Yield Assertion-Reason Anchors**

- **Floral Formulas as Definitive Identifiers:** In Solanaceae, the ovary is characteristically **obliquely placed** with a swollen placenta bearing many ovules; in Cruciferae, the ovary becomes bilocular due to a false septum called **replum**.
- **Staminal Adaptations:** Diadelphous stamens [(9)+1] are unique to Fabaceae; monadelphous stamens with monothecous anthers are diagnostic of Malvaceae; tetradynamous stamens $(2 + 4)$ are exclusive to Cruciferae; and syngenesious stamens are hallmark to Asteraceae.
- **Placentation Anchor:** Marginal (Fabaceae/Pea), Parietal with replum (Cruciferae/Mustard), Axile (Solanaceae, Malvaceae, Liliaceae), Basal (Asteraceae/Sunflower, Poaceae/Wheat).

# About the Author & Errata Submissions

**Satya Prakash** is an educator, curriculum researcher, and author dedicated to creating rigorous, student-first learning resources for competitive examinations in India. His focus is on eliminating cognitive overload, dismantling obsolete legacy syllabi, and restoring conceptual rigor and mathematical transparency to competitive pre-medical preparation.

## Official Project Repository & Continuous Improvement

Science education thrives on continuous scrutiny and collective refinement. If you identify a typo, biological discrepancy, or pedagogical ambiguity, or if you wish to suggest an enhancement for future editions, please submit your feedback:

- **Direct Author Email:** spdedsec@hotmail.com
- **GitHub Open-Source Repository:** `https://github.com/spdedsec/neet-study-materials`
- **Errata Tracker:** Verified errata and updates are posted directly to the repository.

---



Authored by Satya Prakash · All Rights Reserved