NEET UG BIOLOGY

MASTER TEXTBOOK

NEET UG 2027 Master Reference Series · Complete Authoritative Edition

NEET UG BIOLOGY

MASTER TEXTBOOK

Complete Botany & Zoology: Genetics, Plant Physiology, Reproduction, Ecology, Cell Biology, Human Physiology, Health, Evolution & Quantitative Matrices

Authored by

Satya Prakash

Strictly Calibrated to the Revised NMC Syllabus & Rationalized NCERT Textbooks

Target Examination: National Eligibility cum Entrance Test (NEET UG 2027)

Published for Independent Medical Aspirants, Educators & Academic Study Circles

Comprehensive Academic Edition · Global Print-on-Demand

NEET UG BIOLOGY MASTER TEXTBOOK

Complete Botany, Zoology, Plant Families, Human Physiology & Quantitative Matrices

Author: Satya Prakash
Editorial Correspondence & Academic Feedback: spdedsec@hotmail.com
Source Repository & Official Errata: https://github.com/spdedsec/neet-study-materials

© 2026–2027 Satya Prakash. All rights reserved.

First Edition: October 2026
Publishing Imprint: Independent Academic Publishing / Satya Prakash Editorial Imprint
Target Cycle: NEET UG 2027 (Undergraduate Medical Admissions Entrance Examination)
ISBN-13 (Paperback): 979-8-XXXXXXXXX (Placeholder for Official Assigned ISBN)
ISBN-13 (Hardcover): 979-8-XXXXXXXXX (Placeholder for Official Assigned ISBN)

All Rights Reserved. No part of this publication may be reproduced, stored in a retrieval system, or transmitted in any form or by any means—electronic, mechanical, photocopying, recording, scanning, or otherwise—without the prior written permission of the author, except for brief quotations embodied in critical reviews or scholarly analysis as permitted under copyright law.

Academic & Legal Disclaimer: This textbook is an independent preparation manual authored by Satya Prakash to assist candidates preparing for the National Eligibility cum Entrance Test (NEET UG). While every effort has been made to verify biological accuracy and strict concordance with the National Medical Commission (NMC) curriculum and rationalized NCERT textbooks, the author and publisher make no official claims of affiliation with or endorsement by the National Testing Agency (NTA), the National Medical Commission (NMC), or the Ministry of Education, Government of India. The author disclaims any liability for errors or omissions.

Typesetting & Typography: Typeset in LaTeX 2ε with the Tectonic XeTeX engine using Latin Modern typefaces. Biological taxonomic nomenclature follows the International Code of Nomenclature for algae, fungi, and plants (ICN) and the International Code of Zoological Nomenclature (ICZN). Produced and bound under rigorous academic publishing standards.

Printed in the United States of America / United Kingdom / India.

To my parents, whose selfless dedication and love illuminated every step of my journey;

To the biologists and physicians whose tireless investigations into the living world revealed the exquisite unity of cellular and molecular life;

And to you, future doctor—
may the biological wonders detailed in these pages inspire not only your success in NEET UG,
but a lifelong reverence for the human body and the living systems you will soon heal.

Acknowledgements

Writing an exhaustive, 360-mark reference textbook for NEET UG Biology requires balancing two essential qualities: absolute textual loyalty to rationalized NCERT benchmarks, and clinical conceptual depth that renders a student completely immune to tricky Assertion-Reasoning and Statement-based questions.

I express my deepest gratitude to my family for their unending understanding, encouragement, and sacrifices throughout the compilation of this manuscript.

I thank the academic biology faculty, medical researchers, and botanists whose critical discussions helped ensure that every floral formula, enzyme kinetic relation, and endocrine loop was represented with anatomical and physiological precision.

I am especially grateful to the hundreds of medical aspirants whose questions, doubts, and mock examination analyses directly shaped the high-yield tables and "NEET Trap" callouts throughout this book. Your pursuit of excellence was the primary catalyst for this work.

Finally, I acknowledge the developers of LaTeX 2_ε and Tectonic, whose typesetting ecosystem allowed us to integrate comprehensive taxonomic matrices, biochemical pathways, and color-coded diagnostic summaries into a publication-ready authoritative manuscript.

Satya Prakash
October 2026

Author's Preface: Straight Talk on NEET Biology

Let's cut straight to the chase: Biology carries exactly **360 marks out of 720** in NEET UG. Exactly half the examination.

Because of this, you have probably heard the common advice a thousand times: *"Just read NCERT ten times and you will get 360/360."*

Let me tell you what actually happens in the examination hall. Every year, thousands of brilliant, hard-working students read NCERT ten times, walk into the exam, and come out with 310 or 320 instead of 360. Why? Because they memorized the lines passively without understanding the underlying physiological mechanisms, without analyzing the exceptions in comparative tables, and without realizing how NTA twists two unrelated NCERT sentences into a lethal Assertion-Reasoning pair.

Furthermore, the National Medical Commission (NMC) significantly reorganized the syllabus:

- **Deleted Chapters:** Entire legacy units were deleted—including *Digestion and Absorption*, *Transport in Plants*, *Mineral Nutrition*, and *Strategies for Enhancement in Food Production*. If you are spending hours memorizing nitrogenase enzyme mechanisms from Mineral Nutrition or digestive enzymes in the duodenum, you are burning study hours on zero-yield topics.
- **Newly Mandated Topics:** The NMC explicitly mandated **seven distinct angiosperm families** (Malvaceae, Cruciferae/Brassicaceae, Fabaceae/Leguminosae, Asteraceae/Compositae, Poaceae/Gramineae, Solanaceae, and Liliaceae) and reinstated detailed animal morphology for both **Frog** and **Cockroach**.

I wrote this master book to give you the ultimate, zero-compromise Biology textbook for NEET UG 2027:

1. **Complete Integration of Botany and Zoology:** Part I gives you the complete Botany curriculum—from photosynthesis and molecular genetics to reproductive biology, ecology, and all seven prescribed plant families. Part II delivers complete Zoology—cell biology, mitosis/meiosis, animal taxonomy, frog and cockroach morphology, integrated human physiology, immunology, and evolutionary biology.
2. **Quantitative Biology & Mathematical Formulations (Chapter 8):** Most biology books treat genetics and physiology as purely descriptive. But NEET regularly tests dihybrid and trihybrid probability calculations, recombination frequencies and gene mapping (centimorgan calculations), Hardy-Weinberg equilibrium algebra ($p^2 + 2pq + q^2 = 1$), pulmonary spirometry volumes, cardiac output calculations, and renal net filtration pressure. Chapter 8 brings all these quantitative principles together with full derivations.
3. **The Five-Color Callout Architecture:**
   - **Concept Boxes (`conceptbox`):** Structural definitions, anatomical relationships, and cellular pathways.
   - **Formula Boxes (`formulabox`):** Exact biochemical equations, genetic ratios, and physiological measurements.
   - **Use-Case Boxes (`usecasebox`):** Clinical correlations, ecological applications, and biotechnology techniques.
   - **NEET Trap Boxes (`trapbox`):** Tricky exceptions, easily confused terms (e.g., tracheids vs vessels, juxtamedullary vs cortical nephrons, homologous vs analogous structures), and Assertion-Reason traps.
   - **Direct NCERT Lines (`ncertbox`):** Verbatim high-yield sentences and table extractions that NTA loves to test.

My advice to you as a friend: do not just highlight sentences. When you study a physiological loop (like the Renin-Angiotensin-Aldosterone System or the photolysis of water), close the book and draw the flowchart from memory. When you study genetics, work out the Punnett squares by hand.

You have the discipline within you to master all 360 marks of Biology. Treat this book as your constant companion, review your error logs honestly, and stay committed to the dream that drove you to start this journey.

Satya Prakash
Author & Dedicated Mentor
spdedsec@hotmail.com
October 2026

Syllabus Blueprint & NMC Rationalization Guide

This section details the official NEET UG 2027 Biology curriculum prescribed by the National Medical Commission (NMC), covering both Botany and Zoology domains, examination weightage, and critical syllabus deletions.

Official Biology Examination Architecture

Subject Domain	Section A	Section B	Total Questions	Marks	Question Typology
Botany	35	15 (Attempt 10)	45	180	Single Choice, MCQs,
Zoology	35	15 (Attempt 10)	45	180	Assertion-Reason, Statement I/II,
Master Biology Total	**70**	**30 (Attempt 20)**	**90**	**360**	Match the Column (Matrix)

Table 1: NEET UG Biology Examination Scheme (+4 for correct response, −1 for incorrect response, 0 for unattempted).

Official 10-Unit Biology Syllabus Structure

1. **Unit 1: Diversity in Living World** (What is living? Biodiversity, taxonomic hierarchy, binomial nomenclature, five-kingdom classification: Monera, Protista, Fungi, Plantae, Animalia; Salient features of non-chordates up to phyla level and chordates up to class level).
2. **Unit 2: Structural Organisation in Animals and Plants** (Morphology, anatomy and functions of different parts of flowering plants: Root, stem, leaf, inflorescence, flower, fruit and seed; Description of families: Malvaceae, Cruciferae/Brassicaceae, Fabaceae/Leguminosae, Asteraceae/Compositae, Poaceae/Gramineae, Solanaceae, Liliaceae; Animal tissues; Morphology, anatomy and functions of digestive, circulatory, respiratory, nervous and reproductive systems of Frog and Cockroach).
3. **Unit 3: Cell Structure and Function** (Cell theory, prokaryotic and eukaryotic cells, plant and animal cells, cell envelope, organelles: nucleus, mitochondria, chloroplast, endomembrane system; Biomolecules: carbohydrates, proteins, nucleic acids, enzymes; Cell cycle: interphase, mitosis and meiosis).
4. **Unit 4: Plant Physiology** (Photosynthesis in higher plants: light reactions, photolysis, cyclic and non-cyclic photophosphorylation, Calvin cycle, C_4 pathway, photorespiration; Respiration in plants: glycolysis, TCA cycle, electron transport system, oxidative phosphorylation; Plant growth and development: auxins, gibberellins, cytokinins, ethylene, abscisic acid).
5. **Unit 5: Human Physiology** (Breathing and respiration: pulmonary volumes, alveolar gas exchange, oxygen dissociation; Body fluids and circulation: blood composition, ABO/Rh grouping, cardiac cycle, ECG; Excretory products and elimination: nephron function, counter-current mechanism, RAAS; Locomotion and movement: sliding filament theory, skeletal joints; Neural control: action potentials, synapses; Chemical coordination: endocrine hormones and mechanisms).

6. **Unit 6: Reproduction** (Sexual reproduction in flowering plants: microsporogenesis, megasporogenesis, double fertilization, apomixis; Human reproduction: male and female reproductive systems, gametogenesis, menstrual cycle, fertilization, embryogeny; Reproductive health: contraception, sexually transmitted infections, assisted reproductive technologies).
7. **Unit 7: Genetics and Evolution** (Heredity and variation: Mendelian ratios, incomplete dominance, codominance, multiple alleles, linkage and crossing over, sex determination, sex-linked disorders; Molecular basis of inheritance: DNA structure, replication, transcription, genetic code, translation, lac operon, DNA fingerprinting; Evolution: origin of life, Darwinian selection, evidence for evolution, Hardy-Weinberg equilibrium, human evolution).
8. **Unit 8: Biology and Human Welfare** (Health and disease: pathogens causing typhoid, pneumonia, malaria, amoebiasis, ringworm, filariasis; Basic principles of immunology: innate and acquired immunity, vaccines; AIDS and cancer: etiology, diagnosis, treatment; Microbes in human welfare: household products, industrial processing, sewage treatment, biofertilizers).
9. **Unit 9: Biotechnology and Its Applications** (Principles and processes: recombinant DNA technology, restriction enzymes, cloning vectors, PCR; Applications in agriculture and medicine: genetically modified crops, Bt cotton, RNA interference, human insulin, gene therapy, transgenic animals).
10. **Unit 10: Ecology and Environment** (Organisms and populations: population growth models, species interactions; Ecosystem: structure, productivity, decomposition, energy flow, ecological pyramids; Biodiversity and conservation: patterns of biodiversity, red data book, in-situ and ex-situ conservation).

Zero-Tolerance NMC Exclusions: Topics Strictly Deleted from Biology

Under revised NMC regulations, the following topics and chapters have been **completely eliminated**:

- **Digestion and Absorption** (Entire human digestion chapter eliminated).
- **Transport in Plants** (Xylem/phloem transport mechanisms, water potential, transpiration pull eliminated from Plant Physiology).
- **Mineral Nutrition** (Macro/micronutrients, nitrogen metabolism, deficiency symptoms eliminated).
- **Strategies for Enhancement in Food Production** (Animal husbandry and plant breeding eliminated).
- **Environmental Issues** (Pollution, solid waste, global warming eliminated from Ecology).
- **Anatomy of Flowering Plants (legacy sections)** (Secondary growth, anomalous structures eliminated).

Important Additions to Note:

- **Prescribed 7 Plant Families:** Malvaceae, Cruciferae/Brassicaceae, Fabaceae/Leguminosae, Asteraceae/Compositae, Poaceae/Gramineae, Solanaceae, and Liliaceae must be studied thoroughly.
- **Frog & Cockroach Morphology:** Both are explicitly in the syllabus.

Empirical Question Weightage Matrix

Biology Domain	Typical Questions	Marks Weightage
Genetics & Molecular Basis of Inheritance (Units 7A, 7B)	15–18	60–72 Marks
Human Physiology (Respiration, Circulation, Excretion, Locomotion, Neuro-Endocrine)	12–15	48–60 Marks
Plant Physiology (Photosynthesis, Respiration, Plant Growth Regulators)	6–8	24–32 Marks
Cell Biology & Cell Division (Cell Structure, Biomolecules, Mitosis, Meiosis)	7–9	28–36 Marks
Reproduction & Biotechnology (Plant/Human Reproduction, Health, rDNA, Biotech)	14–16	56–64 Marks
Ecology & Environment (Populations, Ecosystems, Biodiversity)	8–10	32–40 Marks
Diversity in Living World & Systematic Plant Families (Taxonomy, 7 Families, Frog/Cockroach)	10–12	40–48 Marks
Immunology, Human Health & Evolutionary Biology	6–8	24–32 Marks
Total Biology Paper	**90 Attempted / 100**	**360 Marks**

Table 2: Empirical Question Distribution in Recent NEET UG Papers.

Contents

Acknowledgements ix

Author's Preface: Straight Talk on NEET Biology xi

Syllabus Blueprint & NMC Rationalization Guide xiii

I Botany: Plant Biology, Genetics, Ecology & Angiosperm Families 1

1 Plant Physiology: Photosynthesis, Cellular Respiration & Phytohormones 3

1.1 Photosynthesis in Higher Plants . . . 3

1.2 Respiration in Plants . . . 5

1.3 Plant Growth Regulators (Phytohormones) . . . 6

2 Genetics & Molecular Basis of Inheritance 9

2.1 Principles of Inheritance & Variation (Mendelian Genetics) . . . 9

2.2 Molecular Basis of Inheritance (DNA, Replication, Transcription & Translation) . . . 11

3 Sexual Reproduction & Applied Biotechnology 17

3.1 Sexual Reproduction in Flowering Plants (Angiosperms) . . . 17

3.2 Human Reproduction & Reproductive Health . . . 18

3.3 Biotechnology: Principles, Tools & Modern Applications . . . 20

4 Ecology, Environmental Biology & Systematic Plant Families 23

4.1 Organisms & Populations . . . 23

4.2 Ecosystem: Structure, Function & Ecological Succession . . . 24

4.3 Biodiversity & Conservation . . . 25

4.4 Systematic Morphology of Plant Families (Prescribed NMC Syllabus) . . . 26

II Zoology: Cellular Life, Animal Diversity, Human Physiology & Health 27

5 Cell Biology, Cell Division & Animal Morphology 29

5.1 Cell Structure, Function & Biomolecules . . . 29

5.2 Cell Cycle & Cell Division: Mitosis & Meiosis . . . 31

5.3 Animal Kingdom: Non-Chordates & Chordates Blueprint . . . 31

5.4 Structural Organisation in Animals: Frog & Cockroach Anatomy . . . 32

6 Human Physiology: Integrated Organ Systems **35**
6.1 Breathing & Exchange of Gases . . . 35
6.2 Body Fluids & Circulation . . . 36
6.3 Excretory Products & Their Elimination . . . 37
6.4 Locomotion & Movement: Musculoskeletal Mechanics . . . 38
6.5 Neural Control & Chemical Coordination . . . 39

7 Immunology, Human Pathogens, Disease & Evolutionary Biology **41**
7.1 Principles of Immunology & Host Defense Mechanisms . . . 41
7.2 Major Human Pathogens, Transmission & Diagnostic Tests . . . 43
7.3 Evolutionary Biology: Origins, Evidence & Mechanisms . . . 44

III High-Yield Quantitative Biology & Diagnostic Matrices **47**

8 Quantitative Biology, Genetic Formulations & High-Yield Diagnostic Matrices **49**
8.1 Quantitative Genetics & Probability Calculus . . . 49
8.2 Recombination Frequency & Chromosome Mapping Calculus . . . 50
8.3 Population Genetics: Hardy-Weinberg Mathematical Model . . . 51
8.4 Quantitative Human Physiology Matrices . . . 51
8.5 Master Systematic Comparison of All 7 Prescribed Plant Families . . . 52

About the Author & Errata Submissions **53**

PART I

Botany: Plant Biology, Genetics, Ecology & Angiosperm Families

Chapter 1

Plant Physiology: Photosynthesis, Cellular Respiration & Phytohormones

Plant Physiology forms the biochemical core of botany. In this chapter, we explore chloroplast thylakoid architectures, the Z-scheme of light reactions, cyclic and non-cyclic photophosphorylation, the Calvin C_3 cycle, the C_4 dicarboxylic acid pathway with Kranz anatomy, photorespiration, mitochondrial cellular respiration (glycolysis, link reaction, Krebs cycle, and the electron transport system), and the physiological roles of the five major phytohormone classes.

A Personal Note from Satya: Active Recall vs. The Highlighting Trap

Before you turn another page, let's talk about the biggest mistake students make in Biology.

You open NCERT. You pick up a neon yellow highlighter. You read a paragraph about the Hatch-Slack C_4 pathway, highlight three sentences, and move on. By the end of the chapter, your textbook looks like a coloring book. Your brain feels satisfied and says: *"I know this!"*

That is the **illusion of competence**. What happened in your brain was simple *recognition*, not *retrieval*. When you see a familiar sentence on a printed page, your brain recognizes the words. But on exam day, NTA gives you an empty bubble sheet and a tricky question asking: *"In C_4 plants, the primary carboxylation reaction takes place in the mesophyll cytoplasm, catalyzed by which enzyme?"* Suddenly, your mind draws a blank.

How to Build Real 360/360 Retrieval Strength:

1. **The Blank Sheet Test:** Close this book. Take a plain white piece of paper. Draw the Z-scheme of light reactions, or write down the floral formula of Solanaceae entirely from memory. The exact moment your hand freezes is your real preparation level. That is the blind spot you must fix today.
2. **Differentiating Similar Concepts:** Never just read facts in isolation. Group them: Compare C_3 vs C_4 directly; compare Gramineae vs Malvaceae floral parts side-by-side.

Active recall feels uncomfortable. It makes your head hurt a little. But that slight discomfort is literally the physical sensation of memory circuits forming. Test yourself relentlessly, write things out by hand, and you will walk into the biology section with absolute confidence.

1.1 Photosynthesis in Higher Plants

Chloroplast Architecture & Photosynthetic Pigments

Photosynthesis occurs in chloroplasts of green mesophyll cells:

- **Membrane System (Grana & Stroma Lamellae):** Responsible for trapping light energy and synthesizing ATP and NADPH (**Light Reactions / Photochemical Phase**).
- **Fluid Stroma:** Contains soluble enzymes for enzymatic reduction of CO_2 into carbohydrates (**Dark Reactions / Carbon Reactions / Biosynthetic Phase**).
- **Pigment Separation (Paper Chromatography):**
  - **Chlorophyll *a*:** Bright / blue-green; primary reaction center pigment.
  - **Chlorophyll *b*:** Yellow-green; accessory pigment.
  - **Xanthophylls:** Yellow; accessory pigment.
  - **Carotenoids:** Yellow to yellow-orange; accessory pigments; protect chlorophyll *a* from **photo-**

oxidation.

- **Absorption vs Action Spectrum:** Maximum absorption by chlorophyll *a* occurs in the **blue** (400 − 450 nm) and **red** (650 − 700 nm) regions, showing maximal rate of photosynthesis (first demonstrated by T.W. Engelmann using *Cladophora* and aerobic bacteria).

Photochemical Phase: Z-Scheme & Photophosphorylation

Light-harvesting complexes (LHC) comprise hundreds of pigment molecules bound to proteins:

Photosystem I (PS I): Reaction center chlorophyll *a* has absorption peak at **700** nm (P_{700}).

Photosystem II (PS II): Reaction center chlorophyll *a* has absorption peak at **680** nm (P_{680}).

Photolysis of Water: $2H_2O \xrightarrow[Mn^{2+},Cl^-]{OEC} 4H^+ + O_2 \uparrow + 4e^-$

(Associated with PS II on inner thylakoid membrane).

Comparison of Non-Cyclic vs Cyclic Photophosphorylation:

Parameter	Non-Cyclic Photophosphorylation (Z-Scheme)	Cyclic Photophosphorylation
Photosystems	Both **PS II and PS I** operate in series	**Only PS I** is active
Electron Pathway	Non-cyclic (electrons from water reduce $NADP^+$)	Cyclic (electrons cycle back to P_{700})
End Products	**ATP, NADPH and** O_2 evolved	**ATP alone** (no NADPH, no O_2 evolution)
Site in Chloroplast	Grana thylakoids (appressed membrane)	**Stroma lamellae** (lack PS II and NADP reductase)
Wavelength Needed	Normal sunlight ($\lambda \leq 680$ nm)	Active when light of $\lambda > 680$ nm is available

Chemiosmotic Hypothesis (Peter Mitchell)

ATP synthesis in chloroplasts is coupled to the generation of a **proton gradient across the thylakoid membrane**. Protons accumulate in high concentration inside the **thylakoid lumen**, dropping lumenal pH:

1. **Splitting of water** on inner side of membrane releases H^+ into the lumen.
2. **Plastoquinone (PQ)** acts as a mobile hydrogen carrier; removes protons from stroma and dumps them into lumen.
3. **NADP reductase enzyme** on stroma side removes protons from stroma to reduce $NADP^+$ to $NADPH + H^+$.

Proton Motive Force: Protons break the gradient by passing through the transmembrane channel of F_0 **of ATP Synthase** into stroma $\implies$ drives conformational changes in F_1 **headpiece** to catalyze $ADP + P_i \rightarrow ATP$.

Biosynthetic Phase: C_3 Calvin Cycle Energetics

Discovered by Melvin Calvin using radioactive ^{14}C in algal photosynthesis (*Chlorella*).

Phase 1: Carboxylation: $\text{RuBP (5C)} + CO_2 \xrightarrow{\textbf{RuBisCO}} 2 \times \text{3-PGA (3C)}$

Phase 2: Reduction: $2 \times \text{3-PGA} + 2\,\text{ATP} + 2\,\text{NADPH} \rightarrow 2\,\text{Triose-P} + 2\,\text{ADP} + 2\,NADP^+$

Phase 3: Regeneration: $\text{Triose-P} + 1\,\text{ATP} \longrightarrow \text{RuBP} + 1\,\text{ADP} + P_i$

$$\textbf{Per Single } CO_2 \textbf{ Fixed:} \quad 3\,\text{ATP} + 2\,\text{NADPH} \tag{1.1}$$

$$\textbf{Per Glucose (6 } CO_2\textbf{):} \quad 6\,CO_2 + \textbf{18}\,\text{ATP} + \textbf{12}\,\text{NADPH} \longrightarrow C_6H_{12}O_6 + 18\,\text{ADP} + 12\,NADP^+ \tag{1.2}$$

Hatch-Slack Pathway (C_4 Plants) & Photorespiration (C_2 Cycle)

1. Kranz Anatomy of C_4 Leaves (Maize, Sugarcane, Sorghum):

- Bundle sheath cells surround vascular bundles in large layers; possess thick walls impermeable to gas exchange, no intercellular spaces, and large agranal chloroplasts with abundant **RuBisCO** (lack PEPcase).
- Mesophyll cells have granal chloroplasts containing **PEPcase** (lack RuBisCO).
- Primary CO_2 acceptor is **Phosphoenolpyruvate (PEP, 3C)**, forming **Oxaloacetic acid (OAA, 4C)**.
- Malic acid / Aspartic acid transported to bundle sheath cells $\rightarrow$ decarboxylated to release CO_2 and pyruvate (3C).
- **Advantage:** RuBisCO operates at saturated CO_2 concentration, completely **preventing photorespiration**!
- **Energetic Cost of C_4 Pathway:** Requires 30 ATP + 12 NADPH per glucose molecule (extra 12 ATP to regenerate PEP).

2. Photorespiration (C_2 Cycle / Glycolate Pathway): Occurs under high O_2, low CO_2, and high temperature when RuBisCO acts as an **oxygenase**:

$$\text{RuBP (5C)} + O_2 \xrightarrow{\text{RuBisCO}} \text{Phosphoglycerate (3C)} + \textbf{Phosphoglycolate (2C)} \tag{1.3}$$

Operates sequentially in three organelles: **Chloroplast $\rightarrow$ Peroxisome $\rightarrow$ Mitochondria**. A wasteful process that yields **neither ATP nor sugar**, but consumes ATP and releases CO_2!

NEET Trap: Blackman's Law & Factors Affecting Photosynthesis

- **Blackman's Law of Limiting Factors (1905):** If a chemical process is affected by more than one factor, its rate is determined by the factor which is nearest to its minimal value.
- **CO_2 Saturation Levels:** C_4 plants saturate at 360 μL/L; C_3 plants saturate only beyond 450 μL/L. Because current atmospheric CO_2 is $\approx$ 400 ppm, CO_2 is the major limiting factor for C_3 plants (greenhouse crops like tomatoes and bell peppers show enhanced productivity in CO_2-enriched atmospheres).
- **Light Saturation:** Occurs at **10% of full sunlight**. Except for plants in shade or dense forests, light is rarely a limiting factor in nature.

1.2 Respiration in Plants

Glycolysis (EMP Pathway) & Link Reaction

Discovered by Gustav Embden, Otto Meyerhof, and J. Parnas. Occurs in the **cytoplasm** of all living cells; universal to both aerobic and anaerobic organisms.

Phase 1: Preparatory: $\text{Glucose} \xrightarrow{1\,\text{ATP}} \text{G-6-P} \rightarrow \text{F-6-P} \xrightarrow{1\,\text{ATP}} \text{F-1,6-BP}$ (Invests 2 ATP)

Phase 2: Cleavage: $\text{F-1,6-BP (6C)} \xrightarrow{\text{Aldolase}} \text{PGAL (3C)} \underset{}{\overset{\text{TPI}}{\rightleftharpoons}} \text{DHAP (3C)}$

Phase 3: Payoff Phase (x2):

$$2\,\text{PGAL} + 2\,NAD^+ \rightarrow 2 \times 1,3\text{-BPGA} + \mathbf{2}\,\text{NADH} + \mathbf{2}\,H^+$$

$$2 \times 1,3\text{-BPGA} + 2\,\text{ADP} \rightarrow 2 \times \text{3-PGA} + \mathbf{2}\,\text{ATP (SLP)}$$

$$2\,\text{PEP} + 2\,\text{ADP} \xrightarrow{\text{PK}} 2\,\text{Pyruvate} + \mathbf{2}\,\text{ATP (SLP)}$$

Net Glycolysis Yield: $\mathbf{2}\,\text{Pyruvate} + \mathbf{2}\,\text{ATP} + \mathbf{2}\,\text{NADH} + \mathbf{2}\,H^+$ (No CO_2 or O_2) (1.4)

The Link Reaction (Oxidative Decarboxylation of Pyruvate): Occurs in the **mitochondrial matrix**, catalyzed by *pyruvate dehydrogenase* multienzyme complex (requires Mg^{2+}, TPP, Lipoamide, NAD^+, CoA):

$$2\,\text{Pyruvate (3C)} + 2\,\text{CoA} + 2\,NAD^+ \longrightarrow \mathbf{2}\,\text{Acetyl-CoA (2C)} + \mathbf{2}\,CO_2 \uparrow + \mathbf{2}\,\text{NADH} + \mathbf{2}\,H^+ \tag{1.5}$$

Krebs Cycle (TCA Cycle) & Electron Transport System (ETS)

1. Tricarboxylic Acid (TCA) Cycle (Hans Krebs, 1937): Occurs in the **mitochondrial matrix**:

Per Acetyl-CoA (2C): $\text{Acetyl-CoA} + \text{OAA (4C)} + \text{H}_2\text{O} \xrightarrow{\text{Citrate Synthase}} \text{Citrate (6C)} + \text{CoA}$

Decarboxylations: $\text{Isocitrate} \rightarrow \alpha\text{-ketoglutarate (5C)} + \text{CO}_2 + \text{NADH}$

$\alpha\text{-ketoglutarate} \rightarrow \text{Succinyl-CoA (4C)} + \text{CO}_2 + \text{NADH}$

Substrate-Level: $\text{Succinyl-CoA} + \text{GDP} + \text{P}_\text{i} \rightarrow \text{Succinate} + \mathbf{1}\,\text{GTP/ATP} + \text{CoA}$

Oxidations: $\text{Succinate} \rightarrow \text{Fumarate} + \mathbf{1}\,\text{FADH}_2$ (Succinate dehydrogenase in inner membrane)

$\text{Malate} + \text{NAD}^+ \rightarrow \text{OAA} + \mathbf{1}\,\text{NADH} + \text{H}^+$

Net Yield of TCA Cycle (2 Acetyl-CoA per Glucose): $\mathbf{4}\,\text{CO}_2 + \mathbf{6}\,\text{NADH} + \mathbf{2}\,\text{FADH}_2 + \mathbf{2}\,\text{ATP/GTP}$ (1.6)

2. Inner Mitochondrial Membrane ETS Complexes:

- **Complex I: NADH dehydrogenase** (contains FMN and Fe-S centers); transfers electrons from NADH to Ubiquinone (CoQ).
- **Complex II: Succinate dehydrogenase** (contains FAD and Fe-S centers); transfers electrons from succinate to Ubiquinone.
- **Complex III: Cytochrome bc_1 complex** (Cytochrome b, Fe-S, Cytochrome c_1).
- **Mobile Electron Carriers: Ubiquinone** (in lipid bilayer) and **Cytochrome** c (small soluble protein attached to outer surface of inner membrane).
- **Complex IV: Cytochrome** c **oxidase** (contains Cytochrome a, a_3 and two copper centers, Cu_A and Cu_B); reduces terminal electron acceptor O_2:

$$\frac{1}{2}\text{O}_2 + 2\text{H}^+ + 2e^- \longrightarrow \text{H}_2\text{O} \quad \text{(Metabolic water)} \tag{1.7}$$

- **Complex V: ATP Synthase ($F_0 - F_1$ complex).** Pass of 2H^+ through F_0 produces 1 ATP.

Total Yield per Glucose: **36** or **38** ATP (or $30 - 32$ ATP by modern P/O) (1.8)

Respiratory Quotient (RQ) Formulations

$$\text{RQ} = \frac{\text{Volume of } \text{CO}_2 \text{ evolved}}{\text{Volume of } \text{O}_2 \text{ consumed}} \tag{1.9}$$

Carbohydrates (Glucose): $\text{C}_6\text{H}_{12}\text{O}_6 + 6\text{O}_2 \rightarrow 6\text{CO}_2 + 6\text{H}_2\text{O} \implies \text{RQ} = \mathbf{1.0}$

Fats (Tripalmitin): $2\text{C}_{51}\text{H}_{98}\text{O}_6 + 145\text{O}_2 \rightarrow 102\text{CO}_2 + 98\text{H}_2\text{O} \implies \text{RQ} = \dfrac{\mathbf{102}}{\mathbf{145}} = \mathbf{0.7}$

Proteins: $\text{RQ} \approx \mathbf{0.9}$

Organic Acids (Malic Acid): $\text{C}_4\text{H}_6\text{O}_5 + 3\text{O}_2 \rightarrow 4\text{CO}_2 + 3\text{H}_2\text{O} \implies \text{RQ} = \dfrac{\mathbf{4}}{\mathbf{3}} = \mathbf{1.33} > \mathbf{1}$

Anaerobic Respiration: $\text{C}_6\text{H}_{12}\text{O}_6 \rightarrow 2\text{C}_2\text{H}_5\text{OH} + 2\text{CO}_2 \implies \text{RQ} = \dfrac{\mathbf{2}}{\mathbf{0}} = \infty$

1.3 Plant Growth Regulators (Phytohormones)

Five Classic Phytohormone Classes: Functions & Applications

Hormone Class	Discovery / Chemistry	Key Physiological Functions & Agricultural Applications
Auxins	Charles & Francis Darwin (*Phalaris* coleoptile); isolated by F.W. Went from *Avena* coleoptile tip. Natural: IAA, IBA; Synthetic: NAA, 2,4-D.	• Promotes **apical dominance** (growing apical bud inhibits lateral buds; removal / decapitation induces bushy hedge). • Initiates root formation in stem cuttings. • Promotes flowering in pineapples. Prevents premature fruit/leaf drop, promotes abscission of older leaves. • **2,4-D** is widely used as a selective weedicide / herbicide to eradicate dicotyledonous weeds without affecting monocot plants. • Induces parthenocarpy in tomatoes.
Gibberellins	E. Kurosawa discovered in *Gibberella fujikuroi* (foolish seedling / *bakanae* disease of rice). $>$ 100 known (GA_1, GA_2, GA_3 most studied). Terpene nature.	• Causes elongation of internodes; increases length of grape stalks and elongates apples. • **Bolting** (internode elongation prior to flowering) in rosette plants like cabbage and beet. • Increases sugarcane stem length, boosting yield by up to 20 tonnes per acre. • Accelerates malting process in brewing industry.
Cytokinins	F. Skoog discovered kinetin from autoclaved herring sperm DNA; Letham isolated **Zeatin** from corn kernels. Aminopurine derivative.	• Promotes rapid cell division; high cytokinin-to-auxin ratio promotes shoot differentiation. • Overcomes apical dominance; promotes nutrient mobilization and delays leaf senescence (**Richmond-Lang effect**). • Promotes adventitious shoot formation and lateral shoot growth.
Ethylene	H.H. Cousins established volatile substance from ripened oranges ripened bananas. Only **gaseous hormone**.	• Accelerates fruit ripening accompanied by a sudden rise in respiration rate (**Respiratory Climacteric**). • Source compound: **Ethephon** (releases ethylene slowly). • Promotes senescence and abscission of plant organs (leaves, flowers). • Promotes root growth and root hair formation, increasing absorption surface area. • Breaks seed and bud dormancy (initiates germination in peanut seeds, sprouting of potato tubers). • Promotes female flowers in cucumbers to increase yield.
Abscisic Acid (ABA)	Discovered as Inhibitor-β, Abscission II, and Dormin. Sesquiterpene derivative; acts as general growth inhibitor.	• Known as the **Stress Hormone**: stimulates rapid closure of stomata during water stress / drought. • Induces and maintains **seed dormancy**, preventing vivipary and premature sprouting. • Acts as a direct antagonist to gibberellins (anti-GA).

Photoperiodism & Vernalization Highlights

- **Photoperiodism (Garner & Allard):** Response of plants to periods of day/night.
  - **Short-Day Plants (SDP):** Flower when day length is less than critical photoperiod (e.g., Tobacco, Soyabean, Chrysanthemum).
  - **Long-Day Plants (LDP):** Flower when day length exceeds critical photoperiod (e.g., Wheat, Radish, Spinach).
  - **Day-Neutral Plants (DNP):** No correlation between light exposure and flowering (e.g., Tomato, Sunflower, Pea).
  - **Site of Perception:** The site of perception of light/dark duration is the **leaves**. The hypothetical hormonal substance responsible for flowering is **Florigen** (migrates from leaves to shoot apices).
- **Vernalization:** Quantitative or qualitative dependence of flowering on low-temperature exposure ($1 - 6^{\circ}$C); prevents precocious reproductive development late in the growing season (e.g., winter varieties of wheat, barley, rye; biennials like sugarbeet, cabbage, carrot). Perceived by the **shoot apical meristem**.

Chapter 2

Genetics & Molecular Basis of Inheritance

Genetics is the highest-weightage domain in the entire NEET examination, typically generating between 15 and 18 questions. This chapter covers classical Mendelian principles of segregation and independent assortment, incomplete dominance, codominance, chromosomal theories of linkage, pedigree analysis, DNA structure, semi-conservative replication, prokaryotic and eukaryotic transcription, the universal genetic code, translation, the lac operon, and DNA fingerprinting.

2.1 Principles of Inheritance & Variation (Mendelian Genetics)

Mendel's Experimental Architecture & Contrasting Traits

Gregor Johann Mendel conducted hybridization experiments on garden peas (*Pisum sativum*) for 7 years (1856–1863). He investigated **7 pairs of contrasting traits** located across 4 chromosomes (Chromosomes 1, 4, 5, 7):

Character	Dominant Trait	Recessive Trait	Chromosome
Stem Height	Tall (T)	Dwarf (t)	4
Flower Color	Violet / Purple (W)	White (w)	1
Flower Position	Axial (A)	Terminal (a)	4
Pod Shape	Inflated / Full (I)	Constricted (i)	4
Pod Color	Green (G)	Yellow (g)	5
Seed Shape	Round (R)	Wrinkled (r)	7
Seed Color	Yellow (Y)	Green (y)	1

Mendelian Ratios & Mathematical Formulations

For a cross involving heterozygous parents ($Aa \times Aa$ for monohybrid, $AaBb \times AaBb$ for dihybrid):

Monohybrid Phenotypic Ratio: $3:1$ (Genotypic: $1:2:1 = 1\,AA:2\,Aa:1\,aa$)

Dihybrid Phenotypic Ratio: $9:3:3:1$ (Genotypic: $1:2:1:2:4:2:1:2:1$)

Trihybrid Phenotypic Ratio: $(3+1)^3 = 27:9:9:9:3:3:3:1$

Number of Phenotypes: 2^n (n = number of heterozygous gene pairs)

Number of Genotypes: 3^n

Number of Gamete Types: 2^n

Number of Zygotes / Offspring: 4^n

Test Cross Ratio: Heterozygote crossed with homozygous recessive:

- Monohybrid Test Cross ($Aa \times aa$): Ratio $= 1:1$.
- Dihybrid Test Cross ($AaBb \times aabb$): Ratio $= 1:1:1:1$.

Deviations from Pure Mendelian Inheritance

1. **Incomplete Dominance:** Neither allele is dominant; the heterozygote displays an intermediate phenotype.
   - *Examples:* Flower color in *Mirabilis jalapa* (4 o'clock plant) and *Antirrhinum majus* (Snap-

dragon / Dog flower).
- Crossing Red (RR) × White (rr) gives Pink (Rr) in F_1.
- F_2 Phenotypic ratio = 1 Red : 2 Pink : 1 White (1 : 2 : 1).
- **Crucial Rule: Phenotypic ratio equals Genotypic ratio** (1 : 2 : 1).

2. **Codominance:** Both alleles express themselves fully and simultaneously in the heterozygote without blending.
   - *Examples:* ABO Blood grouping in humans, Sickle cell trait (Hb^AHb^S), MN blood group.
3. **Multiple Allelism:** More than two alleles govern a single gene locus in a population (can only be studied in population studies, not in an individual).
   - *ABO Blood Group Gene (I):* Located on Chromosome 9; 3 alleles (I^A, I^B, i).
   - Number of possible genotypes for n multiple alleles:

$$\text{Total Genotypes} = \frac{n(n+1)}{2} = \frac{3(3+1)}{2} = 6 \text{ genotypes} \implies 4 \text{ phenotypes (A, B, AB, O)} \tag{2.1}$$

4. **Pleiotropy:** A single gene influences multiple unrelated phenotypic traits.
   - *Phenylketonuria (PKU):* Mutation in gene coding for enzyme *phenylalanine hydroxylase* causes mental retardation, reduction in hair, and skin pigmentation.
   - *Starch Synthesis in Pea Seeds:* Gene B controls starch grain size (incomplete dominance: BB large, Bb intermediate, bb small) and seed shape (complete dominance: BB, Bb round, bb wrinkled).
5. **Polygenic Inheritance:** Quantitative inheritance where multiple additive genes control a continuous trait, influenced by the environment (e.g., Human skin color, human height, kernel color in wheat).
   - Governed by 3 genes (A, B, C): $AABBCC$ (darkest) to $aabbcc$ (lightest).
   - F_2 phenotypic distribution follows a bell-shaped Gaussian curve: 1 : 6 : 15 : 20 : 15 : 6 : 1.

Chromosomal Theory, Linkage & Genetic Mapping

1. Chromosomal Theory of Inheritance (Sutton & Boveri, 1902): Behavior of chromosomes parallels the behavior of genes during meiosis. Chromosomes segregate during anaphase I and assort independently during anaphase I/metaphase I.

2. Morgan's Linkage Experiments (Thomas Hunt Morgan, 1910): Used the fruit fly *Drosophila melanogaster* because:
- Can be grown on simple synthetic medium in the laboratory.
- Short life cycle (about 2 weeks / 14 days).
- A single mating produces a large number of progeny flies.
- Clear morphological dimorphism (female is distinctly larger than male).
- Low chromosome number ($2n = 8$) easily observable under low power microscope.

Morgan's Crosses on X-Chromosome:
- **Cross A (Tightly Linked):** Yellow body (y), white eye (w) crossed with wild brown body (y^+), red eye (w^+). Recombination frequency = **1.3%** (Parental types = **98.7%**).
- **Cross B (Loosely Linked):** White eye (w), miniature wing (m) crossed with wild red eye (w^+), normal wing (m^+). Recombination frequency = **37.2%** (Parental types = **62.8%**).

3. Chromosome Mapping (Alfred Sturtevant): Recombination frequency is directly proportional to the physical distance between genes on a chromosome:

$$\text{Distance between genes (in centimorgans / map units)} = \frac{\text{Total Recombinants}}{\text{Total Progeny}} \times 100\% \tag{2.2}$$

$$1 \text{ Map Unit (cM)} = 1\% \text{ Recombination frequency} \tag{2.3}$$

Boundary Limit: Maximum observable recombination frequency for two linked genes cannot exceed **50%** (at 50%, genes behave as if assorting independently).

Sex Determination Mechanisms in Animals & Humans

- **XX–XY Type (Male Heterogamety):** Humans and *Drosophila*. Male produces two types of sperms: 50% with X, 50% with Y. Female is homogametic (XX).
- **XX–XO Type (Male Heterogamety):** Grasshopper and certain insects. Male has only one X-chromosome ($2A + XO$); female has two ($2A + XX$).
- **ZZ–ZW Type (Female Heterogamety):** Birds, fowls, butterflies. Female produces two types of eggs: 50% with Z, 50% with W ($2A + ZW$). Male is homogametic ($2A + ZZ$).
- **Haplodiploidy (Honeybee):** Female (Queen/Worker) develops from fertilized diploid egg ($2n = 32$) via sexual reproduction. Male (Drone) develops from unfertilized haploid egg ($n = 16$) by **parthenogenesis**. Drones have no father and cannot have sons, but have a grandfather and can have grandsons!

NEET Trap: Genetic Disorders & Pedigree Distinctions

1. **Mendelian Disorders (Single-Gene Mutations):**

- **Autosomal Dominant:** *Myotonic dystrophy*, Huntington's chorea. Trait does not skip generations; affected child always has at least one affected parent.
- **Autosomal Recessive:** *Sickle-cell Anemia, Phenylketonuria, Thalassemia, Cystic Fibrosis.* Trait frequently skips generations; parents of affected individuals are usually unaffected heterozygous carriers (Aa). Both sexes equally affected.
  - **Sickle-cell Anemia:** Point mutation in β-globin gene on chromosome 11. GAG codon mutates to GUG $\implies$ substitution of Glutamic acid (Glu) by **Valine (Val) at position 6** of the β-chain. Under low O_2 tension, Hb^S polymerizes into elongated sickle shapes.
  - **Thalassemia:** Quantitative disorder (reduced rate of globin chain synthesis), whereas Sickle-cell anemia is a qualitative disorder (defective globin produced). α-Thalassemia is controlled by two closely linked genes $HBA1$ and $HBA2$ on chromosome 16. β-Thalassemia is controlled by a single gene HBB on chromosome 11.
- **Sex-Linked (X-Linked) Recessive:** *Hemophilia, Color Blindness.*
  - Show **Criss-cross inheritance** (passed from carrier mother to son, or father to daughter who becomes a carrier).
  - Far more prevalent in males (X^cY) than females (X^cX^c, requiring carrier mother and affected father).
  - **Hemophilia A:** Deficiency of blood clotting Factor VIII; **Hemophilia B:** Factor IX (Christmas disease).

2. **Chromosomal Disorders (Aneuploidy & Polyploidy):**

- **Down's Syndrome (Trisomy 21):** First described by Langdon Down (1866). Karyotype = $47, XX + 21$ or $47, XY + 21$. Short stature, small round head, furrowed tongue, partially open mouth, broad palm with characteristic palm crease, physical and mental retardation.
- **Klinefelter's Syndrome (Trisomy of Sex Chromosome):** Karyotype = $47, XXY$. Overall masculine development with feminine features (e.g., development of breasts / **Gynaecomastia**), sterile male.
- **Turner's Syndrome (Monosomy of Sex Chromosome):** Karyotype = $45, XO$. Sterile female, rudimentary ovaries, lack of secondary sexual characters, short stature, webbed neck.

2.2 Molecular Basis of Inheritance (DNA, Replication, Transcription & Translation)

Watson-Crick B-DNA Structural Dimensions

Proposed by James Watson and Francis Crick in 1953, based on X-ray diffraction data of Rosalind Franklin and Maurice Wilkins.

Helix Type: Right-handed double helix with antiparallel strands ($5' \to 3'$ and $3' \to 5'$)

Helix Pitch (P) :$3.4\,\text{nm} = 34\,\text{Å}$ (one complete turn of helix)

Base Pairs per Turn: $10\,\text{bp}$

Distance between bp: $0.34\,\text{nm} = 3.4\,\text{Å} = 0.34 \times 10^{-9}\,\text{m}$

Helix Diameter: $2.0\,\text{nm} = 20\,\text{Å}$

Hydrogen Bonds: A = T (2 H-bonds), G ≡ C (3 H-bonds)

Human DNA Length: $6.6 \times 10^{9}\,\text{bp} \times 0.34 \times 10^{-9}\,\text{m} \approx \mathbf{2.2}\,\text{m}$

Packaging of DNA: Nucleosome Architecture

Because a 2.2 m long negatively charged DNA must fit inside a microscopic nucleus ($\sim 10^{-6}\,\text{m}$), it is tightly packaged around basic proteins called **histones**:

- **Histones:** Rich in basic amino acids **Lysine** and **Arginine** (both carry positive charges in their side chains).
- **Histone Octamer:** Core consists of two molecules each of H_2A, H_2B, H_3, and H_4.
- **Nucleosome Core:** Negatively charged DNA is wrapped around the octamer:

$$\text{Length of DNA wrapped around single nucleosome} \approx \mathbf{200}\,\text{bp} \tag{2.4}$$

- **Linker Histone (H_1):** Seals the DNA at the entry and exit points of the nucleosome core.
- **Chromatin States:**
  - **Euchromatin:** Loosely packed, stains light, transcriptionally active.
  - **Heterochromatin:** Densely packed, stains dark, transcriptionally inactive.

Historic Experiments Proving DNA is the Genetic Material

1. **Griffith's Transforming Principle (Frederick Griffith, 1928):** Used *Streptococcus pneumoniae* (Pneumococcus).
   - Smooth (S) strain has mucous (polysaccharide) coat $\implies$ Virulent (mice die of pneumonia).
   - Rough (R) strain lacks capsule $\implies$ Non-virulent (mice survive).
   - Heat-killed S strain → Mice survive.
   - Heat-killed S strain + Living R strain → **Mice die**, and living S strain bacteria were recovered from blood!
   - *Conclusion:* R strain was transformed by a heat-stable transforming principle from heat-killed S.
2. **Biochemical Characterization (Avery, MacLeod & McCarty, 1944):** Purified biochemicals (proteins, RNA, DNA) from heat-killed S cells.
   - Proteases and RNases did **not** inhibit transformation.
   - Digestion with **Deoxyribonuclease (DNase) completely inhibited transformation**.
   - *Conclusion:* DNA is the substance causing transformation!
3. **Unequivocal Proof: Hershey-Chase Experiment (1952):** Used Bacteriophage T2 and *Escherichia coli*.
   - Phages grown on radioactive sulfur (^{35}S) labeled protein coat (cysteine and methionine contain S, DNA has no S).
   - Phages grown on radioactive phosphorus (^{32}P) labeled DNA core (DNA has P, proteins have no P).
   - Steps: (i) Infection → (ii) Blending (agitation in kitchen blender detached coats) → (iii) Centrifugation.
   - Bacteria infected with ^{32}P phages were radioactive (pellet had radioactivity). Supernatant

with ^{35}S had viral ghosts.
- *Conclusion:* DNA is the genetic material passed from virus to bacteria!

DNA Replication Machinery & Enzymology

Replication is **semiconservative** and **bidirectional**, originating at specific sites called *Ori* (Origin of Replication).

Meselson & Stahl: Grew *E. coli* in $^{15}NH_4Cl$ → all heavy ($^{15}N - ^{15}N$).
In $^{14}NH_4Cl$: Gen 1 (20 min): 100% Hybrid ($^{15}N - ^{14}N$).
Gen 2 (40 min): 50% Hybrid ($^{15}N - ^{14}N$) + 50% Light ($^{14}N - ^{14}N$).
Taylor: Tritiated thymidine in *Vicia faba* root cells.

Core Enzymes and Functional Roles:

Enzyme	Specific Biological Action
Helicase	Unwinds the parental DNA double helix by breaking hydrogen bonds, forming replication fork.
SSB Proteins	Single-Strand Binding proteins stabilize single strands and prevent premature re-annealing.
Topoisomerase / Gyrase	Relieves positive supercoiling strain ahead of the progressing replication fork by cutting and resealing.
RNA Primase	Synthesizes short complementary RNA primers (5 → 10 nucleotides) providing free 3′-OH for DNA polymerase.
DNA Polymerase III	Main replicating enzyme in prokaryotes; synthesizes in strictly 5′ → 3′ **direction**; high catalytic speed (≈ 2000 bp/sec) with proofreading (3′ → 5′ exonuclease activity).
DNA Polymerase I	Removes RNA primers (via 5′ → 3′ exonuclease activity) and fills gaps with deoxynucleotides (Kornberg enzyme).
DNA Ligase	Catalyzes formation of phosphodiester bonds between discontinuous Okazaki fragments on the lagging strand.

Transcription: Prokaryotes vs Eukaryotes

A transcription unit comprises: (1) **Promoter** (located towards 5′ end of coding strand / upstream), (2) **Structural gene**, (3) **Terminator** (located towards 3′ end of coding strand / downstream).

$$\text{Template Strand: } 3' \longrightarrow 5' \qquad \text{Coding Strand (Non-template): } 5' \longrightarrow 3' \tag{2.5}$$

Convention: All reference points (promoter, terminator) are defined with respect to the **coding strand**.

1. Transcription in Prokaryotes: Single RNA polymerase conducts all phases:
- Initiation: RNA Polymerase binds to promoter with transient assistance of **Sigma (σ) factor**.
- Elongation: RNA Polymerase synthesizes RNA in 5′ → 3′ direction using ribonucleoside triphosphates.
- Termination: Termination factor **Rho (ρ)** associates with polymerase to terminate transcription.
- Translation can begin **coupled with transcription** even before mRNA is fully synthesized (no nuclear membrane separation).

2. Transcription in Eukaryotes (Three Polymerases & Post-Transcriptional Processing):
- **RNA Polymerase I:** Transcribes rRNAs ($28S, 18S$, and $5.8S$).
- **RNA Polymerase II:** Transcribes precursor of mRNA, called **heterogeneous nuclear RNA (hnRNA)**.
- **RNA Polymerase III:** Transcribes **tRNA**, $5S$ **rRNA**, and **snRNAs** (small nuclear RNAs).

Post-Transcriptional Modifications of hnRNA → mature mRNA:
1. **Capping:** An unusual nucleotide, **methyl guanosine triphosphate (m^7Gppp)**, is added to the 5′-end.
2. **Tailing (Polyadenylation):** Adenylate residues (200 – 300) are added at 3′-end in a template-

independent manner.

3. **Splicing:** Non-coding intervening sequences (**Introns**) are excised, and coding sequences (**Exons**) are joined by spliceosomes (complex of snRNPs).

The Genetic Code: Salient Features & Wobble Phenomenon

Deciphered by George Gamow, Har Gobind Khorana, Marshall Nirenberg, and Severo Ochoa (polynucleotide phosphorylase).

Triplet Nature: 64 codons (4^3); 61 code for 20 amino acids; 3 stop codons.
Stop Codons: **UAA** (Ochre), **UAG** (Amber), **UGA** (Opal).
Dual Function: **AUG** codes for Met AND acts as initiator codon.
Degenerate: Amino acids coded by > 1 codon (Leu, Ser, Arg have 6).
Exceptions: AUG (Met) and UGG (Trp) have single codons alone.
Unambiguous: One specific codon codes for strictly one amino acid.
Universal: From bacteria to humans, UUU codes for Phe.

Wobble Hypothesis (Crick, 1966): The third base ($3'$ position of codon) does not pair as strictly with the first base ($5'$ position of anticodon) on tRNA. Non-Watson-Crick base pairing allows a single tRNA species to recognize multiple degenerate codons, reducing the minimum required tRNAs to 31.

Translation: Mechanism of Polypeptide Synthesis

1. **Charging of tRNA (Aminoacylation):**

$$\text{Amino Acid} + \text{ATP} \xrightarrow{\text{Aminoacyl-tRNA synthetase}} \text{Aminoacyl-AMP} + \text{PP}_\text{i} \tag{2.6}$$

$$\text{Aminoacyl-AMP} + \text{tRNA} \longrightarrow \text{Aminoacyl-tRNA} + \text{AMP} \tag{2.7}$$

2. **Initiation:** Small ribosomal subunit ($30S$ in prokaryotes, $40S$ in eukaryotes) binds to mRNA at $5'$-end (Shine-Dalgarno sequence in prokaryotes). Initiator tRNA binds to P-site (Peptidyl site).
3. **Elongation:** Incoming charged tRNA binds to the A-site (Aminoacyl site). Peptide bond formation is catalyzed by the ribozyme:

$$\text{Peptidyl transferase} = \textbf{23S}\ \text{rRNA (in prokaryotes)} \quad \text{or} \quad \textbf{28S}\ \text{rRNA (in eukaryotes)} \tag{2.8}$$

   Ribosome advances codon by codon along mRNA in $5' \to 3'$ direction (**Translocation**, requires EF-G and GTP).
4. **Termination:** Stop codon arrives at A-site; release factors (RF1, RF2, RF3) bind to stop codon, hydrolyzing polypeptide from tRNA.

Gene Regulation: The Operon Model of Jacob & Monod

The **Lac Operon** of *E. coli* is an example of an **inducible, negative regulation system**:

- **Gene i (Inhibitor / Regulatory):** Constitutively expresses the active Lac repressor protein.
- **Promoter (p):** Binding site for RNA polymerase.
- **Operator (o):** Binding site for the Lac repressor protein.
- **Structural Genes:**
  - **Gene z:** Codes for **β-galactosidase** (hydrolyzes lactose into glucose and galactose).
  - **Gene y:** Codes for **Permease** (increases cellular membrane permeability to β-galactosides).
  - **Gene a:** Codes for **Transacetylase** (transfers acetyl group from acetyl-CoA to β-galactosides).
- **Regulation States:**
  - *In Absence of Inducer (Lactose):* Repressor binds tightly to operator $\implies$ RNA polymerase blocked $\implies$ Operon switched OFF.
  - *In Presence of Inducer (Lactose / Allolactose):* Inducer binds to repressor, inducing confor-

mational change $\Longrightarrow$ repressor inactivated and detaches $\Longrightarrow$ RNA polymerase transcribes z, y, a $\Longrightarrow$ Operon switched ON.

Human Genome Project (HGP) & DNA Fingerprinting High-Yield Facts

1. **Human Genome Project Core Metrics:**
- Total base pairs: 3.1647×10^9 bp; average gene contains **3000 bases**.
- Largest known human gene: **Dystrophin** on X-chromosome (2.4×10^6 bases).
- Total estimated genes: $\approx 30,000$ (drastically lower than previous estimates of 140,000).
- Less than **2% of the genome** codes for proteins.
- Chromosome 1 has the most genes (**2968**); the Y chromosome has the fewest (**231**).
- Scientists have identified about **1.4 million locations** where single-base DNA differences (**SNPs** – single nucleotide polymorphisms) occur.

2. **DNA Fingerprinting (Alec Jeffreys, 1984):**
- Explores **DNA polymorphism** in non-coding repetitive DNA sequence (**Satellite DNA**).
- **VNTRs (Variable Number of Tandem Repeats):** Belong to a class of satellite DNA called *minisatellites*. The copy number varies from chromosome to chromosome in an individual (size varies from 0.1 to 20 kb).
- **Protocol Steps:** Isolation of DNA → Digestion by restriction endonucleases → Separation by agarose gel electrophoresis → Southern blotting (transfer to nylon membrane) → Hybridization using labeled VNTR probes → Detection by autoradiography.

Chapter 3

Sexual Reproduction & Applied Biotechnology

This chapter bridges plant and human reproduction with modern molecular biotechnology. We analyze microsporogenesis, megasporogenesis, pollen-pistil interactions, double fertilization, endosperm development, apomixis, human gametogenesis and hormonal regulation, contraceptive methods, and the principles and tools of recombinant DNA technology including restriction endonucleases, plasmid vectors, PCR amplification, Bt transgenic crops, and gene therapy.

3.1 Sexual Reproduction in Flowering Plants (Angiosperms)

Microsporogenesis & Male Gametophyte (Pollen Grain)

- **Anther Structure:** Typically bilobed, dithecous (two thecae per lobe), and tetrasporangiate (four microsporangia).
- **Wall Layers (from outside to inside):**
  1. **Epidermis:** Protective single outer layer.
  2. **Endothecium:** Layer with fibrous hygroscopic α-cellulose bands; assists in anther dehiscence.
  3. **Middle Layers:** $1-3$ transient ephemeral layers; degenerate to provide nourishment.
  4. **Tapetum:** Innermost nutritive layer; cells possess dense cytoplasm and are generally **multinucleate / polyploid**; secretes **sporopollenin**, **callase**, and pollenkitt.
- **Pollen Grain Wall (Sporoderm):**
  - **Exine:** Hard outer layer composed of **Sporopollenin** (the most resistant organic substance known; withstands high temperatures, strong acids, and alkali; no enzyme that degrades sporopollenin is known). Has apertures called **Germ Pores** where sporopollenin is absent.
  - **Intine:** Thin, continuous inner wall composed of **cellulose and pectin**.
- **Pollen Shedding Stage:** In $> 60\%$ of angiosperms, pollen is shed at the **2-celled stage** (large vegetative cell with irregular nucleus + small spindle-shaped generative cell floating in vegetative cytoplasm). In $< 40\%$, generative cell divides mitotically to form 2 male gametes before shedding (**3-celled stage**).

Megasporogenesis & The Monosporic Polygonum-Type Embryo Sac

- **Ovule (Integumented Megasporangium):** Attached to placenta by **funicle**; point of attachment to ovule body is **hilum**. Massive central tissue is **nucellus**.
- **Megasporogenesis:** Megaspore Mother Cell (MMC) undergoes meiosis to produce a linear tetrad of 4 haploid megaspores. 3 micropylar megaspores degenerate; only **1 chalazal megaspore remains functional** (**Monosporic development**).
- **Embryo Sac Organization (7-celled, 8-nucleate at maturity):**
  - **Micropylar End (Egg Apparatus - 3 cells):** 1 central **Egg cell** + 2 flanking **Synergids** possessing **Filiform apparatus** (finger-like wall projections that guide pollen tube entry).
  - **Chalazal End (Antipodals - 3 cells):** Vegetative nutritive cells; degenerate after fertilization.
  - **Central Cell (1 large cell):** Contains **2 Polar Nuclei** that fuse before fertilization to form a diploid **Secondary Nucleus** ($2n$).

Double Fertilization, Endosperm & Apomixis

1. Double Fertilization (Discovered by S.G. Nawaschin, 1898): Unique to angiosperms; involves two distinct nuclear fusion events:

Syngamy: Male Gamete (n) + Egg (n) $\longrightarrow$ Zygote (**2n**) $\longrightarrow$ Embryo

Triple Fusion: Male Gamete (n) + Secondary Nucleus $(2n)$ $\longrightarrow$ PEN (**3n**) $\longrightarrow$ Endosperm

Total Nuclei: 1 (sperm) + 1 (egg) + 1 (sperm) + 2 (polar nuclei) = **5** Nuclei

2. Endosperm Development: Triploid Primary Endosperm Nucleus (PEN) divides to form endosperm before the zygote divides (ensures nutrition for embryo).

- **Free-Nuclear Endosperm:** PEN undergoes repeated free nuclear divisions without cell wall formation (e.g., coconut water is free-nuclear endosperm consisting of thousands of free nuclei; surrounding white kernel is cellular endosperm).

3. Special Reproductive Modes:

- **Apomixis:** Form of asexual reproduction that **mimics sexual reproduction**, producing seeds without fertilization (e.g., *Asteraceae* and grasses). Nucellar cells divide and protrude into embryo sac to form diploid embryos.
- **Polyembryony:** Occurrence of more than one embryo in a seed (common in *Citrus* and mango).

3.2 Human Reproduction & Reproductive Health

Male & Female Reproductive Architecture

1. Male System:

- **Scrotum:** Keeps testes at $2-2.5°C$ lower than internal core body temperature, necessary for spermatogenesis.
- **Seminiferous Tubules:** Lined by (i) **Spermatogonia** (male germ cells) and (ii) **Sertoli cells** (nurse cells providing nutrition, secrete Androgen Binding Protein and Inhibin).
- **Interstitial / Leydig Cells:** Located in interstitial spaces outside tubules; synthesize and secrete androgens (**Testosterone**) under LH stimulation.
- **Accessory Glands:** Paired Seminal vesicles (contribute $60-70\%$ of semen, rich in **fructose, calcium**, and prostaglandins), single Prostate gland (milky alkaline fluid), paired Bulbourethral / Cowper's glands (secrete alkaline mucus for lubricating penis).

2. Female System:

- **Ovaries:** Primary female sex organs producing ovum and steroid hormones (Estrogen, Progesterone).
- **Fallopian Tube (Oviduct, 10–12 cm):** Consists of (i) **Infundibulum** with finger-like **fimbriae** to collect ovum after ovulation, (ii) **Ampulla** (the wider part; **site of fertilization**), (iii) **Isthmus** (narrow lumen joining uterus).
- **Uterine Wall:** Outer thin **perimetrium**, middle thick smooth-muscle **myometrium** (exhibits strong contractions during parturition), and inner glandular **endometrium** (undergoes cyclic changes during menstrual cycle).

Gametogenesis: Spermatogenesis vs Oogenesis Timing

1. Spermatogenesis (Continuous from Puberty through Old Age):

$$\text{Spermatogonia } (2n) \xrightarrow{\text{mitosis}} \text{Primary Spermatocyte } (2n) \xrightarrow{\text{Meiosis I}} 2 \times \text{Secondary Spermatocytes } (n)$$
$$\xrightarrow{\text{Meiosis II}} 4 \times \text{Spermatids } (n) \xrightarrow{\textbf{Spermiogenesis}} 4 \times \text{Spermatozoa } (\mathbf{n})$$

2. Oogenesis (Initiated during Fetal Life, Suspended until Puberty):

$$\text{Oogonia } (2n) \xrightarrow{\text{mitosis}} \text{Primary Oocyte } (2n) \xrightarrow{\text{Arrested at Prophase I}} \text{At Birth}$$

$$\text{At Puberty: Completes Meiosis I} \longrightarrow \text{Secondary Oocyte } (\mathbf{n}) + \mathbf{1}^{\text{st}} \text{ Polar Body } (\mathbf{n})$$

$$\text{Fertilization: Completes Meiosis II} \longrightarrow \text{Ovum } (\mathbf{n}) + \mathbf{2}^{\text{nd}} \text{ Polar Body } (\mathbf{n})$$

Menstrual Cycle Hormonal Regulation & Implantation

Standard 28-day cycle divided into four phases:

1. **Menstrual Phase (Days 1–5):** Breakdown of endometrial lining and blood vessels due to sudden **withdrawal of progesterone**; average blood loss $= 50 - 100$ mL.
2. **Follicular / Proliferative Phase (Days 6–13):** Pituitary **FSH** stimulates maturation of primary follicle into mature **Graafian follicle**; growing follicle secretes **Estrogen**, which regenerates endometrium.
3. **Ovulatory Phase (Day 14):** Peak levels of both LH and FSH; rapid surge of LH (**LH Surge**) induces rupture of Graafian follicle and release of secondary oocyte (**Ovulation**).
4. **Luteal / Secretory Phase (Days 15–28):** Ruptured follicle transforms into endocrine **Corpus Luteum**, which secretes massive amounts of **Progesterone** (essential for maintenance of endometrium). If fertilization fails, corpus luteum degenerates into **Corpus Albicans**, causing progesterone drop and menses.

Fertilization & Pregnancy Endocrinology:

- **Site of Fertilization:** Ampullary-isthmic junction / **Ampulla** of Fallopian tube.
- **Cortical Reaction:** Sperm induces depolarization and hardening of **zona pellucida**, blocking polyspermy.
- **Implantation:** Trophoblast layer of **Blastocyst** attaches to uterine endometrium ≈ 7 days after fertilization.
- **Placental Hormones exclusively in pregnancy:** Human Chorionic Gonadotropin (**hCG**), Human Placental Lactogen (**hPL**), and **Relaxin** (secreted by ovary in late pregnancy).

Reproductive Health: Contraception, MTP & ART Protocols

1. Contraceptive Methodologies:

- **Natural / Traditional:** Periodic abstinence (avoid coitus days $10 - 17$ of cycle), coitus interruptus, **lactational amenorrhea** (intense lactation prevents ovulation for max 6 months postpartum).
- **Barrier Methods:** Condoms (protect against STIs/AIDS), Diaphragms, cervical caps, vaults.
- **Intrauterine Devices (IUDs):**
  - **Non-medicated IUDs:** *Lippes loop* (increases phagocytosis of sperms).
  - **Copper-releasing IUDs: CuT, Cu7, Multiload 375** (Cu^{2+} ions suppress sperm motility and fertilizing capacity).
  - **Hormone-releasing IUDs: Progestasert, LNG-20** (make uterus unsuitable for implantation and cervix hostile to sperms).
- **Oral Contraceptive Pills:** Daily combined progestogen-estrogen pills; or **Saheli** (developed by CDRI Lucknow; non-steroidal, once-a-week pill with high contraceptive value and negligible side effects).
- **Surgical / Permanent Methods: Vasectomy** (cutting/tying vas deferens in males) and **Tubectomy** (fallopian tubes in females); irreversible sterilization.

2. Assisted Reproductive Technologies (ART):

- **IVF-ET (In Vitro Fertilization - Embryo Transfer):** Test-tube baby program.
  - **ZIFT (Zygote Intra-Fallopian Transfer):** Zygote or early embryo **up to 8 blastomeres** transferred into Fallopian tube.
  - **IUT (Intra-Uterine Transfer):** Embryos with **more than 8 blastomeres** transferred directly into Uterus.

- **GIFT (Gamete Intra-Fallopian Transfer):** Transfer of ovum collected from donor into fallopian tube of female who cannot produce one.
- **ICSI (Intra-Cytoplasmic Sperm Injection):** Sperm directly injected into ovum cytoplasm in laboratory.
- **IUI (Intra-Uterine Insemination):** Semen collected from husband/donor introduced artificially into uterus.

3.3 Biotechnology: Principles, Tools & Modern Applications

Tools of Recombinant DNA Technology

1. Restriction Endonucleases ("Molecular Scissors"): Discovered by Arber, Smith, and Nathans (1970). Cut double-stranded DNA at specific palindromic recognition sequences.

- **First Discovered:** *Hind II* (cuts at specific 6 bp sequence).
- **Nomenclature (*Eco*RI):** *E* for genus *Escherichia*, *co* for species *coli*, R for strain RY13, I for first enzyme isolated.
- **Recognition Sequence of *Eco*RI:**

$$5' - \text{G} \downarrow \text{A A T T C} - 3' \qquad 3' - \text{C T T A A} \uparrow \text{G} - 5' \tag{3.1}$$

Produces staggered single-stranded overhangs called **sticky ends**.

2. Cloning Vectors (Features of Ideal Vector pBR322):

- **Origin of Replication (*Ori*):** Specific sequence where replication initiates; controls copy number of linked foreign DNA.
- **Selectable Markers:** Identify and select transformants and eliminate non-transformants. Example: genes conferring resistance to antibiotics (amp^R **and** tet^R).
- **Insertional Inactivation:** Ligation of foreign DNA at *BamHI* site of tet^R gene inactivates tetracycline resistance $\implies$ recombinant grows on ampicillin, dies on tetracycline.
- **Blue-White Screening:** Foreign DNA inserted into gene coding for enzyme **β-galactosidase** (α-complementation). Non-recombinants produce blue colonies with X-gal; recombinants produce **white colonies**.

3. Competent Host Transformation Techniques:

- **Chemical Method:** Treating bacterial cells with divalent cation (Ca^{2+}) followed by **Heat Shock** (42°C for 2 minutes and ice plunge).
- **Microinjection:** Recombinant DNA directly injected into nucleus of animal cell.
- **Biolistics / Gene Gun:** Plant cells bombarded with high-velocity microparticles of **gold or tungsten** coated with DNA.
- **Disarmed Pathogen Vectors:** *Agrobacterium tumefaciens* (Ti plasmid for dicots) and retroviruses for animal cells.

Polymerase Chain Reaction (PCR) & Downstream Processing

Invented by Kary Mullis in 1983. In vitro enzymatic amplification of DNA sequence by a billion-fold (2^n copies):

Step 1: Denaturation: $94 - 96°\text{C}$ (Thermal separation of dsDNA strands by breaking H-bonds)

Step 2: Annealing: $50 - 60°\text{C}$ (Hybridization of two oligonucleotide primers to flanking $3'$-ends)

Step 3: Extension: $72°\text{C}$ (Extension by Taq Polymerase from *Thermus aquaticus*)

Bioreactors & Downstream Processing:

- **Bioreactors ($100 - 1000\,\text{L}$):** Provide optimal growth conditions (pH, temperature, oxygen, substrates, salts). **Stirred-tank** provides continuous mixing; **sparged stirred-tank** bubbles sterile air through broth.
- **Downstream Processing:** Separation and purification of biosynthesized product, followed by

clinical trials, formulation with preservatives, and stringent quality control.

High-Yield Transgenic Applications & Medical Biotechnology

1. **Bt Cotton & Pest Resistance:** Soil bacterium *Bacillus thuringiensis* produces crystalline proteinaceous protoxins (**Cry proteins**). The inactive protoxin is converted into active toxin in the **alkaline pH of the insect midgut**, binding to epithelial cells, creating pores, causing cell lysis and insect death:

   CryIAc and **CryIIAb** control Cotton Bollworms **CryIAb** controls Corn Borer (3.2)

2. **RNA Interference (RNAi):** Cellular defense mechanism in all eukaryotes. Nematode *Meloidogyne incognita* infects roots of tobacco plants. Using *Agrobacterium* vectors, nematode-specific genes are introduced producing both sense and anti-sense RNA $\implies$ forms **double-stranded RNA (dsRNA)** $\implies$ silences specific mRNA of nematode.
3. **Genetically Engineered Insulin (Humulin):** Human insulin consists of chain A (21 amino acids) and chain B (30 amino acids) linked by **disulphide bonds**. Synthesized as **prohormone containing an extra C-peptide** (33 amino acids), which is removed during maturation. In 1983, American company **Eli Lilly** produced two separate DNA sequences for chain A and chain B, introduced into *E. coli*, and combined them by creating disulphide bonds.
4. **Gene Therapy for ADA Deficiency:** First clinical gene therapy performed in 1990 on a 4-year-old girl with **Adenosine Deaminase (ADA) deficiency** (causes Severe Combined Immunodeficiency - SCID). Lymphocytes from patient's blood transduced with functional ADA cDNA using a retroviral vector and returned to patient. (Requires periodic infusion unless embryonic stem cells are engineered).
5. **Transgenic Animals & Biopiracy:**
   - **First Transgenic Cow Rosie (1997):** Produced human protein-enriched milk (2.4 g/litre) containing **human alpha-lactalbumin**, nutritionally more balanced for human babies than natural cow milk.
   - **Biopiracy Case:** In 1997, US company RiceTec was granted patent on Basmati rice cross-bred with semi-dwarf varieties, misappropriating traditional Indian farmers' strains.

Chapter 4

Ecology, Environmental Biology & Systematic Plant Families

Ecology examines the interactions of organisms with their abiotic and biotic environment, population growth kinetics, ecosystem trophic pyramids, and global biodiversity conservation strategies. Furthermore, this chapter provides a systematic morphological analysis of the seven angiosperm families specifically prescribed by the National Medical Commission: Malvaceae, Cruciferae (Brassicaceae), Fabaceae (Leguminosae), Asteraceae (Compositae), Poaceae (Gramineae), Solanaceae, and Liliaceae.

4.1 Organisms & Populations

Population Growth Formulations: Exponential vs Logistic Model

1. **Exponential Growth (Unlimited Resources / Geometric Progression):**

$$\frac{dN}{dt} = (b-d)N = rN \implies N_t = N_0 e^{rt} \quad \textbf{(J-shaped curve)} \tag{4.1}$$

Where N = population size, b = per capita birth rate, d = per capita death rate, r = **intrinsic rate of natural increase**:

- Norway rat: $r = 0.015$ | Flour beetle: $r = 0.12$ | Human population in India (1981): $r = 0.0205$.

2. **Logistic Growth / Verhulst-Pearl Model (Limited Resources / Realistic):**

$$\frac{dN}{dt} = rN\left(\frac{K-N}{K}\right) \quad \textbf{(Sigmoid / S-shaped curve)} \tag{4.2}$$

Where K = **Carrying Capacity** (maximum population density the habitat can support indefinitely).

- Four phases: Lag phase → Acceleration / Log phase → Deceleration → Asymptote ($N = K$).
- When $N \to K$, environmental resistance $\left(\frac{K-N}{K}\right) \to 0$, growth rate drops to zero.

Interspecific Population Interactions Matrix

Interaction Type	Species A	Species B	NCERT Canonical Examples
Mutualism	+	+	Lichens (Algae + Fungi); Mycorrhizae (Fungi + Roots of higher plants); Fig tree and pollinator Fig Wasp; Mediterranean orchid *Ophrys* (**sexual deceit** by mimicking female bee *Colpa*).
Competition	−	−	Abingdon tortoise in Galapagos became extinct within a decade after goats were introduced (browsing efficiency); **Gause's Exclusion Principle** (inferior eliminated if resources limiting); **MacArthur's Warblers** (5 species coexisted via **resource partitioning** and foraging behavior).
Predation	+	−	Biological control: *Cactoblastis* moth controlled invasive prickly pear cactus in Australia (1920); Pisaster starfish removal caused extinction of > 10 invertebrate species; Monarch butterfly accumulates cardiac glycosides from weed in caterpillar stage.
Parasitism	+	−	Ectoparasites: Lice on humans, ticks on dogs, marine copepods on fishes, *Cuscuta* (dodder) on hedge plants; Endoparasites: Liver fluke (requires snail and fish hosts), Malarial parasite; **Brood Parasitism** (Cuckoo/Koel lays eggs in Crow's nest).
Commensalism	+	0	Epiphytic orchid on mango tree branch; Barnacles attached to back of whale; Cattle egret foraging alongside grazing cattle; Clownfish living among stinging tentacles of Sea anemone.
Amensalism	−	0	*Penicillium* secreting penicillin antibiotic inhibiting bacterial growth; Black walnut (*Juglans nigra*) secreting juglone inhibiting understory plants.

4.2 Ecosystem: Structure, Function & Ecological Succession

Productivity Equations & Decomposition Dynamics

1. Primary Productivity Formulations:

Gross Primary Productivity (GPP): Total rate of organic matter synthesis in photosynthesis.

Net Primary Productivity (NPP): GPP $- R$ (R = respiratory loss)

Global Annual Biosphere NPP: $\approx$ **170** billion tons (dry organic matter)

Oceans Productivity: **55** billion tons (Covers 70% but yields 32% NPP)

2. Stages of Decomposition: Detritus (dead leaves, bark, animal remains) undergoes five simultaneous steps:

1. **Fragmentation:** Breakdown of detritus into smaller particles by detritivores (e.g., Earthworm).
2. **Leaching:** Water-soluble inorganic nutrients percolate into soil horizon and precipitate as unavailable salts.
3. **Catabolism:** Bacterial and fungal enzymes degrade detritus into simpler inorganic substances.
4. **Humification:** Accumulation of dark, amorphous, colloidal **Humus** (highly resistant to microbial action, extremely slow decomposition, acts as reservoir of nutrients).
5. **Mineralization:** Slow release of inorganic nutrients by microbial degradation of humus.

Rate Factors: Decomposition is fast if detritus is rich in **nitrogen and water-soluble sugars**; slow if rich in **lignin and chitin**.

Energy Flow & Ecological Pyramids Architecture

- **10% Law of Lindeman:** Only **10**% of energy entering a trophic level is transferred to the next higher level; 90% is lost as metabolic heat.
- **Photosynthetically Active Radiation (PAR):** Out of total incident solar radiation, less than

50% is PAR. Plants capture only **2 – 10**% of PAR (1 – 5% of total incident radiation).

- **Ecological Pyramids Comparison:**
  - **Pyramid of Numbers:** Mostly upright (grassland); **inverted** in tree ecosystem supporting thousands of parasitic insects.
  - **Pyramid of Biomass:** Upright on land (trees/grasses have vast biomass); **Inverted in Aquatic / Marine Ecosystems** (biomass of phytoplankton is drastically smaller than that of zooplankton and predatory fishes).
  - **Pyramid of Energy: Strictly and always upright**; can never be inverted because energy is lost as heat at each transfer (Second Law of Thermodynamics).
- **Succession Trajectory:** Both **Hydrarch** (water) and **Xerarch** (bare rock) successions converge toward identical moderate water conditions called **Mesic conditions**.

4.3 Biodiversity & Conservation

Species-Area Relationship (Alexander von Humboldt)

Alexander von Humboldt observed that within a region, species richness increases with increasing explored area, but only up to a limit:

$$S = CA^Z \implies \log \mathbf{S} = \log \mathbf{C} + \mathbf{Z} \log \mathbf{A} \tag{4.3}$$

Where S = Species richness, A = Area, C = Y-intercept, Z = **Slope of regression line**:

- For small regional areas (plants in Britain, birds in California): Z is strictly 0.1 **to** 0.2.
- For vast continents (frugivorous birds and mammals in tropical forests): Z is much steeper, 0.6 **to** 1.2 (typically 1.15).

Global Biodiversity Metrics & The "Evil Quartet" Causes of Loss

1. Biodiversity Estimates (Robert May): Global species total ≈ **7** million. More than **70% of all recorded species are animals**, and plants comprise ≤ 22%. Among animals, **insects make up** > 70% (7 out of 10 animals on Earth are insects). India has **2.4% of global land area** but accounts for **8.1% of global species diversity** (one of 12 mega-diversity countries).

2. The "Evil Quartet" (Major Drivers of Biodiversity Extinction):

1. **Habitat Loss & Fragmentation:** The single most catastrophic cause. Tropical rainforests once covered 14% of Earth's land, now reduced to < 6%. Amazon rainforest is being cleared for soybean cultivation and cattle pasture.
2. **Over-Exploitation:** Excessive hunting led to extinction of *Steller's sea cow* and *Passenger pigeon*.
3. **Alien Species Invasions:**
   - Nile perch introduced into Lake Victoria caused extinction of > 200 endemic *Cichlid fish* species.
   - Invasive weeds: *Parthenium* (Carrot grass), *Lantana*, and *Eichhornia crassipes* (Water Hyacinth / "Terror of Bengal").
   - Illegal introduction of African catfish *Clarias gariepinus* threatens indigenous catfishes in Indian rivers.
4. **Co-Extinctions:** Obligate co-evolved relationships (e.g., host fish and its specific monogenean parasite; plant-pollinator mutualism).

3. In-Situ vs Ex-Situ Conservation:

- **In-Situ (On-site):** 14 Biosphere Reserves, 90 National Parks, 448 Wildlife Sanctuaries, Sacred Groves (Khasi and Jaintia Hills in Meghalaya, Aravalli Hills in Rajasthan, Western Ghats).
- **Ex-Situ (Off-site):** Zoological parks, Botanical gardens, Wildlife safari parks, Cryopreservation of gametes at −196°C in liquid nitrogen, Seed banks, Tissue culture.

4.4 Systematic Morphology of Plant Families (Prescribed NMC Syllabus)

Diagnostic Master Comparison of Prescribed Angiosperm Families

Family	Floral Formula	Androecium / Gynoecium	Diagnostic Traits & Important Economic Plants
Brassicaceae (Mustard)	$\oplus\ K_{2+2}\ C_4\ A_{2+4}\underline{G}_{(2)}$	**Tetradynamous** (2 outer short, 4 inner long); Bicarpellary, unilocular with **false septum (Replum)**.	Cruciform corolla (clawed petals); Siliqua fruit. *Brassica campestris* (mustard), *Raphanus sativus* (radish).
Fabaceae (Legume)	$\%\ K_{(5)}\ C_{1+2+(2)}\ A_{(9)+1}\underline{G}_1$	**Diadelphous** (9 fused +1 free); Monocarpellary, marginal placentation.	**Vexillary aestivation** (1 standard, 2 wings, 2 keel); Root nodules with *Rhizobium*. Pulses (Gram, Arhar), Oil (Soyabean, Groundnut), Dye (*Indigofera*), Fodder (*Sesbania*), Medicine (*Muliathi*).
Solanaceae (Potato)	$\oplus\ K_{(5)}\ \overbrace{C_{(5)}\ A_5}\ \underline{G}_{(2)}$	Epipetalous stamens; Bicarpellary, syncarpous, **oblique ovary**, **swollen axile placenta**.	Persistent calyx; Berry or capsule. Food (Potato, Tomato, Brinjal), Spice (Chilli), Medicine (*Belladonna*, *Ashwagandha*), Fumigatory (Tobacco), Ornamental (*Petunia*).
Liliaceae (Lily)	$\oplus\ \overbrace{P_{(3+3)}\ A_{3+3}}\ \underline{G}_{(3)}$	Epiphyllous stamens; Tricarpellary, syncarpous, trilocular with axile placentation.	Perianth of 6 tepals in two whorls; Capsule/berry. Ornamental (Tulip, *Gloriosa*), Medicine (*Aloe*), Vegetable (*Asparagus*), Colchicine (*Colchicum autumnale*).
Malvaceae (Cotton)	$\oplus\ K_{(5)}\ C_5\ A_{(\infty)}\underline{G}_{(5)}$	**Monadelphous** staminal tube; **Monothecous, reniform anthers**.	**Epicalyx** present; Mucilaginous sap; Twisted aestivation. Cotton (*Gossypium*), Lady's finger (*Abelmoschus*), China rose (*Hibiscus*).
Compositae (Sunflower)	Ray: $\%\ K_{pappus}\ C_{(5)}\ A_0\underline{G}_{(2)}$ Disc: $\oplus\ K_{pappus}\ \overbrace{C_{(5)}\ A_{(5)}}\ \underline{G}_{(2)}$	**Syngenesious** (anthers fused, filaments free); Bicarpellary, unilocular, **basal placentation**.	**Capitulum inflorescence**; Calyx modified into hairy **Pappus**; Cypsela fruit. Sunflower (*Helianthus*), Marigold (*Tagetes*), *Chrysanthemum*.
Gramineae (Poaceae - Grass)	$\%\ P_2\ A_3$ or $A_{3+3}\underline{G}_1$	Versatile anthers; Monocarpellary, unilocular, basal placentation with **feathery bifid stigma**.	**Spikelet inflorescence**; Perianth represented by 2 scale-like **Lodicules**; Caryopsis fruit. Cereals (Wheat, Rice, Maize, Sugarcane, Bamboo).

PART II

Zoology: Cellular Life, Animal Diversity, Human Physiology & Health

Chapter 5

Cell Biology, Cell Division & Animal Morphology

Every biological inquiry starts at the cellular level. This chapter analyzes prokaryotic and eukaryotic ultrastructure, organelle compartmentalization, enzyme catalytic mechanics, the stages of mitosis and meiosis with synapsis and crossing over, the phylogenetic classification of animal phyla, and the complete anatomical systems of the frog (*Rana tigrina*) and cockroach (*Periplaneta americana*).

A Personal Note from Satya: The Physiology of Memory — Why Sleep is Non-Negotiable for Future Doctors

Since you are studying Human Physiology and Cell Biology, let's talk about your own brain physiology.
Every year, I see bright students ruin months of hard work during the final 30 days before NEET by falling into the "heroic all-nighter" trap. They sleep 4 hours, drink excessive coffee, and brag about studying until 4:00 AM.
Here is what is literally happening inside your skull when you do that:

- **The Glial Waste-Clearance System:** During deep sleep, the interstitial space between your neurons increases by 60%, allowing cerebrospinal fluid (CSF) to flush out toxic metabolic waste products like β-amyloid and adenosine. When you deprive yourself of sleep, these metabolic toxins remain, creating severe cognitive brain fog.
- **Long-Term Potentiation (LTP) in the Hippocampus:** Memory is encoded through synaptic plasticity via NMDA glutamate receptors. In a sleep-deprived state, synaptic fatigue blocks LTP. You can read the respiratory volumes (TV, IRV, ERV, RV) five times, but your brain physically cannot encode them into long-term synaptic tracks.

As a future medical professional, you must respect biological laws. Give yourself 7 to 8 hours of solid sleep. When you wake up, your neurotransmitters are replenished, your focus is sharp, and you will retain the intricate details of animal morphology and human physiology with effortless clarity.

5.1 Cell Structure, Function & Biomolecules

Prokaryotic vs Eukaryotic Cell Architecture

1. **Cell Theory Milestones:**
- **M.J. Schleiden (1838)** observed plants; **Theodore Schwann (1839)** studied animal cells and noted presence of cell wall unique to plant cells.
- **Rudolf Virchow (1855)** modified cell theory: ***Omnis cellula-e cellula*** (all cells arise from pre-existing cells).

2. **Prokaryotic Diagnostic Architecture:**
- **Cell Envelope (3 Layers):** Outermost **Glycocalyx** (loose slime layer or rigid **capsule**) → Middle **Peptidoglycan cell wall** → Innermost semi-permeable **Plasma membrane**.
- **Mesosomes:** Invaginations of plasma membrane in the form of vesicles, tubules, and lamellae; increase surface area for enzymatic respiration, secretion, and replication.
- **Plasmids:** Small, circular, self-replicating extrachromosomal dsDNA; confer specific phenotypes like antibiotic resistance.
- **Inclusion Bodies:** Non-membrane bound storage granules free in cytoplasm (e.g., phosphate

granules, cyanophycean granules, glycogen granules, gas vacuoles in blue-green and purple photosynthetic bacteria).

3. Eukaryotic Organelles & The Endomembrane System: The endomembrane system includes functionally coordinated organelles: **Endoplasmic Reticulum (ER), Golgi Apparatus, Lysosomes, and Vacuoles** (Mitochondria, Chloroplasts, and Peroxisomes are NOT part of it).

- **Fluid Mosaic Model (Singer & Nicolson, 1972):** Quasi-fluid lipid bilayer (phospholipids with hydrophilic heads outside, hydrophobic fatty acid tails inside) with integrated peripheral and integral proteins.
- **Rough ER (RER):** Studded with $80S$ ribosomes; active in **protein synthesis and secretion**.
- **Smooth ER (SER):** Major site for **lipid and steroid hormone synthesis**; detoxifies drugs in liver.
- **Golgi Complex (Camillo Golgi, 1898):** Flattened cisternae ($0.5 - 1.0\,\mu$m); convex forming **cis-face** receives vesicles from ER; concave maturing **trans-face** releases modified vesicles; crucial site for **glycosylation of proteins (glycoproteins)** and **lipids (glycolipids)**.
- **Lysosomes:** Formed by Golgi packaging; contain hydrolytic enzymes (lipases, proteases, carbohydrases) active at **acidic pH** (≈ 5.0).
- **Ribosomes (George Palade, 1953):** Non-membrane bound ribonucleoprotein granules ($70S = 50S + 30S$ in prokaryotes/organelles; $80S = 60S + 40S$ in eukaryotic cytoplasm).
- **Cilia & Flagella:** Possess $9+2$ axoneme microtubules; basal body derived from centrioles ($9+0$ cartwheel).

Enzyme Catalysis, Kinetics & Michaelis-Menten Dynamics

Enzymes are biocatalysts (mostly proteins; ribozymes are RNA enzymes like $23S$ rRNA):

- Lower the **activation energy** barrier without altering standard free energy change (ΔG) or equilibrium constant (K_{eq}).
- **Michaelis Constant (K_m):** Substrate concentration at which the reaction velocity reaches **half-maximal velocity** ($V_{\max}/2$).

$$K_m \propto \frac{1}{\text{Substrate Affinity of Enzyme}} \quad (\text{Lower } K_m \implies \text{Higher catalytic affinity}) \tag{5.1}$$

- **Competitive Inhibition:** Inhibitor closely resembles substrate in molecular structure and competes for active site:

$$\textbf{Competitive Inhibitor Effect:} \quad \mathbf{K_m} \text{ increases} \qquad \mathbf{V}_{\max} \text{ remains unchanged} \tag{5.2}$$

  Canonical NCERT Example: Inhibition of **succinate dehydrogenase by malonate** (resembles succinate).
- **Non-Competitive Inhibition:** Inhibitor binds to allosteric site: $\mathbf{V}_{\max}$ decreases, while $\mathbf{K_m}$ remains unchanged.
- **Enzyme Classes (IUBMB - 6 Classes):** (1) **Oxidoreductases**, (2) **Transferases**, (3) **Hydrolases**, (4) **Lyases** (cleave bonds without water), (5) **Isomerases**, (6) **Ligases** (join bonds with ATP hydrolysis).
- **Cofactors (Non-Protein Moieties):**
  - **Prosthetic Groups:** Tightly and permanently bound organic compounds (e.g., **Haem** in peroxidase and catalase).
  - **Coenzymes:** Transiently bound organic compounds derived from vitamins (e.g., **NAD and NADP** contain niacin).
  - **Metal Ions:** Form coordination bonds (e.g., Zn^{2+} in carboxypeptidase and carbonic anhydrase).

5.2 Cell Cycle & Cell Division: Mitosis & Meiosis

Cell Cycle Progression & Chromosome/DNA Mathematics

Standard eukaryotic cell cycle (human cell divides once in ≈ 24 hours; yeast divides in 90 minutes):

Interphase ($> 95\%$ of cycle): $\mathbf{G_1}$ (growth) $\longrightarrow$ **S** (DNA replication) $\longrightarrow$ $\mathbf{G_2}$ (spindle proteins)
M-Phase ($< 5\%$ of cycle): Prophase $\longrightarrow$ Metaphase $\longrightarrow$ Anaphase $\longrightarrow$ Telophase $\longrightarrow$ Cytokinesis

Quantitative Dynamics of Chromosome Number ($2n$) and DNA Content (C):

Stage	G_1 Phase	S Phase	G_2 Phase	Metaphase	Anaphase	Daughter Cells
Chromosome Count	$2n$	$2n$	$2n$	$2n$	**4n** (split centromeres)	$2n$
DNA Content	$2C$	**4C** (doubles)	$4C$	$4C$	$4C$	$2C$

Meiotic Substages: Prophase I Molecular Events

Meiosis reduces chromosome number by half ($2n \rightarrow n$). Prophase I is prolonged and divided into 5 distinct substages:

1. **Leptotene:** Chromosomes gradually become visible under light microscope; compaction of chromosomes continues.
2. **Zygotene:** Homologous chromosomes pair together (**Synapsis**), forming a nucleoprotein complex called **Synaptonemal Complex**. The paired complex is termed a **Bivalent** or **Tetrad**.
3. **Pachytene:** Non-sister chromatids of homologous chromosomes exchange genetic material (**Crossing Over**) at recombination nodules, catalyzed by the enzyme **Recombinase**.
4. **Diplotene:** Dissolution of synaptonemal complex; homologous chromosomes separate except at crossover sites, forming X-shaped configurations called **Chiasmata**. (Can remain arrested for months/years in oocytes of vertebrates, e.g., Dictyotene).
5. **Diakinesis: Terminalization of chiasmata**; chromosomes are fully condensed; nucleolus disappears and nuclear envelope breaks down.

Crucial Meiotic Checkpoints:

- **Anaphase I:** Homologous chromosomes separate to opposite poles; **sister chromatids remain attached at their centromeres** $\Longrightarrow$ Reduction in chromosome number ($2n \rightarrow n$).
- **Anaphase II: Centromeres split**, allowing sister chromatids to move to opposite poles $\Longrightarrow$ Resembles normal mitotic anaphase.

5.3 Animal Kingdom: Non-Chordates & Chordates Blueprint

Diagnostic Hallmarks of Major Animal Phyla

Phylum	Level of Org. / Symmetry	Coelom / Germ Layers	Diagnostic Features & Canonical Examples
Porifera (Sponges)	Cellular; Asymmetrical	Acoelomate; Diploblastic	**Choanocytes / Collar cells** line spongocoel; Water canal system; Spicules/spongin skeleton. *Sycon, Spongilla* (freshwater), *Euspongia* (bath sponge).
Cnidaria (Coelenterata)	Tissue; Radial symmetry	Acoelomate; Diploblastic	**Cnidoblasts / Nematocysts** on tentacles for defense/anchorage; Polyp and Medusa; **Metagenesis** in *Obelia. Hydra, Physalia* (Portuguese man-of-war), *Adamsia* (sea anemone).
Ctenophora (Comb jellies)	Tissue; Radial symmetry	Acoelomate; Diploblastic	Eight external rows of ciliated **comb plates** for locomotion; **Bioluminescence**; Strictly marine. *Pleurobrachia, Ctenoplana.*
Platyhelminthes (Flatworms)	Organ / Organ-system; Bilateral	**Acoelomate**; Triploblastic	Dorso-ventrally flattened body; Hooks and suckers in parasitic forms; **Flame cells (Protonephridia)** for osmoregulation. *Taenia* (tapeworm), *Fasciola* (liver fluke).
Aschelminthes (Roundworms)	Organ-system; Bilateral	**Pseudocoelomate**; Triploblastic	Circular cross-section; Muscular pharynx; Sexual dimorphism (female longer). *Ascaris, Wuchereria* (filarial worm), *Ancylostoma* (hookworm).
Annelida (Segmented worms)	Organ-system; Bilateral	**Coelomate**; Triploblastic	**Metameric segmentation**; Longitudinal/circular muscles; **Nephridia** for excretion; Closed circulation. *Nereis* (parapodia), *Pheretima* (earthworm), *Hirudinaria* (leech).
Arthropoda (Jointed legs)	Organ-system; Bilateral	Coelomate; Triploblastic	**Largest phylum** ($>2/3$ species); Chitinous exoskeleton; **Malpighian tubules** for excretion; Statocysts; Open circulation. *Apis, Bombyx, Laccifer, Anopheles, Locusta, Limulus* (living fossil).
Mollusca (Soft-bodied)	Organ-system; Bilateral	Coelomate; Triploblastic	**Second largest phylum**; Calcareous shell; Unsegmented with head, muscular foot, and visceral hump; Mantle cavity with ctenidia; Rasping organ **Radula**. *Pila, Pinctada, Sepia, Octopus.*
Echinodermata (Spiny-skinned)	Organ-system; Radial (adult), Bilateral (larva)	Coelomate; Triploblastic	Endoskeleton of calcareous ossicles; **Water vascular system** (ambulacral system) for locomotion, food capture, and respiration; Excretory system absent. *Asterias* (starfish), *Echinus, Antedon.*
Hemichordata	Organ-system; Bilateral	Coelomate; Triploblastic	Worm-like marine; Proboscis, collar, and trunk; **Stomochord** in collar; Open circulation; Proboscis gland for excretion. *Balanoglossus, Saccoglossus.*
Chordata	Organ-system; Bilateral	Coelomate; Triploblastic	Fundamental hallmarks: (1) **Notochord**, (2) **Dorsal hollow nerve cord**, (3) **Paired pharyngeal gill slits**, (4) **Post-anal tail**.

5.4 Structural Organisation in Animals: Frog & Cockroach Anatomy

Cockroach Anatomy (*Periplaneta americana*) Prescribed Scope

- **Morphology:** Body divided into Head, Thorax, and Abdomen; covered by brown chitinous exoskeleton with hardened plates called **Sclerites** (dorsal **Tergites**, ventral **Sternites**, joined by flexible **arthrodial membrane**).
- **Head:** Triangular, hypognathous; formed by fusion of **6 embryonic segments**; bears a pair of compound eyes (each composed of $\approx$ **2000** hexagonal ommatidia providing mosaic vision with high sensitivity but low resolution).
- **Mouthparts (Biting and Chewing Type):** Labrum (upper lip), paired Mandibles (with grinding and incising teeth), paired Maxillae, Labium (lower lip), and median flexible lobe **Hypopharynx** (tongue).
- **Thorax:** Prothorax, Mesothorax, Metathorax. Forewings (**Tegmina** from mesothorax) are opaque, dark, leathery, and protective at rest; Hindwings (from metathorax) are transparent, membranous, and used in **flight**.
- **Abdomen:** Consists of **10 segments** in both sexes.
  - **Sexual Dimorphism:** Males bear a pair of unjointed **Anal Styles** on the 9th sternite (absent in females). Both sexes bear a pair of jointed **Anal Cerci** on the 10th segment.
- **Digestive System:** Alimentary canal: Mouth $\rightarrow$ Pharynx $\rightarrow$ Oesophagus $\rightarrow$ **Crop** (storage) $\rightarrow$ **Gizzard / Proventriculus** (has outer circular muscles and 6 thick chitinous teeth for grinding food).
  - **Hepatic / Gastric Caeca:** $6-8$ blind tubules at the junction of foregut and midgut; secrete digestive juice.
  - **Malpighian Tubules:** $100-150$ thin, yellow, filamentous tubules at the junction of midgut

and hindgut; extract uric acid from hemolymph (**Uricotelic**).

- **Circulatory System:** Open type; heart consists of **13 funnel-shaped chambers** with ostia on either side.
- **Nervous System:** Consists of a series of fused ganglia: **3 in the thorax and 6 in the abdomen**. If head is cut off, cockroach survives for up to a week because the head holds only a bit of the nervous system!

Frog Anatomy (*Rana tigrina*) Prescribed Scope

- **Morphology:** Cold-blooded (**Poikilotherm**); amphibious; smooth, moist, slimy skin with mucus glands; camouflages (**Mimicry**). Body divided into **Head and Trunk** (neck and tail absent). Eyes are protected by a **Nictitating membrane**. Tympanum receives sound vibrations.
  - **Sexual Dimorphism:** Male frogs have sound-producing **Vocal Sacs** and a **Copulatory pad** on the first digit of the forelimb (both absent in female).
- **Respiration:** Cutaneous (skin in water and hibernation/aestivation), Buccopharyngeal cavity, and Pulmonary (lungs on land).
- **Circulatory System:** Closed, 3-chambered heart (**2 Atria and 1 Ventricle**).
  - Triangular **Sinus Venosus** joins the right atrium; receives blood from vena cava.
  - Ventricle opens ventrally into **Conus Arteriosus**.
  - **RBCs:** Nucleated, oval, and contain hemoglobin (unlike non-nucleated human RBCs).
  - Well-developed **Hepatic Portal** and **Renal Portal** systems.
- **Excretory System:** Paired compact, dark red **Mesonephric kidneys**; **Ureotelic** (excretes urea). Functional unit is uriniferous tubule. In males, ureters act as **Urogenital ducts** opening into the **Cloaca** (common chamber for digestive, excretory, and reproductive products).
- **Male Reproductive System:** Paired yellowish ovoid testes adhered to upper kidneys by double fold of peritoneum called **Mesorchium**. $10-12$ **Vasa efferentia** enter kidneys to run into **Bidder's Canal**, opening into urogenital duct.
- **Development:** External fertilization in water; indirect development via swimming aquatic larva called **Tadpole** (undergoes metamorphosis governed by thyroxine).

Chapter 6

Human Physiology: Integrated Organ Systems

Human Physiology represents the core clinical foundation for medical aspirants. In this chapter, we explore the mechanical and biochemical processes of breathing and alveolar gas exchange, body fluids and the cardiac cycle, nephron counter-current exchange and osmoregulation, musculoskeletal excitation-contraction coupling in sliding filament mechanics, and neuro-endocrine coordination via action potentials and peptide/steroid hormone cascades.

6.1 Breathing & Exchange of Gases

Respiratory Volumes & Pulmonary Capacities Matrix

All values represent healthy adult human averages measured by spirometer (residual volume cannot be measured by spirometer):

Tidal Volume (TV): 500 mL ($\approx 6000 - 8000$ mL/min for healthy adult)
Inspiratory Reserve Volume (IRV): $2500 - 3000$ mL
Expiratory Reserve Volume (ERV): $1000 - 1100$ mL
Residual Volume (RV): $1100 - 1200$ mL (air remaining after maximal expiration)
Inspiratory Capacity (IC): TV + IRV = $3000 - 3500$ mL
Expiratory Capacity (EC): TV + ERV = $1500 - 1600$ mL
Functional Residual Capacity (FRC): ERV + RV = $2100 - 2300$ mL
Vital Capacity (VC): ERV + TV + IRV = $3500 - 4500$ mL
Total Lung Capacity (TLC): RV + ERV + TV + IRV = VC + RV = $5000 - 6000$ mL

Partial Pressures & Gas Exchange Gradients

Diffusion of gases across the three-layered respiratory membrane (alveolar squamous epithelium, basement membrane, capillary endothelium) occurs along partial pressure gradients:

Gas	Atmospheric Air	Alveoli	Deoxygenated Blood	Oxygenated Blood	Tissues
O_2	159 mmHg	**104** mmHg	40 mmHg	**95** mmHg	40 mmHg
CO_2	0.3 mmHg	**40** mmHg	45 mmHg	**40** mmHg	45 mmHg

Solubility Factor: The solubility of CO_2 is **20–25 times higher** than that of O_2. Hence, the amount of CO_2 that can diffuse across the diffusion membrane per unit difference in partial pressure is much higher. Total membrane thickness is < 1 mm (around 0.2 mm).

Oxygen Dissociation Curve & Gas Transport Mechanisms

1. **Transport of Oxygen:**
   - 97% bound to hemoglobin as **Oxyhemoglobin** in RBCs (1 g Hb binds 1.34 mL O_2; each Hb molecule binds max 4 molecules of O_2).
   - 3% dissolved in blood plasma.
   - **Delivery Yield:** Under normal physiological conditions, every 100 mL of oxygenated blood de-

livers approximately 5 mL **of** O_2 to the tissues.

2. **Oxygen-Hemoglobin Dissociation Curve (Sigmoid / S-shaped):**

- **Shift to Right (Bohr Effect / O_2 Release at Tissues):** Promoted by:

$$\uparrow pCO_2, \quad \uparrow [H^+] \ (\downarrow \text{pH}), \quad \uparrow \text{Temperature}, \quad \uparrow 2,3\text{-BPG} \tag{6.1}$$

- **Shift to Left (Loading at Alveoli):** Promoted by $\uparrow pO_2, \downarrow pCO_2, \downarrow [H^+]$ ($\uparrow$ pH), $\downarrow$ Temperature.

3. **Transport of Carbon Dioxide:**

- As **Bicarbonate ions** (HCO_3^-)**: 70**% facilitated by RBC enzyme *carbonic anhydrase* and chloride shift (Hamburger phenomenon).
- As **Carbamino-hemoglobin: 20 – 25**% bound to amino groups of hemoglobin.
- In **Dissolved state** in plasma: **7**%.
- **Delivery Yield:** Every 100 mL of deoxygenated blood delivers approximately 4 mL **of** CO_2 to the alveoli.

NEET Trap: Respiratory Regulation & Clinical Pathology

- **Primary Chemical Sensor:** Chemosensitive area located adjacent to the rhythm center in medulla is **highly sensitive to** CO_2 **and** H^+ **ions**. The role of oxygen in the regulation of respiratory rhythm is **quite insignificant**!
- **Respiratory Centers:** Respiratory rhythm center is in **Medulla oblongata**. **Pneumotaxic center** is in **Pons varolii** (moderates rhythm center and reduces duration of inspiration).
- **Asthma:** Allergic inflammation of bronchi and bronchioles causing wheezing.
- **Emphysema:** Chronic disorder in which **alveolar walls are damaged**, decreasing respiratory surface area; majorly caused by cigarette smoking (α-1-antitrypsin used in treatment).
- **Occupational Respiratory Disorders:** Silicosis, asbestosis; chronic exposure leads to **fibrosis** (proliferation of fibrous tissues) in upper parts of lungs.

6.2 Body Fluids & Circulation

Blood Composition, Formed Elements & Coagulation Cascade

Blood is a specialized fluid connective tissue consisting of plasma (55%) and formed elements (45%):

- **Plasma:** 90 – 92% water, 6 – 8% proteins (Fibrinogen for clotting, Globulins for defense/immunity, Albumins for osmotic balance).
- **Erythrocytes (RBCs):** Biconcave, enucleated in adult mammals, $5.0 - 5.5 \times 10^6/\text{mm}^3$. Average lifespan = 120 days; destroyed in the **spleen** (graveyard of RBCs). Hemoglobin count = 12 – 16 g/100 mL.
- **Leukocytes (WBCs):** Nucleated, $6000 - 8000/\text{mm}^3$.
  - *Granulocytes:* **Neutrophils** (60 – 65%, phagocytic), **Eosinophils** (2 – 3%, resist infections, allergic response), **Basophils** (0.5 – 1%, secrete histamine, serotonin, heparin; mediate inflammation).
  - *Agranulocytes:* **Monocytes** (6 – 8%, phagocytic), **Lymphocytes** (20 – 25%, B and T lymphocytes for adaptive immunity).
- **Thrombocytes (Platelets):** Cell fragments produced from **megakaryocytes** in bone marrow ($1.5 - 3.5 \times 10^5/\text{mm}^3$).
- **Blood Coagulation Cascade (Enzyme Cascade Theory):**

$$\text{Thromboplastin (Injured Tissue + Platelets)} \xrightarrow{Ca^{2+}} \text{Prothrombinase / Thrombokinase} \tag{6.2}$$

$$\text{Prothrombin (Inactive)} \xrightarrow[Ca^{2+}]{\text{Thrombokinase}} \text{Thrombin (Active)} \tag{6.3}$$

$$\text{Fibrinogen (Soluble)} \xrightarrow{\text{Thrombin}} \text{Fibrin (Insoluble Mesh)} \xrightarrow{\text{traps dead elements}} \text{Blood Clot} \tag{6.4}$$

Calcium (Ca^{2+}) plays an indispensable cofactor role in almost all clotting steps!

Cardiac Cycle, Hemodynamics & Electrocardiogram (ECG)

1. **Cardiac Cycle Timing (72 beats/min $\implies$ Duration = 0.8 sec):**

Atrial Systole:	0.1 sec	(increases ventricular filling by 30%)
Ventricular Systole:	0.3 sec	(isovolumetric contraction $\rightarrow$ ejection period)
Joint Diastole:	0.4 sec	(passive filling of atria and ventricles)

Stroke Volume (SV): EDV − ESV = 120 mL − 50 mL = **70** mL

Cardiac Output (CO): Stroke Volume × Heart Rate = 70 mL × 72/min ≈ **5040** mL/min ≈ **5** L/min

2. **Heart Sounds:**
- **First Heart Sound (LUBB):** Closure of **atrioventricular valves** (Tricuspid and Bicuspid) at the onset of ventricular systole; low pitched, longer duration.
- **Second Heart Sound (DUBB):** Closure of **semilunar valves** at the onset of ventricular diastole; high pitched, shorter duration.

3. **Standard Electrocardiogram (ECG) Deflections:**
- **P-wave:** Depolarization of atria (leads to atrial contraction).
- **QRS Complex:** Depolarization of ventricles (marks onset of ventricular contraction; counting QRS complexes per unit time gives heart rate).
- **T-wave:** Repolarization of ventricles (return to resting state; end of T-wave marks end of systole).

Double Circulation & Hepatic Portal System

- **Pulmonary Circulation:** Deoxygenated blood from Right Ventricle $\xrightarrow{\text{Pulmonary Artery}}$ Lungs $\xrightarrow{\text{Pulmonary Veins}}$ Left Atrium.
- **Systemic Circulation:** Oxygenated blood from Left Ventricle $\xrightarrow{\text{Aorta}}$ Body Tissues $\xrightarrow{\text{Vena Cava}}$ Right Atrium.
- **Hepatic Portal System:** Unique vascular connection carrying blood from the **digestive tract to the liver** via the Hepatic Portal Vein before reaching systemic circulation.
- **Coronary Circulation:** Exclusive network of blood vessels supplying muscular walls of the heart.

6.3 Excretory Products & Their Elimination

Modes of Nitrogenous Excretion & Nephron Anatomy

- **Ammonotelism:** Excretion of toxic ammonia requiring large volumes of water; diffuses directly across body surfaces or gill surfaces as ammonium ions (e.g., bony fishes, aquatic amphibians, aquatic insects). Kidneys play no significant role in ammonia removal.
- **Ureotelism:** Urea synthesized in the liver via Ornithine cycle; requires moderate water (e.g., mammals, terrestrial amphibians, marine cartilaginous fishes).
- **Uricotelism:** Uric acid excreted as pellet/paste with minimal water loss; least toxic (e.g., reptiles, birds, land snails, insects).
- **Nephron Types:**
  - **Cortical Nephrons** (85%)**:** Henle's loop is short and extends only very little into medulla; vasa recta absent or highly reduced.
  - **Juxtamedullary Nephrons** (15%)**:** Henle's loop is extremely long and runs deep into the inner medulla; well-developed vasa recta; **essential for concentrating urine**.

Renal Physiology: GFR & Counter-Current Mechanism

1. **Glomerular Filtration Rate (GFR):**

$$\text{Renal Blood Flow} \approx 1100 - 1200\,\text{mL/min} \quad (\approx 20 - 25\%\ \text{of Cardiac Output}) \tag{6.5}$$

$$\textbf{GFR} = \textbf{125}\,\text{mL/min} = \textbf{180}\,\text{litres/day} \implies \text{Urine output} \approx \textbf{1.5}\,\text{litres/day}\ (99\%\ \text{reabsorbed!}) \tag{6.6}$$

2. **Net Filtration Pressure (NFP):**

$$\text{NFP} = \text{GBHP} - (\text{BCOP} + \text{CHP}) = 60 - (32 + 18) = \textbf{10}\,\text{mmHg} \tag{6.7}$$

3. **Counter-Current Multiplier & Exchanger:**

- Operated between **Loop of Henle** and **Vasa Recta**.
- Flow of filtrate in two limbs of Henle's loop is in opposite directions; flow of blood in two limbs of vasa recta is also in opposite directions.
- **Descending Limb of Henle:** Permeable to water, almost impermeable to electrolytes $\implies$ filtrate becomes concentrated as it moves down ($300 \rightarrow 1200$ mOsmol/L).
- **Ascending Limb of Henle:** Impermeable to water, actively/passively transports electrolytes (NaCl) into medullary interstitium $\implies$ filtrate becomes diluted as it ascends ($1200 \rightarrow 200$ mOsmol/L).
- Creates an osmolarity gradient in medullary interstitium: 300 mOsmol/L **in cortex** $\rightarrow$ 1200 mOsmol/L **in inner medulla**.
- Human kidneys can produce urine nearly **4 times more concentrated** than the initial filtrate (1200 mOsmol/L vs 300 mOsmol/L).

Hormonal Regulation of Kidney Function

1. **RAAS (Renin-Angiotensin-Aldosterone System):**

$$\downarrow \text{GFR / Blood Pressure} \xrightarrow{\text{detected by JGA}} \text{Juxtaglomerular cells release } \textbf{Renin} \tag{6.8}$$

$$\text{Angiotensinogen (Liver)} \xrightarrow{\text{Renin}} \text{Angiotensin I} \xrightarrow{\text{ACE}} \text{Angiotensin II (Potent Vasoconstrictor)} \tag{6.9}$$

Angiotensin II increases GFR directly by arteriolar constriction AND stimulates **adrenal cortex** to secrete **Aldosterone** $\implies$ promotes active reabsorption of Na^+ and water from DCT $\implies \uparrow$ Blood Volume and BP.

2. **Antidiuretic Hormone (ADH / Vasopressin):** Released by posterior pituitary when hypothalamic osmoreceptors detect $\uparrow$ blood osmolarity (dehydration); makes DCT and Collecting Duct permeable to water via aquaporins, preventing diuresis.
3. **Atrial Natriuretic Factor (ANF):** Released by atrial walls of heart in response to $\uparrow$ blood flow/pressure; causes **vasodilation** and promotes natriuresis (Na^+ excretion). Acts as a negative feedback check on RAAS!

6.4 Locomotion & Movement: Musculoskeletal Mechanics

Sarcomere Architecture & Sliding Filament Theory

A sarcomere is the basic functional contractile unit of a muscle fiber, bounded between **two successive Z-lines**:

- **A-Band (Anisotropic):** Contains both thick (myosin) and thin (actin) filaments; length **remains completely constant** during contraction.
- **I-Band (Isotropic):** Contains only thin actin filaments; bisected by elastic **Z-line**; **shortens** during contraction.
- **H-Zone:** Central region of A-band where thick filaments are not overlapped by thin filaments;

bisected by fibrous **M-line**; **disappears** during maximal contraction.

Myofilament Biochemistry:

- **Thin Filament:** Composed of 2 helically wound polymeric **F-actin** filaments, 2 filamentous **Tropomyosin** running along F-actin grooves, and complex **Troponin** triplets (TnT, TnI, TnC). **Troponin-C (TnC)** binds Ca^{2+}, unmasking active myosin-binding sites on actin.
- **Thick Filament:** Composed of polymerized **Myosin** monomer units (**Meromyosins**):
  - **Heavy Meromyosin (HMM):** Globular head and short arm; projects outwards as cross-bridge; contains actin-binding site and active **ATPase enzyme site**.
  - **Light Meromyosin (LMM):** Tail portion.

Contraction Cycle: Action potential depolarizes sarcolemma → Sarcoplasmic reticulum releases Ca^{2+} → Ca^{2+} binds to Troponin-C → Tropomyosin shifts → Myosin head hydrolyzes ATP ($ADP+P_i$) and binds actin forming cross-bridge → **Power Stroke** pulls thin filaments towards center of A-band → New ATP binds myosin head, causing detachment.

Human Skeletal Framework (206 Bones) & Articulations

- **Axial Skeleton (80 Bones):**
  - **Skull (22 bones):** Cranial (8) + Facial (14). Associated: Hyoid (1) + Ear Ossicles ($2 \times 3 = 6$: Malleus, Incus, Stapes).
  - **Vertebral Column (26 in adult):** Cervical (7, C_1 = Atlas, C_2 = Axis), Thoracic (12), Lumbar (5), Sacral (1 fused), Coccygeal (1 fused).
  - **Thoracic Cage:** Sternum (1) + Ribs (12 pairs = 24). True ribs ($1^{st}-7^{th}$ pair, vertebrosternal), False ribs ($8^{th}, 9^{th}, 10^{th}$ pair, vertebrochondral), Floating ribs ($11^{th}, 12^{th}$ pair, vertebral).
- **Appendicular Skeleton (126 Bones):**
  - **Forelimb ($30 \times 2 = 60$):** Humerus (1), Radius (1), Ulna (1), Carpals (8 wrist), Metacarpals (5 palm), Phalanges (14 digits, formula 2,3,3,3,3).
  - **Pectoral Girdle ($2 \times 2 = 4$):** Clavicle (collar bone) + Scapula (has acromion process, spine, and glenoid cavity for head of humerus).
  - **Hindlimb ($30 \times 2 = 60$):** Femur (thigh, longest/strongest), Patella (kneecap, sesamoid bone), Tibia (1), Fibula (1), Tarsals (7 ankle), Metatarsals (5), Phalanges (14).
  - **Pelvic Girdle (2 coxal bones):** Each formed by fusion of Ilium, Ischium, and Pubis; meet ventrally at pubic symphysis (fibrous cartilage); cavity = **Acetabulum** (articulates with femur).
- **Joint Classifications:**
  - **Fibrous (Synarthroses):** Immovable (sutures between flat skull bones).
  - **Cartilaginous (Amphiarthroses):** Limited movement (pubic symphysis, between adjacent vertebrae).
  - **Synovial (Diarthroses):** Fluid-filled synovial cavity; freely movable.
    - *Ball and Socket:* Shoulder (glenoid-humerus) and Hip (acetabulum-femur).
    - *Hinge Joint:* Knee joint, Elbow joint, Interphalangeal joints.
    - *Pivot Joint:* Between Atlas and Axis ($C_1 - C_2$).
    - *Gliding Joint:* Between adjacent Carpals.
    - *Saddle Joint:* Between Carpal and Metacarpal of human thumb.

6.5 Neural Control & Chemical Coordination

Nerve Impulse: Resting, Depolarization & Repolarization

Resting Membrane Potential: $-70\,\text{mV}$ (Inside negative; permeable to K^+, impermeable to Na^+)

Sodium-Potassium Pump: Transports **3** Na^+ out for **2** K^+ in (Consumes 1 ATP)

Action Potential: $+30\,\text{mV}$ (Voltage-gated Na^+ channels open $\implies$ Na^+ influx)

Repolarization: Na^+ channels close, voltage-gated K^+ channels open $\implies$ K^+ efflux

Synapse: AP at terminal $\rightarrow$ Ca^{2+} influx $\rightarrow$ ACh exocytosis $\rightarrow$ receptors

Endocrine Master Table: Glands, Hormones, Target & Disorders

Endocrine Gland	Hormone	Nature	Major Action & Pathology
Anterior Pituitary	Growth Hormone (GH)	Peptide	Childhood excess: **Gigantism**; adult excess: **Acromegaly**; deficiency: **Pituitary Dwarfism**.
Posterior Pituitary	Oxytocin, Vasopressin	Peptide	Oxytocin: uterine contraction & milk ejection. Vasopressin: water reabsorption; deficiency $\implies$ **Diabetes Insipidus**.
Thyroid	Thyroxine (T_4, T_3)	Iodinated	BMR regulation. Pregnancy deficiency $\implies$ **Cretinism**; adult: **Myxoedema**; excess: **Graves' disease**.
Thyroid / C-cells	Thyrocalcitonin (TCT)	Peptide	**Hypocalcemic**; lowers blood Ca^{2+} via bone deposition.
Parathyroid	Parathormone (PTH)	Peptide	**Hypercalcemic**; bone resorption; antagonist to TCT.
Adrenal Cortex	Aldosterone, Cortisol	Steroid	Na^+ retention. Hyposecretion $\implies$ **Addison's disease**; hypersecretion $\implies$ **Cushing's syndrome**.
Adrenal Medulla	Adrenaline	Catecholamine	Emergency / Fight-or-Flight; $\uparrow$ heart rate, glycogenolysis.
Pancreas (β-cells)	Insulin	Peptide	Hypoglycemic; promotes cellular glucose uptake; deficiency $\implies$ **Diabetes Mellitus**.
Pancreas (α-cells)	Glucagon	Peptide	Hyperglycemic; stimulates glycogenolysis; antagonist to insulin.

Mechanism of Hormone Action: Peptide vs Steroid Pathway

- **Peptide / Protein / Amine Hormones (Water-Soluble):** Insulin, glucagon, pituitary hormones, epinephrine. Cannot cross lipid bilayer; bind to **extracellular membrane-bound receptors** $\implies$ generate second messengers (**cyclic AMP,** IP_3, Ca^{2+}) $\implies$ cascade modifies cellular biochemical metabolism.
- **Steroid & Iodothyronine Hormones (Lipid-Soluble):** Estrogen, progesterone, testosterone, cortisol, thyroxine (T_3/T_4). Cross cell membrane freely; bind to **intracellular / nuclear receptors** $\implies$ hormone-receptor complex interacts with genome/chromosome to regulate gene expression and protein synthesis.

Chapter 7

Immunology, Human Pathogens, Disease & Evolutionary Biology

This chapter explores human health, host-pathogen interactions, defense mechanisms, and the history of life on Earth. Topics include innate vs acquired immunity, antibody structure, active and passive immunization, major bacterial/viral/protozoan/helminthic diseases, oncogenesis, chemical evolution, homologous vs analogous morphological evidence, natural selection, the Hardy-Weinberg equilibrium principle, and the timeline of human evolution.

7.1 Principles of Immunology & Host Defense Mechanisms

Innate Immunity: Non-Specific First & Second Lines of Defense

Present from birth, non-specific, lacks immunological memory. Comprises four distinct anatomical and physiological barrier tiers:

1. **Physical Barriers:** Skin (keratinized stratum corneum prevents microbial ingress); Mucous coating of epithelial lining across respiratory, gastrointestinal, and urogenital tracts traps entrapped microbes.
2. **Physiological Barriers:** Gastric hydrochloric acid (pH $\approx 1.5 - 2.0$ destroys ingested bacteria); Saliva and Tears contain **lysozyme** (hydrolyzes $\beta(1 \to 4)$ glycosidic bonds in bacterial peptidoglycan cell walls).
3. **Cellular Barriers:** Phagocytes that engulf and digest foreign pathogens:
   - **PMNL** (Polymorphonuclear Leukocytes / Neutrophils): Most abundant circulating WBCs $(60 - 65\%)$.
   - **Monocytes** $(6 - 8\%)$ differentiating into tissue **Macrophages** (e.g., Kupffer cells, microglia, alveolar macrophages).
   - **Natural Killer (NK) Cells:** Specialized cytotoxic lymphocytes that destroy virus-infected and malignant tumor cells non-specifically via perforins and granzymes.
4. **Cytokine Barriers:** Virus-infected cells secrete non-viral glycoproteins called **Interferons** (α, β, γ), which bind to uninfected neighboring cells, inducing translation of antiviral proteins (AVPs) that inhibit viral translation and replication.

Acquired (Adaptive) Immunity: Humoral vs Cell-Mediated Immunity (CMI)

Characterized by antigenic specificity, diversity, discrimination between self and non-self, and immunological memory.

Property	Humoral / Antibody-Mediated Immunity (AMI)	Cell-Mediated Immunity (CMI)
Effector Cells	**B-lymphocytes** differentiating into Plasma cells (antibody factories) and Memory B cells.	**T-lymphocytes** maturing into Helper T ($CD4^+T_H$), Cytotoxic T ($CD8^+T_C$), and Regulatory T cells.
Mode of Action	Secretes soluble **Antibodies** (H_2L_2) into blood plasma, lymph, and extracellular fluids.	Direct cell-to-cell contact; T_C cells release perforins creating pores in target membranes; activates macrophages.
Target Pathogens	Extracellular bacteria, viruses in circulation, circulating venoms, toxins.	**Intracellular pathogens** (*Mycobacterium tuberculosis*, viruses inside cells, fungi, parasites) and cancerous cells.
Organ Graft Rejection	Plays a minor secondary role (hyperacute humoral rejection).	**Primary Mediator of Graft Rejection!** Allograft rejection is mediated by T_C cells recognizing non-self MHC/HLA antigens.
Immunosuppression	Intravenous immunoglobulins (IVIg), plasmapheresis.	**Cyclosporin A** (fungal metabolite from *Trichoderma polysporum*) inhibits calcineurin, suppressing T-cell activation.

Antibody Architecture (H_2L_2) & Immunoglobulin Classes

Each antibody is a Y-shaped heterodimer composed of **4 polypeptide chains**: two identical Heavy (H) chains ($\approx 50\,\text{kDa}$ each) and two identical Light (L) chains ($\approx 25\,\text{kDa}$ each), cross-linked by **interchain disulfide bonds** ($-S-S-$).

$$\text{Antibody Formula: } H_2L_2 \quad (\text{Two Heavy Chains H} + \text{Two Light Chains L}) \tag{7.1}$$

Fab (Antigen-Binding Fragment) contains Variable (V_H, V_L) domain; Fc mediates effector binding.

Class	Valency / Form	Diagnostic Role & High-Yield NCERT Facts
IgG	Monomer (Bivalent)	Most abundant ($75-80\%$ of serum Igs); **only antibody class that crosses human placenta** providing natural passive immunity to fetus.
IgA	Dimer (Tetravalent)	Secretory Ig linked by J-chain and secretory piece. Abundant in **Colostrum** (first yellowish milk), saliva, tears, respiratory mucous.
IgM	Pentamer (Decavalent)	Largest molecular weight, linked by J-chain; first antibody synthesized during primary immune response; potent complement activator.
IgE	Monomer (Bivalent)	Mediates **Type I Hypersensitivity (Allergies)**; binds Fc receptors on mast cells and basophils, triggering release of histamine and serotonin.
IgD	Monomer (Bivalent)	Co-expressed with IgM on the surface of naive mature B-cells; functions as antigen receptor initiating B-cell differentiation.

Lymphoid Organs: Primary vs Secondary Tier Architecture

- **Primary Lymphoid Organs (Generative/Central):** Sites where immature lymphocytes differentiate and proliferate into antigen-sensitive lymphocytes.
  - **Bone Marrow:** Primary site of hematopoiesis; produces all blood cells including B- and T-progenitors. B-cells mature completely here.
  - **Thymus:** Lobed organ situated near heart beneath breastbone; large at birth, undergoes progressive atrophy with age. Provides microenvironment for **T-lymphocyte maturation and thymic selection**.
- **Secondary Lymphoid Organs (Peripheral):** Sites where lymphocytes interact with antigens and proliferate into effector/memory cells.
  - **Spleen:** Large bean-shaped organ containing lymphocytes and phagocytes. Acts as a physiological blood filter trapping blood-borne microbes; serves as a massive **reservoir of erythrocytes (RBCs)**.
  - **Lymph Nodes:** Small solid structures along lymphatic vessels; filter lymph fluid by trapping micro-organisms, presenting them to resident dendritic cells and lymphocytes.
  - **MALT (Mucosa-Associated Lymphoid Tissue):** Located within mucosal lining of major tracts (digestive, respiratory, urogenital). Constitutes $\approx 50\%$ **of all lymphoid tissue in the human body!** Includes Peyer's patches (ileum) and palatine tonsils.

7.2 Major Human Pathogens, Transmission & Diagnostic Tests

Bacterial, Viral, Protozoan & Helminthic Infectious Diseases

Disease	Causative Organism	Transmission Vector	Diagnostic Pathology & NCERT Key Hallmarks
Typhoid	*Salmonella typhi* (Gram-negative bacillus)	Contaminated food and water; classic carrier: **Mary Mallon** (Typhoid Mary).	Sustained high fever ($39-40^{\circ}$C), weakness, stomach pain, constipation, headache, anorexia. **Intestinal perforation** in severe cases. Confirmatory test: **Widal Test**.
Pneumonia	*Streptococcus pneumoniae*, *Haemophilus influenzae*	Inhaling droplets/aerosols; sharing glasses/utensils.	**Infects pulmonary alveoli**; alveoli become fluid-filled leading to severe respiratory distress. Lips and fingernails turn **gray to bluish** in severe cases.
Common Cold	**Rhino viruses** (ssRNA)	Inhaling droplets; contaminated pens, books, cups.	**Infects nose and respiratory passage, but NOT lungs!** Nasal congestion, discharge, sore throat, hoarseness, cough; lasts $3-7$ days.
Amoebiasis	*Entamoeba histolytica* (Protozoan parasite)	**Houseflies** act as mechanical carriers from feces to food.	Infects large intestine; constipation, abdominal pain, cramps, stools with **excess mucous and blood clots**.
Ascariasis	*Ascaris lumbricoides* (Intestinal roundworm)	Ingestion of water, vegetables, fruits contaminated with fecal eggs.	Internal bleeding, muscular pain, fever, anemia, and mechanical **blockage of the intestinal passage**.
Filariasis (Elephantiasis)	*Wuchereria bancrofti*, *W. malayi* (Filarial worms)	Bite of infected **female *Culex*** mosquito.	Chronic inflammation of organs, especially **lymphatic vessels of lower limbs and genital organs**, resulting in gross gross deformities.
Ringworm	*Microsporum*, *Trichophyton*, *Epidermophyton*	Direct contact, sharing towels, clothes, comb; heat and moisture favor growth.	Dry, scaly, pruritic (itchy) circular lesions on skin, nails, scalp, and groin folds.

Life Cycle of *Plasmodium* (Malarial Parasite: Digenetic Cycle)

Plasmodium requires two hosts: **Primary host** = Female *Anopheles* mosquito (sexual cycle); **Secondary host** = Human (asexual cycle).

1. **Inoculation:** Female *Anopheles* mosquito injects infective stage **Sporozoites** with saliva into human blood during feeding.
2. **Hepatic Schizogony:** Sporozoites invade liver parenchymal cells, multiply asexually, and burst hepatocytes, releasing merozoites.
3. **Erythrocytic Schizogony:** Merozoites penetrate RBCs, form ring-stage trophozoites $\rightarrow$ schizonts, and burst RBCs synchronously every $48-72$ hours.
   - Rupture of RBCs releases pyrogenic toxin **Haemozoin** (formed from digested hemoglobin iron-porphyrin), producing **chills and high recurring fever**.
4. **Gametocytogenesis:** Some merozoites in RBCs develop into sexual stages: male microgametocytes and female macrogametocytes.
5. **Mosquito Phase (Sexual Cycle):** Female *Anopheles* ingests gametocytes with blood meal. Fertilization and zygote formation occur in the **mosquito gut**. Zygote becomes motile **ookinete**, penetrates gut wall to form **oocyst**. Oocyst divides by sporogony, generating thousands of sporozoites which migrate to and lodge in the **mosquito salivary glands**.

HIV/AIDS Pathogenesis & Viral Replication Mechanism

Caused by **Human Immunodeficiency Virus (HIV)**, an enveloped Lentivirus (Retroviridae) harboring **two identical single-stranded RNA (+ssRNA)** molecules and **Reverse Transcriptase**.

1. **Host Cell Attachment:** Viral envelope glycoprotein gp120 binds to CD4 **receptors** and chemokine coreceptors (CCR5/CXCR4) on macrophages and Helper T (T_H) lymphocytes.
2. **Macrophage Phase ("HIV Factory"):** Viral RNA is converted into viral DNA by Reverse Transcriptase (RNA $\rightarrow$ cDNA $\rightarrow$ dsDNA). Viral DNA integrates into host genome via **Integrase**. Macrophages produce viral particles without immediate cell lysis.
3. T_H **Cell Depletion:** Virus enters $CD4^{+}T_H$ cells, replicates, and lyses host cells. Helper T-cell count plummets from normal ($900-1200/\mu$L) to below **200**/μL.
4. **Opportunistic Infections:** Severe cell-mediated immunodeficiency permits life-threatening infections by *Mycobacterium tuberculosis*, *Toxoplasma gondii*, *Pneumocystis jirovecii*, and cytomegalovirus (CMV).

5. **Diagnosis & Management:** Screening test: **ELISA** (Enzyme-Linked Immunosorbent Assay); Confirmatory test: **Western Blot** (detects viral antigens p24, gp120, gp41). Treatment: Anti-Retroviral Therapy (**ART**) cocktail of nucleoside reverse transcriptase inhibitors (NRTIs, e.g., Zidovudine/AZT) and protease inhibitors.

Cancer Biology: Oncogenesis, Characteristics & Clinical Therapies

- **Transformation Hallmarks:** Normal cells exhibit **contact inhibition** (contact with neighboring cells inhibits uncontrolled proliferation). Cancer cells lose contact inhibition, proliferating relentlessly into **tumors**:
  - **Benign Tumors:** Encapsulated, remain localized to primary site, do not invade adjacent tissues.
  - **Malignant Tumors:** Unencapsulated masses of neoplastic cells exhibiting **Metastasis** (cells slough off into circulation and colonize distant organs; **the most feared property of malignant neoplasms**).
- **Etiological Carcinogens:**
  - **Physical:** Ionizing radiation (X-rays, γ-rays break DNA strands); Non-ionizing radiation (**UV-B** induces cyclobutane pyrimidine/thymine dimers).
  - **Chemical:** Tobacco smoke contains ***N*-nitrosamines** and benzo[*a*]pyrene (major cause of bronchogenic lung carcinoma).
  - **Biological:** Oncogenic viruses carrying viral oncogenes (*v-onc*); cellular proto-oncogenes (*c-onc*) mutated into active oncogenes.
- **Therapeutic Modalities:** Surgery (tumor excision); Radiotherapy (targeted lethal irradiation of malignant tissue); Chemotherapy (antimitotic drugs like Vincristine, Vinblastine from *Catharanthus roseus*); **Immunotherapy:** Biological response modifiers like **α-Interferon** activate the patient's immune system to destroy tumors.

Psychoactive Drugs, Hallucinogens & Addiction Pharmacology

Class	Representative Drugs	Receptors / Site	Pharmacological Action & Botanical Extraction
Opioids	Morphine, Heroin (**Smack** / Diacetylmorphine), Codeine	Specific opioid receptors in **CNS and GIT**.	Extracted from latex of unripe capsules of ***Papaver somniferum*** (Opium poppy). Heroin is synthesized by **acetylation of morphine**; white, odorless, bitter crystalline compound. Acts as a **depressant and slows down body functions**.
Cannabinoids	Marijuana, Hashish, Charas, Ganja	Cannabinoid receptors primarily located in the **brain**.	Obtained from inflorescences, leaves, and resin of ***Cannabis sativa*** (Hemp plant). Inhaled or orally ingested; induces relaxation, alters sensory perception, and severely impacts **cardiovascular system**.
Coca Alkaloids	**Cocaine** (Coke / Crack)	Interferes with reuptake transport of **Dopamine**.	Extracted from leaves of South American shrub ***Erythroxylum coca***. Potent CNS stimulant producing euphoria and increased energy; high doses induce **hallucinations**. Other hallucinogens: *Atropa belladonna*, *Datura*.
Tobacco / Nicotine	Nicotine (Alkaloid)	Stimulates nicotinic acetylcholine receptors.	Stimulates **adrenal medulla** to release adrenaline and noradrenaline into circulation, raising blood pressure and heart rate. Induces lung cancer, emphysema, bronchitis; elevates carbon monoxide (CO) in blood, decreasing oxyhemoglobin concentration.

7.3 Evolutionary Biology: Origins, Evidence & Mechanisms

Origin of Life: Abiogenesis & Miller-Urey Experimental Proof

- **Primordial Earth Atmosphere:** Formed $\approx$ 4.5 billion years ago. Primitive atmosphere was strictly **reducing**: no free O_2, rich in CH_4, NH_3, H_2O (vapor), H_2.
- **Chemical Evolution (Oparin-Haldane Hypothesis):** First forms of life originated from pre-existing non-living organic molecules (RNA, proteins, sugars), preceding biological evolution.
- **Miller-Urey Spark-Discharge Experiment (1953):** Stanley Miller simulated prebiotic atmospheric conditions:

- Closed spark-discharge flask containing gases: $CH_4 : NH_3 : H_2$ in ratio **2 : 1 : 2** along with continuously boiling water vapor (H_2O).
- Tungsten electrodes produced continuous electrical discharge ($\approx 75,000\,V$) maintaining chamber temperature at **800**°C.
- **Experimental Yield:** Condensed effluent contained amino acids: **Glycine, α-Alanine, and Aspartic acid**, conclusively validating chemical evolution.

Comparative Morphological Evidences: Homology vs Analogy

Parameter	Homologous Organs (Divergent Evolution)	Analogous Organs (Convergent Evolution)
Fundamental Basis	**Same anatomical origin and basic structure**, modified to perform different functions in response to different adaptations.	**Different anatomical origin and basic structure**, modified to perform similar functions due to similar selective pressures.
Evolutionary Pattern	**Divergent Evolution** (Radiates from a common ancestor into diverse ecological niches).	**Convergent Evolution** (Unrelated lineages evolve functionally convergent phenotypes).
Vertebrate Forelimb Examples	Forelimb skeletal architecture of **Whale (flipper), Bat (patagium), Cheetah (running), and Human (grasping)** shares identical arrangement: humerus, radius, ulna, carpals, metacarpals, phalanges.	**Wings of Butterfly** (chitinous membranous folds) vs **Wings of Birds** (feather-covered forelimb bones).
Sensory / Functional Organs	Vertebrate heart and brain anatomy across fish, amphibians, reptiles, birds, and mammals.	**Eye of Octopus** (invertebrate retina derived from skin) vs **Eye of Mammals** (derived from embryonic diencephalon). **Flippers of Penguins (bird)** vs **Dolphins (mammal)**.
Botanical Examples	**Thorns of *Bougainvillea*** (axillary bud protection) vs **Tendrils of *Cucurbita*** (axillary bud climbing).	**Sweet Potato** (root tuber modification) vs **Potato** (underground stem tuber modification).

Adaptive Radiation & Convergent Marsupial-Placental Counterparts

Adaptive Radiation: Evolution of divergent species starting from a common ancestral point and radiating into distinct geographical habitats and ecological niches.

- **Classic Examples:** Darwin's Finches on Galapagos archipelago (ancestral seed-eating beak diversified into vegetarian, insectivorous, warbler, and cactus-puncturing beaks); Australian Marsupials radiating from a single ancestral stock within the isolated island continent.
- **Marsupial vs Placental Mammal Convergence:** When adaptive radiation occurs in isolated continents for different lineages occupying similar ecological niches, convergent evolution generates striking morphological analogs:

Ecological Niche	Australian Marsupial Counterpart	Placental Mammal Counterpart
Fossorial Burrower	Marsupial mole (*Notoryctes*)	Placental Mole
Myrmecophage (Ant-eater)	Numbat / Banded anteater (*Myrmecobius*)	Anteater
Small Glider	Flying phalanger / Sugar glider (*Petaurus*)	Flying squirrel
Feline Carnivore	Tasmanian tiger cat (*Dasyurus*)	Bobcat
Apex Carnivore	Tasmanian wolf (*Thylacinus*)	Wolf (*Canis lupus*)

Hardy-Weinberg Equilibrium: Mathematical Formulations & Selection Dynamics

In a large, randomly mating, panmictic diploid population with no evolutionary disruptive forces, allele frequencies and genotype frequencies remain constant across generations:

$$\mathbf{p + q = 1} \qquad \Longrightarrow \qquad \mathbf{p^2 + 2pq + q^2 = 1} \tag{7.2}$$

Where p = frequency of dominant allele A, q = frequency of recessive allele a, p^2 = frequency of homozygous dominant AA, $2pq$ = heterozygous Aa, and q^2 = homozygous recessive aa.

- **Five Evolutionary Forces Disrupting Equilibrium:** (1) Gene migration / Gene flow; (2) **Genetic Drift** (random allele frequency fluctuation in small populations, including **Founder Effect** and **Bottleneck Effect**); (3) Mutation; (4) Genetic Recombination during meiosis; (5) Natural Selection.
- **Three Modes of Natural Selection on Phenotypic Traits:**
  1. **Stabilizing Selection:** Favors intermediate phenotype ($2pq$); phenotypic curve becomes narrower and taller (e.g., human infant birth weight centered at $3 - 3.5\,kg$).
  2. **Directional Selection:** Favors one extreme phenotype; shifts population mean towards one direction (e.g., Industrial Melanism in *Biston betularia*, antibiotic resistance in bacteria).
  3. **Disruptive Selection:** Favors both extreme phenotypes over intermediate; produces a bi-

modal, two-peaked curve.

Chronological Geological Timeline of Human Evolution

Ancestor Stage	Geological Era	Fossil Provenance	Cranial Capacity	Anatomical & Cultural Characteristics
Dryopithecus	$\approx$ 15 mya (Miocene)	Africa and Eurasia	$\approx$ 350 – 400 cc	Arboreal, hairy, arms longer than legs; walked quadrupedally like modern gorillas and chimpanzees.
Ramapithecus	$\approx$ 15 mya (Miocene)	Shivalik hills (India)	$\approx$ 400 – 450 cc	More man-like than *Dryopithecus*; upright posture tendency; walked semi-bipedally.
Australopithecus	$\approx$ 2 – 3 mya (Pliocene)	East African grasslands	$\approx$ 450 – 600 cc	**Bipedal locomotion**, upright posture; hunted with stone weapons but **essentially ate fruit**.
Homo habilis	$\approx$ 1.7 – 2 mya (Pleistocene)	Olduvai Gorge (Tanzania)	**650 – 800** cc	**First human-like hominid** ("Tool maker" / Handy man); constructed crude stone tools; **did not eat meat**.
Homo erectus	$\approx$ 1.5 mya (Pleistocene)	Java (1891 fossils)	**900** cc	Fully erect posture; proficient toolmaker; first to master **fire**; **ate meat**.
Homo neanderthalensis	100,000 – 40,000 ya	Neander Valley (Germany)	**1400** cc	Heavy brow ridges, receded chin; **used animal hides to protect bodies; buried their dead with rituals**.
Homo sapiens	75,000 – 10,000 ya	Emerged in Africa	**1350 – 1450** cc	Modern man; migrated across continents; prehistoric cave art ($\approx$ 18,000 ya at Bhimbetka); **agriculture began** $\approx$ 10,000 **ya**.

High-Frequency NEET Assertion-Reason Traps in Health & Evolution

- **Rhino Virus vs Streptococcus:** Rhino viruses infect the **upper respiratory tract and nasal passage, strictly sparing the pulmonary alveoli**; *Streptococcus pneumoniae* infects the alveoli causing fluid accumulation and cyanosis.
- **Malarial Host Nomenclature:** Sexual reproduction occurs **only in the female *Anopheles* mosquito** (the definitive/primary host). Asexual multiple fission occurs in humans (intermediate host).
- **Morphine vs Heroin:** Morphine is a natural alkaloid; Heroin is a **semi-synthetic derivative produced by acetylation of morphine**.
- **Darwin vs de Vries on Variation:** Darwinian variations are **small, continuous, and directional**; de Vriesian mutations are **single-step large mutations (Saltation), random, and directionless**.
- **Meat Consumption in Hominids:** *Homo habilis* **did not eat meat**; *Homo erectus* **ate meat**!

PART III

High-Yield Quantitative Biology & Diagnostic Matrices

Chapter 8

Quantitative Biology, Genetic Formulations & High-Yield Diagnostic Matrices

To achieve a perfect score of 360 in NEET Biology, students must master both qualitative concepts and quantitative formulations. This chapter provides a unified analytical compendium of genetic probability calculations, recombination frequencies, population genetics models, physiological capacity values, floral formula matrices for all seven prescribed plant families, and high-frequency Assertion-Reasoning anchors.

A Personal Note from Satya: The Stethoscope Waiting at the End of This Tunnel

You are in the final stretch of this master book. Take a moment to remember why you began.
You did not choose this path because it was easy. You chose medicine because deep down, you want to be the person who walks into an emergency room, keeps a cool head when everyone else is panicking, and literally saves a human life.
Right now, in the quiet of your room, you are solving dihybrid crosses, calculating Hardy-Weinberg allele frequencies, and memorizing floral formulas of Compositae and Gramineae. It might feel tedious. It might feel like a never-ending grind. But every single page you conquer is forging the clinical stamina and analytical precision you will need when diagnosing real patients five years from now.
When the days get tough and the mock exam scores fluctuate, do not lose heart. Medicine tests your resilience long before you ever step into a hospital ward. Keep showing up. Keep pushing through the doubt. That white coat with your name stitched onto it, and that stethoscope draped around your neck, is waiting for you. Make yourself proud!

8.1 Quantitative Genetics & Probability Calculus

Foundational Mendelian Probability Principles

Mendelian inheritance patterns obey the fundamental laws of probability:

1. **Rule of Multiplication (Product Rule):** The probability of two independent events occurring simultaneously is the product of their individual probabilities:

$$P(A \cap B) = P(A) \times P(B) \tag{8.1}$$

   Application: Probability of obtaining a homozygous recessive genotype ($aabb$) from a dihybrid cross ($AaBb \times AaBb$) is $P(aa) \times P(bb) = \frac{1}{4} \times \frac{1}{4} = \frac{1}{16}$.
2. **Rule of Addition (Sum Rule):** The probability of either of two mutually exclusive events occurring is the sum of their individual probabilities:

$$P(A \cup B) = P(A) + P(B) \tag{8.2}$$

   Application: In a monohybrid cross ($Aa \times Aa$), the probability of obtaining a heterozygote (Aa) via either maternal or paternal allele donation is $\frac{1}{4} + \frac{1}{4} = \frac{1}{2}$.

Master Combinatorial Formulas for Multi-Gene Crosses

For a cross involving n independently assorting heterozygous gene pairs ($AaBbCc\ldots$):

$$\text{Number of Genetically Distinct Gametes: } N_{\text{gametes}} = 2^n \tag{8.3}$$

$$\text{Total Number of Offspring Genotypes: } N_{\text{genotypes}} = 3^n \tag{8.4}$$

$$\text{Total Number of Offspring Phenotypes: } N_{\text{phenotypes}} = 2^n \quad \text{(assuming complete dominance)} \tag{8.5}$$

$$\text{Number of Squares in Punnett Grid: } N_{\text{grid}} = 2^n \times 2^n = 4^n \tag{8.6}$$

$$\text{Fraction of Completely Homozygous Recessive: } F_{\text{recessive}} = \left(\frac{1}{4}\right)^n \tag{8.7}$$

$$\text{Fraction of Completely Homozygous Dominant: } F_{\text{dominant}} = \left(\frac{1}{4}\right)^n \tag{8.8}$$

$$\text{Fraction of Completely Heterozygous Individuals: } F_{\text{hetero}} = \left(\frac{1}{2}\right)^n \tag{8.9}$$

Chi-Square (χ^2) Goodness-of-Fit Test in Genetic Analysis

To determine whether observed offspring numbers deviate significantly from theoretical Mendelian expectations:

$$\chi^2 = \sum \frac{(O-E)^2}{E} \tag{8.10}$$

where O is the observed frequency, E is the expected frequency based on Mendelian ratio, and degrees of freedom $df = k-1$ (where k is the number of phenotypic classes). If $\chi^2_{\text{calc}} < \chi^2_{\text{critical}}$ at $p = 0.05$, the observed deviation is attributed strictly to random chance and the Mendelian hypothesis is accepted.

8.2 Recombination Frequency & Chromosome Mapping Calculus

Morgan-Sturtevant Gene Mapping Formulations

The recombination frequency (RF) between two linked genes on a chromosome is directly proportional to the physical distance between them:

$$RF = \left(\frac{\text{Total Number of Recombinant Offspring}}{\text{Total Number of Offspring Progeny}}\right) \times 100\% \tag{8.11}$$

Units & Metric Standards:

- 1% Recombination Frequency = 1 Map Unit (m.u.) = 1 centimorgan (cM).
- Maximum observable recombination frequency for two genes on the same chromosome is **50**% (at which point they assort independently, indistinguishable from unlinked genes on different chromosomes).

Three-Point Test Cross & Interference Formulations:

$$\text{Coefficient of Coincidence } (C) = \frac{\text{Observed Frequency of Double Crossovers}}{\text{Expected Frequency of Double Crossovers}} \tag{8.12}$$

$$\text{Interference } (I) = 1 - C \tag{8.13}$$

Interpretation: If $I > 0$ (positive interference), a crossover in one interval suppresses crossover formation in an adjacent chromosomal region. If $I = 1$, double crossovers are completely suppressed.

8.3 Population Genetics: Hardy-Weinberg Mathematical Model

Hardy-Weinberg Equilibrium Equations

For a diploid population with two alleles (A and a) at a single locus with frequencies p and q:

$$\text{Allelic Frequency Identity: } p + q = 1 \tag{8.14}$$

$$\text{Genotypic Frequency Expansion: } (p+q)^2 = p^2 + 2pq + q^2 = 1 \tag{8.15}$$

where:

- p^2: Frequency of homozygous dominant individuals (AA).
- $2pq$: Frequency of heterozygous carrier individuals (Aa).
- q^2: Frequency of homozygous recessive affected individuals (aa).

NEET Trap: Allelic Frequency vs Genotypic Frequency

Pay meticulous attention to the wording in Hardy-Weinberg exam questions:

- If given "frequency of recessive **allele**": that is q.
- If given "frequency of recessive **disease / phenotype / individuals**": that is q^2. You must take the square root $\sqrt{q^2} = q$ first, then find $p = 1 - q$, and finally compute carrier frequency $2pq$.
- **Equilibrium Preconditions:** Strictly valid only for: (1) large population size, (2) random mating, (3) no gene flow / migration, (4) no mutation, and (5) no natural selection.

8.4 Quantitative Human Physiology Matrices

Physiological Parameter	Standard Value	Mathematical Definition / Formula
Tidal Volume (TV)	500 mL	Volume inspired or expired during normal breathing
Inspiratory Reserve Volume (IRV)	2500–3000 mL	Extra volume inspired forcibly above TV
Expiratory Reserve Volume (ERV)	1000–1100 mL	Extra volume expired forcibly after normal expiration
Residual Volume (RV)	1100–1200 mL	Gas remaining in lungs after maximal expiration
Vital Capacity (VC)	3500–4500 mL	VC = TV + IRV + ERV
Total Lung Capacity (TLC)	5000–6000 mL	TLC = VC + RV = TV + IRV + ERV + RV
Cardiac Output (CO)	$\approx$ 5000 mL/min	CO = Stroke Volume (70 mL) $\times$ Heart Rate (72 bpm)
Glomerular Filtration Rate (GFR)	125 mL/min	Equivalent to 180 L/day across both kidneys
Net Filtration Pressure (NFP)	10 mmHg	NFP = GHP (60) − [BCOP (32) + CHP (18)]
Urine Excretion Rate	1.0–1.5 L/day	$>$ 99% of filtrate is reabsorbed by renal tubules

Table 8.1: Master Diagnostic Matrix: Quantitative Human Physiological Constants.

8.5 Master Systematic Comparison of All 7 Prescribed Plant Families

Family	Floral Formula	Androecium	Gynoecium	Key Diagnostic Examples
Malvaceae	$\oplus \text{Epi}_{3-8}\text{K}_{(5)}\text{C}_5\text{A}_{(\infty)}\underline{\text{G}}_{(5-\infty)}$	Monadelphous stamens, monothecous reniform anthers	Syncarpous, superior, axile placentation	*Hibiscus*, Cotton (*Gossypium*), Lady's finger (*Abelmoschus*)
Cruciferae	$\oplus\ \text{K}_{2+2}\ \text{C}_4\ \text{A}_{2+4}\ \underline{\text{G}}_{(2)}$	Tetradynamous stamens (2 short outer + 4 long inner)	Bicarpellary, unilocular becoming bilocular via false septum (replum)	Mustard (*Brassica*), Radish (*Raphanus*), Turnip
Fabaceae	$\%\text{K}_{(5)}\text{C}_{1+2+(2)}\text{A}_{(9)+1}\underline{\text{G}}_1$	Diadelphous stamens $[(9)+1]$, dithecous	Monocarpellary, superior, marginal placentation	Gram, Pea, Soybean, Sunnhemp, Lupin, *Sesbania*
Asteraceae	$\oplus$ or $\%\text{K}_{\text{pappus}}\text{C}_{(5)}\text{A}_{(5)}\overline{\text{G}}_{(2)}$	Syngenesious (anthers fused, filaments free)	Bicarpellary, inferior, basal placentation, single ovule	Sunflower (*Helianthus*), Marigold, *Taraxacum*
Poaceae	$\%\ \text{P}_2\ \text{A}_{3\ \text{or}\ 6}\ \underline{\text{G}}_1$	Stamens versatile, pendulous	Monocarpellary, superior, basal placentation, feathery stigma	Wheat (*Triticum*), Rice, Maize, Bamboo, Sugarcane
Solanaceae	$\oplus\ \text{K}_{(5)}\ \text{C}_{(5)}\ \text{A}_5\ \underline{\text{G}}_{(2)}$	Epipetalous stamens, alternipetalous	Bicarpellary, syncarpous, superior, obliquely placed ovary, swollen placenta	Potato, Tomato, Brinjal, Chilli, Belladonna, Ashwagandha, Tobacco
Liliaceae	$\oplus\ \text{P}_{(3+3)}\ \text{A}_{3+3}\ \underline{\text{G}}_{(3)}$	Epiphyllous stamens in two whorls	Tricarpellary, syncarpous, superior, axile placentation	*Allium* (Onion, Garlic), *Aloe*, *Asparagus*, *Colchicum autumnale*

Table 8.2: Systematic Comparative Diagnostic Matrix for the 7 NMC-Prescribed Angiosperm Families.

Direct NCERT High-Yield Assertion-Reason Anchors

- **Floral Formulas as Definitive Identifiers:** In Solanaceae, the ovary is characteristically **obliquely placed** with a swollen placenta bearing many ovules; in Cruciferae, the ovary becomes bilocular due to a false septum called **replum**.
- **Staminal Adaptations:** Diadelphous stamens $[(9)+1]$ are unique to Fabaceae; monadelphous stamens with monothecous anthers are diagnostic of Malvaceae; tetradynamous stamens $(2+4)$ are exclusive to Cruciferae; and syngenesious stamens are hallmark to Asteraceae.
- **Placentation Anchor:** Marginal (Fabaceae/Pea), Parietal with replum (Cruciferae/Mustard), Axile (Solanaceae, Malvaceae, Liliaceae), Basal (Asteraceae/Sunflower, Poaceae/Wheat).

About the Author & Errata Submissions

Satya Prakash is an educator, curriculum researcher, and author dedicated to creating rigorous, student-first learning resources for competitive examinations in India. His focus is on eliminating cognitive overload, dismantling obsolete legacy syllabi, and restoring conceptual rigor and mathematical transparency to competitive pre-medical preparation.

Official Project Repository & Continuous Improvement

Science education thrives on continuous scrutiny and collective refinement. If you identify a typo, biological discrepancy, or pedagogical ambiguity, or if you wish to suggest an enhancement for future editions, please submit your feedback:

- **Direct Author Email:** spdedsec@hotmail.com
- **GitHub Open-Source Repository:** `https://github.com/spdedsec/neet-study-materials`
- **Errata Tracker:** Verified corrections and updates are posted directly to the project repository.

Authored by Satya Prakash · All Rights Reserved