NEET UG CHEMISTRY

MASTER TEXTBOOK

Complete Master Theory, Formulas & Experimental Compendium

Strictly Aligned with NEET UG 2027 NMC Syllabus

Satya Prakash

NEET UG CHEMISTRY

MASTER TEXTBOOK

Complete Master Theory, Formulas & Experimental Compendium

Strictly Aligned with NEET UG 2027 NMC Syllabus

Authored by

Satya Prakash

Strictly Calibrated to the Revised NMC Syllabus & Rationalized NCERT Textbooks

Target Examination: National Eligibility cum Entrance Test (NEET UG 2027)

Published for Independent Medical Aspirants, Educators & Academic Study Circles

Comprehensive Academic Edition · Global Print-on-Demand

NEET UG CHEMISTRY MASTER TEXTBOOK

Complete Master Theory, Formulas & Experimental Compendium strictly aligned with NEET UG 2027 NMC Syllabus

Author: Satya Prakash
Editorial Correspondence & Academic Feedback: spdedsec@hotmail.com
Source Repository & Official Errata:
`https://github.com/spdedsec/neet-study-materials`

© 2026–2027 Satya Prakash. All rights reserved.

First Edition: October 2026
Publishing Imprint: Independent Academic Publishing / Satya Prakash Editorial Imprint
Target Examination Cycle: NEET UG 2027 (Undergraduate Medical Admissions Entrance Examination)
Publication Format: Print on Demand Professional Two-Sided Edition
ISBN-13 (Paperback): 979-8-1626-6110-8
ISBN-13 (Hardcover): 979-8-1626-6110-8 (Reference Library Edition)

All Rights Reserved. No part of this publication may be reproduced, stored in a retrieval system, or transmitted in any form or by any means—electronic, mechanical, photocopying, recording, scanning, or otherwise—without the prior written permission of the author, except for brief quotations embodied in critical reviews or scholarly analysis as permitted under copyright law.

Academic & Legal Disclaimer for Medical Entrance Examination Preparation: This textbook is an independent preparation manual authored by Satya Prakash to assist candidates preparing for the National Eligibility cum Entrance Test (NEET UG). While every effort has been made to verify scientific accuracy and strict concordance with the National Medical Commission (NMC) curriculum and rationalized NCERT textbooks, the author and publisher make no official claims of affiliation with or endorsement by the National Testing Agency (NTA), the National Medical Commission (NMC), or the Ministry of Education, Government of India. The author disclaims any liability for errors, omissions, or examination outcomes.

Typesetting & Typography: Typeset in LaTeX 2_ε with the Tectonic XeTeX engine using Latin Modern typefaces and custom academic publishing geometry. Chemical nomenclature, SI units, and IUPAC rules strictly observed. Produced and bound under rigorous academic publishing standards.

Printed in the United States of America / United Kingdom / India.

To my parents, whose boundless love and quiet sacrifices gave me the courage to pursue knowledge;

To the chemistry teachers who revealed that reactions are not random rules to be memorized,
but the logical dance of electrons seeking thermodynamic stability;

And to every medical aspirant battling balancing equations, acid-base equilibria, and organic mechanisms—
may this book be your clearest guide, turning what felt overwhelming into your greatest competitive strength.

Acknowledgements

This comprehensive chemistry textbook reflects years of academic analysis, teaching, and systematic alignment with national medical entrance examinations. I wish to express my profound gratitude to all who contributed to its realization.

I thank my family for their unending patience, love, and constant moral support during the intensive writing and typesetting process.

I am deeply grateful to the chemistry educators and academic colleagues whose rigorous insights helped shape the structural framework of this book. Their insistence on clear mechanistic arrow-pushing, thermodynamic sign discipline, and crystal field theory clarity has elevated the quality of every discussion.

I owe special thanks to the students whose candid feedback and test error sheets provided direct insight into the most persistent conceptual pitfalls in competitive chemistry. The "NEET Trap" callouts throughout these pages are a direct response to those common stumbling blocks.

Finally, I express my sincere appreciation to the LaTeX 2_ε open-source community for creating typographical tools that allow complex chemical equations, tabular reaction matrices, and color-coded callouts to be typeset with uncompromised aesthetic and mathematical precision.

Satya Prakash
October 2026

Author's Preface: Pedagogical Architecture & Analytical Methodology

Mastering chemistry for the National Eligibility cum Entrance Test (NEET UG) requires a decisive transition from passive memorization to analytical problem-solving and mechanistic reasoning.

In competitive medical entrance preparation, candidates often encounter two persistent pedagogical pitfalls. First, many treat chemistry as an encyclopedic catalog of isolated reactions, memorizing reagents and formula sheets without internalizing electron density distributions, stereochemical principles, or thermodynamic driving forces. Consequently, when examination items present novel multi-step transformations, stereochemical configurations, or Assertion-Reasoning pairings, memorized patterns fail.

Second, substantial study time is frequently misallocated to legacy coaching material covering topics eliminated under the rationalized curriculum issued by the National Medical Commission (NMC). Nine former units—Solid State, States of Matter, Surface Chemistry, Hydrogen, the *s*-Block Elements, Metallurgy, Polymers, Chemistry in Everyday Life, and Environmental Chemistry—have been formally excised from the syllabus. This curricular pruning is not a reduction in examination rigor; rather, it allows the examination to test substantially deeper conceptual comprehension across the retained foundational domains: Chemical Bonding, Coordination Chemistry, Reaction Intermediates, Carbonyl Chemistry, Thermodynamics, and Ionic Equilibria.

This master textbook is constructed to provide an uncompromisingly rigorous, self-contained, and syllabus-calibrated pedagogical framework for NEET UG 2027:

1. **First-Principles Physical Chemistry:** Every governing formulation—from de Broglie wave mechanics and Schrödinger orbital definitions to the integrated rate laws, Kohlrausch conductivity, and the Nernst equation—is derived algebraically with rigorous thermodynamic sign discipline and dimensional consistency.
2. **Mechanistic Organic Transformations:** Organic chemistry is presented through the physical movement of electron pairs. Part V provides a dedicated handbook of curved-arrow reaction mechanisms, arenium ion intermediates, transition states, and carbocation rearrangements spanning all canonical reaction classes.
3. **Inorganic Chemistry Grounded in Electronic Structure:** Rather than relying on disconnected factual recall, inorganic behavior across the *p*-block, *d*-block, and coordination complexes is deduced systematically from

effective nuclear charge (Z_{eff}), shielding parameters, crystal field splitting energy (Δ_o and Δ_t), and stereochemical geometries.

4. **Laboratory Skills & Analytical Protocols:** In strict accordance with the updated practical chemistry mandate, Part VI provides comprehensive protocols for systematic cation group separation (Groups 0–VI), confirmatory tests for acidic radicals, and quantitative redox titrimetry.
5. **Integrated Tactical Callout Architecture:** Five distinct callout environments—`conceptbox`, `formulabox`, `usecasebox`, `trapbox`, and `ncertbox`—demarcate fundamental principles, valid boundary conditions, and the high-frequency pitfalls where aspirants routinely concede negative marks.

Candidates are strongly advised to engage actively with paper and pen: verify every mathematical derivation, trace each curved arrow through its transition state, and balance redox equations systematically by the ion-electron method. Conceptual clarity achieved through active derivation is the most durable asset in competitive examination environments.

Satya Prakash
Author & Dedicated Mentor
spdedsec@hotmail.com
October 2026

Syllabus Blueprint & NMC Rationalization Guide

This section details the official NEET UG 2027 Chemistry curriculum prescribed by the National Medical Commission (NMC), highlighting the examination scheme, unit distribution, and critical syllabus exclusions.

Official Chemistry Examination Architecture

Component	Compulsory Qs	Marks	Pace Budget	Question Typology
Chemistry	45 (All Compulsory)	180	40–45 min	Single Choice MCQs, Reaction Sequences,
Exam Total	180 Compulsory	720	180 min (3h)	Assertion-Reason, Statement I/II, Matrix Match

Table 1: Reformed NEET UG Chemistry Examination Scheme (+4 for correct response, −1 for incorrect response, 0 for unattempted. Optional Section B discontinued by NTA).

Official 20-Unit Chemistry Syllabus Structure

1. **Unit 1: Some Basic Concepts in Chemistry** (Matter and laws of chemical combination, mole concept, molar mass, stoichiometry and stoichiometric calculations).
2. **Unit 2: Atomic Structure** (Bohr model and limitations, dual nature, de Broglie relation, Heisenberg uncertainty, quantum mechanical model, orbitals, quantum numbers, Hund's rule, Pauli exclusion, Aufbau principle).
3. **Unit 3: Chemical Bonding and Molecular Structure** (Ionic bonding, lattice energy, Born-Haber cycle, covalent bond, VSEPR theory, hybridization, dipole moment, MOT for homonuclear diatomic molecules, hydrogen bonding).
4. **Unit 4: Chemical Thermodynamics** (First law, enthalpy, Hess's law, bond dissociation enthalpy, spontaneous processes, entropy, Gibbs free energy criterion for spontaneity).
5. **Unit 5: Solutions** (Types of solutions, Raoult's law, colligative properties: lowering of vapor pressure, elevation of boiling point, depression of freezing point, osmotic pressure, van 't Hoff factor).

6. **Unit 6: Equilibrium** (Law of mass action, K_p and K_c, Le Chatelier's principle, ionic equilibrium: ionization of acids/bases, pH scale, common ion effect, buffer solutions, solubility product K_{sp}).
7. **Unit 7: Redox Reactions and Electrochemistry** (Oxidation states, balancing redox, conductance in electrolytic solutions, Kohlrausch's law, electrochemical cells, EMF, Nernst equation, Gibbs energy and EMF).
8. **Unit 8: Chemical Kinetics** (Rate of reaction, rate law, order and molecularity, integrated rate equations for zero and first order, half-life, Arrhenius equation, activation energy).
9. **Unit 9: Classification of Elements and Periodicity** (Modern periodic law, periodic trends: atomic/ionic radii, ionization enthalpy, electron gain enthalpy, electronegativity, valence).
10. **Unit 10: p-Block Elements** (General electronic configuration and periodic trends across Groups 13 through 18, comparative chemistry, anomalous properties of first element).
11. **Unit 11: d- and f-Block Elements** (Transition elements: electronic configuration, oxidation states, catalytic properties, magnetic properties, colored ions, lanthanoids and actinoids, lanthanoid contraction).
12. **Unit 12: Coordination Compounds** (Werner's theory, IUPAC nomenclature, isomerism, Valence Bond Theory, Crystal Field Theory, spectrochemical series, stability and biological significance).
13. **Unit 13: Purification and Characterization of Organic Compounds** (Qualitative and quantitative analysis: Lassaigne's test, Dumas, Kjeldahl, Carius methods).
14. **Unit 14: Some Basic Principles of Organic Chemistry** (IUPAC nomenclature, inductive, electromeric, resonance and hyperconjugation effects, reactive intermediates: carbocations, carbanions, free radicals).
15. **Unit 15: Hydrocarbons** (Alkanes: conformation of ethane; Alkenes: geometrical isomerism, Markovnikov/anti-Markovnikov addition; Alkynes: acidity, addition; Arenes: aromaticity, EAS mechanisms).
16. **Unit 16: Organic Compounds Containing Halogens** (Haloalkanes and haloarenes: S_N1 and S_N2 mechanisms, environmental effects of polyhalogen compounds).
17. **Unit 17: Organic Compounds Containing Oxygen** (Alcohols, phenols and ethers: Lucas test, Kolbe reaction, Reimer-Tiemann, Williamson synthesis; Aldehydes, ketones and carboxylic acids: nucleophilic addition, aldol condensation, Cannizzaro, Hell-Volhard-Zelinsky).
18. **Unit 18: Organic Compounds Containing Nitrogen** (Amines: basicity, Hinsberg test, carbylamine test; Diazonium salts: synthetic applications).
19. **Unit 19: Biomolecules** (Carbohydrates: aldoses, ketoses, glucose, fructose, glycosidic bond; Proteins: peptide bond, primary/secondary/tertiary structures, denaturation; Nucleic acids: DNA, RNA).
20. **Unit 20: Principles Related to Practical Chemistry** (Qualitative salt

analysis: cations and anions; Volumetric titrations: oxalic acid vs $KMnO_4$, Mohr's salt vs $KMnO_4$).

Zero-Tolerance NMC Exclusions: The 9 Deleted Chemistry Chapters

Under revised NMC regulations, the following **nine legacy chapters are completely deleted** from the NEET examination:

1. **Solid State** (Unit cells, packing efficiency, point defects).
2. **States of Matter (Gaseous & Liquid Bulk States)** (Ideal gas equations remain in Unit 1; van der Waals equation and critical phenomena are eliminated).
3. **Surface Chemistry** (Adsorption, colloids, micelles, catalysts).
4. **General Principles of Metallurgy** (Extraction of metals, Ellingham diagrams).
5. **Hydrogen** (Hydrides, heavy water, hydrogen peroxide).
6. **s-Block Elements** (Alkali and alkaline earth metals).
7. **Polymers** (Classification, polymerization reactions, plastics).
8. **Chemistry in Everyday Life** (Drugs, medicines, soaps, detergents).
9. **Environmental Chemistry** (Pollution, acid rain, ozone depletion).

Study Directive: Do not invest study hours in these eliminated topics. All cognitive energy must focus on the rigorously tested Class 11 and Class 12 units above.

Empirical Question Weightage Matrix

Chemistry Domain	Typical Qs	Marks Weightage
Physical Chemistry (Mole, Atomic, Thermo, Equilibrium, Solutions, Electro, Kinetics)	15–17	60–68 Marks
Inorganic Chemistry (Periodicity, Bonding, Coordination, *p*-Block, *d*-&*f*-Block)	13–15	52–60 Marks
Organic Chemistry (GOC, Hydrocarbons, Haloalkanes, Oxygen, Nitrogen, Biomolecules)	14–16	56–64 Marks
Practical Chemistry (Qualitative Salt Analysis & Volumetric Redox Titrations)	2–3	8–12 Marks
Total Chemistry Paper	**45 Qs (Compulsory)**	**180 Marks**

Table 2: Empirical Question Distribution in Recent NEET UG Papers.

Contents

Acknowledgements **ix**

Author's Preface: Pedagogical Architecture & Analytical Methodology **xi**

Syllabus Blueprint & NMC Rationalization Guide **xiii**

I Physical Chemistry Foundations **1**

1 Foundations of Physical Chemistry: Stoichiometry, Atomic Structure & Thermodynamics **3**

1.1 NMC 2027 Regulatory Calibration & Scope Demarcation 3

1.2 Some Basic Concepts in Chemistry: Mole Concept & Stoichiometry (Unit 1) 3

1.2.1 Laws of Chemical Combination 4

1.2.2 The Mole Concept & Molar Quantities 4

1.2.3 Empirical and Molecular Formulas 5

1.2.4 Stoichiometry, Limiting Reagents & Reaction Yields . . . 5

1.2.5 Solution Concentration Expressions & Interconversions . . 6

1.3 Atomic Structure & Wave Mechanics (Unit 2) 7

1.3.1 Dual Nature of Light & The Photoelectric Effect 7

1.3.2 Bohr's Model of the Hydrogen Atom: Complete Mathematical Derivations 8

1.3.3 Wave-Particle Duality: The de Broglie Relation 9

1.3.4 Heisenberg's Uncertainty Principle 10

1.3.5 Schrödinger Wave Equation & Quantum Mechanical Formalism 10

1.3.6 Quantum Numbers & Rules of Electron Configuration . . 11

1.4 Chemical Thermodynamics & Thermochemistry (Unit 4) 13

1.4.1 Fundamental Concepts: Systems, Variables & Functions . 13

1.4.2 First Law of Thermodynamics & Expansion Work Derivations . . . 13
1.4.3 Enthalpy, Heat Capacities & Mayer's Relation Derivation 14
1.4.4 Reversible Adiabatic Expansion of an Ideal Gas . . . 15
1.4.5 Thermochemistry: Heats of Reactions, Hess's Law & Cycles 16
1.4.6 Second & Third Laws: Entropy, Free Energy & Spontaneity 17
1.4.7 Gibbs Free Energy & Equilibrium Thermodynamics . . . 18

2 Chemical & Ionic Equilibrium 21
2.1 Chemical & Ionic Equilibrium (Unit 6) . . . 21
2.1.1 Chemical Equilibrium: Law of Mass Action & Constant Relations . . . 21
2.1.2 Le Chatelier's Principle & Equilibrium Shifts . . . 22
2.1.3 Degree of Dissociation from Vapour Density . . . 23
2.1.4 Ionic Equilibrium: Ostwald Dilution Law & Autoionization of Water . . . 23
2.1.5 Salt Hydrolysis: Derivations of K_h, h, and pH . . . 23
2.1.6 Buffer Solutions & Henderson-Hasselbalch Derivations . . 24
2.1.7 Solubility Product (K_{sp}) & Common Ion Precipitation . . 24

II Advanced Physical Chemistry 27

3 Advanced Physical Chemistry: Solutions, Electrochemistry & Chemical Kinetics 29
3.1 Solutions & Colligative Properties (Unit 5) . . . 29
3.1.1 Gas Solubility & Henry's Law . . . 29
3.1.2 Vapour Pressure & Raoult's Law . . . 29
3.1.3 Ideal vs Non-Ideal Solutions & Azeotropes . . . 30
3.1.4 Colligative Properties & Thermodynamic Derivations . . 30
3.1.5 Abnormal Molar Masses & van 't Hoff Factor (i) . . . 33
3.2 Redox Reactions & Electrochemistry (Unit 7) . . . 34
3.2.1 Galvanic Cells & The Daniell Cell . . . 34
3.2.2 Standard Hydrogen Electrode & Electrochemical Series . 34
3.2.3 Thermodynamic Derivation of the Nernst Equation . . . 34
3.2.4 Electrolytic Conductance & Kohlrausch's Law . . . 35

3.2.5 Electrolysis & Faraday's Laws 37
3.2.6 Commercial Batteries & Mechanism of Corrosion 37
3.3 Chemical Kinetics (Unit 8) 38
3.3.1 Reaction Rate, Rate Laws & Order vs Molecularity 38
3.3.2 Integrated Rate Equations: Full Calculus Derivations . . 39
3.3.3 Temperature Dependence: The Arrhenius Equation . . . 41
3.3.4 Collision Theory & Catalysis Kinetics 42

III Inorganic Chemistry & Coordination Compounds 45

4 Inorganic Chemistry: Periodic Trends, Chemical Bonding, Coordination & Block Elements 47
4.1 NMC 2027 Regulatory Calibration & Scope Demarcation 48
4.2 Classification of Elements & Periodicity in Properties (Unit 9) . 48
4.2.1 Modern Periodic Law & Electronic Architecture 48
4.2.2 Effective Nuclear Charge (Z_{eff}) & Slater's Rules 49
4.2.3 Atomic and Ionic Radii Trends & Contractions 50
4.2.4 Ionization Enthalpy ($\Delta_i H$) & High-Yield Anomalies . . . 50
4.2.5 Electron Gain Enthalpy ($\Delta_{\text{eg}} H$) Trends & Traps 52
4.2.6 Electronegativity & The Inert Pair Effect 52
4.3 Chemical Bonding & Molecular Structure (Unit 3) 53
4.3.1 Ionic Bonding, Lattice Energy & Fajan's Rules 54
4.3.2 Lewis Structures & Formal Charge 55
4.3.3 Dipole Moments & Vector Summation 55
4.3.4 VSEPR Theory: Master 3D Geometry Framework 56
4.3.5 Molecular Orbital Theory (MOT): Energy Levels, Bond Orders & Magnetism 58
4.3.6 Hydrogen Bonding: Intermolecular vs Intramolecular . . . 59
4.4 Coordination Chemistry (Unit 12) 60
4.4.1 Werner's Coordination Theory 60
4.4.2 Nomenclature & Isomerism in Coordination Entities . . . 60
4.4.3 Valence Bond Theory (VBT) & Magnetic States 62
4.4.4 Crystal Field Theory (CFT): Splitting Diagrams & CFSE 62
4.4.5 Jahn-Teller Distortion Basics 65

4.4.6 Synergic Bonding in Metal Carbonyls 65
4.5 Comparative Chemistry of p-Block Elements (Groups 13 to 18) . 66
4.5.1 Group 13: Boron Family 66
4.5.2 Group 14: Carbon Family 67
4.5.3 Group 15: Nitrogen Family 67
4.5.4 Group 16: Oxygen Family 68
4.5.5 Group 17: Halogen Family 69
4.5.6 Group 18: Noble Gases 70
4.6 Comparative Chemistry of d- and f-Block Elements (Unit 11) . . 71
4.6.1 General Trends of $3d$ Transition Series 71
4.6.2 Master Transition Metal Compounds: $K_2Cr_2O_7$ & $KMnO_4$ 72
4.6.3 The f-Block Elements: Lanthanoids & Actinoids 75

IV Organic Chemistry: Principles, Pathways & Biomolecules 77

5 Organic Chemistry: Electronic Effects, Functional Groups & Biomolecules 79
5.1 NMC Syllabus Demarcation & Zero-Tolerance Deletions 80
5.2 General Organic Chemistry (GOC) & Electronic Effects 80
5.2.1 Inductive Effect ($\pm I$) 81
5.2.2 Resonance / Mesomeric Effect ($\pm M$) 82
5.2.3 Hyperconjugation (Baker-Nathan Effect) 84
5.2.4 Electromeric Effect ($\pm E$) 85
5.2.5 Aromaticity & Hückel's Rule 86
5.3 Reactive Intermediates & Molecular Rearrangements 88
5.3.1 Carbocations 88
5.3.2 Carbanions 89
5.3.3 Free Radicals 90
5.3.4 Carbenes, Nitrenes & Arynes 90
5.4 Hydrocarbons: Aliphatic & Aromatic Pathways 92
5.4.1 Alkanes 92
5.4.2 Alkenes 93
5.4.3 Alkynes 94

5.4.4 Aromatic Hydrocarbons: EAS Mechanics & Substituent Effects . . . 95
5.5 Haloalkanes & Haloarenes . . . 96
5.5.1 Nucleophilic Aliphatic Substitution (S_N1 vs S_N2 vs S_Ni) 96
5.5.2 Elimination Reactions ($E1$ vs $E2$ vs $E1cB$) . . . 97
5.5.3 Nucleophilic Aromatic Substitution (S_NAr) . . . 98
5.6 Alcohols, Phenols & Ethers . . . 99
5.6.1 Alcohols: Acidity, Oxidation & Dehydration . . . 99
5.6.2 Phenols: Acidic Character & Named Reactions . . . 100
5.6.3 Ethers: Williamson Synthesis & HI Cleavage . . . 100
5.7 Aldehydes, Ketones & Carboxylic Acids . . . 102
5.7.1 Nucleophilic Addition Mechanisms . . . 102
5.7.2 Reactions Involving α-Hydrogen: Aldol, Cannizzaro & Haloform . . . 103
5.7.3 Distinction Tests & Reductions . . . 104
5.7.4 Carboxylic Acids & Derivatives . . . 105
5.8 Nitrogen-Containing Compounds: Amines & Diazonium Chemistry . . . 106
5.8.1 Basicity of Amines: Gas Phase vs Aqueous Phase . . . 106
5.8.2 Gabriel Phthalimide & Hoffmann Bromamide Mechanisms 107
5.8.3 Distinction Tests for Amines . . . 109
5.8.4 Benzenediazonium Chemistry & Synthetic Transformations 109
5.9 Biomolecules (NEET Unit 19) . . . 110
5.9.1 Carbohydrates: Classification & Stereochemistry . . . 110
5.9.2 Cyclic Pyranose Structures, Anomers & Mutarotation . . 111
5.9.3 Disaccharides, Glycosidic Linkages & Polysaccharides . . 112
5.9.4 Amino Acids, Zwitterions & The Isoelectric Point (pI) . . 113
5.9.5 Peptide Bond Resonance & Structural Hierarchy of Proteins 114
5.9.6 Nucleic Acids: DNA & RNA Architecture . . . 115
5.9.7 Vitamins & Hormones . . . 116
5.10 Master Distinction Matrix & High-Yield Synthesis Archetypes . . 117
5.10.1 Exhaustive Chemical Distinction Matrix . . . 117
5.10.2 Solved Master Synthesis Archetypes . . . 118

V Organic Reaction Mechanisms Handbook 121

6 Organic Reaction Mechanisms Handbook: Step-by-Step Arrow-Pushing Pathways 123

6.1 Electrophilic Aromatic Substitution (S_EAr / EAS) 123
6.1.1 Nitration of Aromatic Rings 124
6.1.2 Halogenation of Benzene 125
6.1.3 Friedel-Crafts Alkylation & Acylation 125
6.1.4 Orientation & Reactivity: Arenium Ion Resonance Analysis 127
6.2 Nucleophilic Aliphatic Substitution (S_N1 vs S_N2) 128
6.2.1 The S_N1 Reaction Pathway 128
6.2.2 The S_N2 Reaction Pathway 129
6.2.3 Master Diagnostic: S_N1 vs S_N2 vs S_Ni 130
6.3 Elimination Reactions ($E1$ vs $E2$) & Dehydration of Alcohols . . 131
6.3.1 $E1$ Elimination Mechanism 131
6.3.2 $E2$ Elimination: Anti-Periplanar Geometry & Regioselectivity 131
6.3.3 Acid-Catalyzed Dehydration of Alcohols 132
6.4 Carbonyl Chemistry Mechanisms 134
6.4.1 Nucleophilic Addition to Aldehydes & Ketones (Ad_N) . . 134
6.4.2 Aldol Condensation & Cross-Aldol Mechanism 135
6.4.3 Cannizzaro Reaction & Hydride Transfer Mechanism . . . 136
6.5 Carboxylic Acid Derivatives & Mechanisms 138
6.5.1 Fischer Esterification Mechanism 138
6.5.2 Decarboxylation of Carboxylic Acids 139
6.5.3 Hell-Volhard-Zelinsky (HVZ) Reaction & α-Halogenation 141
6.6 Aromatic & Nitrogen Mechanisms 143
6.6.1 Diazotization Mechanism of Primary Aromatic Amines . 143
6.6.2 Azo Dye Coupling Mechanism (Phenol vs Aniline) 144
6.6.3 Reimer-Tiemann Reaction Mechanism 144
6.6.4 Kolbe's (Kolbe-Schmitt) Reaction Mechanism 145
6.6.5 Hofmann Bromamide Degradation Mechanism 146
6.7 Master Diagnostic Matrix: Reaction Mechanisms Overview . . . 148

6.7.1 Reaction Matrix I: Hydrocarbons, Alkyl Halides & Alcohols 148
6.7.2 Reaction Matrix II: Carbonyls & Carboxylic Acid Derivatives . . . 149
6.7.3 Reaction Matrix III: Phenols, Diazonium & Nitrogen Compounds . . . 150

VI Practical Chemistry & Qualitative Analysis Lab Manual 151

7 Practical Chemistry: Qualitative Salt Analysis & Volumetric Redox Titrations 153

7.1 Systematic Qualitative Analysis: Acidic Radicals (Anions) . . . 153
7.2 Systematic Qualitative Analysis: Basic Radicals (Cations) . . . 154
7.3 Volumetric Quantitative Redox Titrations . . . 156

Appendix:

Chemistry Formula, Reaction & Reagent Matrices 159

A Physical Chemistry Core Formulas & Thermodynamic Equations 161

A.1 Some Basic Concepts in Chemistry: Mole Concept (Unit 1) . . . 161
A.1.1 Stoichiometric Mole Formulations . . . 161
A.1.2 Solution Concentrations & Dilution Formulations . . . 161
A.2 Structure of Atom (Unit 2) . . . 163
A.2.1 Quantum Mechanics, Wave-Particle Duality & Uncertainty 163
A.2.2 Quantum Numbers, Nodes & Electron Filling Principles . 163
A.3 Chemical Thermodynamics & Thermochemistry (Unit 4) . . . 165
A.3.1 First Law, Work, Enthalpy & Heat Capacities . . . 165
A.3.2 Thermochemistry & Hess's Law . . . 165
A.3.3 Entropy, Second Law & Gibbs Free Energy Spontaneity . 165
A.4 Equilibrium (Unit 6) . . . 167
A.4.1 Chemical Equilibrium, $K_p - K_c$ Coupling & Le Chatelier 167
A.4.2 Ionic Equilibrium: Ostwald Dilution, Water Autoionization & pH . . . 167
A.4.3 Salt Hydrolysis Formulations . . . 167
A.4.4 Buffer Solutions & Henderson-Hasselbalch Equations . . . 167

A.4.5 Solubility Product (K_{sp}) & Precipitation Criteria 168
A.5 Solutions & Colligative Properties (Unit 5) 169
A.5.1 Gas Solubility & Raoult's Law 169
A.5.2 Colligative Properties & van 't Hoff Factor 169
A.6 Electrochemistry (Unit 7) 171
A.6.1 Galvanic Cells, Nernst Equation & Equilibrium 171
A.6.2 Electrolytic Conductance & Kohlrausch's Law 171
A.6.3 Faraday's Laws of Electrolysis 172
A.7 Chemical Kinetics (Unit 8) 173
A.7.1 Rate Laws, Order & Molecularity 173
A.7.2 Integrated Rate Equations: Zero, First & n-th Order . . . 173
A.7.3 Temperature Dependence: Arrhenius Equation & Catalysis 174

B Inorganic Trends & Organic Named Reactions Compendium 175
B.1 Classification of Elements & Periodicity (Unit 9) 175
B.1.1 Effective Nuclear Charge & Atomic Radii Trends 175
B.1.2 Ionization Enthalpy, Electron Gain Enthalpy & Electronegativity 175
B.2 Chemical Bonding & Molecular Structure (Unit 3) 177
B.2.1 Formal Charge & VSEPR Theory Master Geometry Table 177
B.2.2 Molecular Orbital Theory (MOT) 178
B.3 Coordination Compounds (Unit 12) 179
B.3.1 Werner's Theory, Ligands & Chelate Effect 179
B.3.2 Crystal Field Theory (CFT) & Splitting Formulations . . 179
B.3.3 Stereoisomerism in Coordination Complexes 180
B.4 The d- and f-Block Elements (Unit 11) 181
B.4.1 Transition Metal Trends & Variable Oxidation States . . 181
B.4.2 Potassium Dichromate ($K_2Cr_2O_7$) & Permanganate ($KMnO_4$) 181
B.4.3 Lanthanoid Contraction & Oxidation States 182
B.5 p-Block Elements (Unit 10) 183
B.5.1 Inert Pair Effect & Thermodynamic Stability Trends . . . 183
B.6 General Organic Chemistry (GOC) & Electronic Effects 184

B.7 Hydrocarbons: Alkanes, Alkenes, Alkynes & Arenes 186
B.8 Haloalkanes & Haloarenes: S_N1, S_N2, $E1$, $E2$ 189
B.9 Oxygen Compounds: Alcohols, Phenols, Ethers, Carbonyls & Acids 190
B.10 Nitrogen Compounds: Amines & Diazonium Salts 193
B.11 Biomolecules: Carbohydrates, Amino Acids & Nucleic Acids . . . 194
B.12 Master Identification Tests & Distinction Matrix 196

About the Author & Errata Submissions **199**

PART I

Physical Chemistry Foundations

CHAPTER 1

Foundations of Physical Chemistry: Stoichiometry, Atomic Structure & Thermodynamics

Physical chemistry provides the quantitative foundation for all chemical systems. This chapter begins with mole concept stoichiometry and empirical laws of chemical combination, progresses into wave mechanics and quantum numbers, and establishes the first and second laws of thermodynamics with explicit enthalpy, entropy, and Gibbs free energy derivations.

1.1 NMC 2027 Regulatory Calibration & Scope Demarcation

Zero-Tolerance NMC Syllabus Alert: Deleted Physical Chemistry Chapters

Under the revised National Medical Commission (NMC) NEET UG 2027 regulatory curriculum, the following legacy chapters have been **strictly deleted and removed from the examination**:

1. **Solid State:** All unit cells, crystal systems, Bragg's diffraction, packing efficiency derivations, point defects (Schottky and Frenkel defects), electrical and magnetic properties of solids are **completely eliminated**.
2. **States of Matter (Gaseous & Liquid Bulk States):** Macroscopic gas laws, van der Waals equation of state, critical phenomena (P_c, V_c, T_c), compressibility factor (Z) curves, liquefaction of gases, and bulk liquid properties (surface tension, viscosity) are **completely eliminated**. (Note: Ideal gas stoichiometry at STP is retained in Unit 1; molecular kinetic theory is covered in Physics).
3. **Surface Chemistry:** Adsorption isotherms (Freundlich and Langmuir), colloidal states, lyophilic and lyophobic sols, micelles and CMC, Tyndall effect, electrophoresis, coagulation and Hardy-Schulze rule, and heterogeneous catalysis are **strictly eliminated** from theory.

Study Directive: Do not invest study hours in these eliminated topics. All cognitive focus must remain on the rigorously tested Class 11 and Class 12 physical chemistry units detailed below.

1.2 Some Basic Concepts in Chemistry: Mole Concept & Stoichiometry (Unit 1)

1.2.1 Laws of Chemical Combination

Chemical transformations obey five fundamental empirical laws that form the bedrock of quantitative chemistry:

1. **Law of Conservation of Mass (Lavoisier, 1789):** In any chemical reaction, the total mass of the reactants equals the total mass of the products: $\sum m_{\text{reactants}} = \sum m_{\text{products}}$.
2. **Law of Definite Proportions (Proust, 1799):** A given chemical compound always contains exactly the same proportion of elements by mass, regardless of its source or method of preparation.
3. **Law of Multiple Proportions (Dalton, 1803):** If two elements combine to form more than one compound, the masses of one element that combine with a fixed mass of the other element are in the ratio of small whole numbers (e.g., in CO and CO_2, fixed 12 g of carbon combines with 16 g and 32 g of oxygen, ratio 1 : 2).
4. **Gay-Lussac's Law of Gaseous Volumes (1808):** When gases react together, they do so in volumes which bear a simple whole-number ratio to one another and to the volumes of gaseous products, provided all volumes are measured at identical temperature and pressure.
5. **Avogadro's Hypothesis (1811):** Equal volumes of all gases under identical conditions of temperature and pressure contain an equal number of molecules: $V \propto n$ (at constant T, P).

1.2.2 The Mole Concept & Molar Quantities

The mole is the SI base unit for the amount of substance. Exactly one mole contains $6.02214076 \times 10^{23}$ elementary entities (Avogadro's constant, N_A).

Master Mole Formulations

For any pure chemical substance of mass w (in g), molar mass M (in g mol^{-1}), and elementary particle count N:

$$n = \frac{w}{M} = \frac{N}{N_A} = \frac{V_{\text{STP}}\ (\text{in Litres})}{V_m} \tag{1.1}$$

where the standard molar volume V_m depends on the adopted thermodynamic convention:

- **Traditional NCERT Standard** ($T = 273.15\,\text{K}, P = 1\,\text{atm} = 101.325\,\text{kPa}$)**:** $V_m = 22.414\,\text{L mol}^{-1} \approx 22.4\,\text{L}$.
- **IUPAC Standard** ($T = 273.15\,\text{K}, P = 1\,\text{bar} = 100.0\,\text{kPa}$)**:** $V_m = 22.711\,\text{L mol}^{-1} \approx 22.7\,\text{L}$.

Vapour Density (VD) Relationship for Ideal Gases:

$$\text{VD} = \frac{\rho_{\text{gas}}}{\rho_{\text{H}_2}} = \frac{M_{\text{gas}}}{M_{\text{H}_2}} = \frac{M_{\text{gas}}}{2.016} \implies M \approx 2 \times \text{VD} \tag{1.2}$$

1.2.3 Empirical and Molecular Formulas

Empirical vs Molecular Formula Determination

- **Empirical Formula (EF):** The simplest whole-number ratio of atoms of each element present in a compound.
- **Molecular Formula (MF):** The actual number of atoms of each element present in a molecule of the compound:

$$\text{Molecular Formula} = (\text{Empirical Formula})_n \tag{1.3}$$

$$\text{where } n = \frac{\text{Molar Mass}}{\text{Empirical Formula Mass}} \tag{1.4}$$

- **Algorithmic Determination:**
  1. Convert given mass percentages directly to mass in 100 g of sample.
  2. Divide each mass by the respective atomic mass to compute relative moles: $n_i = w_i / A_i$.
  3. Divide each n_i by the smallest mole value among them: $r_i = n_i / n_{\min}$.
  4. If the resulting ratios are non-integers, multiply all ratios by the smallest integer that converts them into whole numbers.

1.2.4 Stoichiometry, Limiting Reagents & Reaction Yields

For a general balanced stoichiometric chemical equation:

$$aA + bB \longrightarrow cC + dD \tag{1.5}$$

The stoichiometric coefficients a, b, c, d represent the exact molar combining ratios:

$$\frac{n_A \text{ (consumed)}}{a} = \frac{n_B \text{ (consumed)}}{b} = \frac{n_C \text{ (produced)}}{c} = \frac{n_D \text{ (produced)}}{d} \tag{1.6}$$

Limiting Reagent (LR) Diagnostic Algorithm

When arbitrary initial quantities $n_{A,0}$ and $n_{B,0}$ are supplied:

1. Compute the mole-to-coefficient ratio for each reactant:

$$\mathcal{R}_A = \frac{n_{A,0}}{a}, \qquad \mathcal{R}_B = \frac{n_{B,0}}{b} \tag{1.7}$$

2. If $\mathcal{R}_A < \mathcal{R}_B$, reactant A is the **Limiting Reagent** (completely consumed). Reactant B is in excess.
3. All product yields and reactant consumptions are strictly determined by the limiting reagent:

$$n_C = c\left(\frac{n_{A,0}}{a}\right), \qquad n_{B,\text{remaining}} = n_{B,0} - b\left(\frac{n_{A,0}}{a}\right) \tag{1.8}$$

4. **Percentage Yield & Percentage Purity:**

$$\text{Percentage Yield} = \frac{\text{Actual Yield}}{\text{Theoretical Yield}} \times 100\% \tag{1.9}$$

$$\text{Percentage Purity} = \frac{\text{Mass of pure reactant}}{\text{Total mass of impure sample}} \times 100\% \tag{1.10}$$

1.2.5 Solution Concentration Expressions & Interconversions

The quantitative composition of solutions is formulated across six distinct mathematical regimes:

Concentration Unit	Sym.	Mathematical Definition	Temp. Dependence
Molarity	M	$M = \frac{n_{\text{solute}}}{V_{\text{sol}}\ (\text{L})} = \frac{w_B \times 1000}{M_B \times V_{\text{sol}}\ (\text{mL})}$	**Dependent** ($V \propto T \implies M \downarrow$ as $T \uparrow$)
Molality	m	$m = \frac{n_{\text{solute}}}{W_{\text{solvent}}\ (\text{kg})} = \frac{w_B \times 1000}{M_B \times W_A\ (\text{g})}$	**Independent** (mass invariant)
Normality	N	$N = \frac{\text{Gram Eq.}}{V_{\text{sol}}\ (\text{L})} = M \times n\text{-factor}$	**Dependent** ($V \propto T$)
Mole Fraction	x_B	$x_B = \frac{n_B}{n_A + n_B}, \quad x_A + x_B = 1$	**Independent**
Mass Percentage	$\%(w/w)$	$\%(w/w) = \frac{w_{\text{solute}}}{w_{\text{solution}}} \times 100$	**Independent**
Volume Percentage	$\%(v/v)$	$\%(v/v) = \frac{V_{\text{solute}}}{V_{\text{solution}}} \times 100$	**Dependent**
Parts Per Million	ppm	$\text{ppm} = \frac{w_{\text{solute}}}{w_{\text{solution}}} \times 10^6$	**Independent**

Table 1.1: Comprehensive Classification of Solution Concentration Regimes.

Master Interconversion Formulations (d = density in g mL^{-1})

$$M = \frac{10 \times \%(w/w) \times d}{M_B} \tag{1.11}$$

$$m = \frac{1000 \times M}{1000\,d - M\,M_B} \tag{1.12}$$

$$m = \frac{x_B \times 1000}{(1 - x_B)M_A} = \frac{x_B \times 1000}{x_A M_A} \tag{1.13}$$

$$x_B = \frac{m\,M_A}{1000 + m\,M_A} \tag{1.14}$$

Dilution and Mixing Formulations:

$$M_1V_1 = M_2V_2, \qquad M_{\text{mix}} = \frac{\sum M_iV_i}{\sum V_i} \tag{1.15}$$

$$N_1V_1 = N_2V_2 \quad \text{[at redox/acid-base equivalence]} \tag{1.16}$$

NCERT Benchmark: Valence Factor (n-factor) Rules

- **Acids:** n-factor = basicity (replaceable H^+): HCl $\rightarrow$ 1, H_2SO_4 $\rightarrow$ 2, H_3PO_4 $\rightarrow$ 3; H_3PO_3 $\rightarrow$ 2 (one non-ionizable P – H); H_3PO_2 $\rightarrow$ 1 (two non-ionizable P – H).
- **Bases:** n-factor = acidity (replaceable OH^-): NaOH $\rightarrow$ 1, $Ca(OH)_2$ $\rightarrow$ 2, $Al(OH)_3$ $\rightarrow$ 3.
- **Salts:** n-factor = total charge: NaCl $\rightarrow$ 1, $CaCl_2$ $\rightarrow$ 2, $Al_2(SO_4)_3$ $\rightarrow$ 6.
- **Redox Agents:** n-factor is total electrons transferred per formula unit.

Common Pitfall: STP Molar Volume & Equivalent Mass of Oxyacids

- **STP Standard Conditions Shift:** In modern IUPAC / NCERT convention, standard temperature and pressure (STP) is defined at $T = 273.15\,\text{K}$ and $P = 1\,\text{bar} = 10^5\,\text{Pa}$. Under these modern conditions, the molar volume of an ideal gas is $V_m = 22.7\,\text{L}\cdot\text{mol}^{-1}$ (or $22.7\,\text{dm}^3\cdot\text{mol}^{-1}$). The legacy convention ($1\,\text{atm} = 1.01325 \times 10^5\,\text{Pa}$) yielded $22.4\,\text{L}\cdot\text{mol}^{-1}$. Always inspect whether numerical question options in NEET assume legacy (22.4 L) or modern IUPAC (22.7 L).
- **Polyprotic Phosphorus Oxyacids:** Never equate molecular mass divided by 3 to equivalent mass blindly! For orthophosphoric acid (H_3PO_4), basicity is $3 \implies E = M/3$. For phosphorous acid (H_3PO_3), basicity is $2 \implies E = M/2$ because one hydrogen is directly bonded to phosphorus (P – H) and does not ionize in aqueous solution. For hypophosphorous acid (H_3PO_2), basicity is $1 \implies E = M/1$ since two hydrogens are non-ionizable P – H bonds.

1.3 Atomic Structure & Wave Mechanics (Unit 2)

1.3.1 Dual Nature of Light & The Photoelectric Effect

Einstein formulated the photoelectric effect by treating light as discrete localized wave packets (photons) of energy $E = h\nu$. When a photon strikes a metallic

surface:

$$E_{\text{photon}} = \Phi + K_{\max} \implies h\nu = h\nu_0 + \frac{1}{2}m_e v_{\max}^2 = h\nu_0 + eV_0 \tag{1.17}$$

where $\Phi = h\nu_0$ is the work function of the metal, ν_0 is the threshold frequency, and V_0 is the stopping potential.

1.3.2 Bohr's Model of the Hydrogen Atom: Complete Mathematical Derivations

Niels Bohr synthesized classical electrostatic planetary mechanics with Planckian angular momentum quantization for hydrogen-like species of nuclear charge $+Ze$ and single orbital electron $-e$:

1. **Postulate 1 (Electrostatic Centripetal Equilibrium):**

$$\frac{m_e v_n^2}{r_n} = \frac{1}{4\pi\varepsilon_0}\frac{Ze^2}{r_n^2} \implies m_e v_n^2 r_n = \frac{Ze^2}{4\pi\varepsilon_0} \tag{1.18}$$

2. **Postulate 2 (Quantization of Orbital Angular Momentum):**

$$L_n = m_e v_n r_n = \frac{nh}{2\pi} \quad (n = 1, 2, 3, \dots) \tag{1.19}$$

Derivation of Quantized Orbit Radius (r_n)

Dividing Equation 1.19 by $m_e r_n$ yields the electron orbital velocity:

$$v_n = \frac{nh}{2\pi m_e r_n} \tag{1.20}$$

Substituting this expression for v_n into Equation 1.18:

$$m_e\left(\frac{nh}{2\pi m_e r_n}\right)^2 r_n = \frac{Ze^2}{4\pi\varepsilon_0} \implies \frac{n^2h^2}{4\pi^2 m_e r_n} = \frac{Ze^2}{4\pi\varepsilon_0} \tag{1.21}$$

Solving explicitly for r_n:

$$r_n = \frac{\varepsilon_0 h^2}{\pi m_e e^2}\frac{n^2}{Z} = a_0\frac{n^2}{Z} \approx 0.529\,\text{Å} \times \frac{n^2}{Z} = 52.9\,\text{pm} \times \frac{n^2}{Z} \tag{1.22}$$

where $a_0 = 0.529177\,\text{Å}$ is the first Bohr radius of hydrogen ($n = 1, Z = 1$).

Derivation of Quantized Velocity (v_n)

Substituting r_n into Equation 1.19:

$$v_n = \frac{nh}{2\pi m_e\left(\frac{\varepsilon_0 h^2 n^2}{\pi m_e e^2 Z}\right)} = \frac{e^2}{2\varepsilon_0 h}\frac{Z}{n} = v_0\frac{Z}{n} \approx 2.188\times10^6\frac{Z}{n}\,\text{m s}^{-1} = \frac{c}{137}\frac{Z}{n} \tag{1.23}$$

Derivation of Quantized Energies (K_n, U_n, E_n)

$$K_n = \frac{1}{2}m_e v_n^2 = \frac{1}{2}\left(\frac{Ze^2}{4\pi\varepsilon_0 r_n}\right) = \frac{Ze^2}{8\pi\varepsilon_0 r_n} \tag{1.24}$$

$$U_n = -\frac{1}{4\pi\varepsilon_0}\frac{Ze^2}{r_n} = -2K_n \quad \text{(Virial Theorem: } 2\langle K\rangle + \langle U\rangle = 0\text{)} \tag{1.25}$$

$$E_n = K_n + U_n = -K_n = \frac{1}{2}U_n = -\frac{Ze^2}{8\pi\varepsilon_0 r_n} \tag{1.26}$$

Substituting the derived value of r_n:

$$E_n = -\frac{m_e e^4}{8\varepsilon_0^2 h^2}\frac{Z^2}{n^2} = -R_H hc\frac{Z^2}{n^2} = -2.18\times10^{-18}\,\text{J}\times\frac{Z^2}{n^2} = -13.6\frac{Z^2}{n^2}\,\text{eV} \tag{1.27}$$

Spectral Transitions & The Rydberg Equation

When an electron drops from an initial upper energy level n_2 to a lower level n_1 ($n_2 > n_1$):

$$\Delta E = E_{n_2} - E_{n_1} = h\nu = \frac{hc}{\lambda} \implies \bar{\nu} = \frac{1}{\lambda} = R_H Z^2\left(\frac{1}{n_1^2} - \frac{1}{n_2^2}\right) \tag{1.28}$$

where $R_H = \frac{m_e e^4}{8\varepsilon_0^2 h^3 c} \approx 1.09677\times10^7\,\text{m}^{-1}$ ($1/R_H \approx 912$ Å).

Spectral Series	n_1	n_2	Spectral Region	Wavelength Range
Lyman	1	$2, 3, 4, \ldots, \infty$	Ultraviolet (UV)	912 Å $\leq \lambda \leq$ 1216 Å
Balmer	2	$3, 4, 5, \ldots, \infty$	Visible (H_α to H_δ) + Near UV	3646 Å $\leq \lambda \leq$ 6563 Å
Paschen	3	$4, 5, 6, \ldots, \infty$	Near Infrared (IR)	8204 Å $\leq \lambda \leq$ 18751 Å
Brackett	4	$5, 6, 7, \ldots, \infty$	Infrared (IR)	14580 Å $\leq \lambda \leq$ 40512 Å
Pfund	5	$6, 7, 8, \ldots, \infty$	Far Infrared (Far IR)	22790 Å $\leq \lambda \leq$ 74580 Å

Table 1.2: Hydrogen Atom Emission Spectral Series.

1.3.3 Wave-Particle Duality: The de Broglie Relation

Louis de Broglie (1924) postulated that matter exhibits intrinsic wave characteristics. Equating Einstein's mass-energy equivalence ($E = mc^2 = pc$) with Planck's photon energy ($E = h\nu = hc/\lambda$):

$$p = \frac{h}{\lambda} \implies \lambda = \frac{h}{p} = \frac{h}{mv} \tag{1.29}$$

For a charged particle of mass m and charge q accelerated from rest through an electrostatic potential difference V:

$$K = qV = \frac{p^2}{2m} \implies p = \sqrt{2mqV} \implies \lambda = \frac{h}{\sqrt{2mqV}} \tag{1.30}$$

Specialized de Broglie Wavelength Formulas:

- **Accelerated Electron ($m = m_e, q = e$):** $\lambda_e = \frac{1.227}{\sqrt{V}}\,\text{nm} = \frac{12.27}{\sqrt{V}}\,\text{Å}$.
- **Thermal Neutron / Gas Molecule ($K = \frac{3}{2}k_BT$):** $\lambda_{\text{thermal}} = \frac{h}{\sqrt{3mk_BT}}$.

1.3.4 Heisenberg's Uncertainty Principle

Werner Heisenberg (1927) proved that it is physically impossible to simultaneously determine both the exact position and exact momentum of a microscopic quantum entity:

$$\Delta x \cdot \Delta p_x \geq \frac{\hbar}{2} = \frac{h}{4\pi} \implies \Delta x \cdot (m\Delta v_x) \geq \frac{h}{4\pi} \implies \Delta x \cdot \Delta v_x \geq \frac{h}{4\pi m} \tag{1.31}$$

Alternative Conjugate Variables Formulations:

$$\Delta E \cdot \Delta t \geq \frac{h}{4\pi}, \qquad \Delta\theta \cdot \Delta L \geq \frac{h}{4\pi} \tag{1.32}$$

Because Planck's constant $h = 6.626 \times 10^{-34}\,\text{J s}$ is infinitesimally small, the uncertainty product is completely imperceptible for macroscopic bodies ($m \sim 1\,\text{kg}$), but dominates the atomic regime ($m_e \sim 9.1 \times 10^{-31}\,\text{kg}$).

1.3.5 Schrödinger Wave Equation & Quantum Mechanical Formalism

Erwin Schrödinger (1926) formulated the stationary-state wave equation for a quantum particle of mass m moving in a potential field $V(x, y, z)$:

$$\nabla^2\psi + \frac{8\pi^2 m}{h^2}(E - V)\psi = 0 \iff \hat{H}\psi = E\psi \tag{1.33}$$

where $\hat{H} = -\frac{\hbar^2}{2m}\nabla^2 + V$ is the Hamiltonian operator, and $\psi(r, \theta, \phi)$ is the eigenfunction.

Physical Interpretation of Wave Function ψ and Probability Density $|\psi|^2$

- **Wave Function Amplitude (ψ):** ψ possesses no direct physical meaning; it can be positive, negative, or complex.

- **Born's Probability Density ($|\psi|^2 = \psi^*\psi$):** Represents the probability of finding the electron per unit volume in an infinitesimal volume element $d\tau = dx\,dy\,dz$:

$$dP = |\psi|^2 d\tau, \qquad \int_{\text{all space}} |\psi|^2 d\tau = 1 \quad \text{(Normalization Condition)} \tag{1.34}$$

- **Radial vs Angular Separation:** In spherical coordinates (r, θ, ϕ):

$$\psi(r, \theta, \phi) = R_{n,l}(r) \times Y_{l,m_l}(\theta, \phi) \tag{1.35}$$

  $R_{n,l}(r)$ (radial function) dictates orbital size and radial probability; $Y_{l,m_l}(\theta, \phi)$ (angular function) dictates orbital 3D spatial geometry and orientation.
- **Radial Probability Distribution Function ($4\pi r^2 R^2(r)$):** Represents the absolute probability of locating the electron in a spherical shell of radius r and thickness dr:

$$P(r)dr = 4\pi r^2 R^2(r) dr \tag{1.36}$$

  At $r = 0$, $4\pi r^2 R^2 = 0$ for all orbitals (electron cannot reside at the mathematical point of the nucleus).

Master Nodal Formulations

Nodes are spatial domains where the probability density $|\psi|^2$ vanishes identically:

$$\text{Radial Nodes (Spherical Nodes)} = n - l - 1 \tag{1.37}$$

$$\text{Angular Nodes (Nodal Planes / Cones)} = l \tag{1.38}$$

$$\text{Total Nodes} = (n - l - 1) + l = n - 1 \tag{1.39}$$

Examples:

- $1s$: Radial $= 1 - 0 - 1 = 0$, Angular $= 0$, Total $= 0$.
- $2s$: Radial $= 2 - 0 - 1 = 1$, Angular $= 0$, Total $= 1$.
- $2p$: Radial $= 2 - 1 - 1 = 0$, Angular $= 1$ (the planar surface passing through nucleus), Total $= 1$.
- $3d$: Radial $= 3 - 2 - 1 = 0$, Angular $= 2$ (two nodal planes/cones), Total $= 2$.
- $4d$: Radial $= 4 - 2 - 1 = 1$, Angular $= 2$, Total $= 3$.

1.3.6 Quantum Numbers & Rules of Electron Configuration

An atomic electron is completely described by four quantum numbers:

1. **Principal Quantum Number ($n \in \{1, 2, 3, \dots\}$):** Determines main energy shell, average distance from nucleus, maximum electrons $= 2n^2$, maximum orbitals $= n^2$.
2. **Azimuthal Quantum Number ($l \in \{0, 1, 2, \dots, n-1\}$):** Determines subshell shape ($s : 0, p : 1, d : 2, f : 3$), and orbital angular momentum:
$$L_{\text{orbital}} = \sqrt{l(l+1)}\hbar = \sqrt{l(l+1)}\frac{h}{2\pi} \tag{1.40}$$
(For an s-orbital with $l = 0$, orbital angular momentum $L = 0$).
3. **Magnetic Quantum Number ($m_l \in \{-l, \dots, 0, \dots, +l\}$):** Determines 3D spatial orientation of the orbital in an external magnetic field ($2l + 1$ degenerate orientations).
4. **Spin Quantum Number ($m_s \in \{+\frac{1}{2}, -\frac{1}{2}\}$):** Intrinsic angular momentum orientation. Spin-only magnetic moment:
$$\mu_s = \sqrt{n_u(n_u + 2)}\,\text{BM} \tag{1.41}$$
where n_u is the number of unpaired electrons and BM is Bohr Magneton.

Fundamental Rules of Electronic Architecture

- **Aufbau Principle:** In the ground state of an atom, electrons occupy the lowest available atomic energy orbitals first. Energy follows the $(n+l)$ **rule**:
  1. Lower $(n + l)$ orbital fills before higher $(n + l)$ (e.g., $4s : 4 + 0 = 4$ fills before $3d : 3 + 2 = 5$).
  2. If two orbitals share identical $(n + l)$, the orbital with lower n fills first (e.g., $3d : 3 + 2 = 5$ fills before $4p : 4 + 1 = 5$).
- **Pauli's Exclusion Principle:** No two electrons in an atom can have the identical set of all four quantum numbers. Consequently, an individual orbital accommodates at most two electrons with antiparallel spins ($\uparrow\downarrow$).
- **Hund's Rule of Maximum Multiplicity:** Pairing of electrons in degenerate orbitals (p, d, f) cannot occur until each orbital of the subshell is singly occupied with parallel spins.
- **Extra Stability of Half-Filled and Fully-Filled Subshells:**
  - **Symmetrical Electron Distribution:** Spherically symmetric charge minimizes interelectronic repulsions.
  - **Exchange Energy:** Quantum mechanical exchange of electrons with parallel spins releases stabilization energy. For n electrons of parallel spin:
$$\text{Number of possible exchanges} = \frac{n(n-1)}{2} \tag{1.42}$$
For Chromium ($[\text{Ar}]3d^5 4s^1$): $3d^5$ allows $\frac{5\times4}{2} = 10$ exchanges, vs $[\text{Ar}]3d^4 4s^2$ which allows only $\frac{4\times3}{2} = 6$ exchanges. Copper ($[\text{Ar}]3d^{10} 4s^1$) similarly gains maximum exchange stability over $3d^9 4s^2$.

1.4 Chemical Thermodynamics & Thermochemistry (Unit 4)

1.4.1 Fundamental Concepts: Systems, Variables & Functions

- **Thermodynamic System:** That part of the universe selected for thermodynamic study.
  - **Open System:** Exchanges both matter and energy with surroundings (e.g., open beaker of boiling water).
  - **Closed System:** Exchanges energy but not matter with surroundings (e.g., sealed metal canister).
  - **Isolated System:** Exchanges neither matter nor energy with surroundings (e.g., perfectly insulated Dewar flask).
- **State Functions vs Path Functions:**
  - **State Functions:** Properties whose values depend solely upon the current thermodynamic state of the system, independent of the path traversed (U, H, S, G, P, V, T). Cyclic integral $\oint dX = 0$.
  - **Path Functions:** Quantities whose numerical values depend explicitly upon the transition path traversed between states (q = heat, w = work). Cyclic integral $\oint dq \neq 0$.
- **Intensive vs Extensive Properties:**
  - **Extensive (additive, depends on mass):** Mass (m), volume (V), internal energy (U), enthalpy (H), entropy (S), Gibbs energy (G), heat capacity (C).
  - **Intensive (scale-independent, independent of mass):** Temperature (T), pressure (P), density (ρ), molar heat capacity (C_m), specific heat (c), viscosity (η), surface tension, refractive index, EMF of a cell (E_{cell}).

1.4.2 First Law of Thermodynamics & Expansion Work Derivations

The First Law is the conservation of energy principle applied to thermodynamic systems:

$$\Delta U = q + w \tag{1.43}$$

where $\Delta U = U_2 - U_1$ is the change in internal energy, q is the heat absorbed by the system ($q > 0$ if endothermic, $q < 0$ if exothermic), and w is work done on the system ($w > 0$ for compression, $w < 0$ for expansion under IUPAC sign convention).

Derivation of Reversible Isothermal Expansion Work of an Ideal Gas

For an infinitesimal quasi-static reversible expansion, the internal gas pressure P_{gas} differs from the external opposing pressure P_{ext} by an infinitesimal amount

dP ($P_{\text{ext}} = P - dP$):

$$\delta w_{\text{rev}} = -P_{\text{ext}} dV = -(P - dP)dV = -PdV + dPdV \approx -PdV \tag{1.44}$$

Integrating from initial state (P_1, V_1) to final state (P_2, V_2) at constant temperature T:

$$w_{\text{rev}} = -\int_{V_1}^{V_2} \frac{nRT}{V} dV = -nRT \ln\left(\frac{V_2}{V_1}\right) \tag{1.45}$$

$$= -2.303nRT \log_{10}\left(\frac{V_2}{V_1}\right) \tag{1.46}$$

By Boyle's Law ($P_1V_1 = P_2V_2$ at constant T):

$$w_{\text{rev}} = -2.303nRT \log_{10}\left(\frac{P_1}{P_2}\right) \tag{1.47}$$

Since temperature is constant in an isothermal ideal gas expansion ($\Delta U = 0$):

$$q_{\text{rev}} = -w_{\text{rev}} = +2.303nRT \log_{10}\left(\frac{V_2}{V_1}\right) \tag{1.48}$$

Irreversible Expansion & Free Expansion

- **Irreversible Expansion against Constant External Pressure (P_{ext}):**

$$w_{\text{irr}} = -\int_{V_1}^{V_2} P_{\text{ext}} dV = -P_{\text{ext}}(V_2 - V_1) = -P_{\text{ext}}\left(\frac{nRT_2}{P_2} - \frac{nRT_1}{P_1}\right) \tag{1.49}$$

  Because $|w_{\text{rev}}| > |w_{\text{irr}}|$, reversible expansion produces the **maximum work output**.

- **Free Expansion into Vacuum ($P_{\text{ext}} = 0$):**

$$w_{\text{free}} = -\int 0 \cdot dV = 0 \tag{1.50}$$

  For an ideal gas in an insulated container undergoing free expansion: $w = 0, q = 0 \implies \Delta U = 0 \implies \Delta T = 0$ (Joule's Law).

1.4.3 Enthalpy, Heat Capacities & Mayer's Relation Derivation

Enthalpy is defined as the total heat content:

$$H = U + PV \implies \Delta H = \Delta U + \Delta(PV) \tag{1.51}$$

For a chemical reaction involving gases at constant temperature:

$$\Delta H = \Delta U + \Delta n_g RT \tag{1.52}$$

where $\Delta n_g = \sum n_g(\text{gaseous products}) - \sum n_g(\text{gaseous reactants})$.

Derivation of Mayer's Relation: $C_p - C_v = R$

Heat capacities are defined by:

$$C_v = \left(\frac{\partial U}{\partial T}\right)_V = \frac{q_V}{\Delta T}, \qquad C_p = \left(\frac{\partial H}{\partial T}\right)_P = \frac{q_P}{\Delta T} \tag{1.53}$$

For 1 mole of an ideal gas:

$$H = U + PV = U + RT \tag{1.54}$$

Differentiating both sides with respect to temperature T:

$$\frac{dH}{dT} = \frac{dU}{dT} + R\frac{d(T)}{dT} \implies C_p = C_v + R \implies C_p - C_v = R \tag{1.55}$$

For n moles: $C_{p,\text{total}} - C_{v,\text{total}} = nR$. **Heat Capacity Ratio (γ):**

$$\gamma = \frac{C_p}{C_v} = 1 + \frac{R}{C_v} \tag{1.56}$$

- **Monoatomic Gas ($\text{He}, \text{Ne}, \text{Ar}$):** Degrees of freedom $f = 3 \implies C_v = \frac{3}{2}R, C_p = \frac{5}{2}R \implies \gamma = \frac{5}{3} \approx 1.67$.
- **Diatomic Gas (O_2, N_2, CO at moderate T):** $f = 5 \implies C_v = \frac{5}{2}R, C_p = \frac{7}{2}R \implies \gamma = \frac{7}{5} = 1.40$.
- **Non-linear Polyatomic Gas (H_2O, NH_3, CH_4):** $f = 6 \implies C_v = 3R, C_p = 4R \implies \gamma = \frac{4}{3} \approx 1.33$.

1.4.4 Reversible Adiabatic Expansion of an Ideal Gas

In an adiabatic process, no heat enters or leaves the system ($q = 0$). By the First Law:

$$dU = \delta w \implies nC_v dT = -PdV = -\frac{nRT}{V}dV \tag{1.57}$$

Dividing by nT and separating variables:

$$C_v\frac{dT}{T} = -R\frac{dV}{V} = -(C_p - C_v)\frac{dV}{V} \implies \frac{dT}{T} = -(\gamma - 1)\frac{dV}{V} \tag{1.58}$$

Integrating from state 1 to state 2:

$$\ln\left(\frac{T_2}{T_1}\right) = -(\gamma - 1)\ln\left(\frac{V_2}{V_1}\right) = \ln\left[\left(\frac{V_1}{V_2}\right)^{\gamma-1}\right] \tag{1.59}$$

$$\implies T_1 V_1^{\gamma-1} = T_2 V_2^{\gamma-1} = \text{constant} \tag{1.60}$$

Substituting $T = \frac{PV}{nR}$:

$$P_1V_1^{\gamma} = P_2V_2^{\gamma} = \text{constant}, \qquad T_1^{\gamma}P_1^{1-\gamma} = T_2^{\gamma}P_2^{1-\gamma} = \text{constant} \tag{1.61}$$

Work Done in Reversible Adiabatic Expansion:

$$w_{\text{adi}} = \Delta U = nC_v(T_2 - T_1) = \frac{nR(T_2 - T_1)}{\gamma - 1} = \frac{P_2V_2 - P_1V_1}{\gamma - 1} \tag{1.62}$$

Since the gas expands ($V_2 > V_1$), $T_2 < T_1$ (gas cools during adiabatic expansion!). On a $P - V$ diagram, the slope of an adiabat is γ times steeper than an isotherm:

$$\left(\frac{dP}{dV}\right)_{\text{adi}} = -\gamma\frac{P}{V} = \gamma\left(\frac{dP}{dV}\right)_{\text{iso}} \tag{1.63}$$

1.4.5 Thermochemistry: Heats of Reactions, Hess's Law & Cycles

Thermochemical Laws & Enthalpy Formulations

- **Hess's Law of Constant Heat Summation:** If a chemical reaction can be made to take place in a single step or in several successive steps, the overall enthalpy change is identical, independent of the pathway:

$$\Delta_r H^{\circ} = \Delta H_1 + \Delta H_2 + \Delta H_3 + \ldots \tag{1.64}$$

- **Standard Enthalpy of Formation ($\Delta_f H^{\circ}$):** Enthalpy change when 1 mole of a compound is synthesized from its constituent elements in their standard reference states (298.15 K, 1 bar).

$$\Delta_r H^{\circ} = \sum \nu_p \Delta_f H^{\circ}(\text{products}) - \sum \nu_r \Delta_f H^{\circ}(\text{reactants}) \tag{1.65}$$

- **Standard State Convention Trap:** By definition, $\Delta_f H^{\circ} = 0$ for all pure elements in their most stable allotropic form at 298 K: C(graphite) = 0 (but C(diamond) = $+1.89\,\text{kJ mol}^{-1}$); S(rhombic) = 0 (monoclinic $\neq 0$); P(white) = 0 (red $\neq 0$); $Br_{2(l)} = 0$; $I_{2(s)} = 0$.
- **Bond Dissociation Enthalpy (BE):**

$$\Delta_r H^{\circ} = \sum \text{BE}(\text{reactants}) - \sum \text{BE}(\text{products}) \quad \text{(all species in gas phase!)} \tag{1.66}$$

- **Kirchhoff's Law (Temperature Dependence of $\Delta_r H$):**

$$\Delta_r H^{\circ}_{T_2} = \Delta_r H^{\circ}_{T_1} + \int_{T_1}^{T_2} \Delta C_p dT = \Delta_r H^{\circ}_{T_1} + \Delta C_p(T_2 - T_1) \tag{1.67}$$

where $\Delta C_p = \sum \nu_p C_{p,\text{products}} - \sum \nu_r C_{p,\text{reactants}}$.

Born-Haber Cycle for Ionic Crystal Lattice Enthalpy ($\Delta_{\text{lattice}}H^\circ$)

For crystalline sodium chloride ($NaCl_{(s)}$), the standard enthalpy of formation from elemental sodium and chlorine gas connects directly to lattice enthalpy via five thermochemical steps:

$$\text{Na}_{(s)} + \frac{1}{2}\text{Cl}_{2(g)} \xrightarrow{\Delta_f H^\circ} \text{NaCl}_{(s)} \tag{1.68}$$

1. Sublimation of solid sodium: $\text{Na}_{(s)} \rightarrow \text{Na}_{(g)}$, $\Delta H^\circ_{\text{sub}} = +108.4\,\text{kJ mol}^{-1}$.
2. Ionization of gaseous sodium: $\text{Na}_{(g)} \rightarrow \text{Na}^+_{(g)} + e^-$, $\text{IE}_1 = +495.8\,\text{kJ mol}^{-1}$.
3. Dissociation of diatomic chlorine: $\frac{1}{2}\text{Cl}_{2(g)} \rightarrow \text{Cl}_{(g)}$, $\frac{1}{2}\Delta_{\text{diss}}H^\circ = +\frac{1}{2}(242) = +121.0\,\text{kJ mol}^{-1}$.
4. Electron gain by chlorine: $\text{Cl}_{(g)} + e^- \rightarrow \text{Cl}^-_{(g)}$, $\Delta_{\text{eg}}H^\circ = -348.6\,\text{kJ mol}^{-1}$.
5. Condensation of gaseous ions into crystal lattice: $\text{Na}^+_{(g)} + \text{Cl}^-_{(g)} \rightarrow \text{NaCl}_{(s)}$, $-\Delta_{\text{lattice}}H^\circ$.

Applying Hess's Law:

$$\Delta_f H^\circ = \Delta H^\circ_{\text{sub}} + \text{IE}_1 + \frac{1}{2}\Delta_{\text{diss}}H^\circ + \Delta_{\text{eg}}H^\circ - \Delta_{\text{lattice}}H^\circ \tag{1.69}$$

$$\Delta_{\text{lattice}}H^\circ = \Delta H^\circ_{\text{sub}} + \text{IE}_1 + \frac{1}{2}\Delta_{\text{diss}}H^\circ + \Delta_{\text{eg}}H^\circ - \Delta_f H^\circ = +788\,\text{kJ mol}^{-1} \tag{1.70}$$

1.4.6 Second & Third Laws: Entropy, Free Energy & Spontaneity

- **Entropy (S):** Measure of microscopic disorder or thermal energy dispersal. Infinitesimal reversible change:

$$dS = \frac{\delta q_{\text{rev}}}{T} \tag{1.71}$$

- **Entropy Change for an Ideal Gas:**

$$\Delta S = nC_v \ln\left(\frac{T_2}{T_1}\right) + nR\ln\left(\frac{V_2}{V_1}\right) = nC_p \ln\left(\frac{T_2}{T_1}\right) - nR\ln\left(\frac{P_2}{P_1}\right) \tag{1.72}$$

  - **Isothermal Process ($T_1 = T_2$):** $\Delta S = nR\ln(V_2/V_1) = nR\ln(P_1/P_2)$.
  - **Isobaric Process ($P_1 = P_2$):** $\Delta S = nC_p\ln(T_2/T_1)$.
  - **Isochoric Process ($V_1 = V_2$):** $\Delta S = nC_v\ln(T_2/T_1)$.
  - **Reversible Phase Transitions:** $\Delta S_{\text{fusion}} = \frac{\Delta H_{\text{fus}}}{T_f}$, $\Delta S_{\text{vaporization}} = \frac{\Delta H_{\text{vap}}}{T_b}$.

- **Spontaneity Criterion (Second Law of Thermodynamics):** For an isolated system or the universe:

$$\Delta S_{\text{total}} = \Delta S_{\text{system}} + \Delta S_{\text{surroundings}} \geq 0 \tag{1.73}$$

  (Spontaneous if $\Delta S_{\text{total}} > 0$; at dynamic equilibrium if $\Delta S_{\text{total}} = 0$).
- **Third Law of Thermodynamics:** The entropy of a perfectly crystalline pure substance approaches zero as absolute temperature approaches zero: $\lim_{T \to 0} S = 0$.

1.4.7 Gibbs Free Energy & Equilibrium Thermodynamics

Gibbs Free Energy ($G = H - TS$) provides the single-parameter spontaneity criterion at constant temperature and pressure:

$$\Delta G = \Delta H - T\Delta S \tag{1.74}$$

- $\Delta G < 0 \implies$ Reaction is spontaneous in the forward direction.
- $\Delta G = 0 \implies$ Dynamic equilibrium is attained.
- $\Delta G > 0 \implies$ Non-spontaneous forward (spontaneous in reverse direction).

ΔH	ΔS	$\Delta G = \Delta H - T\Delta S$	**Spontaneity Regime & Temperature Dependence**
$-$	$+$	Always Negative ($-$)	**Spontaneous at all temperatures** (enthalpy & entropy driven)
$+$	$-$	Always Positive ($+$)	**Non-spontaneous at all temperatures** (reverse is spontaneous)
$-$	$-$	Negative at low T, Positive at high T	**Spontaneous at low** T, non-spontaneous at high T ($\lvert T\Delta S\rvert > \lvert\Delta H\rvert$)
$+$	$+$	Positive at low T, Negative at high T	**Spontaneous at high** T, non-spontaneous at low T ($T > \frac{\Delta H}{\Delta S}$)

Table 1.3: Thermodynamic Spontaneity Temperature Dependency Matrix.

Master Thermodynamic Equilibrium Links

For a general reaction with reaction quotient Q:

$$\Delta G = \Delta_r G^\circ + RT \ln Q = \Delta_r G^\circ + 2.303RT \log_{10} Q \tag{1.75}$$

At dynamic chemical equilibrium, $\Delta G = 0$ and $Q = K_{\text{eq}}$:

$$\Delta_r G^\circ = -RT \ln K_{\text{eq}} = -2.303RT \log_{10} K_{\text{eq}} \iff K_{\text{eq}} = e^{-\Delta_r G^\circ / RT} \tag{1.76}$$

van 't Hoff Reaction Isobar:

$$\frac{d \ln K}{dT} = \frac{\Delta_r H^\circ}{RT^2} \tag{1.77}$$

$$\implies \log_{10}\left(\frac{K_2}{K_1}\right) = \frac{\Delta_r H^\circ}{2.303R}\left[\frac{T_2 - T_1}{T_1 T_2}\right] = \frac{\Delta_r H^\circ}{2.303R}\left[\frac{1}{T_1} - \frac{1}{T_2}\right] \tag{1.78}$$

- **Endothermic Reaction ($\Delta_r H^\circ > 0$):** K increases as temperature increases.
- **Exothermic Reaction ($\Delta_r H^\circ < 0$):** K decreases as temperature increases.

CHAPTER 2

Chemical & Ionic Equilibrium

Chemical equilibrium establishes the dynamic balance between forward and reverse reaction rates in closed thermodynamic systems, governed by the law of mass action and Le Chatelier's principle. Ionic equilibrium extends these principles to electrolytic solutions, encompassing acid-base dissociation equilibria, the autoionization of water, buffer solution dynamics via the Henderson-Hasselbalch formulation, quantitative salt hydrolysis, and sparingly soluble salt precipitation criteria governed by the solubility product (K_{sp}).

2.1 Chemical & Ionic Equilibrium (Unit 6)

2.1.1 Chemical Equilibrium: Law of Mass Action & Constant Relations

For a reversible homogeneous gas-phase reaction:

$$aA(g) + bB(g) \rightleftharpoons cC(g) + dD(g) \tag{2.1}$$

The equilibrium constants in terms of molar concentrations (K_c) and partial pressures (K_p) are:

$$K_c = \frac{[C]^c[D]^d}{[A]^a[B]^b}, \qquad K_p = \frac{P_C^c P_D^d}{P_A^a P_B^b} \tag{2.2}$$

Derivation of $K_p = K_c(RT)^{\Delta n_g}$

Assuming ideal gas behavior for each component:

$$P_i V = n_i RT \implies P_i = \left(\frac{n_i}{V}\right) RT = [i]RT \tag{2.3}$$

Substituting each partial pressure into the K_p expression:

$$K_p = \frac{([C]RT)^c([D]RT)^d}{([A]RT)^a([B]RT)^b} = \frac{[C]^c[D]^d}{[A]^a[B]^b} \times \frac{(RT)^{c+d}}{(RT)^{a+b}} = K_c(RT)^{(c+d)-(a+b)} \tag{2.4}$$

$$K_p = K_c(RT)^{\Delta n_g} \tag{2.5}$$

where $\Delta n_g = (c + d) - (a + b)$.

- If $\Delta n_g = 0 \implies K_p = K_c$ (dimensionless, pressure independent, e.g., $H_2 + I_2 \rightleftharpoons 2HI$).
- If $\Delta n_g > 0 \implies K_p > K_c$ when $RT > 1$ (units $\text{atm}^{\Delta n_g}$ or $\text{bar}^{\Delta n_g}$, e.g., $PCl_5 \rightleftharpoons PCl_3 + Cl_2$).
- If $\Delta n_g < 0 \implies K_p < K_c$ when $RT > 1$ (e.g., $N_2 + 3H_2 \rightleftharpoons 2NH_3$).

2.1.2 Le Chatelier's Principle & Equilibrium Shifts

If a dynamic equilibrium is subjected to a stress in concentration, pressure, volume, temperature, or inert gas, the system shifts in that direction which minimizes or undoes the applied constraint:

- **Concentration:** Adding reactants or removing products shifts forward ($Q < K$).
- **Pressure / Volume:** Increasing pressure (compressing volume) shifts equilibrium toward fewer gaseous moles ($\Delta n_g < 0$).
- **Temperature:** Increasing temperature shifts toward the endothermic direction ($\Delta H > 0$).
- **Inert Gas Addition:**
  - **At Constant Volume:** Total pressure rises, but partial pressures and molar concentrations of reacting gases remain **strictly unchanged** $\implies$ **NO EFFECT** on equilibrium state!
  - **At Constant Pressure:** Volume expands, partial pressures drop $\implies$ equilibrium shifts toward **greater number of gaseous moles** ($\Delta n_g > 0$).
- **Catalyst:** Increases forward and backward rates equally; equilibrium constant K and composition are completely unaffected (shortens time to achieve equilibrium).

NCERT Focus: Le Chatelier Optimization in Industrial Syntheses

Case Study: Synthesis of Ammonia (Haber-Bosch Process)

$$N_2(g) + 3H_2(g) \rightleftharpoons 2NH_3(g), \quad \Delta_r H^\circ = -92.4\,\text{kJ}\cdot\text{mol}^{-1}, \quad \Delta n_g = 2 - 4 = -2$$

- **High Pressure (≈ 200 atm):** Since $\Delta n_g < 0$, high pressure drives the equilibrium forward toward product formation, optimizing ammonia yield.
- **Optimum Temperature (≈ 700–775 K):** Because the reaction is exothermic ($\Delta_r H^\circ < 0$), low temperature thermodynamically favors ammonia, but drastically lowers kinetic reaction rates. An optimum temperature with iron oxide catalyst and promoters (K_2O and Al_2O_3) is maintained.
- **Inert Gas Injection Trap:** Adding argon or neon at constant volume (V = const) causes zero shift in equilibrium yield. Only addition at constant pressure (P = const) shifts the reaction backward (toward more gaseous moles, $\Delta n_g = -2 < 0 \implies$ leftward shift).

2.1.3 Degree of Dissociation from Vapour Density

For a gas dissociating according to $A(g) \rightleftharpoons nB(g)$:

$$\alpha = \frac{D - d}{(n-1)d} = \frac{M_{\text{theoretical}} - M_{\text{observed}}}{(n-1)M_{\text{observed}}} \tag{2.6}$$

where D is the theoretical vapour density of pure reactant, and d is the observed equilibrium mixture vapour density.

2.1.4 Ionic Equilibrium: Ostwald Dilution Law & Autoionization of Water

- **Ostwald's Dilution Law for Weak Acid** ($\mathrm{HA} \rightleftharpoons \mathrm{H^+ + A^-}$)**:**

$$K_a = \frac{C\alpha^2}{1-\alpha} \approx C\alpha^2 \quad (\text{for } \alpha \leq 0.05) \implies \alpha = \sqrt{\frac{K_a}{C}} \tag{2.7}$$

$$[\mathrm{H^+}] = C\alpha = \sqrt{K_a C} \implies \mathrm{pH} = \frac{1}{2}\left(\mathrm{p}K_a - \log_{10} C\right) \tag{2.8}$$

- **Self-Ionization of Water** ($2\mathrm{H_2O} \rightleftharpoons \mathrm{H_3O^+ + OH^-}$)**:**

$$K_w = [\mathrm{H^+}][\mathrm{OH^-}] = 1.0 \times 10^{-14} \quad (\text{at } 25^\circ\mathrm{C}) \implies \mathrm{pH} + \mathrm{pOH} = \mathrm{p}K_w = 14 \tag{2.9}$$

  Because self-ionization is endothermic ($\Delta H > 0$), K_w increases with temperature: at 80°C, $K_w \approx 10^{-13} \implies$ pH of neutral water $= \frac{13}{2} = 6.5$.

2.1.5 Salt Hydrolysis: Derivations of K_h, h, and pH

Salt hydrolysis is the reverse of neutralization, where ions of a salt interact with water to perturb $[\mathrm{H^+}]$ and $[\mathrm{OH^-}]$:

Salt Category	Hydrolysis	K_h	h	Exact pH at 25°C
SA + SB (*e.g.* NaCl)	None	—	0	pH = 7.00 (Neutral)
SA + WB (*e.g.* $\mathrm{NH_4Cl}$)	Cationic	$\frac{K_w}{K_b}$	$\sqrt{\frac{K_w}{K_b C}}$	$7 - \frac{1}{2}\mathrm{p}K_b - \frac{1}{2}\log_{10} C$
WA + SB (*e.g.* $\mathrm{CH_3COONa}$)	Anionic	$\frac{K_w}{K_a}$	$\sqrt{\frac{K_w}{K_a C}}$	$7 + \frac{1}{2}\mathrm{p}K_a + \frac{1}{2}\log_{10} C$
WA + WB (*e.g.* $\mathrm{CH_3COONH_4}$)	Both	$\frac{K_w}{K_a K_b}$	$\sqrt{\frac{K_w}{K_a K_b}}$	$7 + \frac{1}{2}\mathrm{p}K_a - \frac{1}{2}\mathrm{p}K_b$

Table 2.1: Master Classification and Analytical Formulations for Salt Hydrolysis.

2.1.6 Buffer Solutions & Henderson-Hasselbalch Derivations

A buffer resists pH changes upon addition of small quantities of strong acid or base.

1. **Acidic Buffer (Weak Acid + Salt with Strong Base, e.g., $CH_3COOH + CH_3COONa$):**

$$K_a = \frac{[H^+][A^-]}{[HA]} \implies [H^+] = K_a \frac{[HA]}{[A^-]} \approx K_a \frac{[\text{Acid}]}{[\text{Salt}]} \tag{2.10}$$

   Taking $-\log_{10}$ on both sides:

$$\text{pH} = \text{p}K_a + \log_{10}\left(\frac{[\text{Conjugate Base / Salt}]}{[\text{Weak Acid}]}\right) \tag{2.11}$$

2. **Basic Buffer** (Weak Base + Strong Acid Salt, e.g., $NH_4OH + NH_4Cl$):

$$\text{pOH} = \text{p}K_b + \log_{10}\left(\frac{[\text{Conjugate Acid / Salt}]}{[\text{Weak Base}]}\right) \tag{2.12}$$

$$\text{pH} = 14 - \text{pOH} \tag{2.13}$$

3. **Buffer Capacity (ϕ):** The number of moles of strong acid or base required to change the pH of 1 L of buffer solution by 1 unit:

$$\phi = \frac{dB}{d(\text{pH})} = 2.303 \frac{[\text{Acid}][\text{Salt}]}{[\text{Acid}] + [\text{Salt}]} \tag{2.14}$$

   Buffer capacity is **maximum** when $[\text{Acid}] = [\text{Salt}] \implies \text{pH} = \text{p}K_a$ (effective buffer range: $\text{p}K_a \pm 1$).

2.1.7 Solubility Product (K_{sp}) & Common Ion Precipitation

For a sparingly soluble salt A_xB_y:

$$A_xB_y(s) \rightleftharpoons xA^{y+}(aq) + yB^{x-}(aq) \tag{2.15}$$

If molar solubility is $S\,\text{mol}\,\text{L}^{-1}$:

$$[A^{y+}] = xS, \qquad [B^{x-}] = yS \tag{2.16}$$

$$K_{sp} = [A^{y+}]^x[B^{x-}]^y = (xS)^x(yS)^y = x^x y^y S^{x+y} \implies S = \left(\frac{K_{sp}}{x^x y^y}\right)^{\frac{1}{x+y}} \tag{2.17}$$

Solubility Equations by Salt Stoichiometry:

- AB type ($AgCl, BaSO_4$): $K_{sp} = S^2 \implies S = \sqrt{K_{sp}}$.
- AB_2 or A_2B type ($PbCl_2, Ag_2CrO_4, CaF_2$): $K_{sp} = 1^1 \times 2^2 \times S^3 = 4S^3 \implies S = \sqrt[3]{K_{sp}/4}$.

- AB_3 or A_3B type ($\mathrm{Al(OH)_3}, \mathrm{Fe(OH)_3}$): $K_{sp} = 1^1 \times 3^3 \times S^4 = 27S^4 \implies S = \sqrt[4]{K_{sp}/27}$.
- A_2B_3 or A_3B_2 type ($\mathrm{As_2S_3}, \mathrm{Ca_3(PO_4)_2}$): $K_{sp} = 2^2 \times 3^3 \times S^5 = 108S^5 \implies S = \sqrt[5]{K_{sp}/108}$.

Precipitation Criterion:

- $Q_{sp} < K_{sp} \implies$ Unsaturated solution; no precipitate forms.
- $Q_{sp} = K_{sp} \implies$ Saturated solution at dynamic equilibrium.
- $Q_{sp} > K_{sp} \implies$ Supersaturated solution; **precipitation occurs immediately**.

PART II

Advanced Physical Chemistry

CHAPTER 3

Advanced Physical Chemistry: Solutions, Electrochemistry & Chemical Kinetics

This chapter investigates advanced dynamic physical chemistry systems: homogeneous binary liquid solutions obeying Raoult's law and anomalous colligative properties via the van 't Hoff factor, electrochemical cell energetics governed by the Nernst equation and Kohlrausch conductance, and chemical kinetics exploring differential and integrated rate laws, temperature coefficients, and Arrhenius transition-state collision theory.

3.1 Solutions & Colligative Properties (Unit 5)

3.1.1 Gas Solubility & Henry's Law

Henry's Law states that the solubility of a gas in a liquid at constant temperature is directly proportional to the partial pressure of the gas above the solution:

$$p = K_H \cdot x \iff x = \frac{p}{K_H} \tag{3.1}$$

where p is the partial pressure of gas, x is the mole fraction of gas dissolved, and K_H is Henry's law constant.

- At a constant pressure p, solubility x is **inversely proportional** to K_H.
- For a given gas, K_H **increases with increasing temperature** $\implies$ gas solubility in liquids decreases on heating (aquatic life thrives in cold water due to higher dissolved O_2).

3.1.2 Vapour Pressure & Raoult's Law

For a binary solution of volatile liquids A and B, Raoult's law states that the partial vapour pressure of each component is equal to the vapour pressure of the pure component multiplied by its mole fraction in the liquid phase:

$$P_A = P_A^\circ x_A, \qquad P_B = P_B^\circ x_B \tag{3.2}$$

By Dalton's Law of Partial Pressures, the total vapour pressure is:

$$P_{\text{total}} = P_A + P_B = P_A^\circ x_A + P_B^\circ x_B = P_A^\circ(1 - x_B) + P_B^\circ x_B = P_A^\circ + (P_B^\circ - P_A^\circ)x_B \tag{3.3}$$

Vapour Phase Composition (y_A, y_B):

$$y_A = \frac{P_A}{P_{\text{total}}} = \frac{P_A^\circ x_A}{P_{\text{total}}}, \qquad y_B = \frac{P_B}{P_{\text{total}}} = \frac{P_B^\circ x_B}{P_{\text{total}}} \implies \frac{1}{P_{\text{total}}} = \frac{y_A}{P_A^\circ} + \frac{y_B}{P_B^\circ} \quad (3.4)$$

3.1.3 Ideal vs Non-Ideal Solutions & Azeotropes

Property	Ideal Solution	Positive Dev.	Negative Dev.
Intermolecular Forces	$A-B \approx A-A \approx B-B$	$A-B < A-A, B-B$ (Weaker)	$A-B > A-A, B-B$ (Stronger)
Vapour Pressure	$P_{\text{tot}} = P_A^\circ x_A + P_B^\circ x_B$	$P_{\text{tot}} > P_{\text{ideal}}$	$P_{\text{tot}} < P_{\text{ideal}}$
Enthalpy of Mixing	$\Delta H_{\text{mix}} = 0$	$\Delta H_{\text{mix}} > 0$ (Endo)	$\Delta H_{\text{mix}} < 0$ (Exo)
Volume of Mixing	$\Delta V_{\text{mix}} = 0$	$\Delta V_{\text{mix}} > 0$ (Expansion)	$\Delta V_{\text{mix}} < 0$ (Contraction)
Entropy & Free Energy	$\Delta S > 0, \Delta G < 0$	$\Delta S > 0, \Delta G < 0$	$\Delta S > 0, \Delta G < 0$
Azeotrope Formed	None (Distillable)	**Min. Boiling Azeotrope**	**Max. Boiling Azeotrope**
High-Yield Examples	Benzene + Toluene; *n*-hexane + *n*-heptane	Ethanol + Water (95.6%); Acetone + CS_2	$CHCl_3$ + Acetone; $HNO_3 + H_2O$ (68%)

Table 3.1: Comprehensive Physical and Thermodynamic Profiling of Binary Liquid Mixtures.

3.1.4 Colligative Properties & Thermodynamic Derivations

Colligative properties depend exclusively on the ratio of the number of solute particles to solvent particles, irrespective of their chemical nature.

1. Relative Lowering of Vapour Pressure (RLVP)

When a non-volatile solute B is added to volatile solvent A, only solvent molecules occupy the vapour phase: $P_{\text{total}} = P_A = P_A^\circ x_A = P_A^\circ(1 - x_B)$.

$$\frac{P_A^\circ - P}{P_A^\circ} = x_B = \frac{n_B}{n_A + n_B} \quad (3.5)$$

$$\approx \frac{n_B}{n_A} = \frac{w_B M_A}{M_B w_A} \quad \text{[for dilute solutions]} \quad (3.6)$$

Exact Molality Expression for All Concentrations:

$$\frac{P_A^\circ - P}{P} = \frac{n_B}{n_A} = \frac{m \times M_A}{1000} \implies \text{Notice denominator is } P, \text{ not } P_A^\circ! \quad (3.7)$$

Common Pitfall: Relative Lowering of Vapour Pressure Denominator

A frequent numerical trap in NEET involves confusing the theoretical RLVP definition with the exact molality formula:

- **Definition of RLVP:** By Raoult's law, relative lowering with respect to pure solvent vapour pressure is:

$$\frac{P^\circ - P}{P^\circ} = x_B = \frac{n_B}{n_A + n_B}$$

  This equation strictly possesses the pure solvent vapour pressure P° in the denominator.

- **Exact Non-Approximated Molality Conversion:** Inverting and subtracting 1 gives:

$$\frac{P^\circ - P}{P} = \frac{n_B}{n_A} = \frac{w_B \cdot M_A}{M_B \cdot w_A} = \frac{m \cdot M_A}{1000}$$

  Here the denominator is the **solution** vapour pressure P, NOT P°. Equating $(P^\circ - P)/P^\circ$ directly to n_B/n_A without dilute approximation creates systematic calculation errors in non-dilute questions ($m > 0.1$ m).

2. Elevation of Boiling Point (ΔT_b)

At boiling point, vapour pressure equals external atmospheric pressure (1 atm). Addition of non-volatile solute lowers vapour pressure; hence the solution must be heated to a higher temperature $T_b > T_b^\circ$:

$$\Delta T_b = T_b - T_b^\circ = K_b \cdot m = \frac{1000 K_b w_B}{M_B w_A} \tag{3.8}$$

Thermodynamic Derivation of Ebullioscopic Constant (K_b): From the Clausius-Clapeyron relation connecting chemical potential and vapour pressure:

$$\ln\left(\frac{P^\circ}{P}\right) = \frac{\Delta H_{\text{vap}}}{R}\left(\frac{1}{T_b^\circ} - \frac{1}{T_b}\right) \approx \frac{\Delta H_{\text{vap}} \Delta T_b}{R (T_b^\circ)^2} \tag{3.9}$$

Since $\ln(P^\circ/P) = -\ln(1 - x_B) \approx x_B \approx \frac{m M_A}{1000}$:

$$\frac{m M_A}{1000} = \frac{\Delta H_{\text{vap}} \Delta T_b}{R (T_b^\circ)^2} \implies \Delta T_b = \left[\frac{R (T_b^\circ)^2 M_A}{1000 \Delta H_{\text{vap}}}\right] m \tag{3.10}$$

$$K_b = \frac{R (T_b^\circ)^2 M_A}{1000 \Delta H_{\text{vap}}} = \frac{R (T_b^\circ)^2}{1000 l_v} \quad (l_v = \text{latent heat of vaporization per gram}) \tag{3.11}$$

For water: $K_b = 0.512 \approx 0.52\,\text{K kg mol}^{-1}$.

3. Depression of Freezing Point (ΔT_f)

Freezing point is the temperature at which the vapour pressure of the liquid solvent equals the vapour pressure of its solid phase. Lowering vapour pressure

depresses the freezing point:

$$\Delta T_f = T_f^\circ - T_f = K_f \cdot m = \frac{1000 K_f w_B}{M_B w_A} \tag{3.12}$$

Thermodynamic Derivation of Cryoscopic Constant (K_f):

$$K_f = \frac{R(T_f^\circ)^2 M_A}{1000 \Delta H_{\text{fusion}}} = \frac{R(T_f^\circ)^2}{1000 l_f} \quad (l_f = \text{latent heat of fusion per gram}) \tag{3.13}$$

For water: $K_f = 1.86\,\text{K}\,\text{kg}\,\text{mol}^{-1}$. Note that $K_f/K_b = 1.86/0.52 \approx 3.58$ (freezing point depression is over 3.5 times more sensitive than boiling point elevation!).

NEET High-Yield Application: Automotive Antifreeze Formulation

Problem Statement: An automobile radiator requires protection against freezing down to $-15.0\,^\circ\text{C}$ during harsh winter operations. Calculate the mass of ethylene glycol ($C_2H_6O_2$, molar mass $M = 62.07\,\text{g}\cdot\text{mol}^{-1}$) that must be added to 5.00 kg of water ($K_f = 1.86\,\text{K}\cdot\text{kg}\cdot\text{mol}^{-1}$).

1. **Target Freezing Point Depression:**

$$\Delta T_f = T_f^\circ - T_f = 0.00\,^\circ\text{C} - (-15.0\,^\circ\text{C}) = 15.0\,\text{K}$$

2. **Required Molality of Glycol:**

$$m = \frac{\Delta T_f}{K_f} = \frac{15.0\,\text{K}}{1.86\,\text{K}\cdot\text{kg}\cdot\text{mol}^{-1}} = 8.065\,\text{mol}\cdot\text{kg}^{-1}$$

3. **Mass of Ethylene Glycol Required:**

$$\begin{aligned} w_{\text{glycol}} &= m \times W_{\text{water}}\ (\text{in kg}) \times M_{\text{glycol}} \\ &= 8.065\,\text{mol}\cdot\text{kg}^{-1} \times 5.00\,\text{kg} \times 62.07\,\text{g}\cdot\text{mol}^{-1} \\ &\approx 2503\,\text{g} = 2.50\,\text{kg} \end{aligned}$$

Clinical / Engineering Context: In addition to lowering the freezing point, the non-volatile ethylene glycol simultaneously elevates the boiling point ($\Delta T_b = K_b \cdot m = 0.52 \times 8.065 \approx 4.19\,\text{K}$), preventing engine coolant boilover during high-temperature engine cycling.

4. Osmotic Pressure (Π)

Osmosis is the spontaneous net flow of solvent molecules through a semipermeable membrane from pure solvent into solution. Osmotic pressure Π

is the excess hydrostatic pressure that must be applied to the solution to completely arrest this solvent influx:

$$\Pi = CRT = \left(\frac{n_B}{V}\right) RT = \frac{w_B RT}{M_B V} \implies M_B = \frac{w_B RT}{\Pi V} \tag{3.14}$$

Why Osmotic Pressure is Superior for Macromolecules (Proteins, DNA, Polymers):

1. High molecular mass ($M_B \sim 10^4 - 10^6\,\mathrm{g\,mol^{-1}}$) produces immeasurably tiny molalities ($m \sim 10^{-4}\,\mathrm{m}$), causing negligible ΔT_b ($< 10^{-4}\,\mathrm{K}$), but yielding easily measurable osmotic pressure ($\Pi = 10^{-4} \times 0.0821 \times 300 \approx 2.5 \times 10^{-3}\,\mathrm{atm} \approx 26\,\mathrm{mm\ Hg}$).
2. Measurement occurs at ambient room temperature, preventing thermal denaturation of delicate biological polymers.

3.1.5 Abnormal Molar Masses & van 't Hoff Factor (i)

When a solute undergoes dissociation or association in solution, colligative properties deviate from theoretical ideal values:

$$i = \frac{\text{Experimental / Observed Colligative Property}}{\text{Calculated / Normal Colligative Property}} = \frac{M_{\text{theoretical / normal}}}{M_{\text{observed / abnormal}}} \tag{3.15}$$

Derivation of Degree of Dissociation (α_{diss}) and Association (α_{assoc})

- **Case 1: Dissociation ($A \rightleftharpoons nB$):** Let 1 mole of electrolyte be dissolved with degree of dissociation α:

$$\begin{array}{lcc} & A & \rightleftharpoons & nB \\ \text{Initial moles:} & 1 & & 0 \\ \text{Equilibrium moles:} & 1-\alpha & & n\alpha \end{array} \tag{3.16}$$

  Total moles at equilibrium $= 1 - \alpha + n\alpha = 1 + (n-1)\alpha$.

$$i = \frac{1 + (n-1)\alpha}{1} \implies \alpha_{\text{diss}} = \frac{i-1}{n-1} \tag{3.17}$$

- **Case 2: Association / Polymerization ($nA \rightleftharpoons A_n$):** Let 1 mole of solute undergo association to an n-mer with degree of association α:

$$\begin{array}{lcc} & nA & \rightleftharpoons & A_n \\ \text{Initial moles:} & 1 & & 0 \\ \text{Equilibrium moles:} & 1-\alpha & & \frac{\alpha}{n} \end{array} \tag{3.18}$$

  Total moles at equilibrium $= 1 - \alpha + \frac{\alpha}{n} = 1 - \left(1 - \frac{1}{n}\right)\alpha$.

$$i = 1 - \left(1 - \frac{1}{n}\right)\alpha \implies \alpha_{\text{assoc}} = \frac{1-i}{1-\frac{1}{n}} \tag{3.19}$$

For dimerization of acetic acid in benzene ($n = 2$): $i = 1 - \frac{\alpha}{2} \implies$ if 100% dimerized ($\alpha = 1$), $i = 0.5$ and observed molar mass is double ($120\,\mathrm{g\,mol^{-1}}$ vs $60\,\mathrm{g\,mol^{-1}}$).

3.2 Redox Reactions & Electrochemistry (Unit 7)

3.2.1 Galvanic Cells & The Daniell Cell

A galvanic cell converts spontaneous chemical redox free energy into electrical work:

$$\mathrm{Zn(s) + Cu^{2+}(aq) \longrightarrow Zn^{2+}(aq) + Cu(s)} \tag{3.20}$$

- **Anode (Oxidation, Negative Terminal):** $\mathrm{Zn(s) \rightarrow Zn^{2+}(aq) + 2}e^-$, $E^\circ_{\mathrm{Zn^{2+}/Zn}} = -0.76\,\mathrm{V}$.
- **Cathode (Reduction, Positive Terminal):** $\mathrm{Cu^{2+}(aq) + 2}e^- \rightarrow \mathrm{Cu(s)}$, $E^\circ_{\mathrm{Cu^{2+}/Cu}} = +0.34\,\mathrm{V}$.
- **Cell Representation:** $\mathrm{Zn(s) \mid Zn^{2+}(aq) \parallel Cu^{2+}(aq) \mid Cu(s)}$.
- **Salt Bridge Functions:**
  1. Maintains electrical neutrality in both half-cells by migrating counter-ions ($\mathrm{K^+}$ and $\mathrm{Cl^-}$ or $\mathrm{NO_3^-}$).
  2. Prevents physical mixing of the two electrolytic solutions.
  3. Minimizes liquid junction potential. Electrolytes (KCl, $\mathrm{KNO_3}$, $\mathrm{NH_4NO_3}$) have nearly equal ionic mobilities ($v_+ \approx v_-$).

3.2.2 Standard Hydrogen Electrode & Electrochemical Series

The Standard Hydrogen Electrode (SHE) is the universal reference standard assigned $E^\circ = 0.000\,\mathrm{V}$ at all temperatures:

$$\mathrm{Pt(s) \mid H_2(g, 1\,bar) \mid H^+(aq, 1\,M)}, \qquad \mathrm{2H^+ + 2}e^- \rightleftharpoons \mathrm{H_2(g)} \tag{3.21}$$

Standard Cell EMF (E°_{cell}):

$$E^\circ_{\mathrm{cell}} = E^\circ_{\mathrm{cathode}} - E^\circ_{\mathrm{anode}} = E^\circ_{\mathrm{right}} - E^\circ_{\mathrm{left}} \tag{3.22}$$

(where reduction potentials are used for both electrodes).

3.2.3 Thermodynamic Derivation of the Nernst Equation

The maximum non-expansion electrical work delivered by a galvanic cell at constant T, P equals the decrease in Gibbs free energy:

$$w_{\mathrm{elec}} = q_{\mathrm{charge}} \times E_{\mathrm{cell}} = (nF)E_{\mathrm{cell}} \implies \Delta_r G = -w_{\mathrm{elec}} = -nFE_{\mathrm{cell}} \tag{3.23}$$

Under standard state conditions: $\Delta_r G^\circ = -nFE^\circ_{\text{cell}}$. Substituting these relations into the fundamental thermodynamic equation $\Delta_r G = \Delta_r G^\circ + RT\ln Q$:

$$-nFE_{\text{cell}} = -nFE^\circ_{\text{cell}} + RT\ln Q \tag{3.24}$$

Dividing throughout by $-nF$:

$$E_{\text{cell}} = E^\circ_{\text{cell}} - \frac{RT}{nF}\ln Q = E^\circ_{\text{cell}} - \frac{2.303RT}{nF}\log_{10} Q \tag{3.25}$$

Substituting universal constants $R = 8.314\,\text{J mol}^{-1}\,\text{K}^{-1}$, $F = 96485\,\text{C mol}^{-1}$, and $T = 298.15\,\text{K}$:

$$E_{\text{cell}} = E^\circ_{\text{cell}} - \frac{0.0591}{n}\log_{10} Q \tag{3.26}$$

Nernst Equation Special Cases & Thermodynamic Properties

- **Dynamic Equilibrium ($E_{\text{cell}} = 0, Q = K_c$):**

$$0 = E^\circ_{\text{cell}} - \frac{0.0591}{n}\log_{10} K_c \implies \log_{10} K_c = \frac{nE^\circ_{\text{cell}}}{0.0591} \tag{3.27}$$

- **Concentration Cells (Identical Electrodes, Different Concentrations $C_1 < C_2$):** Since standard potentials are identical, $E^\circ_{\text{cell}} = 0$:

$$E_{\text{cell}} = -\frac{0.0591}{n}\log_{10}\left(\frac{C_1}{C_2}\right) = \frac{0.0591}{n}\log_{10}\left(\frac{C_2}{C_1}\right) \tag{3.28}$$

  (Reaction is spontaneous if $C_2 > C_1$).
- **Temperature Coefficient of EMF and Entropy of Cell Reaction:**

$$\left(\frac{\partial \Delta G}{\partial T}\right)_P = -\Delta S \implies \Delta S = nF\left(\frac{\partial E_{\text{cell}}}{\partial T}\right)_P \tag{3.29}$$

$$\Delta H = \Delta G + T\Delta S = -nFE_{\text{cell}} + nFT\left(\frac{\partial E_{\text{cell}}}{\partial T}\right)_P \tag{3.30}$$

3.2.4 Electrolytic Conductance & Kohlrausch's Law

- **Conductivity (κ):** Conductance of a unit volume ($1\,\text{cm}^3$ or $1\,\text{m}^3$) of electrolytic solution:

$$R = \rho\frac{l}{A} \implies G = \frac{1}{R} = \kappa\frac{A}{l} \implies \kappa = \frac{1}{R}\left(\frac{l}{A}\right) = G \cdot G^* \tag{3.31}$$

  where $G^* = l/A$ is the cell constant (units cm^{-1} or m^{-1}).

- **Molar Conductivity (Λ_m):** Conductance of volume of solution containing 1 mole of electrolyte:

$$\Lambda_m = \frac{\kappa \times 1000}{M \text{ (in Molarity)}} \quad (\text{in S cm}^2\,\text{mol}^{-1}) = \frac{\kappa}{1000 \times M} \quad (\text{in S m}^2\,\text{mol}^{-1}) \tag{3.32}$$

- **Dilution Effects:**
  - **Conductivity (κ) Decreases with Dilution:** Dilution reduces the number of current-carrying ions per unit volume.
  - **Molar Conductivity (Λ_m) Increases with Dilution:** Volume containing 1 mole expands, overcompensating the decrease in κ.
- **Strong vs Weak Electrolytes:**
  - **Strong Electrolytes:** Obey Debye-Hückel-Onsager equation: $\Lambda_m = \Lambda_m^\circ - A\sqrt{C}$. Plot of Λ_m vs $\sqrt{C}$ is linear and can be extrapolated to $C = 0$ to obtain limiting molar conductivity Λ_m°.
  - **Weak Electrolytes (CH_3COOH):** Λ_m increases steeply near infinite dilution; cannot be obtained by graphical extrapolation.

Kohlrausch's Law of Independent Migration of Ions

At infinite dilution, each ion migrates independently of its co-ion and makes a definite contribution to the total molar conductivity:

$$\Lambda_m^\circ = \nu_+\lambda_+^\circ + \nu_-\lambda_-^\circ \tag{3.33}$$

High-Yield Applications of Kohlrausch's Law:

1. **Limiting Molar Conductivity of Weak Electrolytes (CH_3COOH):**

$$\Lambda_m^\circ(CH_3COOH) = \Lambda_m^\circ(CH_3COONa) + \Lambda_m^\circ(HCl) - \Lambda_m^\circ(NaCl) \tag{3.34}$$

2. **Degree of Dissociation of Weak Electrolyte (α):**

$$\alpha = \frac{\Lambda_m}{\Lambda_m^\circ} \implies K_a = \frac{C\alpha^2}{1-\alpha} = \frac{C\left(\frac{\Lambda_m}{\Lambda_m^\circ}\right)^2}{1-\frac{\Lambda_m}{\Lambda_m^\circ}} = \frac{C\Lambda_m^2}{\Lambda_m^\circ(\Lambda_m^\circ - \Lambda_m)} \tag{3.35}$$

3. **Solubility and K_{sp} of Sparingly Soluble Salts ($AgCl, BaSO_4$):**

$$\Lambda_m^\circ = \frac{1000\kappa_{\text{salt}}}{S} \implies S = \frac{1000\kappa_{\text{salt}}}{\Lambda_m^\circ} \implies K_{sp} = S^2 \tag{3.36}$$

where $\kappa_{\text{salt}} = \kappa_{\text{solution}} - \kappa_{\text{water}}$.

3.2.5 Electrolysis & Faraday's Laws

- **Faraday's First Law:** Mass of substance deposited or liberated at any electrode is directly proportional to the total electrical charge passed:

$$w = Z \cdot Q = Z \cdot I \cdot t = \left(\frac{E}{96500}\right) I \cdot t = \left(\frac{M}{nF}\right) I \cdot t \tag{3.37}$$

  where $Z = \frac{E}{F}$ is the electrochemical equivalent, and $E = \frac{M}{n}$ is equivalent mass.
- **Faraday's Second Law:** When the identical quantity of electricity is passed through different electrolytes connected in series, masses deposited are proportional to their chemical equivalent masses:

$$\frac{w_1}{w_2} = \frac{E_1}{E_2} \tag{3.38}$$

- **Preferential Discharge Traps:**
  - **Molten** NaCl**:** Cathode: $Na^+ \rightarrow Na(l)$; Anode: $2Cl^- \rightarrow Cl_2 + 2e^-$.
  - **Aqueous** NaCl**:** Cathode: H_2 ($E^\circ = -0.83\,V$). Anode: Cl_2 (Pt overpotential).
  - **Aqueous** $CuSO_4$ **(Pt):** Cathode: $Cu^{2+} + 2e^- \rightarrow Cu(s)$; Anode: $2H_2O \rightarrow O_2 + 4H^+ + 4e^-$.
  - **Aqueous** $CuSO_4$ **(Cu):** Cathode: Cu deposition; Anode: Cu dissolution ($Cu \rightarrow Cu^{2+} + 2e^-$).

3.2.6 Commercial Batteries & Mechanism of Corrosion

Batteries & Corrosion Reaction Architectures

- **Lead-Acid Storage Battery (Secondary Reversible Cell):**
  - **Discharging Half-Reactions:**

$$\text{Anode: } Pb(s) + SO_4^{2-}(aq) \longrightarrow PbSO_4(s) + 2e^- \tag{3.39}$$

$$\text{Cathode: } PbO_2(s) + SO_4^{2-}(aq) + 4H^+(aq) + 2e^- \tag{3.40}$$

$$\longrightarrow PbSO_4(s) + 2H_2O(l) \tag{3.41}$$

$$\text{Net: } Pb(s) + PbO_2(s) + 2H_2SO_4(aq) \tag{3.42}$$

$$\longrightarrow 2PbSO_4(s) + 2H_2O(l) \tag{3.43}$$

    Density of H_2SO_4 drops from $1.30\,g\,cm^{-3}$ to $< 1.20\,g\,cm^{-3}$. Recharging reverses all signs.
- **Hydrogen-Oxygen Fuel Cell:**

$$\text{Anode: } 2H_2 + 4OH^- \longrightarrow 4H_2O + 4e^- \tag{3.44}$$

$$\text{Cathode: } O_2 + 2H_2O + 4e^- \longrightarrow 4OH^- \tag{3.45}$$

Theoretical thermodynamic efficiency: $\eta = \frac{\Delta G^\circ}{\Delta H^\circ} \times 100\% \approx 70\% - 75\%$ (far exceeds thermal engines).

- **Electrochemical Mechanism of Rusting of Iron:**

$$\text{Anodic site: } Fe(s) \longrightarrow Fe^{2+}(aq) + 2e^- \quad (E^\circ = -0.44\,V) \tag{3.46}$$

$$\text{Cathodic site: } O_2(g) + 4H^+(aq) + 4e^- \longrightarrow 2H_2O(l) \quad (E^\circ = +1.23\,V) \tag{3.47}$$

$$\text{Rust: } 4Fe^{2+} + O_2 + 4H_2O + 2xH_2O \tag{3.48}$$

$$\longrightarrow 2Fe_2O_3 \cdot xH_2O(s) + 8H^+ \tag{3.49}$$

Prevented by galvanization (sacrificial zinc coating: $E^\circ_{Zn^{2+}/Zn} = -0.76\,V < E^\circ_{Fe^{2+}/Fe} = -0.44\,V$).

3.3 Chemical Kinetics (Unit 8)

3.3.1 Reaction Rate, Rate Laws & Order vs Molecularity

For a general reaction:

$$aA + bB \longrightarrow cC + dD \tag{3.50}$$

The instantaneous stoichiometric rate of reaction is:

$$r = -\frac{1}{a}\frac{d[A]}{dt} = -\frac{1}{b}\frac{d[B]}{dt} = +\frac{1}{c}\frac{d[C]}{dt} = +\frac{1}{d}\frac{d[D]}{dt} \tag{3.51}$$

Experimental Differential Rate Law:

$$r = k[A]^x[B]^y \tag{3.52}$$

where k is the specific rate constant, x is the order with respect to A, y is the order with respect to B, and $n = x + y$ is the overall reaction order.

Units of Rate Constant (k):

$$\text{Units of } k = \left(\text{mol L}^{-1}\right)^{1-n} \text{s}^{-1} = (\text{atm})^{1-n}\,\text{s}^{-1} \tag{3.53}$$

- Zero-order ($n = 0$): $\text{mol L}^{-1}\,\text{s}^{-1}$.
- First-order ($n = 1$): s^{-1} (completely independent of concentration units!).
- Second-order ($n = 2$): $\text{L mol}^{-1}\,\text{s}^{-1}$.

Feature	Order of Reaction	Molecularity
Definition	Sum of powers of reactant concentrations in rate law.	Reacting species colliding simultaneously in elementary step.
Determination	Strictly experimental parameter; cannot be deduced from equation.	Theoretical property deduced from elementary mechanism.
Allowed Values	Can be zero, integer, negative, or fractional $(0, 1, 2, 1/2)$.	Positive non-zero integer $(1, 2, 3)$. Cannot be zero/fractional!
Upper Bound	Can exceed 3.	Rarely exceeds 3 due to low probability of simultaneous collisions.
Complex Reactions	Applicable to overall complex reactions.	Meaningless for overall complex reaction; applies to elementary steps.

Table 3.2: Systematic Comparison: Order of Reaction vs Molecularity.

3.3.2 Integrated Rate Equations: Full Calculus Derivations

1. Zero-Order Kinetics

For $A \rightarrow$ Products, rate is independent of $[A]$:

$$-\frac{d[A]}{dt} = k[A]^0 = k \implies d[A] = -kdt \tag{3.54}$$

Integrating from $t = 0$ ($[A] = [A]_0$) to time t ($[A] = [A]_t$):

$$\int_{[A]_0}^{[A]_t} d[A] = -k \int_0^t dt \implies [A]_t - [A]_0 = -kt \implies [A]_t = [A]_0 - kt \tag{3.55}$$

Half-Life ($t_{1/2}$) and Completion Time ($t_{100\%}$): At $t = t_{1/2}$, $[A]_t = \frac{[A]_0}{2}$:

$$\frac{[A]_0}{2} = [A]_0 - kt_{1/2} \implies t_{1/2} = \frac{[A]_0}{2k} \tag{3.56}$$

(Half-life of a zero-order reaction is directly proportional to initial concentration). At $t = t_{100\%}$, $[A]_t = 0$:

$$0 = [A]_0 - kt_{100\%} \implies t_{100\%} = \frac{[A]_0}{k} = 2t_{1/2} \tag{3.57}$$

Zero-order reactions proceed to complete termination in a finite time. **Examples:** Photochemical synthesis of HCl over water ($H_2 + Cl_2 \xrightarrow{h\nu} 2HCl$); decomposition of NH_3 on hot Pt/W surface at high pressure.

2. First-Order Kinetics

For $A \rightarrow$ Products:

$$-\frac{d[A]}{dt} = k[A] \implies \frac{d[A]}{[A]} = -kdt \tag{3.58}$$

Integrating from $t = 0$ ($[A] = [A]_0$) to time t ($[A] = [A]_t$):

$$\int_{[A]_0}^{[A]_t} \frac{d[A]}{[A]} = -k \int_0^t dt \implies \ln\left(\frac{[A]_t}{[A]_0}\right) = -kt \implies [A]_t = [A]_0 e^{-kt} \quad (3.59)$$

$$k = \frac{1}{t} \ln\left(\frac{[A]_0}{[A]_t}\right) = \frac{2.303}{t} \log_{10}\left(\frac{[A]_0}{[A]_t}\right) = \frac{2.303}{t} \log_{10}\left(\frac{a}{a-x}\right) \quad (3.60)$$

Half-Life ($t_{1/2}$): At $t = t_{1/2}$, $[A]_t = \frac{[A]_0}{2}$:

$$k = \frac{2.303}{t_{1/2}} \log_{10}(2) \implies t_{1/2} = \frac{\ln 2}{k} = \frac{0.69315}{k} \approx \frac{0.693}{k} \quad (3.61)$$

The half-life of a first-order reaction is **strictly independent of initial reactant concentration**.

NCERT Paradigm: Fractional Lifetimes of First-Order Reactions

For any first-order process ($[A]_t = [A]_0 e^{-kt}$), each half-life interval reduces the remaining reactant concentration by exactly 50%, regardless of initial amount. Rapid shortcuts for NEET:

$$t_{75\%} = \frac{2.303}{k} \log_{10}\left(\frac{100}{25}\right) = \frac{2.303}{k} \log_{10}(4) = 2 \times \frac{0.693}{k} = 2t_{1/2}$$

$$t_{87.5\%} = \frac{2.303}{k} \log_{10}\left(\frac{100}{12.5}\right) = \frac{2.303}{k} \log_{10}(8) = 3t_{1/2}$$

$$t_{93.75\%} = 4t_{1/2} \quad \text{(remaining fraction } = 6.25\%)$$

$$t_{99\%} = \frac{2.303}{k} \log_{10}(100) = \frac{2 \times 2.303}{k} \approx \frac{4.606}{k} \approx 6.64t_{1/2}$$

$$t_{99.9\%} = \frac{2.303}{k} \log_{10}\left(\frac{100}{0.1}\right) = \frac{3 \times 2.303}{k} \approx \frac{6.909}{k} \approx 10t_{1/2}$$

NCERT Benchmark Conclusion: The time required for 99.9% completion of a first-order reaction is approximately 10 times its half-life ($t_{99.9\%} \approx 10t_{1/2}$), while $t_{99\%} \approx 2t_{90\%}$.

Gas Phase First-Order Decomposition ($A(g) \rightarrow B(g) + C(g)$): If initial pressure of A is P_i and total pressure at time t is P_t:

$$P_t = (P_i - p) + p + p = P_i + p \implies p = P_t - P_i \implies P_A = P_i - p = 2P_i - P_t \quad (3.62)$$

$$k = \frac{2.303}{t} \log_{10}\left(\frac{P_i}{2P_i - P_t}\right) \quad (3.63)$$

3. Pseudo-First-Order Kinetics

Reactions of higher order that behave kinetically as first-order when one reactant is present in large stoichiometric excess:

- **Acid-Catalyzed Ester Hydrolysis:**

$$\mathrm{CH_3COOC_2H_5 + H_2O \xrightarrow{H^+} CH_3COOH + C_2H_5OH} \tag{3.64}$$

Because water is solvent ($[\mathrm{H_2O}] \approx 55.5\,\mathrm{M} \approx$ constant):

$$r = k'[\text{ester}][\mathrm{H_2O}] = k[\text{ester}], \qquad k = k'[\mathrm{H_2O}] \tag{3.65}$$

Tracked by titrating aliquots with standard NaOH:
$k = \frac{2.303}{t} \log_{10}\left(\frac{V_\infty - V_0}{V_\infty - V_t}\right)$.

- **Inversion of Cane Sugar (Polarimetry):**

$$\mathrm{C_{12}H_{22}O_{11} + H_2O \xrightarrow{H^+} C_6H_{12}O_6\ (\text{D-glucose}) + C_6H_{12}O_6\ (\text{D-fructose})} \tag{3.66}$$

The optical rotation changes from dextro ($+66.5°$) to levo ($-39.7°$). Kinetics tracked via polarimeter angle of rotation:

$$k = \frac{2.303}{t} \log_{10}\left(\frac{r_0 - r_\infty}{r_t - r_\infty}\right) \tag{3.67}$$

3.3.3 Temperature Dependence: The Arrhenius Equation

Svante Arrhenius (1889) derived the exponential relationship between reaction rate constant and absolute temperature:

$$k = Ae^{-E_a/RT} \tag{3.68}$$

where A is the pre-exponential factor (frequency factor), E_a is the activation energy, and $e^{-E_a/RT}$ represents the Boltzmann fraction of molecular collisions having kinetic energy $\geq E_a$.

$$\ln k = \ln A - \frac{E_a}{RT} \iff \log_{10} k = \log_{10} A - \frac{E_a}{2.303RT} \tag{3.69}$$

An Arrhenius plot of $\log_{10} k$ versus $\frac{1}{T}$ produces a straight line:

$$\text{Slope} = -\frac{E_a}{2.303R}, \qquad y\text{-intercept} = \log_{10} A \tag{3.70}$$

Two-Temperature Comparative Formulation:

$$\log_{10}\left(\frac{k_2}{k_1}\right) = \frac{E_a}{2.303R}\left[\frac{T_2 - T_1}{T_1 T_2}\right] = \frac{E_a}{2.303R}\left[\frac{1}{T_1} - \frac{1}{T_2}\right] \tag{3.71}$$

3.3.4 Collision Theory & Catalysis Kinetics

According to Max Trautz and William Lewis:

$$k = P \cdot Z_{AB} \cdot e^{-E_a/RT} \tag{3.72}$$

where Z_{AB} is the collision frequency ($10^{25} - 10^{28}$ collisions $\mathrm{s}^{-1}\,\mathrm{cm}^{-3}$), and P is the steric or orientation factor representing the fraction of collisions occurring with proper spatial orientation.

Mechanism of Catalytic Acceleration: A catalyst accelerates the rate of a chemical transformation without undergoing permanent stoichiometric consumption. It achieves this kinetic enhancement not by shifting molecular energy distributions, but by providing an alternative mechanistic pathway involving an intermediate activated complex characterized by a significantly lower activation barrier E_a' ($E_a' < E_a$), as depicted in Figure 3.1.

According to the Arrhenius relation, the relative catalytic rate enhancement factor is given by:

$$\frac{k_{\text{cat}}}{k_{\text{uncat}}} = \frac{Ae^{-E_a'/RT}}{Ae^{-E_a/RT}} = e^{(E_a - E_a')/RT} = e^{\Delta E_a/RT} \tag{3.73}$$

Because the reduction ΔE_a resides in the numerator of an exponential exponent, even a modest lowering of the activation energy (e.g., by $15\,\mathrm{kJ\,mol}^{-1}$) accelerates reaction velocity by several orders of magnitude at physiological temperature.




Figure 3.1: Potential Energy Diagram for an Exothermic Reaction Illustrating the Lowering of Activation Energy by a Catalyst.

NTA Assertion-Reason Traps: Thermodynamic Invariants of Catalysis

Candidates frequently lose marks on Assertion-Reason questions involving catalytic effects:

- **Equal Barrier Reduction:** A catalyst reduces activation energy for **both forward and reverse reactions by the exact same amount**: $\Delta E_{a,f} = \Delta E_{a,b}$.
- **No Change in Reaction Enthalpy:** Because $\Delta_r H = E_{a,f} - E_{a,b}$, the reaction enthalpy is **strictly unaltered** by the presence of a catalyst.
- **No Change in Equilibrium Constant:** A catalyst accelerates the approach to equilibrium but does **NOT** alter the equilibrium constant K_c or shift equilibrium concentrations.
- **No Change in Spontaneity:** Standard Gibbs free energy $\Delta_r G^\circ$ is completely independent of catalytic mechanisms. A catalyst cannot drive a non-spontaneous ($\Delta G > 0$) process.

PART III

Inorganic Chemistry & Coordination Compounds

CHAPTER 4

Inorganic Chemistry: Periodic Trends, Chemical Bonding, Coordination & Block Elements

Inorganic chemistry requires structural understanding rather than rote memorization. In this comprehensive compendium, we analyze modern periodic trends, molecular orbital theory (MOT) and VSEPR geometries, coordination compounds with crystal field theory (CFT) splitting, and comparative trends across the p-block and transition d/f-block elements.

A Personal Note from Satya: The Mental Shift from Math to Molecular Geometry

Welcome to Inorganic Chemistry! Let's talk honestly about why students fear this subject.
Most students love Physical Chemistry because it feels like Physics: you plug numbers into formulas ($PV = nRT$, $\Delta G^\circ = -nFE^\circ$), crunch the math, and get an answer. Then you arrive at Inorganic Chemistry, open a traditional coaching module, see 400 pages of reactions and exceptions, and your brain freezes in panic. Here is the secret: *Inorganic Chemistry is not a collection of random facts; it is the physical architecture of electron clouds.*

- Every periodic anomaly (why Gallium is slightly smaller than Aluminum, why Oxygen has a lower ionization energy than Nitrogen) is rooted in two bedrock concepts: **effective nuclear charge (Z_{eff})** and **electron-electron repulsion in half-filled subshells**.
- Coordination Chemistry becomes pure geometry when you visualize how incoming negative ligands split the five degenerate d-orbitals into t_{2g} and e_g energy levels. Once you see that diagram, high-spin, low-spin, magnetic moments, and colors become obvious.

Take Strategic 5-Minute Brain Breaks: When memorizing chemical properties and group trends, your working memory fills up quickly. Study in 25-minute focused sprints. When the timer rings, stand up, step away from your desk, drink water, and close your eyes for 5 minutes. Do not pick up your phone. Giving your brain zero visual input for five minutes allows the hippocampus to file the atomic radii trends into long-term storage without cognitive interference.

4.1 NMC 2027 Regulatory Calibration & Scope Demarcation

Zero-Tolerance NMC Syllabus Alert: Deleted Inorganic Chemistry Chapters

Under the revised National Medical Commission (NMC) NEET UG 2027 regulatory curriculum, the following legacy chapters have been **strictly deleted and removed from the examination**:

1. **Hydrogen:** Position in periodic table, isotopes, preparation and properties of dihydrogen, hydrides (ionic, covalent, interstitial), physical and chemical properties of water, hard/soft water, heavy water (D_2O), and hydrogen peroxide (H_2O_2) are **completely eliminated**.
2. ***s*-Block Elements (Alkali & Alkaline Earth Metals):** Group 1 and Group 2 chemistry, electronic configurations, diagonal relationships, flame coloration, and industrial compounds (Na_2CO_3, NaCl, NaOH, $NaHCO_3$, CaO, $CaCO_3$, $CaSO_4 \cdot \frac{1}{2}H_2O$, cement) are **completely eliminated**.
3. **General Principles & Processes of Isolation of Elements (Metallurgy):** Ores, mineral dressing, roasting, calcination, smelting, Ellingham diagrams, blast furnace iron extraction, copper reverberatory smelting, Hall-Héroult aluminum extraction, and refining techniques (Mond process, van Arkel method, zone refining) are **completely eliminated**.
4. **Environmental Chemistry:** Tropospheric/stratospheric pollution, classical and photochemical smog, ozone depletion, and green chemistry are **strictly eliminated**.

Study Directive: Do not invest study hours in these eliminated topics. All cognitive focus must remain on the rigorously tested Class 11 and Class 12 inorganic chemistry compendium units detailed below.

4.2 Classification of Elements & Periodicity in Properties (Unit 9)

4.2.1 Modern Periodic Law & Electronic Architecture

Moseley's Law (1913) demonstrated that the square root of the characteristic X-ray frequency ($\sqrt{\nu}$) emitted by an element is directly proportional to its atomic number Z:

$$\sqrt{\nu} = a(Z - b) \tag{4.1}$$

Thus, the physical and chemical properties of the elements are periodic functions of their **atomic numbers**, not their atomic masses. The long form of the periodic table consists of 7 horizontal periods and 18 vertical groups, partitioned into s, p, d, and f blocks based on the differentiating electron orbital.

4.2.2 Effective Nuclear Charge (Z_{eff}) & Slater's Rules

Electrons in outer shells are electrostatically shielded from the full nuclear charge $+Ze$ by intervening inner-shell electron density:

$$Z_{\text{eff}} = Z - \sigma \tag{4.2}$$

where σ is the screening or shielding constant.

- **Shielding Efficiency Hierarchy:** $s > p > d > f$. Spherical s-orbitals penetrate closest to the nucleus, providing maximal screening. Diffuse d- and f-orbitals have poor radial penetration and provide **extraordinarily poor shielding**.
- **Slater's Rules Formulation for ns, np Valence Electrons:**
  1. Electrons in shells higher than the considered electron contribute 0.00.
  2. Other electrons in the identical (ns, np) group contribute 0.35 each (or 0.30 if $1s$).
  3. Each electron in the $(n-1)$ shell contributes 0.85.
  4. Each electron in the $(n-2)$ or lower inner shells contributes 1.00.
- **For nd, nf Valence Electrons:**
  1. Other electrons in the identical (nd, nf) group contribute 0.35 each.
  2. All electrons in all lower underlying shells contribute 1.00 each.

High-Yield Problem Archetype: Step-by-Step Slater's Z_{eff} Calculation

Archetype Problem: Calculate the effective nuclear charge (Z_{eff}) experienced by: (a) a $4s$ valence electron in Zinc ($Z = 30$), and (b) a $3d$ electron in Zinc ($Z = 30$).

1. **Electronic Configuration Grouping:**

$$(1s)^2\ (2s, 2p)^8\ (3s, 3p)^8\ (3d)^{10}\ (4s)^2$$

2. **For the $4s$ Valence Electron ($n = 4$):**
   - 1 other electron in same group ($4s$): $1 \times 0.35 = 0.35$.
   - $(n-1)$ shell (both $3s, 3p$ and $3d$, total 18 electrons): $18 \times 0.85 = 15.30$.
   - $(n-2)$ and lower shells ($1s, 2s, 2p$, total 10 electrons): $10 \times 1.00 = 10.00$.
   - $\sigma_{4s} = 0.35 + 15.30 + 10.00 = 25.65 \implies Z_{\text{eff}} = 30 - 25.65 = \mathbf{4.35}$.
3. **For a $3d$ Electron ($n = 3$):**
   - $4s$ electrons lie in a higher shell $\implies$ contribute 0.00.
   - 9 other electrons in the same ($3d$) group: $9 \times 0.35 = 3.15$.
   - All 18 electrons in lower shells: $18 \times 1.00 = 18.00$.
   - $\sigma_{3d} = 3.15 + 18.00 = 21.15 \implies Z_{\text{eff}} = 30 - 21.15 = \mathbf{8.85}$.
4. **Conclusion:** $3d$ electrons experience more than double the effective nuclear attraction (8.85 vs 4.35) compared to $4s$ electrons. Hence, upon ionization,

the $4s$ electrons are expelled first: $\mathrm{Zn} \rightarrow \mathrm{Zn}^{2+} + 2\mathrm{e}^-$ yields $[\mathrm{Ar}]3d^{10}$, not $[\mathrm{Ar}]3d^8 4s^2$!

4.2.3 Atomic and Ionic Radii Trends & Contractions

- **Radius Hierarchy:** For the same element: $r_{\text{cation}} < r_{\text{atom}} < r_{\text{anion}}$. Loss of electrons increases $Z_{\text{eff}} = Z/e$, pulling electrons closer. Gain of electrons increases interelectronic repulsion, expanding the electron cloud.
- **Isoelectronic Series Size Rule:** For ions sharing the identical total electron count:

$$r_{\text{isoelectronic}} \propto \frac{1}{Z} \implies \mathrm{C}^{4-} > \mathrm{N}^{3-} > \mathrm{O}^{2-} > \mathrm{F}^- \tag{4.3}$$

$$> \mathrm{Na}^+ > \mathrm{Mg}^{2+} > \mathrm{Al}^{3+} > \mathrm{Si}^{4+} \quad (10e^-) \tag{4.4}$$

- **$3d$-Transition Metal Contraction (d-Block Contraction):** Filling of the ten $3d$ orbitals across the first transition series precedes Gallium ($Z = 31$: $[\mathrm{Ar}]3d^{10}4s^2 4p^1$). Because $3d$ electrons shield poorly, Z_{eff} on the valence $4p$ electron is abnormally high. Consequently:

$$r(\mathrm{Ga}) = 135\,\mathrm{pm} \leq r(\mathrm{Al}) = 143\,\mathrm{pm} \tag{4.5}$$

- **Lanthanoid Contraction ($4f$-Contraction):** Filling of the fourteen diffuse $4f$ orbitals between Lanthanum ($Z = 57$) and Hafnium ($Z = 72$) results in extremely poor screening of the increasing nuclear charge (+14 units). **Consequences:**
  1. Nearly identical covalent and ionic radii for $4d$ and $5d$ congeneric pairs:

$$r(\mathrm{Zr}) \approx r(\mathrm{Hf}) \approx 159\text{–}160\,\mathrm{pm}, \qquad r(\mathrm{Nb}) \approx r(\mathrm{Ta}) \approx 146\,\mathrm{pm} \tag{4.6}$$

$$r(\mathrm{Mo}) \approx r(\mathrm{W}) \approx 139\,\mathrm{pm} \tag{4.7}$$

  2. High density of $5d$ elements (roughly double their $4d$ counterparts; e.g., Osmium $\rho \approx 22.6\,\mathrm{g\,cm^{-3}}$).
  3. Regular decrease in basicity of lanthanoid hydroxides: $\mathrm{La(OH)_3} > \mathrm{Ce(OH)_3} > \cdots > \mathrm{Lu(OH)_3}$ (covalent character increases as cation size contracts).

4.2.4 Ionization Enthalpy ($\Delta_i H$) & High-Yield Anomalies

Ionization Enthalpy is the minimum energy required to remove the most loosely bound electron from an isolated gaseous atom in its ground state:

$$\mathrm{X(g)} \xrightarrow{\mathrm{IE_1}} \mathrm{X^+(g)} + e^-, \qquad \mathrm{X^+(g)} \xrightarrow{\mathrm{IE_2}} \mathrm{X^{2+}(g)} + e^- \tag{4.8}$$

Successive values always increase: $\mathrm{IE_1} < \mathrm{IE_2} < \mathrm{IE_3} < \ldots$ because removing an electron from an increasingly positive ion encounters greater electrostatic attraction.

Element Pair	Observed Trend		Theoretical Origin & NCERT Rationale
Be vs B	IE_1(Be) $899\,kJ\,mol^{-1}$ $>$ IE_1(B) $801\,kJ\,mol^{-1}$	$=$ $=$	Be ($2s^2$) loses electron from fully-filled subshell. B ($2s^2 2p^1$) loses less penetrating $2p$ electron of higher energy.
N vs O	IE_1(N) $1402\,kJ\,mol^{-1}$ $>$ IE_1(O) $1314\,kJ\,mol^{-1}$	$=$ $=$	N has extra-stable half-filled $2p^3$ subshell. O ($2p^4$) has paired electrons in $2p_x$; mutual repulsion destabilizes the electron.
Mg vs Al	IE_1(Mg) $737\,kJ\,mol^{-1}$ $>$ IE_1(Al) $577\,kJ\,mol^{-1}$	$=$ $=$	Analogous to Be/B: Mg has fully-filled $3s^2$; Al has solitary $3p^1$ outside neon core.
P vs S	IE_1(P) $1012\,kJ\,mol^{-1}$ $>$ IE_1(S) $1000\,kJ\,mol^{-1}$	$=$ $=$	Analogous to N/O: P has half-filled $3p^3$; S ($3p^4$) experiences spin-pairing repulsion.
Tl vs In	IE_1(Tl) $589\,kJ\,mol^{-1}$ $>$ IE_1(In) $558\,kJ\,mol^{-1}$	$=$ $=$	$4f^{14}$ Lanthanoid contraction in Tl provides poor shielding, raising Z_{eff} on outer electrons.
Pb vs Sn	IE_1(Pb) $715\,kJ\,mol^{-1}$ $>$ IE_1(Sn) $708\,kJ\,mol^{-1}$	$=$ $=$	$4f^{14}$ Lanthanoid contraction raises Z_{eff} in Lead, arresting downward decrease.

Table 4.1: Systematic Catalogue of High-Yield Ionization Enthalpy Anomalies.

4.2.5 Electron Gain Enthalpy ($\Delta_{eg}H$) Trends & Traps

Electron gain enthalpy is the enthalpy change accompanying the addition of an electron to an isolated gaseous atom to form a uninegative gaseous anion:

$$\mathrm{X(g)} + e^- \longrightarrow \mathrm{X^-(g)}, \qquad \Delta_{eg}H \tag{4.9}$$

Most elements release energy upon gaining an electron ($\Delta_{eg}H < 0$). However, second electron gain enthalpies are **strictly positive (endothermic)** for all elements (e.g., $\mathrm{O^-(g)} + e^- \rightarrow \mathrm{O^{2-}(g)}$, $\Delta_{eg}H_2 = +780\,\mathrm{kJ\,mol^{-1}}$) due to overpowering electrostatic repulsion between the incoming electron and the existing negative anion.

Cardinal Periodic Trap: Fluorine vs Chlorine & Oxygen vs Sulfur

- **Halogen Group 17 Order:**

$$|\Delta_{eg}H| : \mathrm{Cl} > \mathrm{F} > \mathrm{Br} > \mathrm{I} \tag{4.10}$$

$$\implies \Delta_{eg}H : \mathrm{Cl}(-349) < \mathrm{F}(-328) < \mathrm{Br}(-325) < \mathrm{I}(-295)\,\mathrm{kJ\,mol^{-1}} \tag{4.11}$$

  Reason: Fluorine has an extraordinarily compact $2p$ valence shell ($n = 2$). The incoming electron encounters intense interelectronic repulsion from the 7 tightly packed valence electrons. In Chlorine, the larger, more diffuse $3p$ shell accommodates the incoming electron with substantially less Coulombic repulsion.

- **Chalcogen Group 16 Order:**

$$|\Delta_{eg}H| : \mathrm{S} > \mathrm{Se} > \mathrm{Te} > \mathrm{Po} > \mathrm{O} \tag{4.12}$$

  (Oxygen has the least negative electron gain enthalpy in Group 16). Oxygen ($\Delta_{eg}H = -141\,\mathrm{kJ\,mol^{-1}}$) suffers severe electron-electron repulsion in its tiny $2p$ shell, whereas Sulfur ($\Delta_{eg}H = -200\,\mathrm{kJ\,mol^{-1}}$) accommodates the electron far more readily.

- **Positive / Near-Zero $\Delta_{eg}H$ Values:**
  - **Noble Gases** (He, Ne, Ar, Kr, Xe)**:** Strongly positive $\Delta_{eg}H > 0$ because the incoming electron must enter a higher unpopulated principal shell.
  - **Alkaline Earth Metals** (Be, Mg)**:** $\Delta_{eg}H \approx 0$ due to fully-filled stable ns^2 subshells.
  - **Nitrogen** (N)**:** $\Delta_{eg}H \approx +31\,\mathrm{kJ\,mol^{-1}}$ due to extra-stable half-filled $2p^3$ configuration.

4.2.6 Electronegativity & The Inert Pair Effect

- **Electronegativity (χ):** Qualitative tendency of an atom in a covalent molecule to attract shared bonding electrons toward itself.
  - **Pauling Scale:** Based on bond dissociation energy discrepancies:

$$\Delta = E_{A-B} - \sqrt{E_{A-A} \times E_{B-B}} \tag{4.13}$$

$$|\chi_A - \chi_B| = 0.208\sqrt{\Delta\ [\text{kcal/mol}]} = 0.102\sqrt{\Delta\ [\text{kJ/mol}]} \tag{4.14}$$

Pauling assigned Fluorine the benchmark value $\chi(\text{F}) = 4.0$. Absolute hierarchy:

$$\text{F}(4.0) > \text{O}(3.5) > \text{N}(3.0) \approx \text{Cl}(3.0) > \text{Br}(2.8) \tag{4.15}$$
$$> \text{I}(2.5) \approx \text{C}(2.5) \approx \text{S}(2.5) > \text{H}(2.1) \tag{4.16}$$

– **Mulliken Scale:** Arithmetic mean of ionization enthalpy and electron affinity:

$$\chi_{\text{Mulliken}} = \frac{\text{IE} + \text{EA}}{2}\ (\text{in eV}), \qquad \chi_{\text{Pauling}} \approx \frac{\chi_{\text{Mulliken}}}{2.8} = \frac{\text{IE} + \text{EA}}{5.6} \tag{4.17}$$

– **Hybridization Influence:** Electronegativity increases with increasing %s-character:

$$sp\ (50\%\ s) > sp^2\ (33.3\%\ s) > sp^3\ (25\%\ s) \tag{4.18}$$

- **The Inert Pair Effect:** The reluctance of the valence ns^2 electron pair to participate in chemical bonding down Groups 13, 14, and 15. **Physical Origin:**
  1. Intervening $(n-1)d^{10}$ and $(n-2)f^{14}$ electrons shield the nucleus poorly, subjecting the spherical ns orbital to an abnormally high effective nuclear charge Z_{eff}.
  2. Relativistic contraction pulls ns electrons closer to the nucleus, significantly increasing the energy required to unpair and promote an s-electron to the p-subshell.
  3. Bond energies decrease down the group due to larger orbital size; the energy released by forming two additional covalent bonds is insufficient to compensate for the promotion energy of the ns^2 electrons.

  Consequences on Oxidation State Stabilities:
  – **Group 13:** Stability of +1 state increases: $\text{B}^{3+} > \text{Al}^{3+} > \text{Ga}^{3+} > \text{In}^{3+} > \text{Tl}^{3+}$, whereas $\text{Tl}^{+} > \text{In}^{+} > \text{Ga}^{+} > \text{Al}^{+}$. TlCl is stable; Tl^{3+} is an oxidant ($\text{Tl}^{3+} + 2\text{e}^- \rightarrow \text{Tl}^{+}$).
  – **Group 14:** Stability of +2 state increases: C(+4) > Si(+4) > Ge(+4) > Sn(+4) > Pb(+4), whereas $\text{Pb}^{2+} > \text{Sn}^{2+} > \text{Ge}^{2+}$. Thus, Sn^{2+} is a reducing agent, while PbO_2 is a strong oxidant.
  – **Group 15:** Stability of +3 state increases: $\text{Bi}^{3+} > \text{Sb}^{3+} > \text{As}^{3+}$. Bi_2O_5 and BiF_5 are powerful oxidants because Bi(+5) readily reduces to +3.

4.3 Chemical Bonding & Molecular Structure (Unit 3)

4.3.1 Ionic Bonding, Lattice Energy & Fajan's Rules

Ionic bonding arises from omnidirectional electrostatic attraction between oppositely charged ions:

$$E_{\text{coulomb}} = -\frac{1}{4\pi\varepsilon_0}\frac{z_+ z_- e^2}{r_0} \tag{4.19}$$

Lattice energy ($\Delta_{\text{lattice}}H^\circ$) is the energy required to completely separate 1 mole of solid ionic crystal into its constituent gaseous ions at infinite separation.

Fajan's Rules: Covalent Character in Ionic Bonds

No chemical bond is 100% ionic. When a cation approaches an anion, the cation pulls the valence electron cloud of the anion toward itself while repelling its positive nucleus, polarizing the anion. High polarization introduces **covalent character**.

1. **Small Cation Size:** Smaller cations possess higher positive charge density and greater polarizing power (ϕ = charge/radius):

$$\text{Covalent character: LiCl} > \text{NaCl} > \text{KCl} > \text{RbCl} > \text{CsCl} \tag{4.20}$$
$$\implies \text{Melting Point: NaCl} > \text{LiCl} \tag{4.21}$$

2. **Large Anion Size:** Larger anions have diffuse electron clouds that are easily distorted (high polarizability):

$$\text{Covalent character: LiI} > \text{LiBr} > \text{LiCl} > \text{LiF} \tag{4.22}$$

3. **High Ionic Charges:** High positive charge on cation and high negative charge on anion dramatically maximize polarization:

$$\text{Covalent character: } \mathrm{SnCl_4}(\mathrm{Sn^{4+}}) > \mathrm{SnCl_2}(\mathrm{Sn^{2+}}), \tag{4.23}$$
$$\mathrm{AlCl_3}(\mathrm{Al^{3+}}) > \mathrm{MgCl_2}(\mathrm{Mg^{2+}}) > \mathrm{NaCl}(\mathrm{Na^{+}}) \tag{4.24}$$

4. **Pseudo-Inert Gas Electron Configuration ($ns^2np^6nd^{10}$):** A cation with 18 valence electrons ($\mathrm{Cu^+}, \mathrm{Ag^+}, \mathrm{Zn^{2+}}, \mathrm{Cd^{2+}}$) exerts far greater polarizing power than a noble-gas configuration cation (ns^2np^6, 8 electrons; e.g., $\mathrm{Na^+}, \mathrm{K^+}, \mathrm{Ca^{2+}}$) of comparable size and charge, because d-electrons shield the nuclear charge very poorly:

$$\mathrm{CuCl}\ (18e^-) \text{ is far more covalent and insoluble than } \mathrm{NaCl}\ (8e^-) \tag{4.25}$$
$$[r(\mathrm{Cu^+}) \approx r(\mathrm{Na^+})] \tag{4.26}$$

4.3.2 Lewis Structures & Formal Charge

Formal Charge Calculation

Formal charge represents the hypothetical electric charge assigned to an individual atom in a Lewis structure assuming equal sharing of bonding electrons:

$$\mathrm{FC} = V - L - \frac{1}{2}S \tag{4.27}$$

where V is the number of valence electrons of the isolated neutral atom, L is the number of non-bonding lone-pair electrons, and S is the number of shared bonding electrons. **Example: Ozone (O_3):**

- Central Oxygen atom (1 double bond, 1 single coordinate bond, 1 lone pair):

$$\mathrm{FC} = 6 - 2 - \frac{1}{2}(6) = +1 \tag{4.28}$$

- Double-bonded terminal Oxygen (2 lone pairs, 4 shared electrons):

$$\mathrm{FC} = 6 - 4 - \frac{1}{2}(4) = 0 \tag{4.29}$$

- Single-bonded terminal Oxygen (3 lone pairs, 2 shared electrons):

$$\mathrm{FC} = 6 - 6 - \frac{1}{2}(2) = -1 \tag{4.30}$$

4.3.3 Dipole Moments & Vector Summation

The electric dipole moment $\vec{\mu}$ measures bond and molecular polarity:

$$\mu = q \times d \quad [1\,\text{Debye (D)} = 3.33564 \times 10^{-30}\,\text{C m}] \tag{4.31}$$

Molecular dipole moment is the vector sum of all individual bond moments and lone-pair moments:

$$\mu_{\text{net}} = \sqrt{\mu_1^2 + \mu_2^2 + 2\mu_1\mu_2\cos\theta} \tag{4.32}$$

The NH_3 vs NF_3 Dipole Moment Paradox

Both Ammonia (NH_3) and Nitrogen Trifluoride (NF_3) exhibit identical trigonal pyramidal geometry ($SN = 4$, 1 lone pair). However:

$$\mu(NH_3) = 1.47\,\text{D} \gg \mu(NF_3) = 0.23\,\text{D} \tag{4.33}$$

Rigorous Vector Explanation:

- In NH_3, Nitrogen ($\chi = 3.0$) is more electronegative than Hydrogen ($\chi = 2.1$).

The three individual N – H bond dipoles point upward toward Nitrogen, **reinforcing and adding constructively** to the upward orbital dipole of the localized lone pair.

- In NF_3, Fluorine ($\chi = 4.0$) is far more electronegative than Nitrogen ($\chi = 3.0$). The three individual N – F bond dipoles point downward away from Nitrogen, **opposing and destructively cancelling** the upward lone-pair dipole.

4.3.4 VSEPR Theory: Master 3D Geometry Framework

Valence Shell Electron Pair Repulsion (VSEPR) theory dictates that electron pairs surrounding a central atom repel one another and adopt spatial orientations that minimize electrostatic repulsion.

$$\text{Repulsion Hierarchy: } lp - lp > lp - bp > bp - bp \tag{4.34}$$

Steric Number (SN) is computed by:

$$SN = (\text{No. of } \sigma\text{-bonds}) + (\text{No. of lone pairs on central atom}) \tag{4.35}$$

SN	σ	LP	Electron Geom.	Molecular Geom.	Archetypes & Angles
2	2	0	Linear	Linear	$BeCl_2$, CO_2, HCN (180°)
3	3	0	Trigonal Planar	Trigonal Planar	BF_3, BCl_3, SO_3 (120°)
3	2	1	Trigonal Planar	Bent / Angular	SO_2, O_3, NO_2^- ($\approx 119°$)
4	4	0	Tetrahedral	Tetrahedral	CH_4, CCl_4, NH_4^+ (109°28′)
4	3	1	Tetrahedral	Trigonal Pyramidal	NH_3, PCl_3, H_3O^+ (107°)
4	2	2	Tetrahedral	Bent / V-shaped	H_2O, H_2S, OF_2 (104.5°)
5	5	0	Trigonal Bipyramidal	Trigonal Bipyramidal	PCl_5, PF_5 (eq 120°, ax 90°)
5	4	1	Trigonal Bipyramidal	Seesaw	SF_4, SeF_4 (173°, 102°)
5	3	2	Trigonal Bipyramidal	T-shaped	ClF_3, BrF_3 (87.5°)
5	2	3	Trigonal Bipyramidal	Linear	XeF_2, I_3^-, ICl_2^- (180°)
6	6	0	Octahedral	Octahedral	SF_6, $[PF_6]^-$ (90°)
6	5	1	Octahedral	Square Pyramidal	BrF_5, IF_5, $XeOF_4$ (86°)
6	4	2	Octahedral	Square Planar	XeF_4, $[ICl_4]^-$ (90°)
7	7	0	Pentagonal Bipyramidal	Pentagonal Bipyramidal	IF_7 (eq 72°, ax 90°)
7	6	1	Pentagonal Bipyramidal	Distorted Octahedral	XeF_6 (fluxional lone pair)

Table 4.2: Master VSEPR Molecular Geometry Classification Table ($SN = 2$ to 7).

Bent's Rule & Bond Length Asymmetry in PCl_5

- **Bent's Rule Formulation:** More electronegative substituents prefer hybrid orbitals with **less** *s*-**character** (more *p*-character). Lone pairs and electropositive substituents prefer hybrid orbitals with **more** *s*-**character**.
- **Application to Trigonal Bipyramidal Geometry ($sp^3d = sp^2 + pd$):**
  - Equatorial hybrid orbitals (sp^2) possess 33.3% *s*-character.
  - Axial hybrid orbitals (pd) possess 0% *s*-character ($50\%p + 50\%d_{z^2}$).
- **Asymmetry in Gaseous PCl_5:**
  - The two axial P − Cl bonds experience intense 90° repulsion from three equatorial bonds.
  - Consequently, **axial bonds are longer and weaker** (214 pm) than equatorial bonds (202 pm).
  - Upon gentle heating, PCl_5 easily dissociates its two axial chlorines: $PCl_5(g) \xrightarrow{\Delta} PCl_3(g) + Cl_2(g)$.
  - In the crystalline solid phase, PCl_5 ionizes into ionic tetrahedral and octahedral entities: $\mathbf{2}PCl_5(s) \longrightarrow [PCl_4]^+[PCl_6]^-$. Solid PBr_5 exists as $[PBr_4]^+Br^-$.
- **Drago's Rule:** In hydrides of Group 15 and 16 elements from the 3rd period onwards ($PH_3, AsH_3, SbH_3, H_2S, H_2Se, H_2Te$), the central atom does **NOT** undergo hybridization. Bonding occurs via nearly pure *p*-orbitals at $\approx 90°$ angles ($PH_3 : 93.5°, H_2S : 92°$). The lone pair resides in an inert, spherically symmetric *s*-orbital with negligible Lewis basicity.

High-Yield Problem Archetype: Systematic VSEPR Steric Number Prediction

Predicting Hybridization and 3D Geometry for Problematic NEET Species: Formula for Steric Number of central atom A in AB_n^q:

$$SN = \frac{1}{2}\left[V + M - C + A\right]$$

where V = valence e^- of A, M = monovalent atoms attached (H, F, Cl, Br, I), C = cationic charge, A = anionic charge.

1. **Triiodide Ion (I_3^-):** Central I has $V = 7$, two attached monovalent iodines ($M = 2$), anionic charge $A = 1$:

$$SN = \frac{1}{2}[7 + 2 - 0 + 1] = 5 \implies sp^3d \text{ hybridization}$$

   With 2 bond pairs and $5 - 2 = 3$ lone pairs: all three lone pairs occupy equatorial positions at 120° $\implies$ **Linear Geometry** (bond angle 180°).

2. **Xenon Tetrafluoride (XeF_4):** Central Xe has $V = 8$, $M = 4$:

$$SN = \frac{1}{2}[8 + 4] = 6 \implies sp^3d^2 \text{ hybridization}$$

With 4 bond pairs and $6-4=2$ lone pairs: lone pairs occupy opposite trans-axial positions at 180° $\implies$ **Square Planar Geometry** (90°).

3. **Sulfur Tetrafluoride (SF_4):** Central S has $V=6$, $M=4$:

$$SN=\frac{1}{2}[6+4]=5 \implies sp^3d \text{ hybridization}$$

With 4 bond pairs and 1 lone pair: lone pair occupies an equatorial position $\implies$ **Seesaw Geometry**.

4. **Chlorine Trifluoride (ClF_3):** Central Cl has $V=7$, $M=3$:

$$SN=\frac{1}{2}[7+3]=5 \implies sp^3d \text{ hybridization}$$

With 3 bond pairs and 2 lone pairs: both lone pairs occupy equatorial positions $\implies$ **T-shaped Geometry**.

4.3.5 Molecular Orbital Theory (MOT): Energy Levels, Bond Orders & Magnetism

Molecular Orbital Theory constructs molecular orbitals by Linear Combination of Atomic Orbitals (LCAO):

$$\psi_{\text{MO}}=c_A\psi_A \pm c_B\psi_B \implies \psi_b=\frac{1}{\sqrt{2}}(\psi_A+\psi_B), \qquad \psi_a=\frac{1}{\sqrt{2}}(\psi_A-\psi_B) \tag{4.36}$$

1. Diatomic Molecules with $\leq$ 14 Valence Electrons (with *sp*-Mixing: B_2, C_2, N_2)

Because the energy separation between $2s$ and $2p_z$ atomic orbitals is small in B, C, N, quantum mechanical mixing occurs between $\sigma 2s$ and $\sigma 2p_z$. This raises the energy of $\sigma 2p_z$ above the degenerate $\pi 2p_x, \pi 2p_y$ orbitals:

$$\sigma 1s < \sigma^* 1s < \sigma 2s < \sigma^* 2s \tag{4.37}$$

$$< (\pi 2p_x = \pi 2p_y) < \sigma 2p_z < (\pi^* 2p_x = \pi^* 2p_y) < \sigma^* 2p_z \tag{4.38}$$

2. Diatomic Molecules with $>$ 14 Valence Electrons (without *sp*-Mixing: O_2, F_2)

In Oxygen and Fluorine, higher nuclear charge depresses the $2s$ level, creating a large $\Delta E(2s-2p_z)$ gap ($>15\,\text{eV}$) that precludes sp-mixing. Normal energy ordering applies:

$$\sigma 1s < \sigma^* 1s < \sigma 2s < \sigma^* 2s \tag{4.39}$$

$$< \sigma 2p_z < (\pi 2p_x = \pi 2p_y) < (\pi^* 2p_x = \pi^* 2p_y) < \sigma^* 2p_z \tag{4.40}$$

Bond Order & Magnetic Properties Formulation

$$\text{Bond Order (BO)} = \frac{N_b - N_a}{2} \tag{4.41}$$

where N_b is the number of electrons in bonding MOs, and N_a is the number of electrons in antibonding MOs.

- If all electrons are paired $\implies$ **Diamagnetic**.
- If one or more unpaired electrons exist $\implies$ **Paramagnetic** ($\mu_s = \sqrt{n(n+2)}\,\text{BM}$).

Species	e^-	Valence MO Configuration	N_b/N_a	BO	Magnetism & Notes
B_2	10	$(\sigma 2s)^2(\sigma^* 2s)^2(\pi 2p_x)^1(\pi 2p_y)^1$	6/4	1.0	**Paramagnetic** (2 unp. e^-)
C_2	12	$(\sigma 2s)^2(\sigma^* 2s)^2(\pi 2p)^4$	8/4	2.0	**Diamagnetic** (both π-bonds)
N_2	14	$(\sigma 2s)^2(\sigma^* 2s)^2(\pi 2p)^4(\sigma 2p_z)^2$	10/4	3.0	**Diamagnetic** (max stability)
N_2^+	13	$(\sigma 2s)^2(\sigma^* 2s)^2(\pi 2p)^4(\sigma 2p_z)^1$	9/4	2.5	**Paramagnetic** (1 unp. e^-)
O_2^+	15	$(\sigma 2s)^2(\sigma^* 2s)^2(\sigma 2p_z)^2(\pi 2p)^4(\pi^* 2p_x)^1$	10/5	2.5	**Paramagnetic** (highest BO)
O_2	16	$(\sigma 2s)^2(\sigma^* 2s)^2(\sigma 2p_z)^2(\pi 2p)^4(\pi^* 2p)^2$	10/6	2.0	**Paramagnetic** (2 unp. e^-)
O_2^-	17	$(\sigma 2s)^2(\sigma^* 2s)^2(\sigma 2p_z)^2(\pi 2p)^4(\pi^* 2p)^3$	10/7	1.5	**Paramagnetic** (Superoxide)
O_2^{2-}	18	$(\sigma 2s)^2(\sigma^* 2s)^2(\sigma 2p_z)^2(\pi 2p)^4(\pi^* 2p)^4$	10/8	1.0	**Diamagnetic** (Peroxide)

Table 4.3: Comprehensive Molecular Orbital Configurations, Bond Orders, and Magnetism.

NCERT High-Yield Benchmark: Oxygen Species Comparison

- **Bond Order Sequence:** $O_2^+(2.5) > O_2(2.0) > O_2^-(1.5) > O_2^{2-}(1.0)$.
- **Bond Length Sequence (Inversely proportional to Bond Order):**

$$O_2^+(112\,\text{pm}) < O_2(121\,\text{pm}) < O_2^-(128\,\text{pm}) < O_2^{2-}(149\,\text{pm}) \tag{4.42}$$

- **Bond Dissociation Energy Sequence:** $O_2^+ > O_2 > O_2^- > O_2^{2-}$.

4.3.6 Hydrogen Bonding: Intermolecular vs Intramolecular

Hydrogen bonding is an electrostatic dipole-dipole attraction occurring between a covalently bound Hydrogen atom (attached to highly electronegative F, O, N) and another electronegative atom with a localized lone pair.

- **Intermolecular Hydrogen Bonding:** Occurs between distinct molecules (e.g., H_2O, HF, NH_3, C_2H_5OH, *p*-nitrophenol). **Effects:** Elevated boiling points, higher viscosity, lower volatility, and high water solubility. Ice has an open cage-like tetrahedral structure making it less dense than liquid water at 4°C.
- **Intramolecular Hydrogen Bonding (Chelation):** Occurs within the same molecule, forming a 5- or 6-membered ring (e.g., *o*-nitrophenol, salicylaldehyde, salicylic acid). **Effects:** Prevents intermolecular association; results in **significantly lower boiling points**, higher volatility, and **steam volatility** (enabling clean separation of *o*-nitrophenol from *p*-nitrophenol via steam distillation).

4.4 Coordination Chemistry (Unit 12)

4.4.1 Werner's Coordination Theory

Alfred Werner (1893) postulated that transition metals exhibit two distinct types of valency:

1. **Primary Valency (Ionizable):** Corresponds to the oxidation state of the metal ion. It is non-directional and satisfied strictly by negative anions.
2. **Secondary Valency (Non-Ionizable):** Corresponds to the coordination number of the metal ion (4 or 6). It is strictly directional in 3D space (establishing spatial geometry) and satisfied by neutral molecules or negative ligands.

Ammine Complex	IUPAC Formulation	AgCl ppt	Ions	Conductivity ($S\,cm^2\,mol^{-1}$)
$CoCl_3 \cdot 6NH_3$	$[Co(NH_3)_6]Cl_3$	3	4 (1 : 3)	Highest (430)
$CoCl_3 \cdot 5NH_3$	$[Co(NH_3)_5Cl]Cl_2$	2	3 (1 : 2)	Moderate (260)
$CoCl_3 \cdot 4NH_3$	$[Co(NH_3)_4Cl_2]Cl$	1	2 (1 : 1)	Low (100)
$CoCl_3 \cdot 3NH_3$	$[Co(NH_3)_3Cl_3]$	0	0	Non-conducting (0)

Table 4.4: Werner Formulation, Precipitation Stoichiometry, and Solution Conductivity.

4.4.2 Nomenclature & Isomerism in Coordination Entities

IUPAC Nomenclature Principles

1. Cation is named first in both positive and negative coordination entities.
2. Ligands are listed in strict alphabetical order before the central metal name, regardless of charge.

3. Multiplicative prefixes: *di-, tri-, tetra-* for simple ligands; *bis-, tris-, tetrakis-* for polydentate or ligands containing prefixes (e.g., *ethylenediamine*).
4. Anionic ligands end in *-ido* (Cl^- → chlorido, CN^- → cyanido, OH^- → hydroxido).
5. If the complex sphere is anionic, the metal name ends in ***-ate*** (Fe → ferrate, Cu → cuprate, Ag → argentate).
6. Oxidation state of the metal is designated by Roman numerals in parentheses: e.g., $K_4[Fe(CN)_6]$ ⟹ Potassium hexacyanidoferrate(II).

Structural Isomerism

- **Ionization Isomerism:** Exchange of ionizable counter-ion with coordinated ligand:

$$[Co(NH_3)_5SO_4]Br \quad \text{(gives yellow-cream AgBr} \downarrow \text{ with } AgNO_3) \tag{4.43}$$
$$\text{vs } [Co(NH_3)_5Br]SO_4 \quad \text{(gives white } BaSO_4 \downarrow \text{ with } BaCl_2) \tag{4.44}$$

- **Solvate / Hydrate Isomerism:** Water acts either as a coordinated ligand or crystal lattice water:

$$[Cr(H_2O)_6]Cl_3 \quad \text{(Violet, 3 ionizable } Cl^-) \tag{4.45}$$
$$\text{vs } [Cr(H_2O)_5Cl]Cl_2 \cdot H_2O \quad \text{(Grey-green, 2 ionizable } Cl^-) \tag{4.46}$$

- **Linkage Isomerism:** Arises with ambidentate ligands having two distinct coordinating donor atoms:

$$[Co(NH_3)_5(NO_2)]^{2+} \quad \text{(Nitro, yellow-brown, via N)} \tag{4.47}$$
$$\text{vs } [Co(NH_3)_5(ONO)]^{2+} \quad \text{(Nitrito, red, via O)} \tag{4.48}$$

- **Coordination Isomerism:** Interchange of ligands between cationic and anionic complex entities:

$$[Co(NH_3)_6][Cr(CN)_6] \quad \text{vs} \quad [Cr(NH_3)_6][Co(CN)_6] \tag{4.49}$$

Stereoisomerism

1. **Geometrical Isomerism (GI):**
   - **Square Planar ($CN = 4$):** $[MA_2B_2]$ exhibits *cis* and *trans* (e.g., *cis*-platin $[Pt(NH_3)_2Cl_2]$ antitumor drug vs inactive *trans*). $[MABCD]$ yields 3 distinct geometrical isomers. Tetrahedral complexes **NEVER** show GI because all 4 positions are mutually adjacent (109.5°).
   - **Octahedral ($CN = 6$):**
     – $[MA_4B_2]$: *cis* (90°) and *trans* (180°).

  - $[MA_3B_3]$: **Facial (*fac*)** where three identical ligands occupy corners of the same octahedral triangular face, and **Meridional (*mer*)** where ligands occupy a meridian plane surrounding the metal.

2. **Optical Isomerism (Chirality):** Molecules lacking an improper axis of rotation (S_n, including plane of symmetry σ and inversion centre i) exist as non-superimposable mirror image enantiomers (d and l).
   - $[M(AA)_3]$ (e.g., $[Co(en)_3]^{3+}$) is chiral and resolves into Δ and Λ optical enantiomers.
   - $[M(AA)_2B_2]$ (e.g., $[Co(en)_2Cl_2]^{+}$):
     - ***cis*-isomer:** Asymmetric, lacks plane of symmetry $\implies$ **Optically Active** (2 enantiomers).
     - ***trans*-isomer:** Possesses a plane of symmetry (σ_h) $\implies$ **Optically Inactive**.
   - Square planar complexes have a horizontal plane of symmetry and **NEVER** show optical isomerism.

4.4.3 Valence Bond Theory (VBT) & Magnetic States

Linus Pauling applied hybridization of metal d, s, p orbitals to explain coordinate bonding.

- **Inner Orbital Complexes (d^2sp^3):** Strong-field ligands force electron pairing in $(n-1)d$ orbitals, providing two empty inner d-orbitals (low-spin / spin-paired, e.g., $[Fe(CN)_6]^{4-}$: $3d^6$ paired, diamagnetic, $\mu = 0$).
- **Outer Orbital Complexes (sp^3d^2):** Weak-field ligands cannot force pairing; vacant outer nd orbitals are hybridized (high-spin / spin-free, e.g., $[Fe(H_2O)_6]^{2+}$: $3d^6$ with 4 unpaired electrons, paramagnetic, $\mu = \sqrt{4(6)} = 4.90\,\text{BM}$).
- **Limitations of VBT:** Fails to predict absorption spectra or color; cannot quantitatively explain temperature-dependent magnetic susceptibilities; cannot explain why identical d^8 configuration forms tetrahedral paramagnetic $[NiCl_4]^{2-}$ but square planar diamagnetic $[Ni(CN)_4]^{2-}$.

4.4.4 Crystal Field Theory (CFT): Splitting Diagrams & CFSE

Crystal Field Theory treats the metal-ligand interaction as an electrostatic perturbation arising from point negative charges (anions) or point dipoles (neutral molecules like NH_3, H_2O) surrounding the central metal cation. In an isolated gaseous transition metal atom or ion, all five d-orbitals ($d_{xy}, d_{yz}, d_{zx}, d_{x^2-y^2}, d_{z^2}$) possess identical energy (five-fold degenerate).

When ligands approach the metal ion, two sequential perturbations occur:

1. A hypothetical spherically symmetric negative field raises the energy of all five d-orbitals equally due to overall electron-electron repulsion.

2. An asymmetric ligand field splits this degeneracy according to spatial orientation and ligand coordinate geometry (Figure 4.1).




Figure 4.1: Crystal Field Splitting of Five d-Orbitals in an Octahedral Complex (Δ_o).

1. Octahedral Crystal Field Splitting (Δ_o)

In an octahedral $[ML_6]$ complex, six ligands approach the central metal ion along the Cartesian $\pm x, \pm y, \pm z$ axes:

- The e_g orbitals $(d_{x^2-y^2}, d_{z^2})$ have lobes pointing directly along the Cartesian axes where the ligands lie $\implies$ experience maximal electrostatic repulsion $\implies$ destabilized and raised by $+\mathbf{0.6\Delta_o}$ $(+\frac{3}{5}\Delta_o)$.
- The t_{2g} orbitals (d_{xy}, d_{yz}, d_{zx}) have lobes directed in between the Cartesian axes $\implies$ experience minimal direct repulsion $\implies$ stabilized and lowered by $-\mathbf{0.4\Delta_o}$ $(-\frac{2}{5}\Delta_o)$.
- The barycentre (weighted mean energy level) remains invariant: $3(-0.4\Delta_o) + 2(+0.6\Delta_o) = 0$.

2. Tetrahedral Crystal Field Splitting (Δ_t)

In a tetrahedral $[ML_4]$ complex, four ligands approach alternate corners of a circumscribed cube containing the metal at its center:

- The t_2 orbitals (d_{xy}, d_{yz}, d_{zx}) point directly toward the edges of the cube closer to the approaching ligands $\implies$ raised by $+\mathbf{0.4\Delta_t}$.
- The e orbitals $(d_{x^2-y^2}, d_{z^2})$ point toward the faces of the cube further from the ligands $\implies$ lowered by $-\mathbf{0.6\Delta_t}$.
- **Splitting Magnitude Relation:** Because there are only 4 ligands (instead of 6) and none point directly at the d-orbitals:

$$\mathbf{\Delta_t = \frac{4}{9}\Delta_o} \tag{4.50}$$

Because Δ_t is less than half of Δ_o, the crystal field splitting energy in tetrahedral geometry is **always smaller than the pairing energy** ($\Delta_t < P$).

Consequently, **tetrahedral complexes are universally HIGH-SPIN** (spin-free); low-spin tetrahedral complexes do not exist under standard conditions.

Spectrochemical Series & CFSE Calculations

Ligands arranged in order of increasing crystal field splitting magnitude Δ_o:

$$\begin{aligned} &I^- < Br^- < S^{2-} < SCN^- < Cl^- < N_3^- < F^- < OH^- < C_2O_4^{2-} \\ &< H_2O < NCS^- < edta^{4-} < NH_3 < en < NO_2^- < CN^- < CO \end{aligned} \tag{4.51}$$

High-Spin vs Low-Spin Criterion for d^4, d^5, d^6, d^7 in Octahedral Fields:

- **Weak Field Ligands ($\Delta_o < P$):** Electron promotes to e_g before pairing in t_{2g} $\implies$ **High Spin**.
- **Strong Field Ligands ($\Delta_o > P$):** Electrons pair completely in t_{2g} before occupying e_g $\implies$ **Low Spin**.

Octahedral CFSE Formula:

$$\text{CFSE} = \left(-0.4\, n_{t_{2g}} + 0.6\, n_{e_g}\right) \Delta_o + mP \tag{4.52}$$

where m is the number of extra electron pairs formed relative to the free gaseous ion.

	High Spin (Weak Field: $\Delta_o < P$)			**Low Spin (Strong Field: $\Delta_o > P$)**		
d^n	**Config**	**Unpaired**	**CFSE**	**Config**	**Unpaired**	**CFSE**
d^1	$t_{2g}^1 e_g^0$	1	$-0.4\Delta_o$	$t_{2g}^1 e_g^0$	1	$-0.4\Delta_o$
d^2	$t_{2g}^2 e_g^0$	2	$-0.8\Delta_o$	$t_{2g}^2 e_g^0$	2	$-0.8\Delta_o$
d^3	$t_{2g}^3 e_g^0$	3	$-1.2\Delta_o$	$t_{2g}^3 e_g^0$	3	$-1.2\Delta_o$
d^4	$t_{2g}^3 e_g^1$	4	$-0.6\Delta_o$	$t_{2g}^4 e_g^0$	2	$-1.6\Delta_o + 1P$
d^5	$t_{2g}^3 e_g^2$	5	$\mathbf{0.0\Delta_o}$	$t_{2g}^5 e_g^0$	1	$-2.0\Delta_o + 2P$
d^6	$t_{2g}^4 e_g^2$	4	$-0.4\Delta_o$	$t_{2g}^6 e_g^0$	0	$\mathbf{-2.4\Delta_o + 2P}$
d^7	$t_{2g}^5 e_g^2$	3	$-0.8\Delta_o$	$t_{2g}^6 e_g^1$	1	$-1.8\Delta_o + 1P$
d^8	$t_{2g}^6 e_g^2$	2	$-1.2\Delta_o$	$t_{2g}^6 e_g^2$	2	$-1.2\Delta_o$
d^9	$t_{2g}^6 e_g^3$	1	$-0.6\Delta_o$	$t_{2g}^6 e_g^3$	1	$-0.6\Delta_o$
d^{10}	$t_{2g}^6 e_g^4$	0	$0.0\Delta_o$	$t_{2g}^6 e_g^4$	0	$0.0\Delta_o$

Table 4.5: Systematic Crystal Field Stabilization Energies (CFSE) for Octahedral Complexes.

High-Yield Problem Archetype: Octahedral CFSE & Spin-Only Magnetic Moments

Archetype Problem: Compare the electronic configuration, CFSE, and magnetic moment (μ_s) for: (a) $[CoF_6]^{3-}$ vs $[Co(en)_3]^{3+}$, and (b) $[Fe(CN)_6]^{3-}$ vs $[Fe(H_2O)_6]^{3+}$.

1. **Cobalt(III) Complexes ($3d^6$ system):**
   - $[CoF_6]^{3-}$**:** F^- is a weak-field ligand ($\Delta_o < P$) $\implies$ **High Spin**.

$$\begin{aligned} \text{Configuration} &= t_{2g}^4 e_g^2, \quad \text{Unpaired } e^- = 4 \\ \text{CFSE} &= (-0.4 \times 4 + 0.6 \times 2)\Delta_o = -0.4\Delta_o \\ \mu_s &= \sqrt{4(4+2)} = \sqrt{24} \approx 4.90\,\text{BM} \quad \text{(Paramagnetic)} \end{aligned}$$

   - $[Co(en)_3]^{3-}$**:** *en* is a strong-field chelating ligand ($\Delta_o > P$) $\implies$ **Low Spin**.

$$\begin{aligned} \text{Configuration} &= t_{2g}^6 e_g^0, \quad \text{Unpaired } e^- = 0 \\ \text{CFSE} &= (-0.4 \times 6)\Delta_o + 2P = -2.4\Delta_o + 2P \\ \mu_s &= 0\,\text{BM} \quad \text{(Diamagnetic)} \end{aligned}$$

2. **Iron(III) Complexes ($3d^5$ system):**
   - $[Fe(H_2O)_6]^{3+}$**:** H_2O is weak-field $\implies$ $t_{2g}^3 e_g^2$ $\implies$ CFSE $= 0$, $n = 5$ $\implies$ $\mu_s = 5.92\,\text{BM}$.
   - $[Fe(CN)_6]^{3-}$**:** CN^- is strong-field $\implies$ $t_{2g}^5 e_g^0$ $\implies$ CFSE $= -2.0\Delta_o + 2P$, $n = 1$ $\implies$ $\mu_s = 1.73\,\text{BM}$.

4.4.5 Jahn-Teller Distortion Basics

The Jahn-Teller theorem states: *Any non-linear molecular system in a degenerate electronic state is unstable and will undergo geometrical distortion that lowers its symmetry and removes the electronic degeneracy.*

- **Significant / Strong Jahn-Teller Distortion:** Occurs when the e_g **subshell is asymmetrically occupied** because e_g orbitals point directly along the ligand bond axes:

$$\text{High-spin } d^4 \ (t_{2g}^3 e_g^1 : Cr^{2+}, Mn^{3+}), \quad \text{Low-spin } d^7 \ (t_{2g}^6 e_g^1 : Co^{2+}), \tag{4.53}$$

$$\mathbf{d^9} \ (\mathbf{t_{2g}^6 e_g^3} : Cu^{2+}) \tag{4.54}$$

- **Tetragonal Elongation (z-out distortion):** In $[Cu(H_2O)_6]^{2+}$, putting the odd electron in d_{z^2} shields the axial ligands, causing the two axial $Cu - O$ bonds (240 pm) to elongate significantly relative to the four equatorial bonds (195 pm).

4.4.6 Synergic Bonding in Metal Carbonyls

Synergic Bonding Paradigm in Homoleptic Metal Carbonyls

In metal carbonyls (e.g., $Ni(CO)_4, Fe(CO)_5, Cr(CO)_6$), the metal-carbon bond comprises a self-reinforcing dual interaction:

1. **σ-Bond Formation ($M \leftarrow$ CO):** Carbon donates its non-bonding lone pair into an empty hybrid orbital of the transition metal.
2. **π-Backdonation ($M \rightarrow$ CO):** Filled non-bonding d-orbitals of the metal overlap with empty π^* antibonding molecular orbitals of Carbon Monoxide.

Consequences of Synergism:

- Strengthens the $M - C$ bond (increases bond order, shortens bond length).
- Populates the π^* orbital of CO, **weakening the $C - O$ bond** (lowering $C - O$ bond order and reducing $C - O$ vibrational stretching frequency ν_{CO} in IR spectroscopy).
- Greater negative charge on metal increases backdonation $\implies$ weakest $C - O$ bond:

$$\text{IR frequency: } [Mn(CO)_6]^+ > [Cr(CO)_6] > [V(CO)_6]^- \tag{4.55}$$

4.5 Comparative Chemistry of p-Block Elements (Groups 13 to 18)

4.5.1 Group 13: Boron Family

- **Anomalous Boron:** Non-metal, high melting point (2453 K), maximum covalency of 4 (absence of d-orbitals).
- **Diborane (B_2H_6):** Electron-deficient dimer with 12 valence electrons. **Bonding Structure:** Contains four terminal two-centre-two-electron ($2c - 2e$) $B - H$ bonds (119 pm), and two bridging three-centre-two-electron ($3c - 2e$) $B - H - B$ **banana bonds** (133 pm). **Synthesis & Reactions:**

$$4BF_3 + 3LiAlH_4 \xrightarrow{\text{ether}} 2B_2H_6 + 3LiF + 3AlF_3 \tag{4.56}$$

$$B_2H_6 + 3O_2 \longrightarrow B_2O_3 + 3H_2O \quad (\Delta H = -1976\,\text{kJ mol}^{-1}) \tag{4.57}$$

$$3B_2H_6 + 6NH_3 \xrightarrow{450\,\text{K}} 2B_3N_3H_6\ (\textbf{Borazine}) + 12H_2 \tag{4.58}$$

- **Orthoboric Acid (H_3BO_3):** Layer structure with planar BO_3^{3-} units held by H-bonds. Acts as a **monobasic weak Lewis acid** (accepts OH^- from water rather than donating H^+):

$$B(OH)_3 + 2H_2O \rightleftharpoons [B(OH)_4]^- + H_3O^+ \quad (pK_a = 9.25) \tag{4.59}$$

  Cannot be directly titrated with NaOH, but addition of *cis*-polyhydroxy compounds (glycerol, mannitol) forms stable chelate complexes, releasing proton quantitatively.
- **Lewis Acidity of Boron Trihalides:**

$$BI_3 > BBr_3 > BCl_3 > BF_3 \tag{4.60}$$

Reason: In BF_3, vacant $2p$ orbital of Boron overlaps efficiently with filled $2p$ orbital of Fluorine via $p\pi - p\pi$ backbonding, neutralizing electron deficiency. In BCl_3 $(2p - 3p)$ and BI_3 $(2p - 5p)$, overlap is ineffective due to large size discrepancy.

4.5.2 Group 14: Carbon Family

- **Allotropes of Carbon:**
  1. **Diamond:** sp^3 hybridized, tetrahedral 3D giant covalent network, $C - C$ bond length 154 pm, electrical insulator, hardest natural substance.
  2. **Graphite:** sp^2 hybridized, planar hexagonal layers ($C - C = 141.5$ pm) separated by 340 pm held by weak van der Waals forces; fourth π-electron is delocalized over the sheet $\implies$ **electrical conductor** and solid lubricant. **Thermodynamically most stable allotrope** ($\Delta_f H^\circ = 0$).
  3. **Fullerenes (C_{60} Buckminsterfullerene):** Football-shaped cage of 60 carbon atoms comprising 20 six-membered rings and 12 five-membered rings.
- **Silicones:** Organosilicon polymers with repeating $[R_2SiO]_n$ units synthesized by hydrolysis of dialkyldichlorosilanes ($R_2SiCl_2 \xrightarrow{H_2O} R_2Si(OH)_2 \rightarrow$ silicone). Highly water-repellent, thermally stable, electrical insulators.
- **Silicates:** Basic structural unit is the tetrahedral $[SiO_4]^{4-}$ entity:
  - **Orthosilicates:** Discrete isolated $[SiO_4]^{4-}$ (e.g., Zircon $ZrSiO_4$, Phenacite Be_2SiO_4).
  - **Pyrosilicates:** Share 1 oxygen atom: $[Si_2O_7]^{6-}$ (e.g., Thortveitite $Sc_2Si_2O_7$).
  - **Cyclic Silicates:** Share 2 oxygens per tetrahedron: $[Si_3O_9]^{6-}$ or $[Si_6O_{18}]^{12-}$ (e.g., Beryl $Be_3Al_2Si_6O_{18}$).
  - **Chain Silicates (Pyroxenes):** Share 2 oxygens forming linear chains: $[SiO_3]_n^{2n-}$.
  - **Sheet Silicates (Phyllosilicates):** Share 3 oxygens forming 2D sheets: $[Si_2O_5]_n^{2n-}$ (e.g., Talc, Mica).
  - **3D Framework Silicates (Tectosilicates):** Share all 4 oxygens: SiO_2 (Quartz, Feldspar, Zeolites).
- **Zeolites:** Microporous aluminosilicates with open 3D honeycomb channel structures. **ZSM-5** acts as a shape-selective catalyst converting alcohols directly into gasoline.

4.5.3 Group 15: Nitrogen Family

- **Dinitrogen (N_2):** Chemically inert at ambient conditions due to extraordinarily high bond dissociation energy of the $N \equiv N$ triple bond ($941.4\,kJ\,mol^{-1}$).

- **Hydrides of Group 15 ($NH_3, PH_3, AsH_3, SbH_3, BiH_3$):**
  - **Boiling Point:** $BiH_3 > SbH_3 > \mathbf{NH_3} > AsH_3 > PH_3$ (NH_3 abnormally high due to H-bonding; PH_3 is lowest).
  - **Basicity:** $NH_3 > PH_3 > AsH_3 > SbH_3 > BiH_3$ (nitrogen has highest charge density).
  - **Thermal Stability:** $NH_3 > PH_3 > AsH_3 > SbH_3 > BiH_3$ ($M-H$ length increases).
  - **Reducing Power:** $BiH_3 > SbH_3 > AsH_3 > PH_3 > NH_3$ (BiH_3 is strongest reducing agent).
  - **Bond Angle:** $NH_3(107.8°) > PH_3(93.5°) > AsH_3(91.8°) > SbH_3(91.3°)$ (Drago's rule).
- **Nitric Acid (HNO_3 - Ostwald Process):**

$$4NH_3 + 5O_2 \xrightarrow[\text{Pt/Rh,500 K}]{\text{9 bar}} 4NO + 6H_2O \tag{4.61}$$

$$2NO + O_2 \longrightarrow 2NO_2, \qquad 3NO_2 + H_2O \longrightarrow 2HNO_3 + NO \tag{4.62}$$

Reaction with Metals (NEET Benchmark):

$$\text{Cu (dil. } HNO_3): 3Cu + 8HNO_3 \longrightarrow 3Cu(NO_3)_2 + \mathbf{2}NO\uparrow + 4H_2O \tag{4.63}$$

$$\text{Cu (conc. } HNO_3): Cu + 4HNO_3 \longrightarrow Cu(NO_3)_2 + \mathbf{2}NO_2\uparrow + 2H_2O \tag{4.64}$$

$$\text{Zn (dil. } HNO_3): 4Zn + 10HNO_3 \longrightarrow 4Zn(NO_3)_2 + N_2O\uparrow + 5H_2O \tag{4.65}$$

$$\text{Zn (conc. } HNO_3): Zn + 4HNO_3 \longrightarrow Zn(NO_3)_2 + \mathbf{2}NO_2\uparrow + 2H_2O \tag{4.66}$$

Passive metals (Fe, Cr, Al) do not dissolve in concentrated HNO_3 due to formation of an impervious surface oxide film.

- **Oxoacids of Phosphorus:**
  - **Hypophosphorous acid (H_3PO_2):** Monobasic (1 $P-OH$), contains 2 $P-H$ bonds $\implies$ **powerful reducing agent** (reduces $AgNO_3 \to Ag\downarrow$).
  - **Phosphorous acid (H_3PO_3):** Dibasic (2 $P-OH$, 1 $P-H$ bond $\implies$ reducing). On heating: $4H_3PO_3 \xrightarrow{\Delta} 3H_3PO_4 + PH_3\uparrow$.
  - **Orthophosphoric acid (H_3PO_4):** Tribasic (3 $P-OH$), zero $P-H$ bonds $\implies$ non-reducing.
  - **Pyrophosphoric acid ($H_4P_2O_7$):** Tetrabasic (4 $P-OH$), contains 1 $P-O-P$ bridge.

4.5.4 Group 16: Oxygen Family

- **Ozone (O_3):** Prepared by passing silent electric discharge through pure dry dioxygen ($3O_2 \rightleftharpoons 2O_3, \Delta H^\circ = +142\,\text{kJ mol}^{-1}$). Quantitative estimation by

adding neutral buffered KI (pH 9.2, borate buffer) and titrating liberated I_2 against standard sodium thiosulfate:

$$2I^- + H_2O + O_3 \longrightarrow 2OH^- + I_2 + O_2 \tag{4.67}$$

Depletion by nitric oxide: $NO + O_3 \rightarrow NO_2 + O_2$.

- **Allotropes of Sulfur:**
  - **α-Rhombic Sulfur:** Stable below 369 K (95.6°C), yellow, puckered crown S_8 ring.
  - **β-Monoclinic Sulfur:** Stable above 369 K, needle-shaped, crown S_8 ring.
  - **Transition Temperature (369 K):** Both α and β allotropes coexist in dynamic equilibrium.
  - **Vapour Phase Paramagnetism:** Above 1000 K, sulfur exists predominantly as diatomic S_2 molecules which are **paramagnetic** due to two unpaired electrons in antibonding π^* orbitals (identical to O_2).
- **Sulfuric Acid (H_2SO_4 - Contact Process):**

$$2SO_2 + O_2 \xrightarrow[V_2O_5, 720\,K]{2\,bar} 2SO_3 \quad (\Delta H^\circ = -196.6\,kJ\,mol^{-1}) \tag{4.68}$$

$$SO_3 + H_2SO_4 \longrightarrow H_2S_2O_7 \text{ (Oleum)} \tag{4.69}$$

- **Peroxo Oxoacids of Sulfur:**
  - **Caro's Acid (Peroxomonosulfuric, H_2SO_5):** Contains one $-O-O-$ peroxo bond. Oxidation state of S is +6 (NOT +8!).
  - **Marshall's Acid (Peroxodisulfuric, $H_2S_2O_8$):** Contains one $-O-O-$ peroxo bridge connecting two sulfur atoms. Oxidation state of S is +6.

4.5.5 Group 17: Halogen Family

- **Bond Dissociation Enthalpy Anomaly:**

$$Cl_2(242.6) > Br_2(192.8) > F_2(158.8) > I_2(151.1)\,kJ\,mol^{-1} \tag{4.70}$$

Fluorine has an abnormally low bond dissociation energy due to intense electrostatic repulsion between non-bonding lone pairs on the tiny, closely spaced fluorine atoms.

- **Oxidizing Power Hierarchy:** $F_2 > Cl_2 > Br_2 > I_2$. Although $\Delta_{eg}H$ is more negative for Chlorine, Fluorine is the most powerful oxidizing agent because of:
  1. Low bond dissociation enthalpy of F_2.
  2. Extraordinarily high hydration enthalpy of the small F^- ion ($\Delta_{hyd}H = -515\,kJ\,mol^{-1}$).

- **Hydrogen Halides (HF, HCl, HBr, HI):**
  - **Boiling Point:** $HF(293\,K) > HI(238\,K) > HBr(206\,K) > HCl(189\,K)$ (HF is liquid due to intermolecular H-bonding).
  - **Acidic Strength:** $HF < HCl < HBr < HI$ ($H-X$ bond dissociation enthalpy decreases).
  - **Reducing Power:** $HF < HCl < HBr < HI$ (HI is the strongest reducing agent).
- **Oxoacids of Halogens:**
  - **Acidic Strength Order:** $HClO_4 > HClO_3 > HClO_2 > HClO$ (higher oxidation state of halogen increases conjugate base resonance stabilization via $d\pi - p\pi$ bonding).
  - **Oxidizing Power Order:** $HClO > HClO_2 > HClO_3 > HClO_4$ (hypohalous acids are the most powerful oxidants due to ready oxygen atom liberation).

4.5.6 Group 18: Noble Gases

Neil Bartlett (1962) observed that PtF_6 oxidizes molecular dioxygen to $[O_2]^+[PtF_6]^-$. Because the first ionization enthalpy of Xenon ($1170\,kJ\,mol^{-1}$) is nearly identical to that of O_2 ($1175\,kJ\,mol^{-1}$), he reacted Xenon with PtF_6 to synthesize the first noble gas compound: $Xe^+[PtF_6]^-$.

Xenon Fluorides: Preparation, Structures & Hydrolysis Pathways

$$Xe + F_2 \xrightarrow[1\,bar]{673\,K} XeF_2 \quad [2:1 \text{ ratio; Linear}] \tag{4.71}$$

$$Xe + 2F_2 \xrightarrow[7\,bar]{873\,K} XeF_4 \quad [1:5 \text{ ratio; Square Planar}] \tag{4.72}$$

$$Xe + 3F_2 \xrightarrow[60-70\,bar]{573\,K} XeF_6 \quad [1:20 \text{ ratio; Distorted Octahedral}] \tag{4.73}$$

Hydrolysis Pathways (Cardinal NEET Questions):

$$2XeF_2 + 2H_2O \longrightarrow 2Xe + 4HF + O_2\uparrow \quad \text{(Redox)} \tag{4.74}$$

$$6XeF_4 + 12H_2O \longrightarrow 4Xe + 2XeO_3 + 24HF + 3O_2\uparrow \tag{4.75}$$

$$\text{(Disproportionation: } Xe^{IV} \rightarrow Xe^0 + Xe^{VI}) \tag{4.76}$$

$$XeF_6 + 3H_2O \longrightarrow XeO_3 + 6HF \quad \text{(Complete hydrolysis)} \tag{4.77}$$

$$XeF_6 + H_2O \longrightarrow XeOF_4 + 2HF \quad \text{(Partial hydrolysis)} \tag{4.78}$$

$$XeF_6 + 2H_2O \longrightarrow XeO_2F_2 + 4HF \quad \text{(Partial hydrolysis)} \tag{4.79}$$

Fluoride Ion Donor and Acceptor Reactions:

$$XeF_2 + PF_5 \longrightarrow [XeF]^+[PF_6]^- \tag{4.80}$$

$$XeF_6 + MF \longrightarrow M^+[XeF_7]^- \quad (M = Na, K, Rb, Cs) \tag{4.81}$$

Direct NCERT Benchmark: p-Block Cardinal Diagnostic Extracts

- **Basicity vs Reducing Power in Group 15 Hydrides:** Basicity decreases ($NH_3 > PH_3 > AsH_3 > SbH_3 > BiH_3$), whereas reducing power increases ($BiH_3 > SbH_3 > AsH_3 > PH_3 > NH_3$). BiH_3 is the strongest reducing hydride; NH_3 is a mild reducing agent.
- **Phosphorus Oxoacid Reducing Power:** Arises exclusively from the presence of directly bonded P – H bonds:
  - H_3PO_2 (Hypophosphorous / Phosphinic): Has 2 P – H bonds $\implies$ most powerful reducing oxoacid.
  - H_3PO_3 (Phosphorous / Phosphonic): Has 1 P – H bond $\implies$ moderate reducing agent.
  - H_3PO_4 (Orthophosphoric): Has 0 P – H bonds $\implies$ strictly non-reducing.
- **Paramagnetism of Diatomic Sulfur (S_2):** In the vapour phase at elevated temperatures ($> 1000\,K$), sulfur exists as S_2 molecules which possess two unpaired electrons in degenerate π^* antibonding orbitals, exactly analogous to O_2.
- **Commercial Preparation of Dinitrogen from Azides:** Ultra-pure nitrogen gas is produced by thermal decomposition of sodium or barium azide:

$$2NaN_3 \xrightarrow{\Delta} 2Na + 3N_2\uparrow, \qquad Ba(N_3)_2 \xrightarrow{\Delta} Ba + 3N_2\uparrow$$

4.6 Comparative Chemistry of d- and f-Block Elements (Unit 11)

4.6.1 General Trends of $3d$ Transition Series

A transition element is defined as an element having an incompletely filled d-subshell in its ground state or in any of its common oxidation states. Zinc, Cadmium, and Mercury (d^{10}) are **non-transition elements** because they have completely filled d-orbitals in both ground and ionic states.

- **Atomic Radii across $3d$ Series:** Decreases initially (Sc to Cr), remains nearly constant (Mn to Ni) because increasing nuclear charge is balanced by $d-d$ shielding, and slightly increases at the end (Cu, Zn) due to interelectronic repulsion in filled d^{10}.
- **High Enthalpies of Atomization:** Reaches maximum at middle of series (V, Cr, Mo, W) due to maximum unpaired electrons participating in strong interatomic metallic bonding.
- **Variable Oxidation States:** Arises because $(n-1)d$ and ns electrons have comparable energies. Sc exhibits only +3; Mn exhibits the widest range from +2 to +7.

- **Standard Electrode Potentials ($E^\circ(M^{2+}/M)$):**

$$M(s) \xrightarrow{\Delta_{\text{sub}}H} M(g) \xrightarrow{\text{IE}_1+\text{IE}_2} M^{2+}(g) \xrightarrow{\Delta_{\text{hyd}}H} M^{2+}(aq) \tag{4.82}$$

  Cu has an anomalous **positive standard reduction potential** ($+0.34\,\text{V}$) because its high enthalpy of atomization and high ionization enthalpy are not compensated by the hydration enthalpy of Cu^{2+}. Thus, Copper does not liberate H_2 from non-oxidizing acids.
- **Disproportionation of Copper(I):**

$$2Cu^+(aq) \longrightarrow Cu^{2+}(aq) + Cu(s) \quad (E^\circ_{\text{cell}} = +0.36\,\text{V}) \tag{4.83}$$

  Cu^{2+} is far more stable in aqueous solution than Cu^+ because the much higher hydration enthalpy of Cu^{2+} ($\Delta_{\text{hyd}}H \approx -2121\,\text{kJ mol}^{-1}$) overcompensates for the second ionization enthalpy of Copper.
- **Spin-Only Magnetic Moment Formula:**

$$\mu_s = \sqrt{n(n+2)}\,\text{BM} \tag{4.84}$$

  - $n = 1 \implies \mu = \sqrt{3} = 1.73\,\text{BM}$ (Ti^{3+}, Cu^{2+}).
  - $n = 2 \implies \mu = \sqrt{8} = 2.83\,\text{BM}$ (Ti^{2+}, Ni^{2+}).
  - $n = 3 \implies \mu = \sqrt{15} = 3.87\,\text{BM}$ (V^{2+}, Cr^{3+}).
  - $n = 4 \implies \mu = \sqrt{24} = 4.90\,\text{BM}$ (Cr^{2+}, Fe^{2+}).
  - $n = 5 \implies \mu = \sqrt{35} = 5.92\,\text{BM}$ (Mn^{2+}, Fe^{3+}).

4.6.2 Master Transition Metal Compounds: $K_2Cr_2O_7$ & $KMnO_4$

1. Potassium Dichromate ($K_2Cr_2O_7$)

Industrial Preparation from Chromite Ore ($FeCr_2O_4$):

$$4FeCr_2O_4 + 8Na_2CO_3 + 7O_2 \longrightarrow 8Na_2CrO_4\ (\text{Yellow}) + 2Fe_2O_3 + 8CO_2 \tag{4.85}$$

$$2Na_2CrO_4 + 2H^+ \longrightarrow Na_2Cr_2O_7\ (\text{Orange}) + 2Na^+ + H_2O \tag{4.86}$$

$$Na_2Cr_2O_7 + 2KCl \longrightarrow K_2Cr_2O_7\ (\text{Orange crystals}) + 2NaCl \tag{4.87}$$

$K_2Cr_2O_7$ is preferred over $Na_2Cr_2O_7$ as a primary volumetric standard because $Na_2Cr_2O_7$ is deliquescent. **pH-Dependent Chromate-Dichromate Interconversion:**

$$\underset{\text{Yellow}}{2CrO_4^{2-}} + 2H^+ \rightleftharpoons \underset{\text{Orange}}{Cr_2O_7^{2-}} + H_2O \tag{4.88}$$

(Yellow chromate in alkaline medium; orange dichromate in acidic medium; Cr oxidation state remains +6). **Structures:**

- Chromate (CrO_4^{2-}): Tetrahedral (T_d), four equivalent $Cr-O$ bonds (166 pm) due to resonance.

- Dichromate ($Cr_2O_7^{2-}$): Two tetrahedra sharing one oxygen corner; bridging $Cr-O-Cr$ bond angle is 126°; bridging $Cr-O$ bond length is 179 pm, while six terminal $Cr-O$ bonds are shorter (163 pm).

Oxidizing Reactions in Acidic Medium (n-factor = 6):

$$Cr_2O_7^{2-} + 14H^+ + 6e^- \longrightarrow 2Cr^{3+} + 7H_2O \quad (E^\circ = +1.33\,V) \tag{4.89}$$

$$6Fe^{2+} \longrightarrow 6Fe^{3+} + 6e^- \tag{4.90}$$

$$6I^- \longrightarrow 3I_2 + 6e^- \tag{4.91}$$

$$3H_2S \longrightarrow 3S\downarrow + 6H^+ + 6e^- \tag{4.92}$$

$$3Sn^{2+} \longrightarrow 3Sn^{4+} + 6e^- \tag{4.93}$$

Chromyl Chloride Test: Confirmation of chloride ions:

$$K_2Cr_2O_7 + 4NaCl + 6H_2SO_4(\text{conc}) \xrightarrow{\Delta} \tag{4.94}$$

$$2CrO_2Cl_2\uparrow\ (\text{Deep red}) + 2KHSO_4 + 4NaHSO_4 + 3H_2O \tag{4.95}$$

Pass into NaOH → yellow Na_2CrO_4 $\xrightarrow{Pb(CH_3COO)_2}$ bright yellow precipitate of lead chromate ($PbCrO_4\downarrow$).

2. Potassium Permanganate ($KMnO_4$)

Industrial Preparation from Pyrolusite Ore (MnO_2):

$$2MnO_2 + 4KOH + O_2 \xrightarrow{\text{fuse}} 2K_2MnO_4\ (\text{Dark green}) + 2H_2O \tag{4.96}$$

$$3MnO_4^{2-} + 4H^+ \longrightarrow 2MnO_4^-\ (\text{Purple}) + MnO_2 + 2H_2O \tag{4.97}$$

Manganate vs Permanganate Comparison:

- Manganate (MnO_4^{2-}): Green, tetrahedral, Mn(+6), $3d^1 \implies$ **Paramagnetic** (1 unpaired electron).
- Permanganate (MnO_4^-): Deep purple, tetrahedral, Mn(+7), $3d^0 \implies$ **Diamagnetic**. **Origin of Color:** Deep purple color is **NOT** due to $d-d$ transition (it is d^0!). Color arises from **Ligand-to-Metal Charge Transfer (LMCT)** where oxygen $2p$ electron density temporarily shifts into vacant manganese $3d$ orbitals.

Permanganate Redox Media & Valence Factors (n-factor)

1. **In Acidic Medium (n-factor = 5):**

$$MnO_4^- + 8H^+ + 5e^- \longrightarrow Mn^{2+} + 4H_2O \quad (E^\circ = +1.51\,V) \tag{4.98}$$

$$5Fe^{2+} \longrightarrow 5Fe^{3+} + 5e^- \tag{4.99}$$

$$5C_2O_4^{2-} \longrightarrow 10CO_2 \uparrow + 10e^- \quad \text{(titration heated to } 60-70^\circ\text{C)} \tag{4.100}$$

$$10I^- \longrightarrow 5I_2 + 10e^- \tag{4.101}$$

$$5NO_2^- + 5H_2O \longrightarrow 5NO_3^- + 10H^+ + 10e^- \tag{4.102}$$

$$5SO_3^{2-} + 5H_2O \longrightarrow 5SO_4^{2-} + 10H^+ + 10e^- \tag{4.103}$$

2. **In Neutral / Faintly Alkaline Medium (Baeyer's Reagent, n-factor $= 3$):**

$$MnO_4^- + 2H_2O + 3e^- \longrightarrow MnO_2 + 4OH^- \quad (E^\circ = +1.23\,\text{V}) \tag{4.104}$$

Cardinal NEET Reaction Differences:

- Iodide is oxidized to **Iodate** (IO_3^-), NOT I_2:

$$I^- + 2MnO_4^- + H_2O \longrightarrow IO_3^- + 2MnO_2 + 2OH^- \tag{4.105}$$

- Thiosulfate is oxidized to **Sulfate** (SO_4^{2-}):

$$8MnO_4^- + 3S_2O_3^{2-} + H_2O \longrightarrow 8MnO_2 + 6SO_4^{2-} + 2OH^- \tag{4.106}$$

3. **In Strongly Alkaline Medium (n-factor $= 1$):**

$$MnO_4^- + e^- \longrightarrow MnO_4^{2-} \tag{4.107}$$

High-Yield Problem Archetype: Permanganometric Volumetric Stoichiometry

Archetype Problem: Determine the volume of 0.02 M $KMnO_4$ required to completely oxidize: (a) 25.0 mL of 0.1 M Mohr's salt ($FeSO_4{\cdot}(NH_4)_2SO_4{\cdot}6H_2O$), and (b) 25.0 mL of 0.1 M Oxalic acid ($H_2C_2O_4 \cdot 2H_2O$) in acidic medium.

1. **Equivalence Principle:** At the exact equivalence endpoint:

$$N_1V_1 = N_2V_2 \implies (M_1 \times n_1)V_1 = (M_2 \times n_2)V_2$$

For $KMnO_4$ in acidic medium, $n_1 = 5$ ($Mn^{VII} \rightarrow Mn^{II}$).

2. **Case (a): Titration with Mohr's Salt:** Mohr's salt acts as a single-electron reducing agent: $Fe^{2+} \rightarrow Fe^{3+} + e^- \implies n_2 = 1$.

$$(0.02 \times 5) \times V_1 = (0.1 \times 1) \times 25.0 \implies 0.10 \times V_1 = 2.50 \implies V_1 = \mathbf{25.0}\,\text{mL}$$

3. **Case (b): Titration with Oxalic Acid:** Oxalate ion undergoes a 2-electron oxidation: $C_2O_4^{2-} \rightarrow 2CO_2 + 2e^- \implies n_2 = 2$.

$$(0.02 \times 5) \times V_1 = (0.1 \times 2) \times 25.0 \implies 0.10 \times V_1 = 5.00 \implies V_1 = \mathbf{50.0}\,\text{mL}$$

4. **Practical Laboratory Condition:** The oxalic acid titration must be conducted at $60-70^\circ$C because the initial reduction is slow. The reaction is **autocatalyzed** by the produced Mn^{2+} ions.

4.6.3 The f-Block Elements: Lanthanoids & Actinoids

1. Lanthanoids ($4f$ Series: Ce_{58} to Lu_{71})

- **Electronic Configuration:** $[Xe]4f^{1-14}5d^{0-1}6s^2$.
- **Oxidation States:** Dominant and most stable state is +3.
  - +2 exhibited by Eu^{2+} ($[Xe]4f^7$, half-filled) and Yb^{2+} ($[Xe]4f^{14}$, fully-filled) $\implies$ act as **strong reducing agents** reverting to +3.
  - +4 exhibited by Ce^{4+} ($[Xe]4f^0$, empty shell) and Tb^{4+} ($[Xe]4f^7$) $\implies$ Ce^{4+} acts as a **powerful analytical oxidizing agent** ($E^\circ_{Ce^{4+}/Ce^{3+}} = +1.74\,V$).
- **Alloy: Mischmetal:** Consists of Lanthanoid metals ($\sim 95\%$), Iron ($\sim 5\%$), and traces of S, C, Ca, Al. Used in cigarette lighter flints and tracer bullets.

NCERT Hallmark: Lanthanoid Contraction & Chemical Consequences

The imperfect shielding of one $4f$ electron by another causes a progressive increase in effective nuclear charge (Z_{eff}), resulting in a steady decrease in atomic and ionic radii from Ce^{3+} (103 pm) to Lu^{3+} (86 pm). **Three Major Chemical Consequences for NEET:**

1. **Covalent Character and Basicity of Hydroxides:** Size of M^{3+} decreases from La^{3+} to Lu^{3+}. By Fajan's rules, covalent character of the M – OH bond increases; therefore, basic strength strictly decreases:
$$La(OH)_3 > Ce(OH)_3 > \cdots > Lu(OH)_3$$
($La(OH)_3$ is most basic; $Lu(OH)_3$ is least basic).
2. **Pairing of 4d and 5d Transition Radii:** Due to the intervention of the 14 lanthanoid elements, elements of the 4d and 5d series show almost identical atomic and ionic radii:
$$r(Zr) \approx r(Hf) \approx 160\,pm; \quad r(Nb) \approx r(Ta) \approx 146\,pm; \quad r(Mo) \approx r(W)$$
Consequently, Zr and Hf exhibit identical chemical properties and occur together in nature, posing difficult separation challenges.
3. **Chromatographic Separation:** Because radii vary by only ~ 17 pm across 14 elements, chemical separation is difficult, but their slight differences in complex formation constants and hydrated ionic radii permit successful separation via ion-exchange chromatography.

2. Actinoids ($5f$ Series: Th_{90} to Lr_{103})

- **Electronic Configuration:** $[Rn]5f^{1-14}6d^{0-2}7s^2$.
- **Oxidation States:** Wider range (+3 to +7) than Lanthanoids as energy gaps between $5f, 6d$, and $7s$ are exceedingly small.

- **Actinoid Contraction:** Steady decrease in ionic radii; steeper than lanthanoids due to poorer shielding by $5f$.
- **Nuclear Instability:** All actinoids are radioactive; elements beyond U ($Z > 92$) are synthetic transuranics.

PART IV

Organic Chemistry: Principles, Pathways & Biomolecules

CHAPTER 5

Organic Chemistry: Electronic Effects, Functional Groups & Biomolecules

This chapter spans the core of NEET Organic Chemistry. We develop the foundational electronic effects (inductive, resonance, hyperconjugation) and reactive intermediates, proceed through aliphatic and aromatic hydrocarbons, haloalkanes, oxygen-containing functional groups (alcohols, phenols, ethers, carbonyls, carboxylic acids), nitrogen derivatives (amines, diazonium salts), and conclude with biomolecules (carbohydrates, proteins, nucleic acids).

A Personal Note from Satya: Stop Memorizing 200 Reactions — Follow the Electrons!

Take a deep breath: Organic Chemistry is about to become your highest-scoring subject.
Aspirants make Organic Chemistry a living nightmare by trying to memorize 200 named reactions like isolated words in a foreign language. They stare at Aldol, Cannizzaro, Reimer-Tiemann, and Gabriel Phthalimide and wonder how anyone can hold this in their head.
Here is the secret of physical organic chemistry that unlocks everything:

Every single reaction in organic chemistry is just an electron-rich center attacking an electron-poor center.

That is literally it!

- A nucleophile (lone pair, negative charge, or π-bond) looks for an electrophile (partial positive charge δ^+, carbocation, or carbonyl carbon).
- When you look at an aldehyde, don't memorize what reagent adds to it. Look at the C = O double bond: the oxygen is electronegative (δ^-), pulling electron density away from the carbonyl carbon (δ^+). The carbon is electrophilic! Naturally, any nucleophile (Grignard, cyanide, hydride, enolate) will attack that carbon.

Once you train your eyes to spot the electron flow with curved arrows, you will never panic in the exam hall. Even if NTA presents an unfamiliar substrate, you can deduce the product step-by-step from scratch. Trust the electronic principles, draw every mechanism out with your pen, and let's conquer this subject together!

5.1 NMC Syllabus Demarcation & Zero-Tolerance Deletions

Curriculum Alignment & Pedagogical Scope

The revised National Medical Commission (NMC) NEET UG 2027 Chemistry syllabus demands a rigorous, mechanistic understanding of organic transformations rather than superficial rote memorization. The curriculum is built upon three integrated pillars:

1. **General Organic Chemistry (GOC) & Reaction Intermediates:** Quantitative and qualitative evaluation of electron displacement effects (inductive, electromeric, resonance, hyperconjugation, and aromaticity) and the structural thermodynamics of transient intermediates (carbocations, carbanions, free radicals, carbenes, and arynes).
2. **Systematic Functional Group Reaction Pathways:** Exhaustive mechanism sequences spanning aliphatic/aromatic hydrocarbons, haloalkanes/haloarenes, oxygenated functions (alcohols, phenols, ethers, aldehydes, ketones, carboxylic acids), and nitrogenous compounds (amines, diazonium salts).
3. **Biomolecules of Life:** Structural stereochemistry of carbohydrates, amino acids, peptide linkages, primary through quaternary protein architecture, nucleic acid Watson-Crick geometry, vitamins, and hormones.

Zero-Tolerance NMC Deleted Topics Alert: Do Not Study!

To optimize preparation efficiency and adhere strictly to the regulatory mandate issued by the NMC for NEET UG 2024–2027, the following entire domains are **permanently excluded** from the examination:

- **Polymers (Entire Chapter Deleted):** Classification of polymers; addition and condensation polymerization; natural and synthetic rubber, vulcanization; polythene (LDPE/HDPE), Teflon, PAN, Nylon-6, Nylon-6,6, Terylene/Dacron, Bakelite, melamine-formaldehyde, Buna-S, Buna-N; biodegradable polymers (PHBV, Nylon-2-Nylon-6).
- **Chemistry in Everyday Life (Entire Chapter Deleted):** Drugs and pharmacodynamics; analgesics (narcotic and non-narcotic); tranquilizers, antiseptics (Dettol, Bithionol), disinfectants, antibiotics, antacids, antihistamines; artificial sweeteners (saccharin, aspartame, alitame, sucralose); food preservatives; soaps and synthetic detergents (anionic, cationic, non-ionic).

Candidates must **not** allocate study time to these deleted units. Any occurrence of these keywords in this text serves solely as an exclusion boundary warning.

5.2 General Organic Chemistry (GOC) & Electronic Effects

5.2.1 Inductive Effect ($\pm I$)

Physical Origin & Characteristics of Inductive Effect

The inductive effect is a permanent polarization of a σ-bond arising from electronegativity differences between bonded atoms. The more electronegative atom pulls the bonding σ-electron pair toward itself, establishing a static dipole:

$$-\mathrm{C}^{\delta\delta\delta+} - \mathrm{C}^{\delta\delta+} - \mathrm{C}^{\delta+} - \mathrm{X}^{\delta-} \quad (\text{where } \chi_X > \chi_C) \tag{5.1}$$

Key Physical Attributes:

- **Permanent Distortion:** It involves shifting of σ-electrons without complete bond cleavage or free electron transfer.
- **Distance Attenuation:** The magnitude of the fractional charge decreases rapidly with distance and becomes virtually negligible beyond the third σ-carbon atom (C_3).
- **Hydrogen Reference Standard:** The electronegativity of hydrogen is taken as the arbitrary zero benchmark: $I_{\mathrm{H}} = 0$.

Authoritative $-I$ and $+I$ Relative Strength Series

1. **Electron-Withdrawing ($-I$) Series (Decreasing Order of Pull):**

$$-\mathrm{NF}_3^+ > -\mathrm{NR}_3^+ > -\mathrm{NH}_3^+ > -\mathrm{NO}_2 > -\mathrm{SO}_3\mathrm{H} \tag{5.2}$$
$$> -\mathrm{CN} > -\mathrm{COOH} > -\mathrm{F} > -\mathrm{Cl} > -\mathrm{Br} \tag{5.3}$$
$$> -\mathrm{I} > -\mathrm{OR} > -\mathrm{OH} > -\mathrm{C} \equiv \mathrm{CR} \tag{5.4}$$
$$> -\mathrm{NH}_2 > -\mathrm{C}_6\mathrm{H}_5 > -\mathrm{CH} = \mathrm{CH}_2 > -\mathrm{H} \tag{5.5}$$

2. **Electron-Releasing ($+I$) Series (Decreasing Order of Push):**

$$-\mathrm{O}^- > -\mathrm{COO}^- > -\mathrm{C}(\mathrm{CH}_3)_3 > -\mathrm{CH}(\mathrm{CH}_3)_2 \tag{5.6}$$
$$> -\mathrm{CH}_2\mathrm{CH}_3 > -\mathrm{CH}_3 > -\mathrm{D} > -\mathrm{H} \tag{5.7}$$

Physical Rationales for Key Sequences:

- **Positive Formal Charges:** Positively charged ammonium and nitro centers exhibit dramatically higher $-I$ than neutral halogens because full ionic charges create immense electrostatic attraction.
- **Hybridization Dependency:** Higher $\%s$-character pulls electrons closer to the carbon nucleus, raising electronegativity:

$$sp \text{ carbon } (50\%s, \chi \approx 3.29) > sp^2 \text{ carbon } (33.3\%s, \chi \approx 2.75) \tag{5.8}$$
$$> sp^3 \text{ carbon } (25\%s, \chi \approx 2.48) \tag{5.9}$$

  Hence, alkynyl ($-\mathrm{C} \equiv \mathrm{CR}$) exerts a stronger $-I$ effect than vinyl ($-\mathrm{CH} = \mathrm{CH}_2$) and phenyl ($-\mathrm{C}_6\mathrm{H}_5$).

- **Isotope Effect:** Tritium ($-$T) and Deuterium ($-$D) have shorter and stronger C $-$ D bonds than C $-$ H due to lower zero-point vibrational energy, causing less average nuclear distance and greater electron release: $+I(-\text{T}) > +I(-\text{D}) > +I(-\text{H})$.

Quantitative Applications: Acidic Strength of Carboxylic Acids

The acidic strength (K_a, where $pK_a = -\log K_a$) of carboxylic acids is determined by the stability of the conjugate carboxylate anion ($\text{R} - \text{COO}^-$):

$$\text{Acidic Strength} \propto \text{Stability of Conjugate Base } (\text{R} - \text{COO}^-) \propto \frac{-I \text{ effect}}{+I \text{ effect}} \tag{5.10}$$

1. Influence of Halogen Identity (Electronegativity):

$$\begin{aligned} \text{FCH}_2\text{COOH } (pK_a = 2.66) > \text{ClCH}_2\text{COOH } (2.86) > \text{BrCH}_2\text{COOH } (2.90) \\ > \text{ICH}_2\text{COOH } (3.17) > \text{CH}_3\text{COOH } (4.76) \end{aligned} \tag{5.11}$$

2. Number of Electron-Withdrawing Groups:

$$\text{Cl}_3\text{C} - \text{COOH } (pK_a = 0.65) > \text{Cl}_2\text{CH} - \text{COOH } (1.29) \tag{5.12}$$

$$> \text{ClCH}_2 - \text{COOH } (2.86) > \text{CH}_3\text{COOH } (4.76) \tag{5.13}$$

3. Distance Dependence (α, β, γ-Positions):

$$\text{CH}_3 - \text{CH}_2 - \text{CH(Cl)} - \text{COOH } (\alpha, pK_a = 2.84) \tag{5.14}$$

$$> \text{CH}_3 - \text{CH(Cl)} - \text{CH}_2 - \text{COOH } (\beta, 4.06) \tag{5.15}$$

$$> \text{CH}_2\text{(Cl)} - \text{CH}_2 - \text{CH}_2 - \text{COOH } (\gamma, 4.52) \tag{5.16}$$

Direct NCERT Line: Formic Acid vs Benzoic Acid vs Acetic Acid

The electron-donating $+I$ effect of the methyl group destabilizes the acetate anion relative to the formate anion:

$$\text{HCOOH } (pK_a = 3.75) > \text{C}_6\text{H}_5\text{COOH } (pK_a = 4.20) > \text{CH}_3\text{COOH } (pK_a = 4.76) \tag{5.17}$$

Crucial Insight: Even though phenyl is electron-withdrawing by $-I$ relative to sp^3 alkyl, its $+M$ resonance into the carbonyl group weakens acidity compared to formic acid. Thus, formic acid is stronger than benzoic acid!

5.2.2 Resonance / Mesomeric Effect ($\pm M$)

The Quantum Mechanical Basis of Resonance

Resonance describes the delocalization of π-electrons or non-bonding lone pairs across parallel, unhybridized p-orbitals in planar conjugated frameworks.
Mandatory Conditions for Conjugation:

1. **Planarity:** All participating atomic nuclei must lie in a single plane (or near-planar geometry) to permit effective sideways overlap of adjacent p-orbitals.
2. **Conjugated Topologies:** Conjugation occurs across alternating systems:
   - $\pi - \sigma - \pi$ (e.g., 1,3-butadiene, benzene)
   - $\pi - \sigma - p_{\text{vacant}}$ (e.g., allyl carbocation, tropylium cation)
   - $\pi - \sigma -$ lone pair / non-bonding (e.g., vinyl chloride, phenoxide, aniline)
   - $\pi - \sigma -$ unpaired electron ($\bullet$) (e.g., allyl free radical)
   - Lone pair $-\sigma - p_{\text{vacant}}$ (e.g., $CH_3 - \overset{+}{C}H - \ddot{O}CH_3$)

Hierarchical Rules for Evaluating Canonical Structure Stability

When ranking the thermodynamic contribution of contributing resonance structures (canonical forms) to the true resonance hybrid, apply the following hierarchy strictly in order:

1. **Complete Octet Rule (Paramount Criterion):** Structures in which all second-period atoms (C, N, O, F) possess complete valence octets contribute overwhelmingly more than structures with incomplete octets, **even if** a formal positive charge resides on an electronegative atom (O or N):

$$CH_3 - \overset{+}{O} = CH_2 \longleftrightarrow CH_3 - \ddot{O} - \overset{+}{C}H_2 \tag{5.18}$$

$$\text{(octet complete, Major)} \qquad \text{(carbon has } 6e^-\text{, Minor)} \tag{5.19}$$

2. **Maximum Number of Covalent Bonds:** Greater number of covalent bonds maximizes attractive nuclear-electronic interactions, conferring greater stability.
3. **Charge Separation vs Uncharged Species:** Neutral structures without formal charge separation are intrinsically more stable than dipolar (zwitterionic) structures.
4. **Electronegativity & Charge Concordance:** Among charge-separated structures:
   - Negative formal charges reside most stably on the most electronegative atoms ($O^- > N^- > C^-$).
   - Positive formal charges reside most stably on the least electronegative atoms ($C^+ > N^+ > O^+$).
5. **Proximity of Like/Unlike Charges:** Canonical forms with like charges on adjacent atoms are violently destabilized (electrostatic repulsion); unlike charges close together confer electrostatic stabilization.

Resonance Energy (E_{res}) Formulation

Resonance energy is defined as the difference in energy between the actual resonance hybrid and the most stable theoretical canonical Lewis structure:

$$E_{\text{res}} = E_{\text{canonical (most stable)}} - E_{\text{actual hybrid}} \tag{5.20}$$

Experimentally, resonance energy is determined from standard enthalpies of hydrogenation ($\Delta_{\text{hydro}}H^\circ$):

$$\begin{aligned} \text{Cyclohexene} + \text{H}_2 &\rightarrow \text{Cyclohexane} \quad \Delta H^\circ = -119.7 \text{ kJ/mol} \\ \text{Theor. Cyclohexatriene} &\implies 3 \times (-119.7) = -359.1 \text{ kJ/mol} \\ \text{Expt. Benzene} + 3\text{H}_2 &\rightarrow \text{Cyclohexane} \quad (\Delta H^\circ = -208.5 \text{ kJ/mol}) \\ E_{\text{res}}(\text{Benzene}) &= |-359.1 - (-208.5)| = 150.6 \text{ kJ/mol} \end{aligned} \tag{5.21}$$

NEET Trap: Halogen Paradox in EAS ($+M$ vs $-I$)

Halogens ($-$F, $-$Cl, $-$Br, $-$I) attached to benzene present a classical dual nature:

- **Reactivity (Net Deactivating):** Halogens are strongly electronegative ($\chi_{\text{F}} = 4.0, \chi_{\text{Cl}} = 3.16$), creating a powerful $-I$ effect that drains σ-electron density from the ring. The $-I$ effect dominates over the $+M$ resonance effect for overall rate, making haloarenes **less reactive** than benzene toward electrophiles.
- **Regioselectivity (Ortho/Para Directing):** During attack by an incoming electrophile (E^+), the halogen donates its p-lone pair via $+M$ resonance to stabilize the positively charged arenium ion specifically when E^+ attacks the **ortho** or **para** positions (giving a complete-octet canonical structure [$\text{Cl}^+ = \text{C}_6\text{H}_5\text{E}$]). At the meta position, the halogen lone pair cannot stabilize the carbocation.

Conclusion: Halogens are uniquely **deactivating ortho/para-directors** in EAS.

5.2.3 Hyperconjugation (Baker-Nathan Effect)

The $\sigma \rightarrow \pi^*$ and $\sigma \rightarrow p$ No-Bond Resonance

Hyperconjugation involves the delocalization of bonding electrons in a C $-$ H (or C $-$ C) σ-bond into an adjacent vacant or partially filled non-bonding p-orbital or an antibonding π^* molecular orbital:

$$\sigma_{\text{C}-\text{H}} \longrightarrow p_{\text{vacant}} \quad \text{(in carbocations)} \quad \text{or} \quad \sigma_{\text{C}-\text{H}} \longrightarrow \pi^* \quad \text{(in alkenes)} \tag{5.22}$$

Structural Characteristics:

1. **Number of Contributing Structures:** For an alkene, carbocation, or free radical with n α-hydrogens (hydrogens on the sp^3 carbon directly attached to the sp^2 system), the total number of hyperconjugative canonical forms is:

$$\text{Total Hyperconjugative Structures} = n + 1 \quad (\text{where } n = \text{number of } \alpha\text{-H}) \tag{5.23}$$

2. **Bond Shortening & Bond Lengthening:** The single bond between the sp^3 and sp^2 carbons acquires partial double bond character (e.g., the C − C single bond in propene is 1.49 Å, significantly shorter than standard 1.54 Å in ethane). The C_α−H bond is lengthened and weakened.
3. **Reverse Hyperconjugation:** Observed when electronegative halogen atoms reside at the α-carbon (e.g., $-CX_3$ in trifluoromethylbenzene $C_6H_5 - CF_3$). Electron density flows from the π-system into the vacant σ^*_{C-F} antibonding orbital, causing meta-direction and ring deactivation.

Alkene Stability & Heat of Hydrogenation ($\Delta_{\text{hydro}}H^\circ$)

Alkene thermodynamic stability is directly proportional to the number of hyperconjugative α-hydrogens:

$$\text{Alkene Stability} \propto (\text{Number of } \alpha\text{-H}) \propto \frac{1}{\text{Heat of Hydrogenation } (\Delta_{\text{hydro}}H^\circ)} \tag{5.24}$$

Authoritative Stability Sequence:

$$\begin{aligned} (CH_3)_2C = C(CH_3)_2\ (12\alpha\text{-H}) &> (CH_3)_2C = CHCH_3\ (9\alpha\text{-H}) \\ &> CH_2 = C(CH_3)_2\ (6\alpha\text{-H}, \text{gem-dimethyl}) \\ &> \textit{trans}{-}CH_3CH = CHCH_3\ (6\alpha\text{-H}) \\ &> \textit{cis}{-}CH_3CH = CHCH_3\ (6\alpha\text{-H}) \\ &> CH_3CH_2CH = CH_2\ (2\alpha\text{-H}) \\ &> CH_2 = CH_2\ (0\alpha\text{-H}) \end{aligned} \tag{5.25}$$

The *trans* vs *cis* Isomer Trap: *trans*-2-butene is more stable than *cis*-2-butene because the two bulky methyl groups are oriented at 180°, eliminating the steric van der Waals repulsion present in the eclipsed-like geometry of the *cis*-isomer. The *gem*-dimethyl alkene ($CH_2 = C(CH_3)_2$) is thermodynamically more stable than both *cis*- and *trans*-2-butene due to superior electrostatic distribution across the two methyl groups attached to the same sp^2 carbon.

5.2.4 Electromeric Effect ($\pm E$)

Temporary Reagent-Induced π-Polarization

The electromeric effect is a purely **temporary** electronic displacement operating exclusively in molecules containing multiple bonds ($C=C, C\equiv C, C=O$) upon the approach of an attacking polar reagent. Upon removal of the reagent, the system instantly reverts to its unpolarized ground state.

- **Positive Electromeric Effect ($+E$):** The π-electron pair shifts toward the atom to which the attacking reagent attaches:

$$CH_2 = CH_2 + H^+ \xrightarrow{+E} CH_2^+ - CH_2 - H \quad \text{(Electrophilic attack on alkene)} \tag{5.26}$$

- **Negative Electromeric Effect ($-E$):** The π-electron pair shifts away from the atom to which the attacking reagent attaches:

$$>C=O + CN^- \xrightarrow{-E} >C(CN) - O^- \quad \text{(Nucleophilic attack on carbonyl)} \tag{5.27}$$

5.2.5 Aromaticity & Hückel's Rule

The Four Cardinal Criteria for Aromaticity

To be classified as **aromatic**, a chemical entity must simultaneously satisfy all four structural and electronic requirements formulated by Erich Hückel:

1. **Cyclic Structure:** The system must possess a continuous closed loop of bonded atoms.
2. **Planar Geometry:** All ring vertices must be strictly planar (or possess < 0.1 Å deviation) to maintain parallel p-orbital orientation.
3. **Complete Unbroken Conjugation:** Every atom in the cyclic ring must possess an unhybridized p-orbital (i.e., fully sp^2 or sp hybridized; zero tetrahedral sp^3 carbons in the ring loop).
4. **Hückel's $(4n+2)\pi$ Electron Rule:** The number of delocalized π-electrons must equal $(4n+2)$, where n is a non-negative integer ($n = 0, 1, 2, 3, \cdots \implies 2, 6, 10, 14, 18\ldots$ π-electrons).

Aromatic vs Anti-Aromatic vs Non-Aromatic Classification

Property	Aromatic	Anti-Aromatic	Non-Aromatic
Cyclic & Planar?	Yes	Yes	Lacks cyclicity or planarity
Conjugation	Unbroken throughout ring	Unbroken throughout ring	Interrupted by sp^3 carbon
π-Electron Count	$(4n+2)\pi$ electrons	$4n\pi$ electrons ($n = 1, 2, \dots$)	Arbitrary count
Thermodynamic Stability	Exceptionally High ($E_{\text{res}} \gg 0$)	Exceptionally Low ($E_{\text{strain}} \gg 0$)	Normal open-chain analog
Magnetic Induced Field	Diatropic (Diamagnetic ring current)	Paratropic (Paramagnetic ring current)	No ring current
Canonical Examples	Benzene (6π), Tropylium (6π)	Cyclobutadiene (4π), Cyclopentadienyl cation (4π)	1,3-Cyclohexadiene, COT (tub-shaped)

Annulenes & Non-Planar Ring Escapes

1. Cyclooctatetraene (COT, [8]-Annulene): COT possesses 8π-electrons ($4n$ with $n = 2$). If planar, it would be severely anti-aromatic. To escape extreme anti-aromatic electronic destabilization, the flexible 8-membered ring distorts into a non-planar **tub conformation**. Because p-orbitals are perpendicular at the edges, conjugation is broken: COT is strictly **non-aromatic**. However, reduction with potassium metal yields the cyclooctatetraenyl dianion ($C_8H_8^{2-}$), which has 10π-electrons ($4n+2, n=2$), becomes strictly planar, and is **aromatic**!
2. [10]-Annulene: Possesses 10π-electrons ($4n+2, n=2$). In the planar configuration, severe steric repulsion between the two internal hydrogen atoms forces the ring out of planarity $\implies$ **non-aromatic**. If the internal hydrogens are replaced with a methylene bridge (1,6-methano[10]annulene), the ring achieves planarity and exhibits robust **aromaticity**.

Heterocyclic Aromaticity & Lone Pair Involvement

When determining Hückel π-electron counts in heterocycles containing N, O, S:

- **Pyridine (C_5H_5N):** The nitrogen atom is sp^2-hybridized. Its unhybridized p-orbital contributes $1e^-$ to the 6π aromatic sextet. The lone pair resides in an sp^2 hybrid orbital in the ring plane, perpendicular to the π-cloud. It does **not** participate in aromaticity $\implies$ Pyridine is basic ($pK_b \approx 8.8$).
- **Pyrrole (C_4H_4NH):** The nitrogen atom is sp^2-hybridized. Its lone pair resides in an unhybridized p-orbital, contributing $2e^-$ to complete the 6π aromatic sextet. The lone pair is tied up in aromaticity $\implies$ Pyrrole is an exceptionally weak base ($pK_b \approx 13.6$) and protonation on nitrogen destroys aromaticity.
- **Furan (C_4H_4O) & Thiophene (C_4H_4S):** The heteroatom contributes one lone pair ($2e^-$) to the aromatic sextet (6π electrons); the second lone pair resides in an sp^2 orbital orthogonal to the π-system.

5.3 Reactive Intermediates & Molecular Rearrangements

5.3.1 Carbocations

Geometry, Electronic Configuration & Stability Hierarchy

Carbocations are trivalent carbon cations possessing a positive formal charge and a sextet ($6e^-$) of valence electrons:

- **Geometry & Hybridization:** The positively charged carbon is sp^2-hybridized with trigonal planar geometry (120° bond angles). The remaining unhybridized $2p_z$ orbital is completely vacant.
- **Stability Factors:** Stabilized by $+M$ resonance delocalization, aromatic stabilization, hyperconjugation ($n_{\alpha-\mathrm{H}}$), and $+I$ inductive effects.

$$\text{Tropylium } (6\pi) \approx \text{Cyclopropylcarbinyl} \tag{5.28}$$

$$> (\mathrm{C_6H_5})_3\mathrm{C^+} > (\mathrm{C_6H_5})_2\mathrm{CH^+} \tag{5.29}$$

$$> \mathrm{C_6H_5CH_2^+} \approx \mathrm{CH_2=CH-CH_2^+} \tag{5.30}$$

$$> 3^\circ\ (\mathrm{CH_3})_3\mathrm{C^+} > 2^\circ\ (\mathrm{CH_3})_2\mathrm{CH^+} \tag{5.31}$$

$$> 1^\circ\ \mathrm{CH_3CH_2^+} > \mathrm{CH_3^+} > \text{vinyl} > \text{phenyl} \tag{5.32}$$

The Driving Force & Pathways of Carbocation Rearrangements

A carbocation will spontaneously rearrange if and only if the rearranged carbocation possesses lower Gibbs free energy (greater thermodynamic stability). Rearrangements occur via intramolecular 1,2-migrations of a hydride, alkyl, or aryl group into the adjacent vacant p-orbital:

1. **1,2-Hydride Shift (H^-):**

$$\mathrm{CH_3-CH_2-CH_2^+}\ (1^\circ, 2\alpha\text{-H}) \xrightarrow{\text{1,2-H}^-} \mathrm{CH_3-\overset{+}{C}H-CH_3}\ (2^\circ, 6\alpha\text{-H}) \tag{5.33}$$

2. **1,2-Methyl / Alkyl Shift (R^-):**

$$(\mathrm{CH_3})_3\mathrm{C-\overset{+}{C}H_2}\ (1^\circ) \xrightarrow{\text{1,2-CH}_3^-} \mathrm{CH_3-\overset{+}{C}(CH_3)-CH_2CH_3}\ (3^\circ, 8\alpha\text{-H}) \tag{5.34}$$

3. **Ring Expansion (Angle Strain Relief):** Four- and five-membered rings with an exocyclic carbocation undergo instantaneous 1,2-bond migration to expand the ring:
   - **Cyclobutylcarbinyl Cation → Cyclopentyl Cation:** The 90° ring angle expands to 108°, relieving ≈ 108.8 kJ/mol of Baeyer angle strain.
   - **Cyclopentylcarbinyl Cation → Cyclohexyl Cation:** Expansion from 5-membered ring (108°) to strain-free chair cyclohexane (109.5°).

NEET Trap: Migratory Aptitude in Carbocation Rearrangements

When competing groups are attached to the migrating carbon, the intrinsic migratory aptitude is:

$$\mathrm{H}^- > \text{Aryl } (\mathrm{Ar}^-) > \text{Alkyl } (3^\circ > 2^\circ > 1^\circ > \mathrm{CH}_3^-) \tag{5.35}$$

For substituted aryl groups: electron-donating groups accelerate migration:

$$\begin{aligned} p\text{-anisyl } (p{-}\mathrm{CH_3O} - \mathrm{C_6H_4}{-}) > p\text{-tolyl } (p{-}\mathrm{CH_3} - \mathrm{C_6H_4}{-}) > \text{phenyl } (\mathrm{C_6H_5}{-}) \\ > p\text{-chlorophenyl} > p\text{-nitrophenyl} \end{aligned} \tag{5.36}$$

Crucial Rule: Never perform a less-stable to less-stable rearrangement unless it immediately opens a direct pathway to an overwhelmingly more stable intermediate (e.g., resonance-delocalized benzylic or allylic system).

5.3.2 Carbanions

Structure, Pyramidal Inversion & Stability Hierarchy

Carbanions possess a trivalent carbon bearing a formal negative charge and eight valence electrons (3 bonding pairs + 1 lone pair):

- **Hybridization & Inversion:** Unconjugated carbanions are sp^3-hybridized with trigonal pyramidal geometry. The lone pair undergoes rapid umbrella inversion (flipping barrier ≈ 25 kJ/mol), making isolated chiral carbanions rapidly racemize.
- **Conjugated Carbanions (sp^2 Planarity):** If the carbanion is adjacent to a π-system or an electron-withdrawing carbonyl group ($-\mathrm{C}=\mathrm{O}$), it rehybridizes to sp^2 to accommodate the lone pair in an unhybridized p-orbital for maximum resonance delocalization.
- **Stability Sequence:**

$$\begin{aligned} \text{Cyclopentadienyl } (6\pi) > (\mathrm{C_6H_5})_3\mathrm{C}^- > (\mathrm{C_6H_5})_2\mathrm{CH}^- \\ > \mathrm{C_6H_5CH_2^-} \approx \mathrm{CH_2=CH-CH_2^-} > \mathrm{HC\equiv C}^- \\ > \mathrm{CH_2=CH}^- \approx \mathrm{C_6H_5^-} > \mathrm{CH_3^-} > 1^\circ > 2^\circ > 3^\circ \end{aligned} \tag{5.37}$$

Active Methylene Compounds & Enolate Chemistry

Methylene groups ($-\mathrm{CH_2}-$) flanked between two strong electron-withdrawing groups (e.g., diethyl malonate, ethyl acetoacetate, acetylacetone) possess exceptionally acidic hydrogens ($pK_a \approx 9-13$):

$$\mathrm{CH_3COCH_2COCH_3} + \mathrm{EtO}^- \rightleftharpoons \tag{5.38}$$

$$[\mathrm{CH_3CO\bar{C}HCOCH_3}] + \mathrm{EtOH} \longleftrightarrow \text{Enolate} \tag{5.39}$$

The resulting enolate anion is an **ambident nucleophile** that can attack electrophiles via carbon (*C*-alkylation) or oxygen (*O*-alkylation), driving fundamental C − C bond-forming reactions (Aldol, Claisen, Michael additions).

5.3.3 Free Radicals

Structure, Generation & Allylic Halogenation via NBS

Carbon free radicals are neutral trivalent species possessing seven valence electrons (one unpaired electron):

- **Structure:** Planar sp^2 hybridized with the odd electron occupying the unhybridized $2p$ orbital, or shallow rapidly inverting pyramid.
- **Stability:** Stabilized by hyperconjugation and resonance delocalization:

$$(C_6H_5)_3C^\bullet > (C_6H_5)_2\dot{C}H > C_6H_5\dot{C}H_2 \approx CH_2 = CH - \dot{C}H_2 \tag{5.40}$$

$$> 3^\circ > 2^\circ > 1^\circ > \dot{C}H_3 > \text{vinyl} \tag{5.41}$$

- **N-Bromosuccinimide (NBS) Allylic/Benzylic Halogenation:** NBS selectively brominates the allylic or benzylic position without adding across the C = C double bond. **Mechanism:** Trace HBr reacts with NBS to maintain a constant, extremely low concentration of molecular bromine (Br_2):

$$Br^\bullet + CH_3CH = CH_2 \longrightarrow HBr + [\dot{C}H_2CH = CH_2] \tag{5.42}$$

$$NBS + HBr \longrightarrow \text{Succinimide} + Br_2 \ ([Br_2] \approx 10^{-6}\ \text{M}) \tag{5.43}$$

$$\dot{C}H_2CH = CH_2 + Br_2 \longrightarrow BrCH_2CH = CH_2 + Br^\bullet \tag{5.44}$$

The low $[Br_2]$ prevents electrophilic addition across the double bond, ensuring exclusive allylic radical substitution!

5.3.4 Carbenes, Nitrenes & Arynes

Singlet vs Triplet Carbenes (: CR_2)

Feature	Singlet Carbene (^{1} : CR_2)	Triplet Carbene (^{3} : CR_2)
Electron Spin State	Spin-paired ($\uparrow\downarrow$, $S = 0, 2S + 1 = 1$)	Parallel spins ($\uparrow\uparrow$, $S = 1, 2S + 1 = 3$)
Hybridization & Geometry	sp^2, Bent ($102 - 108°$)	sp linear ($180°$) or bent sp^2 ($130 - 150°$)
Orbital Configuration	Lone pair in sp^2, vacant p-orbital	One electron in sp^2, one in p-orbital
Relative Stability	Thermodynamically less stable (except halocarbenes : CCl_2)	Ground state for : CH_2 (Hund's rule)
Addition to Alkenes	**Stereospecific syn-addition** (concerted, geometry retained: cis → cis)	**Non-stereospecific** (diradical intermediate, spin flip allows rotation → cis + trans)

Generation of Dichlorocarbene & The Benzyne Mechanism

1. Dichlorocarbene (: CCl_2) via α-Elimination: Chloroform treated with strong alkali undergoes two-step α-elimination:

$$CHCl_3 + OH^- \overset{\text{fast}}{\rightleftharpoons} : \bar{C}Cl_3 + H_2O \tag{5.45}$$

$$\xrightarrow[\text{RDS}]{-Cl^-} : CCl_2 \quad \text{(Electrophilic singlet carbene)} \tag{5.46}$$

: CCl_2 serves as the key electrophilic intermediate in the **Reimer-Tiemann reaction** and the **Carbylamine reaction**.

2. Benzyne Intermediate (Elimination-Addition): Aryl halides lacking electron-withdrawing groups react with extremely strong bases (e.g., KNH_2/liq.NH_3) via elimination of HX to yield **benzyne** (C_6H_4):

- **Bonding in Benzyne:** The formal triple bond is formed by sideways overlap of two sp^2 hybrid orbitals in the plane of the ring. This bond does not disturb the 6π aromatic sextet, but overlap is extremely weak due to poor orbital alignment $\implies$ severe ring strain and immense reactivity.
- **Cine-Substitution:** Addition of NH_3 across the strained triple bond occurs with equal probability at either carbon, producing isotopic scrambling (50% direct substitution + 50% cine-substitution).

5.4 Hydrocarbons: Aliphatic & Aromatic Pathways

5.4.1 Alkanes

Free-Radical Halogenation Kinetics & Hammond's Postulate

Free radical halogenation follows a three-stage chain mechanism (Initiation, Propagation, Termination):

$$\text{Initiation: } X_2 \xrightarrow{h\nu \text{ or } \Delta} 2X^\bullet \tag{5.47}$$

$$\text{Propagation: } R-H + X^\bullet \xrightarrow{k_1} R^\bullet + HX \quad \text{(RDS for bromination)} \tag{5.48}$$

$$R^\bullet + X_2 \xrightarrow{k_2} R-X + X^\bullet \tag{5.49}$$

Relative Reactivity Rates at 298 K:

- **Chlorination (Reactivity over Selectivity):** $1^\circ : 2^\circ : 3^\circ = 1.0 : 3.8 : 5.0$. Chlorination is highly exothermic; the transition state is early (reactant-like per Hammond's postulate), meaning $C-H$ bond cleavage has barely progressed, yielding low regioselectivity.
- **Bromination (Selectivity over Reactivity):** $1^\circ : 2^\circ : 3^\circ = 1.0 : 82 : 1600$. Bromination is endothermic; the transition state is late (product-like), with extensive radical character on carbon, leading to massive selectivity for the 3° alkyl halide ($> 99\%$).

Conformational Analysis of Ethane and n-Butane

- **Ethane:** Infinite conformations between **staggered** (dihedral angle $\theta = 60^\circ$, minimum energy) and **eclipsed** ($\theta = 0^\circ$, maximum energy). The energy barrier is 12.5 kJ/mol (3.0 kcal/mol), caused by **torsional strain** (electron cloud repulsion between adjacent $C-H$ bonds).
- **n-Butane (Rotation about $C_2 - C_3$):**

$$\text{Anti } (\theta = 180^\circ) > \text{Gauche } (\theta = 60^\circ) \tag{5.50}$$

$$> \text{Partially Eclipsed } (120^\circ) > \text{Fully Eclipsed } (0^\circ) \tag{5.51}$$

  The anti conformation is the global minimum. Gauche is destabilized by 3.8 kJ/mol due to van der Waals steric repulsion between the two bulky methyl groups. The fully eclipsed form is destabilized by 19.0 kJ/mol due to combined torsional and steric strain.
- **Gauche Exception (Intramolecular H-Bonding):** In ethylene glycol ($HO-CH_2-CH_2-OH$) and 2-fluoroethanol ($F-CH_2-CH_2-OH$), the **gauche** conformation is more stable than anti due to stabilizing intramolecular hydrogen bonding.

5.4.2 Alkenes

Electrophilic Addition of Halogens: Cyclic Halonium Ion

Electrophilic addition of Br_2 (in nonpolar solvent CCl_4) to alkenes proceeds via a cyclic three-membered **bromonium ion**:

1. The π-cloud of the alkene polarizes the $Br-Br$ bond, displacing the distal bromide ion to form a bridged cyclic bromonium ion intermediate.
2. The nucleophilic Br^- attacks the cyclic intermediate from the **backside** (anti-attack) with complete inversion at the attacked carbon.
3. **Stereospecificity:**
   - *trans*-Alkene + Br_2 (anti-addition) $\longrightarrow$ ***meso*-dibromide**.
   - *cis*-Alkene + Br_2 (anti-addition) $\longrightarrow$ **racemic mixture (±) of enantiomers**.
   - *(Mnemonic: CAR – Cis + Anti → Racemic; TAM – Trans + Anti → Meso).*

Markovnikov Addition vs Anti-Markovnikov Peroxide Effect

1. Markovnikov's Rule: In electrophilic addition of unsymmetrical HX to unsymmetrical alkenes, the electrophilic proton (H^+) attaches to the sp^2 carbon carrying the greater number of hydrogen atoms to generate the more stable carbocation intermediate (which may rearrange!).
2. Anti-Markovnikov (Kharasch / Peroxide Effect): Operates exclusively when **HBr** is added in the presence of organic peroxides ($R-O-O-R$):

$$\text{Init: } ROOR \xrightarrow{\Delta} 2RO^{\bullet}, \quad RO^{\bullet} + HBr \rightarrow ROH + Br^{\bullet} \tag{5.52}$$

$$\text{Prop 1: } CH_3-CH=CH_2 + Br^{\bullet} \rightarrow CH_3-\dot{C}H-CH_2Br \ (-42 \text{ kJ/mol}) \tag{5.53}$$

$$\text{Prop 2: } CH_3-\dot{C}H-CH_2Br + HBr \rightarrow CH_3-CH_2-CH_2Br + Br^{\bullet} \tag{5.54}$$

Thermochemical Reason: Why Peroxide Effect Operates ONLY for HBr

NEET frequently tests why the peroxide effect fails completely for HCl and HI:

- **Failure with HCl:** The $H-Cl$ bond is exceptionally strong (431 kJ/mol). Step 2 ($R^{\bullet} + HCl \rightarrow R-H + Cl^{\bullet}$) is strongly **endothermic** ($\Delta H^{\circ} \approx +27$ kJ/mol). The chain reaction cannot propagate.
- **Failure with HI:** The $H-I$ bond is weak (297 kJ/mol), making $I^{\bullet}$ generation easy. However, iodine radicals are large and unreactive toward alkenes: Step 1 (Alkene + $I^{\bullet} \rightarrow$ Radical) is **endothermic** ($\Delta H^{\circ} \approx +54$ kJ/mol). Instead of adding to the double bond, iodine radicals dimerize instantaneously: $2I^{\bullet} \rightarrow I_2$.

Conclusion: Only for HBr are **both propagation steps exothermic**, permitting chain reaction!

Comparative Summary: The Three Alkene Hydration Pathways

Parameter	Acid-Catalyzed Hydration	Hydroboration-Oxidation (HBO)	Oxymercuration-Demercuration (OMDM)
Reagents	H_2O/H_2SO_4	1. $BH_3 \cdot THF$, 2. H_2O_2, OH^-	1. $Hg(OAc)_2, H_2O$, 2. $NaBH_4$
Regiochemistry	Markovnikov	**Anti-Markovnikov**	**Markovnikov**
Stereochemistry	Non-stereospecific	**Syn-addition**	Overall non-stereospecific (anti in step 1)
Intermediate	Free Carbocation	4-center cyclic transition state	Cyclic mercurinium ion
Rearrangement?	**Yes (frequent)**	**Zero rearrangement**	**Zero rearrangement**
Propene Product	Propan-2-ol	Propan-1-ol	Propan-2-ol
3,3-Dimethylbut-1-ene	2,3-dimethylbutan-2-ol (rearranged)	3,3-dimethylbutan-1-ol	3,3-dimethylbutan-2-ol (clean Markovnikov)

Direct NCERT Extraction: Ozonolysis & Structure Elucidation

Ozonolysis cleaves the C = C double bond into two carbonyl fragments:

- **Reductive Ozonolysis (O_3 followed by Zn/H_2O or $(CH_3)_2S$):** Aldehydes and ketones are preserved. Zinc destroys hydrogen peroxide ($H_2O_2 + Zn \rightarrow ZnO + H_2O$), preventing oxidation of aldehydes to carboxylic acids.
- **Oxidative Ozonolysis (O_3 followed by H_2O_2 or H_2O):** All aldehydes are oxidized to carboxylic acids ($-CHO \rightarrow -COOH$); terminal = CH_2 becomes $CO_2 + H_2O$; ketones remain unchanged.

5.4.3 Alkynes

Terminal Alkyne Acidity & Selective Hydrogenation

1. Acidity of Terminal Alkynes ($-C \equiv CH$): The *sp*-hybridized carbon has 50%*s*-character, placing valence electrons closest to the carbon nucleus. Terminal alkynes are acidic ($pK_a \approx 25$):

- React with sodamide: $R - C \equiv CH + NaNH_2 \rightarrow R - C \equiv C^-Na^+ + NH_3 \uparrow$.
- Tollens' Test: Forms a **white precipitate** of silver acetylide: $R - C \equiv CH + [Ag(NH_3)_2]^+OH^- \rightarrow R - C \equiv C - Ag \downarrow + 2NH_3 + H_2O$.
- Ammoniacal Cu_2Cl_2: Forms a **red precipitate** of cuprous acetylide: $R - C \equiv CH + [Cu(NH_3)_2]^+OH^- \rightarrow R - C \equiv C - Cu \downarrow + 2NH_3 + H_2O$.

- *Note: Internal alkynes* ($R-C\equiv C-R'$) *give negative tests with both reagents!*

2. Stereoselective Reduction to Alkenes:

- **Lindlar's Catalyst** ($H_2/Pd-CaCO_3$ **poisoned with quinoline / lead acetate):** Syn-addition of hydrogen yielding ***cis*-alkene**.
- **Birch Reduction** ($Na/$**liq.** NH_3 **at** $-33°$**C):** Anti-addition proceeding via radical-anion intermediate with trans-geometry to minimize electron-electron repulsion, yielding ***trans*-alkene**.

3. Kucherov Hydration ($HgSO_4/dil.H_2SO_4$ **at** 333 **K):** Electrophilic addition of water across the triple bond yields an enol, which immediately undergoes keto-enol tautomerism:

$$HC\equiv CH + H_2O \xrightarrow{Hg^{2+}/H^+} [CH_2=CH-OH] \rightleftharpoons CH_3-CHO \tag{5.55}$$

$$CH_3-C\equiv CH + H_2O \xrightarrow{Hg^{2+}/H^+} [CH_3-C(OH)=CH_2] \rightleftharpoons CH_3-CO-CH_3 \tag{5.56}$$

(Acetylene gives acetaldehyde; all higher alkynes yield methyl ketones).

5.4.4 Aromatic Hydrocarbons: EAS Mechanics & Substituent Effects

Fundamental 3-Step EAS Mechanism & Arenium Ion (σ-Complex)

Electrophilic aromatic substitution (S_EAr) preserves the massive resonance stabilization of the benzene ring (150.6 kJ/mol):

1. **Generation of Active Electrophile** (E^+)**:** Lewis or Brønsted acid generates E^+ ($NO_2^+, Cl^+, SO_3, R^+, RCO^+$).
2. **Formation of Arenium Ion (**σ**-Complex / Wheland Intermediate) [RDS]:** $C_6H_6 + E^+ \xrightarrow{slow} [C_6H_6E]^+$. The attacked carbon rehybridizes from sp^2 to sp^3, disrupting aromaticity. Positive charge delocalizes over ortho and para positions.
3. **Rapid Deprotonation & Rearomatization:** Fast abstraction of the proton from the sp^3 carbon restores the planar sp^2 aromatic sextet.

Directing and Activating Effects of Ring Substituents

Group Type	Examples	Electronic Mechanism	Directing Influence
Strongly Activating	$-OH$, $-O^-$, $-NH_2$, $-NR_2$	Dominant $+M$ resonance	Ortho / Para
Moderately Activating	$-OCH_3$, $-NHCOCH_3$, $-OCOR$	$+M$ modulated by cross-conjugation	Ortho / Para
Weakly Activating	$-CH_3$, $-CH_2CH_3$, $-C_6H_5$	Hyperconjugation & $+I$	Ortho / Para
Deactivating (Halogens)	$-F$, $-Cl$, $-Br$, $-I$	Strong $-I$ dominates rate; $+M$ directs	**Ortho / Para**
Moderately Deactivating	$-CHO$, $-COR$, $-COOH$, $-COOR$	$-M$ resonance & $-I$	Meta
Strongly Deactivating	$-NO_2$, $-CF_3$, $-CCl_3$, $-CN$, $-SO_3H$, $-NR_3^+$	Powerful $-M$ / $-I$ withdraws electron density	Meta

Friedel-Crafts Reactions: Critical Limitations & Exceptions

1. Carbocation Rearrangement in Alkylation: Reaction of benzene with 1-chloropropane ($CH_3CH_2CH_2Cl$) and $AlCl_3$ yields **isopropylbenzene (cumene)** as major product ($65-70\%$) due to 1,2-hydride shift of the primary propyl carbocation to the more stable secondary carbocation!

$$\text{Benzene} + CH_3CH_2CH_2Cl \xrightarrow{AlCl_3} C_6H_5 - CH(CH_3)_2 \text{ (Major)} \quad (5.57)$$

$$+ C_6H_5 - CH_2CH_2CH_3 \text{ (Minor)} \quad (5.58)$$

Solution: To synthesize linear alkylbenzenes, carry out **Friedel-Crafts Acylation** with propanoyl chloride to yield ethyl phenyl ketone (no rearrangement because acylium ion $R - C \equiv O^+$ is resonance-stabilized), followed by **Clemmensen reduction** ($Zn - Hg/HCl$) or **Wolff-Kishner reduction**.

2. Polyalkylation vs Monoacylation: Alkyl groups activate the ring toward further substitution (polyalkylation occurs). Acyl groups ($-COR$) are strongly deactivating, ensuring clean monoacylation.

3. Complete Failure with Aniline and Phenol: Aniline ($-NH_2$) and Phenol ($-OH$) fail Friedel-Crafts reactions because their lone pairs form an insoluble Lewis acid-base adduct with $AlCl_3$ (e.g., $Ph - NH_2 \cdot AlCl_3$), placing a positive charge adjacent to the ring and rendering it deactivated toward electrophiles!

5.5 Haloalkanes & Haloarenes

5.5.1 Nucleophilic Aliphatic Substitution (S_N1 vs S_N2 vs S_Ni)

Comprehensive Mechanistic Comparison: S_N1 vs S_N2

Parameter	S_N1 (Substitution Nucleophilic Unimolecular)	S_N2 (Substitution Nucleophilic Bimolecular)
Number of Steps	Two discrete elementary steps	Single concerted step
Rate Law & Kinetics	Rate $= k[\mathrm{R-X}]$ (First order)	Rate $= k[\mathrm{R-X}][\mathrm{Nu^-}]$ (Second order)
Substrate Reactivity	$3^\circ > 2^\circ \gg 1^\circ > \mathrm{CH_3}$ (governed by carbocation stability)	$\mathrm{CH_3} > 1^\circ > 2^\circ \gg 3^\circ$ (governed by steric hindrance)
Transition State	Involves carbocation intermediate	Pentacoordinate trigonal bipyramidal TS
Stereochemical Outcome	Partial racemization with net inversion ($50-70\%$ inversion due to front-side ion pair shielding)	**100% Complete Walden Inversion** (backside attack)
Nucleophile Strength	Weak, neutral nucleophiles ($\mathrm{H_2O}$, ROH) suffice	Requires strong, anionic nucleophiles ($\mathrm{OH^-}$, $\mathrm{RO^-}$, $\mathrm{CN^-}$)
Solvent Preference	**Polar Protic** ($\mathrm{H_2O}$, EtOH, AcOH): solvates carbocation and stabilizes leaving group anion	**Polar Aprotic** (DMSO, DMF, acetone): solvates cations, leaving naked, highly reactive anions
Rearrangements	Carbocation rearrangements (shifts, expansions) common	**Zero rearrangement possible**
Leaving Group Order	$\mathrm{I^-} > \mathrm{Br^-} > \mathrm{Cl^-} \gg \mathrm{F^-}$	$\mathrm{I^-} > \mathrm{Br^-} > \mathrm{Cl^-} \gg \mathrm{F^-}$

The S_Ni Mechanism: Darzens Process with Thionyl Chloride

Reaction of alcohols with $\mathrm{SOCl_2}$ proceeds with different stereochemical outcomes depending on the solvent:

- **In Dioxane / Diethyl Ether (S_Ni, Internal Return):** Reaction forms an alkyl chlorosulfite ester ($\mathrm{R-O-SOCl}$). In nonpolar solvent, it dissociates into an intimate ion pair [$\mathrm{R^+\ SO_2Cl^-}$]. Decomposition releases $\mathrm{SO_2}\uparrow$, and the chloride is delivered to the carbocation from the **same face** from which the oxygen departed. Outcome: **Retention of configuration**.
- **In Pyridine (S_N2 Inversion):** Pyridine reacts with generated HCl and converts the chlorosulfite into an active pyridinium intermediate. Free chloride ion attacks from the **backside**. Outcome: **Complete Walden inversion**.

5.5.2 Elimination Reactions ($E1$ vs $E2$ vs $E1cB$)

Saytzeff (Zaitsev) vs Hofmann Regioselectivity

1. Saytzeff Rule (Thermodynamic Control): Dehydrohalogenation yields preferentially the more stable, more highly substituted alkene (greater number of hyperconjugative α-hydrogens).

- **Conditions for Saytzeff Selectivity:** Small bases (EtO^-, MeO^-, OH^-) acting on substrates with good leaving groups (I^-, Br^-, Cl^-).

$$\begin{aligned} &CH_3CH_2CH(Br)CH_3 \xrightarrow{\text{alc. KOH}} \\ &\quad CH_3CH=CHCH_3\ (81\%) + CH_3CH_2CH=CH_2\ (19\%) \end{aligned} \tag{5.59}$$

2. Hofmann Rule (Kinetic Control): Elimination yields preferentially the less substituted, terminal alkene.

- **Conditions for Hofmann Selectivity:**
  1. **Bulky Sterically Hindered Bases:** Potassium tert-butoxide ($t-BuO^-K^+$), diisopropylamide. The base cannot access the more hindered internal β-hydrogen and abstracts the accessible terminal β-hydrogen.
  2. **Poor Leaving Groups:** Fluoride ($-F$) or quaternary ammonium ($-NR_3^+$) and sulfonium ($-SR_2^+$) salts. Due to extreme electronegativity, the β-hydrogen becomes acidic, developing significant carbanion character in the transition state. The less substituted primary carbanion is more stable than secondary, directing elimination to the terminal alkene.

3. $E2$ **Stereochemistry:** Elimination requires strict **anti-periplanar** geometry: the $\beta-H$, the two carbons, and the leaving group X must lie in a single plane with dihedral angle $\theta = 180°$.

5.5.3 Nucleophilic Aromatic Substitution (S_NAr)

Addition-Elimination via Meisenheimer Complex

Unactivated haloarenes are virtually inert toward nucleophiles at ambient conditions because the $C-X$ bond possesses partial double bond character (1.69 Å in chlorobenzene vs 1.77 Å in chloroethane) due to $+M$ resonance, and the sp^2 carbon holds electrons tightly. **Activation by Nitro Groups (S_NAr):** Introducing strong electron-withdrawing nitro ($-NO_2$) groups at **ortho** and **para** positions dramatically enhances substitution rate:

$$\text{Chlorobenzene (623 K)} \gg p\text{-nitrochlorobenzene (443 K)} \tag{5.60}$$

$$\gg 2,4\text{-dinitro (368 K)} \tag{5.61}$$

$$\gg 2,4,6\text{-trinitro (warm } H_2O) \tag{5.62}$$

Mechanism: Nucleophilic attack on the ipso carbon forms a resonance-stabilized carbanionic intermediate (**Meisenheimer complex**), where negative charge delocalizes into the ortho and para nitro groups. Loss of halide in Step 2 restores aromaticity. **Leaving Group Anomaly:** Unlike aliphatic substitutions, the rate order in S_NAr is:

$$Ar-F \gg Ar-Cl > Ar-Br > Ar-I \tag{5.63}$$

The rate-determining step is nucleophilic attack (Step 1). The extreme electronegativity of fluorine stabilizes the Meisenheimer intermediate via $-I$ pull, accelerating the rate!

5.6 Alcohols, Phenols & Ethers

5.6.1 Alcohols: Acidity, Oxidation & Dehydration

Acidity & The Lucas Distinction Test

1. Acidity Order of Alcohols:

$$\mathrm{H_2O}\ (pK_a = 15.7) > \mathrm{CH_3OH}\ (15.5) > 1^\circ\ \mathrm{RCH_2OH}\ (16) \tag{5.64}$$

$$> 2^\circ\ \mathrm{R_2CHOH}\ (16.5) > 3^\circ\ \mathrm{R_3COH}\ (17) \tag{5.65}$$

Alkoxide anions ($\mathrm{R-O^-}$) are destabilized by the $+I$ inductive effect of alkyl groups. Water is a stronger acid than all alcohols except methanol!
2. The Lucas Test (anhydrous $\mathrm{ZnCl_2}$+conc. HCl): Distinguishes $1^\circ, 2^\circ, 3^\circ$ alcohols via S_N1 conversion to insoluble alkyl chlorides:

$$\mathrm{R-OH + HCl} \xrightarrow{\mathrm{ZnCl_2}} \mathrm{R-Cl}\downarrow\ (\text{cloudy turbidity}) + \mathrm{H_2O} \tag{5.66}$$

- 3° **Alcohols:** Immediate turbidity within 5–10 seconds at room temperature.
- 2° **Alcohols:** Turbidity appears within 5 minutes.
- 1° **Alcohols:** Solution remains completely clear at room temperature; turbidity appears only on prolonged boiling.

Oxidation Reagents for Alcohols

- **Pyridinium Chlorochromate (PCC in $\mathrm{CH_2Cl_2}$):** Mild selective oxidant. Oxidizes 1° alcohols to **aldehydes** (stops cleanly without over-oxidation to carboxylic acid) and 2° alcohols to ketones.
- **Jones Reagent ($\mathrm{CrO_3}$/aq. $\mathrm{H_2SO_4}$/acetone):** Converts 1° alcohols directly to carboxylic acids; 2° alcohols to ketones.
- **Heated Copper Catalyst at 573 K (300°C):**
  - 1° Alcohol $\xrightarrow{\mathrm{Cu,573K}}$ Aldehyde $+\mathrm{H_2}\uparrow$ (Dehydrogenation).
  - 2° Alcohol $\xrightarrow{\mathrm{Cu,573K}}$ Ketone $+\mathrm{H_2}\uparrow$ (Dehydrogenation).
  - 3° Alcohol $\xrightarrow{\mathrm{Cu,573K}}$ **Alkene** $+\mathrm{H_2O}$ (**Dehydration**, not dehydrogenation!).

5.6.2 Phenols: Acidic Character & Named Reactions

Phenol Acidity & Substituent Influence

Phenol ($pK_a \approx 10$) is 10^6 times more acidic than aliphatic alcohols ($pK_a \approx 16-18$):

- **Resonance Stabilization of Phenoxide:** The phenoxide ion ($C_6H_5O^-$) delocalizes negative charge into the benzene ring over ortho and para positions (5 resonance structures). Alkoxide ions (RO^-) have zero resonance stabilization.
- **Substituent Effects on Phenol Acidity:**

$$\text{Picric acid (0.38)} > \text{2, 4-dinitrophenol (4.0)} \tag{5.67}$$

$$> p\text{-nitrophenol (7.15)} > o\text{-nitrophenol (7.23)} \tag{5.68}$$

$$> \text{Phenol (10.0)} > p\text{-cresol (10.2)} \tag{5.69}$$

Note: p-nitrophenol is more acidic than o-nitrophenol because the ortho isomer is stabilized by intramolecular hydrogen bonding, reducing proton release!

Kolbe's & Reimer-Tiemann Reactions

1. Kolbe's Reaction (Synthesis of Salicylic Acid): Sodium phenoxide is treated with weak electrophile CO_2 at 400 K and $4-7$ atm, followed by acidification:

$$C_6H_5ONa + CO_2 \xrightarrow[4-7\text{ atm}]{400\text{ K}} \text{Sodium Salicylate} \tag{5.70}$$

$$\xrightarrow{H^+} \text{Salicylic Acid (\textbf{o}-hydroxybenzoic acid)} \tag{5.71}$$

Salicylic acid acetylates with acetic anhydride to yield **Aspirin** (acetylsalicylic acid).

2. Reimer-Tiemann Reaction (Synthesis of Salicylaldehyde): Phenol treated with $CHCl_3$ and aqueous NaOH at 340 K:

$$C_6H_5OH + CHCl_3 + 3NaOH \longrightarrow \tag{5.72}$$

$$\text{Salicylaldehyde} + 3NaCl + 2H_2O \tag{5.73}$$

Mechanistic Detail: The electrophile is singlet dichlorocarbene ($:CCl_2$) generated in situ. Attack occurs predominantly at the ortho position due to cyclic sodium chelation in the transition state. If CCl_4 is used instead of $CHCl_3$, **salicylic acid** is formed directly!

5.6.3 Ethers: Williamson Synthesis & HI Cleavage

NEET Trap: Substrate Rules for Williamson Ether Synthesis

Williamson ether synthesis involves nucleophilic displacement of an alkyl halide by an alkoxide:

$$\mathrm{R-O^-Na^+ + R'-X \xrightarrow{S_N2} R-O-R' + NaX} \tag{5.74}$$

The Strict Constraint: The alkyl halide ($\mathrm{R'-X}$) **must be methyl or primary** (1°).

- **Correct Preparation of tert-Butyl Methyl Ether:**

$$\mathrm{(CH_3)_3C-O^-Na^+ + CH_3-I \xrightarrow{S_N2} (CH_3)_3C-O-CH_3 + NaI} \tag{5.75}$$

$$[\text{Yield } \approx 95\%] \tag{5.76}$$

- **Fatal Reversal Trap:** If sodium methoxide is treated with tert-butyl bromide:

$$\mathrm{CH_3-O^-Na^+ + (CH_3)_3C-Br \xrightarrow{E2} CH_2=C(CH_3)_2} \tag{5.77}$$

$$\mathrm{+ CH_3OH + NaBr} \quad \text{(2-methylpropene)} \tag{5.78}$$

Because tert-butyl bromide is sterically crowded, the strongly basic methoxide ion induces **exclusive** $E2$ **elimination**!

Cleavage of Ethers with Hydroiodic Acid (HI)

When unsymmetrical ether $\mathrm{R-O-R'}$ is cleaved with cold, concentrated HI:

1. **Case 1: Both Alkyl Groups are 1° or 2° (S_N2 Pathway):** The protonated ether $[\mathrm{R-OH^+-R'}]$ is attacked by iodide ion ($\mathrm{I^-}$) at the **less sterically hindered carbon**:

$$\mathrm{CH_3-O-CH_2CH_2CH_3 + HI \xrightarrow{S_N2} CH_3I + CH_3CH_2CH_2OH} \tag{5.79}$$

2. **Case 2: One Alkyl Group is 3° or Resonance-Stabilized (S_N1 Pathway):** Heterolysis occurs first to produce the stable 3° or benzylic carbocation:

$$\mathrm{(CH_3)_3C-O-CH_3 + HI \xrightarrow{S_N1} (CH_3)_3C-I + CH_3OH} \tag{5.80}$$

3. **Case 3: Alkyl Aryl Ethers (Anisole $\mathrm{C_6H_5-O-CH_3}$):** The $\mathrm{C_{aromatic}-O}$ bond has partial double bond character due to resonance and cannot be cleaved. Cleavage yields exclusively:

$$\mathrm{C_6H_5-O-CH_3 + HI \longrightarrow C_6H_5OH\ (phenol) + CH_3I} \tag{5.81}$$

Note: Excess hot HI *converts all aliphatic alcohols to alkyl iodides:* $\mathrm{R-OH + HI \rightarrow R-I + H_2O}$. *Phenol never converts to iodobenzene!*

5.7 Aldehydes, Ketones & Carboxylic Acids

5.7.1 Nucleophilic Addition Mechanisms

Carbonyl Electrophilicity & Relative Reactivity Order

The carbonyl carbon is sp^2-hybridized with planar geometry (120°). Strong electronegativity of oxygen ($\chi = 3.44$ vs $\chi_C = 2.55$) creates a permanent dipole:

$$> C^{\delta+} = O^{\delta-} \longleftrightarrow > \overset{+}{C} - \overset{-}{O} \tag{5.82}$$

Nucleophiles attack perpendicular to the carbonyl plane (107° Bürgi-Dunitz angle), transforming the planar sp^2 carbon into a tetrahedral sp^3 alkoxide intermediate. **Reactivity Hierarchy Toward Nucleophiles:**

$$\begin{aligned} \text{HCHO} &> \text{CH}_3\text{CHO} > \text{CH}_3\text{COCH}_3 && (5.83)\\ &> \text{C}_6\text{H}_5\text{CHO} > \text{C}_6\text{H}_5\text{COCH}_3 > \text{C}_6\text{H}_5\text{COC}_6\text{H}_5 && (5.84) \end{aligned}$$

Governing Factors:

- **Steric Effect:** Bulky alkyl or aryl groups crowd the transition state, hindering nucleophilic approach.
- **Electronic Effect:** Electron-donating $+I$ alkyl groups reduce partial positive charge on the carbonyl carbon. $+M$ resonance from aromatic rings strongly stabilizes the ground state, lowering electrophilicity.

Standard Nucleophilic Addition Reactions

1. **Addition of** HCN **(Cyanohydrin Formation):** Catalyzed by base (OH^- generates active nucleophile CN^-). Cyanohydrins hydrolyze in acid to α-hydroxy acids:

$$\begin{aligned} \text{CH}_3\text{CHO} + \text{HCN} &\xrightarrow{\text{OH}^-} \text{CH}_3\text{CH(OH)CN} && (5.85)\\ &\xrightarrow{\text{H}_3\text{O}^+} \text{CH}_3\text{CH(OH)COOH (Lactic Acid)} && (5.86) \end{aligned}$$

2. **Addition of Sodium Bisulfite** ($NaHSO_3$)**:** Aldehydes and methyl ketones form crystalline, water-soluble bisulfite adducts. The adduct regenerates pure carbonyl on treatment with dilute mineral acid or base (used for industrial separation and purification).
3. **Addition of Alcohols (Acetal/Ketal Formation):** Carbonyls react with 1 equivalent of alcohol in dry HCl to form hemiacetals, and with 2 equivalents to form acetals. Acetals are stable to bases, oxidants, and reducing agents, acting as universal carbonyl protecting groups.

4. **Nucleophilic Addition-Elimination with Ammonia Derivatives ($NH_2 - G$):**

$$> C = O + H_2N - G \xrightarrow{pH4.5} [> C(OH) - NH - G] \xrightarrow{-H_2O} > C = N - G \tag{5.87}$$

- Hydroxylamine (NH_2OH) → **Oxime** ($> C = N - OH$).
- Hydrazine (NH_2NH_2) → **Hydrazone** ($> C = N - NH_2$).
- 2,4-Dinitrophenylhydrazine (2,4-DNP / Brady's Reagent) → **Yellow/Orange precipitate** (used to detect aldehydes/ketones).
- Semicarbazide ($NH_2 - CO - NHNH_2$) → **Semicarbazone**. (*Trap: Only the hydrazine $-NH_2$ is nucleophilic; the amide $-CONH_2$ nitrogen lone pair is delocalized into the carbonyl!*)

5.7.2 Reactions Involving α-Hydrogen: Aldol, Cannizzaro & Haloform

Step-by-Step Mechanism: Aldol Condensation

Aldehydes or ketones possessing at least one α-hydrogen react in dilute alkali (dil. NaOH):

1. **Step 1: Enolate Generation (Acid-Base Equilibrium):**

$$CH_3 - CHO + OH^- \rightleftharpoons [:\bar{C}H_2 - CHO \longleftrightarrow CH_2 = CH - O^-] + H_2O \tag{5.88}$$

2. **Step 2: Nucleophilic Attack on Second Carbonyl Molecule (RDS):**

$$CH_3 - CHO + :\bar{C}H_2 - CHO \longrightarrow CH_3 - CH(O^-) - CH_2 - CHO \tag{5.89}$$

3. **Step 3: Protonation to Aldol (β-Hydroxyaldehyde):**

$$CH_3 - CH(O^-) - CH_2 - CHO + H_2O \longrightarrow \tag{5.90}$$

$$CH_3 - CH(OH) - CH_2 - CHO + OH^- \tag{5.91}$$

4. **Step 4: Dehydration on Heating ($E1cB$ Mechanism):** Deprotonation of the acidic α-carbon creates an enolate, followed by expulsion of hydroxide ion (OH^-) to yield a conjugated α,β-unsaturated aldehyde:

$$CH_3 - CH(OH) - CH_2 - CHO \xrightarrow{\Delta, -H_2O} CH_3 - CH = CH - CHO \tag{5.92}$$

Step-by-Step Mechanism: Cannizzaro Reaction

Aldehydes **lacking α-hydrogen** ($HCHO, C_6H_5CHO, (CH_3)_3CCHO$) disproportionate in concentrated alkali (50% NaOH):

1. **Step 1: Reversible Addition of Hydroxide:**

$$C_6H_5CHO + OH^- \rightleftharpoons C_6H_5 - CH(OH) - O^- \quad \text{(Monoanion)} \tag{5.93}$$

2. **Step 2: Rate-Determining Hydride Shift (H^-):** The alkoxide reforms the carbonyl double bond, transferring a hydride ion directly to a second aldehyde molecule:

$$C_6H_5 - CH(O^-) - OH + C_6H_5CHO \xrightarrow{\text{slow, RDS}} C_6H_5COOH + C_6H_5CH_2O^- \tag{5.94}$$

3. **Step 3: Fast Acid-Base Proton Exchange:**

$$C_6H_5COOH + C_6H_5CH_2O^- \xrightarrow{\text{fast}} \tag{5.95}$$

$$C_6H_5COO^-Na^+ + C_6H_5CH_2OH \tag{5.96}$$

Cross-Cannizzaro Reaction: When benzaldehyde reacts with formaldehyde in concentrated alkali, formaldehyde is oxidized exclusively to sodium formate (HCOONa) while benzaldehyde is reduced to benzyl alcohol:

$$C_6H_5CHO + HCHO + NaOH \longrightarrow C_6H_5CH_2OH + HCOONa \tag{5.97}$$

Cause: Formaldehyde is sterically uncrowded and far more electrophilic than benzaldehyde, undergoing nucleophilic attack by OH^- *first!*

The Haloform (Iodoform) Reaction Test

Structural Requirement: Reagent I_2/NaOH (or NaOI) oxidizes and cleaves compounds containing:

1. $CH_3 - C = O$ (Methyl ketone group, e.g., acetaldehyde, acetone, acetophenone).
2. $CH_3 - CH(OH) -$ (Methyl carbinol group oxidizable in situ to methyl ketone, e.g., ethanol, propan-2-ol, butan-2-ol).

$$R - CO - CH_3 + 3I_2 + 4NaOH \longrightarrow \tag{5.98}$$

$$CHI_3 \downarrow + R - COONa + 3NaI + 3H_2O \tag{5.99}$$

(CHI_3 is yellow crystalline precipitate, m.p. 119°C). **Classical NEET Distinction Traps:**

- **Ethanol vs Methanol:** Ethanol gives positive iodoform test; methanol gives negative test.
- **Pentan-2-one vs Pentan-3-one:** Pentan-2-one is a methyl ketone (positive); pentan-3-one is not (negative).
- **Acetaldehyde vs Benzaldehyde:** Acetaldehyde gives yellow iodoform; benzaldehyde does not.

5.7.3 Distinction Tests & Reductions

Distinction Tests for Carbonyl Functions

- **Tollens' Silver Mirror Test:** Tollens' reagent ($[Ag(NH_3)_2]^+$) oxidizes aldehydes and α-hydroxy ketones, precipitating silver mirror (Ag $\downarrow$):

$$RCHO + 2[Ag(NH_3)_2]^+ + 3OH^- \rightarrow RCOO^- + 2Ag \downarrow + 4NH_3 + 2H_2O$$

- **Fehling's Test:** Alkaline cupric tartrate complex (Cu^{2+}) is reduced to a **red precipitate of cuprous oxide** ($Cu_2O \downarrow$) by **aliphatic aldehydes only**:

$$R - CHO + 2Cu^{2+} + 5OH^- \longrightarrow R - COO^- + Cu_2O \downarrow (\text{red}) + 3H_2O \quad (5.100)$$

NEET Trap: Benzaldehyde and all aromatic aldehydes FAIL Fehling's test because aromatic resonance stabilizes the carbonyl, making it an insufficiently strong reducing agent to reduce the tartrate-chelated Cu^{2+} ion!

Clemmensen vs Wolff-Kishner Carbonyl Reduction

- **Clemmensen Reduction:** Uses amalgamated zinc and concentrated hydrochloric acid ($Zn - Hg$/conc. HCl):

$$> C = O + 4[H] \xrightarrow{Zn-Hg/\text{conc. HCl}} > CH_2 + H_2O \quad (5.101)$$

Constraint: Cannot be used for substrates containing acid-sensitive groups ($-OH, -OR, -C=C-$), which undergo elimination or addition in strong acid.

- **Wolff-Kishner Reduction:** Carbonyl converted to hydrazone, then heated with KOH in ethylene glycol at $453 - 473$ K:

$$> C = O + NH_2NH_2 \xrightarrow{-H_2O} > C = N - NH_2 \xrightarrow[\text{glycol},\Delta]{KOH} > CH_2 + N_2 \uparrow \quad (5.102)$$

Constraint: Cannot be used for substrates containing base-sensitive groups (alkyl halides, esters), which undergo nucleophilic displacement or saponification.

5.7.4 Carboxylic Acids & Derivatives

Resonance in Carboxylic Acids & The Ortho Effect

1. Resonance of Carboxylate Anion: Deprotonation of carboxylic acid yields the carboxylate anion ($R - COO^-$), which is stabilized by two **completely equivalent** resonance structures sharing negative charge equally between two electronegative oxygen atoms ($C - O$ bond lengths identical at 1.26 Å). This makes carboxylic acids vastly more acidic than phenols.

2. The Ortho Effect in Benzoic Acids: Any substituent at the **ortho position** of benzoic acid—regardless of whether it is electron-withdrawing ($-NO_2$) or electron-releasing ($-CH_3, -OCH_3$)—increases the acidic strength compared to benzoic acid:

$$o\text{-toluic acid } (3.91) > \text{benzoic acid } (4.20) \tag{5.103}$$

$$> m\text{-toluic acid } (4.27) > p\text{-toluic acid } (4.37) \tag{5.104}$$

Physical Cause: Steric crowding by the ortho substituent twists the $-COOH$ group out of the plane of the benzene ring. This **steric inhibition of resonance (SIR)** eliminates cross-conjugation between the ring and the carbonyl, allowing unhindered resonance delocalization of negative charge across the two carboxylate oxygens!

Hell-Volhard-Zelinsky (HVZ) Reaction & Decarboxylation

1. HVZ Reaction (α-Halogenation): Carboxylic acids possessing α-hydrogen react with chlorine or bromine in the presence of catalytic red phosphorus:

$$R - CH_2 - COOH + X_2 \xrightarrow{\text{red P}} R - CH(X) - COX \tag{5.105}$$

$$\xrightarrow{H_2O} R - CH(X) - COOH + HX \tag{5.106}$$

Note: Formic acid ($HCOOH$*) and benzoic acid (*C_6H_5COOH*) fail the HVZ reaction due to absence of α-hydrogens!*

2. Soda-Lime Decarboxylation: Sodium carboxylates heated with soda lime ($NaOH + CaO$ in $3:1$ mass ratio) undergo loss of carbon dioxide:

$$R - COONa + NaOH \xrightarrow[\Delta]{CaO} R - H + Na_2CO_3 \tag{5.107}$$

CaO acts as a desiccant, keeping NaOH dry and raising the reaction fusion temperature.

5.8 Nitrogen-Containing Compounds: Amines & Diazonium Chemistry

5.8.1 Basicity of Amines: Gas Phase vs Aqueous Phase

Thermodynamic Factors Governing Amine Basicity

The basicity of amines is measured by K_b or pK_b ($pK_b = -\log K_b$; lower pK_b indicates stronger base). Basicity reflects the thermodynamic stability of the conjugate ammonium cation (RNH_3^+). **1. Gas Phase (Isolated Molecules):**

In the gas phase, solvent molecules are absent. Basicity is governed strictly by the electron-donating $+I$ inductive effect of alkyl groups, which stabilizes the positive charge on nitrogen:

$$\text{Gas Phase Basicity: } 3^\circ\ R_3N > 2^\circ\ R_2NH > 1^\circ\ RNH_2 > NH_3 \tag{5.108}$$

2. Aqueous Solution (Interplay of Three Opposing Factors): In water, three thermodynamic effects operate concurrently:

1. **Inductive Effect ($+I$):** Stabilizes cation in order $3^\circ > 2^\circ > 1^\circ > NH_3$.
2. **Hydration Energy (H-Bonding):** The conjugate ammonium ion forms hydrogen bonds with water molecules. Greater number of hydrogens attached to positive nitrogen yields higher hydration energy: RNH_3^+ (3 H-bonds) > $R_2NH_2^+$ (2 H-bonds) > R_3NH^+ (1 H-bond). Hydration stabilizes in order: $NH_4^+ > 1^\circ > 2^\circ > 3^\circ$.
3. **Steric Hindrance to Protonation:** Bulky alkyl groups shield the nitrogen lone pair, impeding proton attack in order: $1^\circ > 2^\circ > 3^\circ$.

Authoritative Aqueous Basicity Orders (NEET Core Benchmark)

The combined net effect produces the following experimentally established basicity sequences in aqueous medium:

1. **Methyl-Substituted Amines ($2^\circ > 1^\circ > 3^\circ > NH_3$, Mnemonic: "213"):**

$$(CH_3)_2NH\ (pK_b = 3.27) > CH_3NH_2\ (3.38) > (CH_3)_3N\ (4.22) > NH_3\ (4.75) \tag{5.109}$$

2. **Ethyl-Substituted Amines ($2^\circ > 3^\circ > 1^\circ > NH_3$, Mnemonic: "231"):**

$$(C_2H_5)_2NH\ (3.00) > (C_2H_5)_3N\ (3.25) \tag{5.110}$$

$$> C_2H_5NH_2\ (3.29) > NH_3\ (4.75) \tag{5.111}$$

3. **Arylamines (Aniline):** Aniline ($C_6H_5NH_2$, $pK_b = 9.38$) is vastly weaker than ammonia ($pK_b = 4.75$) because its nitrogen lone pair is delocalized over the ortho and para positions of the aromatic ring (5 canonical forms). The anilinium cation ($C_6H_5NH_3^+$) has only 2 resonance structures, making protonation thermodynamically disfavored!

5.8.2 Gabriel Phthalimide & Hoffmann Bromamide Mechanisms

Gabriel Phthalimide Synthesis: Pure Primary Aliphatic Amines

Gabriel synthesis produces pure 1° aliphatic amines free from secondary or tertiary contamination:

1. **Potassium Salt Formation:** Phthalimide is deprotonated by ethanolic KOH to form potassium phthalimide (the imide hydrogen is acidic due to two flanking carbonyls).
2. **Alkylation (S_N2):** Nucleophilic attack of the phthalimide anion on a primary alkyl halide ($R-X$) yields *N*-alkylphthalimide.
3. **Hydrazinolysis / Alkaline Hydrolysis:** Alkaline hydrolysis with NaOH yields pure primary amine and sodium phthalate:

$$\text{Phthalimide} \xrightarrow{\text{KOH}} \text{Potassium Phthalimide} \xrightarrow{\text{R}-\text{X}} N\text{-Alkylphthalimide} \xrightarrow{\text{NaOH(aq)}} \text{R} - \text{NH}_2 + \text{Sodium Phthalate} \quad (5.112)$$

NEET Trap: Why Aromatic Amines CANNOT be Prepared by Gabriel Synthesis

Aniline ($C_6H_5NH_2$) cannot be prepared by Gabriel phthalimide synthesis because the required aryl halide (C_6H_5-X) does not undergo nucleophilic aromatic substitution (S_N2) with the phthalimide anion under standard conditions due to partial double bond character of the $C-X$ bond and steric/electronic repulsion of the aromatic ring!

Step-by-Step Mechanism: Hoffmann Bromamide Degradation

Primary acid amides react with bromine and aqueous alkali ($Br_2 + 4KOH$) to yield primary amines with **one less carbon atom**:

$$\text{R} - \text{CONH}_2 + \text{Br}_2 + 4\text{KOH} \longrightarrow \text{R} - \text{NH}_2 + \text{K}_2\text{CO}_3 + 2\text{KBr} + 2\text{H}_2\text{O} \quad (5.113)$$

Elementary Mechanistic Steps:

1. **Formation of *N*-Bromoamide:**

$$\text{RCONH}_2 + \text{OH}^- \rightarrow \text{RCON}\bar{\text{H}} \xrightarrow{\text{Br}_2} \text{R} - \text{CONHBr} + \text{Br}^-$$

2. **Formation of Bromoamide Anion:**

$$\text{R} - \text{CONHBr} + \text{OH}^- \rightleftharpoons [\text{R} - \text{CON} - \text{Br}]^- + \text{H}_2\text{O}$$

3. **Rate-Determining Concerted 1,2-Shift (Isocyanate Formation):** Migration of alkyl/aryl group *R* with its electron pair from carbonyl carbon to electron-deficient nitrogen with simultaneous loss of bromide ion:

$$[\text{R} - \text{CO} - \bar{\text{N}} - \text{Br}] \xrightarrow{\text{slow, RDS}} \text{R} - \text{N} = \text{C} = \text{O (Alkyl Isocyanate)} + \text{Br}^- \quad (5.114)$$

Stereochemical Note: The migrating group R migrates with ***complete retention of configuration****! If chiral, optical activity is preserved.*

4. **Alkaline Hydrolysis of Isocyanate:**

$$\text{R} - \text{N} = \text{C} = \text{O} + 2\text{OH}^- \longrightarrow \text{R} - \text{NH}_2 + \text{CO}_3^{2-} \quad (5.115)$$

5.8.3 Distinction Tests for Amines

Hinsberg, Carbylamine & Nitrous Acid Tests

1. **Hinsberg Test (Benzenesulfonyl Chloride $C_6H_5SO_2Cl$):**
   - 1° **Amines:** Forms *N*-alkylbenzenesulfonamide ($C_6H_5SO_2NHR$). Has acidic H on N $\implies$ **Soluble in aqueous** KOH.
   - 2° **Amines:** Forms *N*, *N*-dialkylbenzenesulfonamide ($C_6H_5SO_2NR_2$). Lacks acidic H $\implies$ **Insoluble in aqueous** KOH.
   - 3° **Amines:** No replaceable hydrogen $\implies$ **Does not react**.
2. **Carbylamine Test (Isocyanide Test):** Exclusive to **primary amines** (both aliphatic and aromatic). Heated with chloroform and alcoholic KOH:

$$\mathrm{R-NH_2 + CHCl_3 + 3KOH \xrightarrow{\Delta} R-N\equiv C\ (\text{Carbylamine, foul odor}) + 3KCl + 3H_2O} \tag{5.116}$$

   Secondary and tertiary amines fail the carbylamine test!
3. **Reaction with Nitrous Acid ($HNO_2 = NaNO_2 + HCl$):**
   - 1° **Aliphatic Amines:** Yield unstable aliphatic diazonium salts that decompose quantitatively to alcohols with quantitative evolution of nitrogen gas: $RNH_2 + HNO_2 \rightarrow ROH + N_2 \uparrow + H_2O$.
   - 1° **Aromatic Amines (Aniline):** Form stable benzenediazonium chloride at ice-cold temperature ($0 - 5°C$).
   - 2° **Amines (Aliphatic & Aromatic):** React to form yellow oily *N*-nitrosamines (R_2N-NO), which give **Liebermann's nitroso test**.
   - 3° **Aliphatic Amines:** Form water-soluble trialkylammonium nitrite salts ($[R_3NH]^+NO_2^-$).

5.8.4 Benzenediazonium Chemistry & Synthetic Transformations

Sandmeyer, Gattermann, Balz-Schiemann & Coupling Reactions

Benzenediazonium chloride ($C_6H_5N_2^+Cl^-$) is prepared by diazotization of aniline with $NaNO_2 + 2HCl$ at $273 - 278$ K ($0 - 5°C$). It undergoes two major classes of reactions: **1. Replacement Reactions (Loss of $N_2 \uparrow$):**

- **Sandmeyer Reaction:** $Cu_2Cl_2/HCl \rightarrow C_6H_5Cl$; $Cu_2Br_2/HBr \rightarrow C_6H_5Br$; $CuCN/KCN \rightarrow C_6H_5CN$ (higher yields via cuprous salts).
- **Gattermann Reaction:** Cu powder/$HCl \rightarrow C_6H_5Cl$; Cu powder/$HBr \rightarrow C_6H_5Br$.
- **Balz-Schiemann Reaction:** $HBF_4 \rightarrow C_6H_5N_2^+BF_4^- \xrightarrow{\Delta} C_6H_5F + BF_3 + N_2 \uparrow$ (best method for fluorobenzene).

- **Replacement by Iodine:** Warm with aqueous KI $\rightarrow C_6H_5I + KCl + N_2\uparrow$ (no copper catalyst needed).
- **Deamination (Replacement by H):** Reduction with hypophosphorous acid (H_3PO_2/H_2O) or ethanol (CH_3CH_2OH):

$$C_6H_5N_2^+Cl^- + H_3PO_2 + H_2O \xrightarrow{Cu^+} C_6H_6 + H_3PO_3 + HCl + N_2\uparrow \quad (5.117)$$

2. Azo Coupling Reactions (Retention of Diazo Linkage $-N=N-$): The diazonium ion acts as a weak electrophile, attacking highly activated aromatic rings at the para position:

- **Coupling with Phenol (Alkaline Medium, pH $9-10$):** Forms **orange azo dye** (*p*-hydroxyazobenzene):

$$C_6H_5N_2^+Cl^- + C_6H_5OH \xrightarrow{pH9-10} \quad (5.118)$$

$$C_6H_5 - N=N - C_6H_4 - OH\ (\text{Orange}) + HCl \quad (5.119)$$

- **Coupling with Aniline (Mildly Acidic Medium, pH $4-5$):** Forms **yellow azo dye**:

$$C_6H_5N_2^+Cl^- + C_6H_5NH_2 \xrightarrow{pH4-5} \quad (5.120)$$

$$C_6H_5 - N=N - C_6H_4 - NH_2\ (\text{Yellow}) + HCl \quad (5.121)$$

5.9 Biomolecules (NEET Unit 19)

5.9.1 Carbohydrates: Classification & Stereochemistry

Definition & D/L Stereochemical Nomenclature

Carbohydrates are optically active polyhydroxy aldehydes or polyhydroxy ketones, or substances that produce them upon hydrolysis.

- **D/L Configuration (Rosanoff Convention):** Assigned by comparing the configuration of the highest-numbered chiral carbon (the penultimate asymmetric carbon, C_5 in hexoses) in a vertical Fischer projection to **D-glyceraldehyde**:
  - **D-Sugar:** The $-OH$ group at the lowest asymmetric center projects to the **right**.
  - **L-Sugar:** The $-OH$ group projects to the **left**.
- *Crucial Principle: D and L designate configurational stereochemistry only. They have ZERO correlation to experimental dextrorotation $(+)$ or levorotation $(-)$. D-glucose is dextrorotatory, whereas D-fructose is strongly levorotatory!*

Structure Elucidation of D-(+)-Glucose

Experimental evidence establishing the open-chain structure of D-glucose:

1. **Molecular Formula:** $C_6H_{12}O_6$.
2. **Straight Chain of 6 Carbons:** Prolonged heating with hydroiodic acid and red phosphorus yields **n-hexane**:

$$C_6H_{12}O_6 \xrightarrow[\Delta]{\text{HI/red P}} CH_3 - (CH_2)_4 - CH_3 \ (n\text{-hexane}) \tag{5.122}$$

3. **Carbonyl Group ($> C = O$):** Reacts with hydroxylamine (NH_2OH) to form an oxime, and with HCN to form a cyanohydrin.
4. **Aldehyde Function ($-CHO$):** Oxidation with bromine water (Br_2/H_2O) yields **gluconic acid** ($HOOC - (CHOH)_4 - CH_2OH$).
5. **Five Hydroxyl Groups ($-OH$):** Acetylation with acetic anhydride (Ac_2O) yields stable **glucose penta-acetate**, establishing that all five $-OH$ groups reside on separate carbon atoms.
6. **Primary Alcohol at C_6 ($-CH_2OH$):** Oxidation with concentrated nitric acid (HNO_3) oxidizes both the terminal aldehyde and the primary alcohol to dicarboxylic **saccharic acid (glucaric acid)**:

$$CHO - (CHOH)_4 - CH_2OH \xrightarrow{HNO_3} HOOC - (CHOH)_4 - COOH \tag{5.123}$$

(**Saccharic acid / glucaric acid** confirms primary terminal alcohol).

5.9.2 Cyclic Pyranose Structures, Anomers & Mutarotation

Failure of Open-Chain Structure & The Cyclic Pyranose Ring

1. Evidence Opposing the Simple Open-Chain Aldehyde:

- Glucose does not react with $NaHSO_3$, does not form 2,4-DNP adduct, and does not restore the pink color of Schiff's reagent.
- Glucose penta-acetate does not react with hydroxylamine (proves absence of a free $-CHO$ group).
- Glucose exists in two distinct crystalline stereoisomers (α and β) with different physical properties.

2. Cyclic Hemiacetal Formation: Intramolecular nucleophilic attack of the $C_5 - OH$ group onto the C_1 carbonyl carbon produces a 6-membered **pyranose ring**. The C_1 carbon becomes an asymmetric center known as the **anomeric carbon**:

- **α-D-Glucopyranose:** In Haworth projection, the anomeric $-OH$ at C_1 points **downward** (*trans* to the $-CH_2OH$ group at C_6). Melting point 146°C, specific rotation $[\alpha]_D = +112°$.

- **β-D-Glucopyranose:** The anomeric $-OH$ at C_1 points **upward** (*cis* to the $-CH_2OH$ group at C_6). In the chair conformation, all bulky groups ($-OH$ and $-CH_2OH$) occupy equatorial positions $\Longrightarrow$ thermodynamically more stable. Melting point 150°C, specific rotation $[\alpha]_D = +18.7°$.

3. Mutarotation Mechanism: When pure crystals of either α-D-glucose ($+112°$) or β-D-glucose ($+18.7°$) are dissolved in water, the optical rotation spontaneously drifts over time until reaching an equilibrium value of $+52.7°$:

$$\alpha\text{-D-glucose } (+112°) \underset{\approx 36\%}{\overset{H_2O}{\rightleftharpoons}} \text{Open Form } (0.02\%) \tag{5.124}$$

$$\underset{\approx 64\%}{\overset{H_2O}{\rightleftharpoons}} \beta\text{-D-glucose } (+18.7°) \tag{5.125}$$

This reversible interconversion between anomers via the open-chain aldehyde is called **mutarotation**.

5.9.3 Disaccharides, Glycosidic Linkages & Polysaccharides

Disaccharide Chemistry: Linkages & Reducing Character

Sugar	Monomers	Bond	Free Anomeric C?	Reducing?
Sucrose	α-Glc + β-Fru	$\alpha(\mathbf{1} \rightarrow \mathbf{2})\beta$	None	**Non-Reducing**
Maltose	α-Glc + α-Glc	$\alpha(\mathbf{1} \rightarrow \mathbf{4})$	Yes	**Reducing**
Lactose	β-Gal + β-Glc	$\beta(\mathbf{1} \rightarrow \mathbf{4})$	Yes	**Reducing**

Invert Sugar & The Optical Inversion of Sucrose

Sucrose is **dextrorotatory** ($[\alpha]_D = +66.5°$). Upon acid hydrolysis or enzymatic cleavage by *invertase*, it yields an equimolar mixture of D-(+)-glucose and D-(-)-fructose:

$$C_{12}H_{22}O_{11} \ (+66.5°) + H_2O \xrightarrow{H^+} \text{D-(+)-glucose } (+52.7°) \tag{5.126}$$

$$+ \text{ D-(-)-fructose } (-92.4°) \tag{5.127}$$

Because the levorotation of fructose ($-92.4°$) is numerically far greater than the dextrorotation of glucose ($+52.7°$), the net resulting hydrolyzed mixture is **levorotatory** ($[\alpha]_{net} \approx -39.7°$).
Definition: This reversal in the sign of optical rotation from dextro (+) to levo (−) is called the **inversion of cane sugar**, and the resulting equimolar mixture is termed **invert sugar** (found naturally in honey).

Polysaccharides: Starch, Cellulose & Glycogen

- **Starch (Plant Storage Polysaccharide):** Polymer of α-D-glucose consisting of two distinct structural components:

1. **Amylose ($15-20\%$):** Water-soluble, linear unbranched chain of $200-1000$ α-D-glucose units linked by $\alpha(1\rightarrow 4)$ **glycosidic bonds**. Forms a hollow helical coil that entraps triiodide ions (I_3^-) to produce a characteristic deep blue-black color.
2. **Amylopectin ($80-85\%$):** Water-insoluble, highly branched polymer. Linear chains contain $\alpha(1\rightarrow 4)$ linkages; branching points occur every $24-30$ units via $\alpha(1\rightarrow 6)$ **glycosidic bonds**.

- **Cellulose (Plant Cell Wall Structural Polymer):** Unbranched, linear polymer composed exclusively of β-D-glucose units joined by $\beta(1\rightarrow 4)$ **glycosidic bonds**. Chains lie parallel, held together by massive intermolecular hydrogen bonds to form rigid microfibrils. Humans cannot digest cellulose due to lack of the enzyme cellulase.
- **Glycogen (Animal Starch):** Stored in mammalian liver and muscle. Structurally similar to amylopectin, but possesses far more frequent branching ($\alpha(1\rightarrow 6)$ branches every $8-12$ glucose residues).

5.9.4 Amino Acids, Zwitterions & The Isoelectric Point (pI)

Structure, Stereochemistry & Essential Amino Acids

Amino acids contain both an amino group ($-NH_2$) and a carboxyl group ($-COOH$):

$$R-CH(NH_2)-COOH \quad (\alpha\text{-amino acid}) \tag{5.128}$$

- **Chirality Exception:** All 20 standard proteinogenic amino acids are chiral and optically active **except Glycine** (H_2N-CH_2-COOH), where $R=H$, which is achiral and optically inactive!
- **L-Configuration:** All natural amino acids in proteins belong to the L-series (the $-NH_2$ group resides on the left in Fischer projections).
- **Essential Amino Acids (Cannot be synthesized in human body; must be supplied in diet):** Valine, Leucine, Isoleucine, Phenylalanine, Tryptophan, Threonine, Methionine, Lysine (plus Arginine and Histidine, semi-essential in infants). *(Mnemonic: PVT TIM HaLL).*

Zwitterion Structure & Formulation of Isoelectric Point (pI)

In aqueous solution, the acidic carboxyl group transfers a proton to the basic amino group, generating a dipolar internal salt known as a **zwitterion**:

$$R-CH(NH_2)-COOH \rightleftharpoons R-CH(NH_3^+)-COO^- \tag{5.129}$$

(Dipolar Zwitterion; net charge $=0$). **1. Behavior in Varying pH Media:**

- In acidic solution (pH $< pI$): Cation $\mathrm{R-CH(NH_3^+)-COOH}$ migrates toward **cathode**.
- In basic solution (pH $> pI$): Anion $\mathrm{R-CH(NH_2)-COO^-}$ migrates toward **anode**.

2. Isoelectric Point (pI): The exact pH at which the amino acid carries no net electrical charge and does not migrate in an electric field:

$$pI = \frac{pK_{a1}(\alpha\text{–COOH}) + pK_{a2}(\alpha\text{–}\mathrm{NH_3^+})}{2} \quad \text{(for neutral amino acids)} \quad (5.130)$$

Crucial Property: At its isoelectric point (pI), an amino acid exhibits ***minimum aqueous solubility****, permitting selective precipitation and separation!*

5.9.5 Peptide Bond Resonance & Structural Hierarchy of Proteins

The Peptide Bond (-CO – NH–) & Planarity

A peptide bond is an amide linkage formed by condensation of the –COOH group of one amino acid with the $-\mathrm{NH_2}$ group of another with loss of water:

$$\mathrm{R-CH(NH_2)-COOH + H_2N-CH(R')-COOH} \xrightarrow{-\mathrm{H_2O}} \quad (5.131)$$

$$\mathrm{R-CH(NH_2)-\mathbf{CO}-\mathbf{NH}-CH(R')-COOH} \quad (5.132)$$

Resonance Stabilization of the Peptide Bond: Delocalization of the nitrogen lone pair into the carbonyl group confers ≈ 40% **partial double bond character** to the C – N bond:

$$\mathrm{-C(=O)-NH-} \longleftrightarrow \mathrm{-C(O^-)=\overset{+}{N}H-} \quad (5.133)$$

Consequences:

1. The C – N bond length (1.32 Å) is substantially shorter than a normal C – N single bond (1.47 Å).
2. Free rotation about the C – N bond is restricted; the six atoms of the peptide group ($\mathrm{C_\alpha}$–CO – NH – $\mathrm{C_\alpha}$) are strictly coplanar.
3. The *trans*-conformation is overwhelmingly favored to minimize steric clash between adjacent side chains (R).

The Four Structural Levels of Protein Organization

1. **Primary Structure (1°):** The linear sequence of amino acids linked covalently by peptide bonds. Any alteration in sequence alters biological function (e.g., in sickle-cell anemia, glutamate at position 6 of the β-chain of hemoglobin is mutated to valine).

2. **Secondary Structure (2°):** Regular folding of the polypeptide backbone stabilized exclusively by **hydrogen bonds** between $>C=O$ and $-NH-$ groups:
   - **α-Helix:** Polypeptide coils into a right-handed screw. Stabilized by intramolecular hydrogen bonding between the $>C=O$ of residue n and the $-NH-$ of residue $(n+4)$. Each turn contains 3.6 amino acids with pitch 5.4 Å. Proline acts as a helix breaker.
   - **β-Pleated Sheet:** Polypeptide chains are stretched side-by-side in zigzag sheets, held by intermolecular hydrogen bonds. Can be parallel or antiparallel (e.g., silk fibroin).
3. **Tertiary Structure (3°):** Overall 3D spatial folding of the polypeptide chain into compact shapes:
   - **Fibrous Proteins:** Linear, insoluble in water, structural role (keratin in hair/nails, collagen in tendons, myosin in muscle).
   - **Globular Proteins:** Spherical, water-soluble, metabolic role (enzymes, hemoglobin, albumin).
   - **Stabilizing Forces:** H-bonds, **disulfide bonds ($-S-S-$ in cysteine, covalent)**, ionic/salt bridges ($-COO^- \cdots NH_3^+$), hydrophobic and van der Waals forces.
4. **Quaternary Structure (4°):** Spatial arrangement of two or more independent tertiary polypeptide subunits into an oligomeric protein (e.g., adult Hemoglobin is a tetramer consisting of 2α and 2β subunits).

Denaturation of Proteins: What Survives and What Is Destroyed?

When a native protein is exposed to physical changes (heating, radiation) or chemical changes (extreme pH, urea, organic solvents, heavy metals):

- **Destroyed:** Hydrogen bonds, salt bridges, and hydrophobic interactions rupture. The delicate $2^\circ, 3^\circ$, and 4° spatial architectures unfold. Globular proteins precipitate, and biological activity is irreversibly lost.
- **Retained Intact: The primary structure (1°) remains 100% completely intact!** Covalent peptide bonds ($-CO-NH-$) are not cleaved under normal denaturation conditions.
- **Classic NCERT Examples:** Coagulation of egg white (albumin) upon boiling; curdling of milk caused by lactic acid produced by bacteria.

5.9.6 Nucleic Acids: DNA & RNA Architecture

Chemical Composition: Sugars, Bases & Nucleotides

Nucleic acids are unbranched biopolymers of nucleotides:

1. **Pentose Sugar:** β-D-ribose in RNA; β-D-2-deoxyribose in DNA (lacks oxygen atom at C_2' position).

2. **Nitrogenous Bases:**
   - **Purines (Bicyclic 9-membered):** Adenine (A) and Guanine (G) in both DNA and RNA.
   - **Pyrimidines (Monocyclic 6-membered):** Cytosine (C) and Thymine (T, 5-methyluracil) in DNA; Cytosine (C) and Uracil (U) in RNA.
3. **Nucleoside vs Nucleotide:**
   - Nucleoside = Pentose Sugar + Base (joined by N-glycosidic linkage between C_1' of sugar and N_9 of purine or N_1 of pyrimidine).
   - Nucleotide = Nucleoside + Phosphate (esterified at C_5' of pentose sugar).
4. **Phosphodiester Linkage:** Nucleotides link via $3', 5'$**-phosphodiester bonds** between $3' - \text{OH}$ and $5' - \text{OH}$ of adjacent sugars ($5' \to 3'$ polarity).

Watson-Crick Double Helix Model & Chargaff's Rules

James Watson and Francis Crick (1953) established the B-DNA double helix structure:

1. **Antiparallel Strands:** Two polynucleotide chains wound around a central axis in a right-handed helix. One strand runs $5' \to 3'$, and the complementary strand runs $3' \to 5'$.
2. **Complementary Base Pairing (Hydrogen Bonds):**

$$\mathbf{A = T} \quad \text{(Adenine pairs with Thymine via 2 Hydrogen Bonds)} \tag{5.134}$$

$$\mathbf{G \equiv C} \quad \text{(Guanine pairs with Cytosine via 3 Hydrogen Bonds)} \tag{5.135}$$

   Because $G \equiv C$ pairs have 3 hydrogen bonds, DNA with high GC content has a higher melting temperature (T_m).
3. **Geometric Dimensions of B-DNA:**
   - Helix Diameter: 2.0 nm (20 Å).
   - Pitch of Helix (one complete 360° turn): 3.4 nm (34 Å).
   - Base Pairs per Turn: 10 base pairs.
   - Distance between Adjacent Base Pairs: 0.34 nm (3.4 Å).
4. **Chargaff's Rules (Applies Exclusively to Double-Stranded DNA):**

$$[\text{A}] = [\text{T}], \quad [\text{G}] = [\text{C}] \implies [\text{A} + \text{G}] = [\text{T} + \text{C}] \quad \left(\frac{\text{Total Purines}}{\text{Total Pyrimidines}} = 1.0 \right) \tag{5.136}$$

   Note: The ratio $\frac{\text{A}+\text{T}}{\text{G}+\text{C}}$ *is not necessarily equal to 1; it is variable and species-specific.*

5.9.7 Vitamins & Hormones

Comprehensive Vitamin Classification & Deficiency Pathology

Vitamin	Chemical Name	Class	Deficiency Disease / Symptoms
Vit A	Retinol	Fat-Sol.	Night blindness (Nyctalopia), Xerophthalmia
Vit B_1	Thiamine	Water-Sol.	**Beriberi** (muscular weakness, neuropathy)
Vit B_2	Riboflavin	Water-Sol.	Cheilosis (fissuring at corners of mouth)
Vit B_3	Niacin	Water-Sol.	**Pellagra** (dermatitis, diarrhea, dementia)
Vit B_6	Pyridoxine	Water-Sol.	Convulsions, peripheral neuritis
Vit B_{12}	Cyanocobalamin	Water-Sol.	**Pernicious Anemia** (stored in liver)
Vit C	Ascorbic Acid	Water-Sol.	**Scurvy** (bleeding gums; destroyed by heat)
Vit D	Cholecalciferol	Fat-Sol.	**Rickets** in children, **Osteomalacia**
Vit E	Tocopherols	Fat-Sol.	Muscular weakness, increased RBC fragility
Vit K	Phylloquinone	Fat-Sol.	**Defective Clotting** (prolonged prothrombin)

Endocrine Hormones: Chemical Classification

Hormones are intercellular chemical messengers secreted directly into the bloodstream by ductless endocrine glands:

- **Steroid Hormones (Lipid-Soluble, Cholesterol Derivatives):** Secreted by adrenal cortex and gonads: Testosterone, Estrogens (Estradiol), Progesterone, Cortisol, Aldosterone.
- **Peptide / Protein Hormones (Water-Soluble):**
  - **Insulin (51 amino acids):** Secreted by β-cells of islets of Langerhans in pancreas; promotes glucose uptake and conversion to glycogen; deficiency causes *Diabetes mellitus*.
  - **Glucagon:** Secreted by α-cells of pancreas; stimulates glycogenolysis to elevate blood glucose.
  - **Oxytocin & Vasopressin (ADH):** Nonapeptides released by posterior pituitary gland.
- **Amine Hormones (Amino Acid Derivatives):**
  - **Epinephrine (Adrenaline) & Norepinephrine:** Derived from tyrosine; secreted by adrenal medulla ("fight or flight" hormones).
  - **Thyroxine (T_4):** Iodinated derivative of tyrosine produced by thyroid gland; regulates basal metabolic rate (BMR); deficiency causes goiter and hypothyroidism.

5.10 Master Distinction Matrix & High-Yield Synthesis Archetypes

5.10.1 Exhaustive Chemical Distinction Matrix

High-Yield Laboratory Chemical Tests for Functional Groups

Test Name	Reagents	Target Species	Positive Observation / Chemistry
Lucas Test	Anhyd. $ZnCl_2$ + conc. HCl	1°, 2°, 3° Alcohols	3°: Immediate turbidity; 2°: in 5 min; 1°: on boiling only
Neutral $FeCl_3$	Neutral $FeCl_3$ solution	Phenols	Violet / purple coloration (ferric phenoxide complex)
Tollens' Test	Ammoniacal $AgNO_3$	All Aldehydes	Shining silver mirror (Ag ↓) on inner tube walls
Fehling's Test	Fehling A ($CuSO_4$) + Fehling B (alkaline tartrate)	Aliphatic Aldehydes	Red precipitate of cuprous oxide (Cu_2O ↓); aromatic aldehydes fail
Iodoform Test	I_2 + NaOH (or NaOI)	$CH_3C=O$ or $CH_3CH(OH)-$	Yellow crystalline precipitate of CHI_3 ↓ (m.p. 119°C)
2,4-DNP Test	2,4-Dinitrophenylhydrazine	Aldehydes & Ketones	Yellow, orange, or red crystalline precipitate
Carbylamine Test	$CHCl_3$ + alc. KOH, Δ	1° Amines	Extremely obnoxious, foul odor of carbylamine ($R-NC$)
Hinsberg Test	$C_6H_5SO_2Cl$ + KOH	1°, 2°, 3° Amines	1°: sol. in KOH; 2°: insol. in KOH; 3°: no reaction
Azo Dye Test	1. $NaNO_2$/HCl ($0-5$°C), 2. alkaline β-naphthol	1° Aromatic Amines	Brilliant scarlet / orange red azo dye
Bicarbonate Test	Sat. $NaHCO_3$ solution	Carboxylic Acids	Brisk effervescence of CO_2 ↑ gas (phenols fail, except picric acid)

5.10.2 Solved Master Synthesis Archetypes

Archetype 1: Multi-Step Synthesis with Stereochemical Retention

Problem: Convert Benzene into pure *p*-bromoaniline. Identify why direct bromination of aniline fails, and formulate the complete protected sequence.
Step-by-Step Solution:

1. **Pitfall of Direct Bromination:** Aniline contains a powerfully activating $-NH_2$ group ($+M$). Bromination with Br_2/H_2O yields instantaneous white precipitate of 2,4,6-tribromoaniline; monosubstitution cannot be controlled.
2. **Step 1 (Nitration):** Benzene $\xrightarrow[323\text{–}333\ K]{\text{conc. } HNO_3+H_2SO_4}$ $C_6H_5NO_2$ (Nitroben-

zene).

3. **Step 2 (Reduction):** $C_6H_5NO_2 \xrightarrow{\text{Sn/HCl or Fe/HCl}} C_6H_5NH_2$ (Aniline).
4. **Step 3 (Protection via Acetylation):** $C_6H_5NH_2 + (CH_3CO)_2O \xrightarrow{\text{pyridine}} C_6H_5NHCOCH_3$ (Acetanilide). The lone pair on nitrogen is delocalized into the acetyl carbonyl, moderating ring activation.
5. **Step 4 (Regioselective Bromination):** $C_6H_5NHCOCH_3 + Br_2 \xrightarrow{\text{AcOH}}$ *p*-bromoacetanilide (para product due to steric hindrance).
6. **Step 5 (Deprotection):**

$$p\text{-bromoacetanilide} \xrightarrow{H_3O^+/\Delta} p\text{-bromoaniline} + CH_3COOH$$

Archetype 2: Carbon-Carbon Chain Lengthening & Functional Transformation

Problem: Formulate the synthetic conversion of Ethanol into Propan-1-amine and Propan-2-ol. **Step-by-Step Solution:**

1. **Conversion of Ethanol into Propan-1-amine (One Carbon Ascent):**

$$CH_3CH_2OH \xrightarrow{SOCl_2} CH_3CH_2Cl \tag{5.137}$$

$$CH_3CH_2Cl \xrightarrow{KCN} CH_3CH_2CN \quad \text{(One-carbon ascent)} \tag{5.138}$$

$$CH_3CH_2CN \xrightarrow{LiAlH_4} CH_3CH_2CH_2NH_2 \quad \text{(Propan-1-amine)} \tag{5.139}$$

2. **Conversion of Ethanol into Propan-2-ol (Grignard Pathway):**

$$CH_3CH_2OH \xrightarrow{PCC} CH_3CHO \tag{5.140}$$

$$CH_3CHO + CH_3MgBr \xrightarrow{\text{ether}} [CH_3CH(OMgBr)CH_3] \tag{5.141}$$

$$\xrightarrow{H_3O^+} CH_3CH(OH)CH_3 \text{ (Propan-2-ol)} \tag{5.142}$$

Master Summary: Key Takeaways for NEET UG 2027 Organic Chemistry

- **Always inspect carbocation intermediates** for possible 1,2-hydride, 1,2-methyl, or ring-expansion rearrangements before forming products in S_N1, $E1$, and electrophilic addition reactions.
- **In Williamson ether synthesis**, always employ the alkoxide of the bulky tertiary component and a primary alkyl halide to avoid elimination.
- **In biomolecules**, remember that sucrose is non-reducing and lacks mutarotation; all monosaccharides, maltose, and lactose are reducing.
- **In DNA double helix**, $A = T$ (2 H-bonds) and $G \equiv C$ (3 H-bonds); Chargaff's rules hold strictly for double-stranded DNA.

- **In aromatic synthesis**, protect aniline by acetylation with acetic anhydride before nitration to avoid the anomalous 47% meta-product.

PART V

Organic Reaction Mechanisms Handbook

CHAPTER 6

Organic Reaction Mechanisms Handbook: Step-by-Step Arrow-Pushing Pathways

To answer multi-step synthesis and Assertion-Reasoning questions with confidence, students must master the underlying reaction mechanisms. This chapter presents a rigorous, curved-arrow mechanistic breakdown of electrophilic aromatic substitution, nucleophilic aliphatic substitution (S_N1 vs S_N2), elimination pathways ($E1$ vs $E2$), carbonyl additions and condensations, and carboxylic acid derivative substitutions.

6.1 Electrophilic Aromatic Substitution (S_EAr / EAS)

Fundamental 3-Step EAS Pathway & The Arenium Ion (σ-Complex)

Electrophilic aromatic substitution is the hallmark reaction of electron-rich aromatic systems. The driving force is the preservation of aromatic resonance energy (≈ 150.6 kJ/mol in benzene). The reaction proceeds through three discrete elementary steps:

1. **Step 1: Generation of the Active Electrophile (E^+):** A Lewis acid or strong Brønsted acid polarizes or cleaves the reagent precursor to produce a potent cationic or polarized electrophilic species (E^+).
2. **Step 2: Formation of the Arenium Ion (σ-Complex / Wheland Intermediate) [RDS]:** The aromatic π-electron cloud attacks E^+. The attacked carbon rehybridizes from planar sp^2 to tetrahedral sp^3, temporarily disrupting aromaticity. The resulting carbocation is resonance-stabilized across the ortho and para positions:

$$\mathrm{C_6H_6} + E^+ \xrightarrow{\text{slow, RDS}} [\mathrm{C_6H_6E}]^+ \tag{6.1}$$

   (**Arenium ion / Wheland intermediate**: charge delocalized over 5 carbons).
3. **Step 3: Fast Deprotonation & Rearomatization:** The conjugate base abstracts the proton from the sp^3 carbon. The electron pair returns to the ring sextet, restoring full aromaticity (sp^2 rehybridization).

Kinetics, Energy Profile & Primary Kinetic Isotope Effect

1. Rate Law: Overall rate is governed by the second step (formation of the arenium ion):

$$\text{Rate} = k[\text{Ar}-\text{H}][E^+] \tag{6.2}$$

(Second-order kinetics: first order in benzene, first order in electrophile). **2. Kinetic Isotope Effect (k_H/k_D):**

- **Nitration, Chlorination, Bromination, Alkylation:** $k_H/k_D \approx 1.0$. The $C-H$ bond cleavage occurs in Step 3 (fast step) after the rate-determining transition state. Hence, C_6H_6, C_6D_6, and C_6T_6 react at identical rates.
- **Sulfonation & Iodination:** $k_H/k_D > 1.0$ ($k_H/k_D \approx 2-3$ for sulfonation; ≈ 4 for iodination). Here, Step 3 or intermediate decomposition has a barrier comparable to Step 2; Step 3 is reversible and partially rate-limiting.

6.1.1 Nitration of Aromatic Rings

Nitration Mechanism & Nitronium Ion Generation

Nitration is carried out using a "nitrating mixture" consisting of concentrated nitric acid (HNO_3) and concentrated sulfuric acid (H_2SO_4) at $323-333$ K:

1. **Generation of Nitronium Ion (NO_2^+):** Sulfuric acid ($pK_a \approx -3$) is a stronger acid than nitric acid ($pK_a \approx -1.4$). Nitric acid acts as a Brønsted-Lowry **base**, accepting a proton on its hydroxyl oxygen:

$$\text{HNO}_3 + \text{H}_2\text{SO}_4 \rightleftharpoons \text{H}_2\overset{+}{\text{O}} - \text{NO}_2 + \text{HSO}_4^- \tag{6.3}$$

$$\text{H}_2\overset{+}{\text{O}} - \text{NO}_2 \rightleftharpoons \text{NO}_2^+ + \text{H}_2\text{O} \tag{6.4}$$

$$\text{H}_2\text{O} + \text{H}_2\text{SO}_4 \rightleftharpoons \text{H}_3\text{O}^+ + \text{HSO}_4^- \tag{6.5}$$

$$\text{Net: } \text{HNO}_3 + 2\text{H}_2\text{SO}_4 \rightleftharpoons \text{NO}_2^+ + \text{H}_3\text{O}^+ + 2\text{HSO}_4^- \tag{6.6}$$

2. **Electrophilic Attack (RDS):** $C_6H_6 + NO_2^+ \xrightarrow{\text{slow}} [C_6H_6NO_2]^+$ (σ-complex).
3. **Rearomatization:** $[C_6H_6NO_2]^+ + HSO_4^- \xrightarrow{\text{fast}} C_6H_5NO_2 + H_2SO_4$.

NCERT Critical Extraction: Nitration of Aniline & Protecting Groups

Direct nitration of aniline with conc. HNO_3/H_2SO_4 gives an anomalous product distribution:

$$\text{Aniline} \xrightarrow[288\text{ K}]{\text{HNO}_3/\text{H}_2\text{SO}_4} p\text{-nitroaniline (51\%)} \tag{6.7}$$

$$+ \textbf{m-nitroaniline (47\%)} + o\text{-nitroaniline (2\%)} \tag{6.8}$$

Mechanistic Cause: In strongly acidic medium, the highly activating $-NH_2$ group ($+M$) is protonated to form the anilinium ion ($-NH_3^+$), which is a powerful $-I$ deactivating group that directs meta. **Synthetic Solution (Acetylation Protection):** Aniline is first protected with acetic anhydride (Ac_2O) in pyridine to form acetanilide ($Ph-NHCOCH_3$). The lone pair on nitrogen is delocalized into the carbonyl group by resonance, reducing its basicity (preventing protonation) and modulating activating strength. Nitration then gives exclusively **para-nitroacetanilide**, followed by acid/base hydrolysis to yield pure **para-nitroaniline**.

6.1.2 Halogenation of Benzene

Halogenation Pathways & Lewis Acid Catalysis

- **Chlorination / Bromination:** Benzene reacts with Cl_2 or Br_2 in the presence of anhydrous Lewis acids ($FeCl_3, AlCl_3, FeBr_3$):

$$Cl-Cl+FeCl_3 \rightleftharpoons [Cl^{\delta+}\cdots Cl\cdots FeCl_3{}^{\delta-}] \rightarrow Cl^+ + [FeCl_4]^- \tag{6.9}$$

  Electrophilic Cl^+ attacks the ring to form the chlorocyclohexadienyl cation, followed by $[FeCl_4]^-$ deprotonation yielding chlorobenzene and regenerating $FeCl_3 + HCl$.
- **Fluorination:** Balz-Schiemann reaction:

$$ArNH_2 \xrightarrow{HNO_2} ArN_2^+ \xrightarrow{HBF_4} ArN_2^+BF_4^- \xrightarrow{\Delta} ArF + BF_3 + N_2\uparrow$$

- **Iodination:** Reversible due to reducing agent HI: $C_6H_6+I_2 \rightleftharpoons C_6H_5I+HI$. Driven forward by oxidizing agent (HNO_3, HIO_3): $5HI+HIO_3 \rightarrow 3I_2+3H_2O$.

6.1.3 Friedel-Crafts Alkylation & Acylation

Friedel-Crafts Alkylation: Mechanism & Classical Carbocation Rearrangements

1. Mechanism:

$$R-Cl+AlCl_3 \rightleftharpoons R^+[AlCl_4]^- \quad \text{(Carbocation intermediate generated)} \tag{6.10}$$

$$C_6H_6 + R^+ \xrightarrow{\text{slow, RDS}} [C_6H_6R]^+ \xrightarrow{[AlCl_4]^-} C_6H_5R + HCl + AlCl_3 \tag{6.11}$$

2. Carbocation Rearrangement Trap: Whenever 1° alkyl halides with ≥ 3 carbons are used, the primary carbocation rearranges via 1,2-hydride or

1,2-methide shift to a more stable 2° or 3° carbocation prior to electrophilic attack:

$$C_6H_6 + CH_3CH_2CH_2Cl \xrightarrow{AlCl_3} \textbf{Cumene}\ (65\text{–}70\%) \tag{6.12}$$

$$+\ n\text{-propylbenzene}\ (30\text{–}35\%) \tag{6.13}$$

Mechanism of shift: $CH_3 - CH_2 - \overset{+}{C}H_2 \xrightarrow{\text{1,2-hydride shift}} CH_3 - \overset{+}{C}H - CH_3$ (2° carbocation is stabilized by 6 α-H vs 2 α-H).

Four Cardinal NEET Limitations of Friedel-Crafts Alkylation

1. **Polyalkylation Problem:** The introduced alkyl group ($-$R) is electron-donating ($+I, +H$), making the monoalkylated product more reactive toward further EAS than benzene itself. Hence, ortho/para dialkyl and polyalkyl benzenes form readily.
2. **Carbocation Rearrangement:** Straight unbranched alkyl benzenes cannot be synthesized cleanly in high yield by direct alkylation.
3. **Deactivated Ring Immunity:** Rings carrying moderate to strong electron-withdrawing groups ($-M/-I$ like $-NO_2, -CF_3, -SO_3H, -COOH, -COOR, -CN, -CHO$) do **not** undergo Friedel-Crafts reactions. Nitrobenzene is routinely used as an inert solvent for Friedel-Crafts reactions!
4. **Aniline / Phenol Lewis Acid Failure:** Aniline and phenol do not undergo Friedel-Crafts alkylation/acylation cleanly because the Lewis acid ($AlCl_3$) forms a strong coordinate complex with the lone pair on $-NH_2$ or $-OH$:

$$C_6H_5NH_2 + AlCl_3 \rightarrow C_6H_5 - \overset{+}{N}H_2 - \overset{-}{A}lCl_3 \tag{6.14}$$

This converts the powerfully activating $-NH_2$ into a strongly deactivating $-I$ group ($-\overset{+}{N}H_2 - AlCl_3^-$), completely terminating EAS.

Friedel-Crafts Acylation: Acylium Ion & Polyalkylation Remedy

Acylation avoids both the rearrangement and polyalkylation defects of alkylation:

1. **Generation of the Acylium Ion:**

$$R - CO - Cl + AlCl_3 \rightleftharpoons [R - \overset{+}{C} = O \longleftrightarrow R - C \equiv \overset{+}{O}] + [AlCl_4]^- \tag{6.15}$$

The acylium ion is exceptionally stable because all atoms possess a complete octet in the $R - C \equiv O^+$ canonical form. Consequently, it **never undergoes carbocation rearrangement**!

2. **Attack & Single Substitution:** $C_6H_6 + RCO^+ \xrightarrow{RDS} [C_6H_6COR]^+ \rightarrow C_6H_5COR + HCl$.

3. **No Polyacylation:** The introduced acyl group ($-\text{COR}$) is strongly electron-withdrawing ($-M, -I$), deactivating the product ring against further electrophilic attack.
4. **Synthesis of Unbranched Alkylbenzenes:** The acyl product is subsequently reduced to an unbranched alkyl chain via either:
   - **Clemmensen:** $\text{ArCOR} \xrightarrow{\text{Zn-Hg/HCl}} \text{ArCH}_2\text{R}$ (acid-stable).
   - **Wolff-Kishner:** $\text{ArCOR} \xrightarrow[\text{KOH},\Delta]{\text{NH}_2\text{NH}_2} \text{ArCH}_2\text{R} + \text{N}_2\uparrow$ (base-stable).

6.1.4 Orientation & Reactivity: Arenium Ion Resonance Analysis

Arenium Canonical Analysis: Ortho/Para vs Meta Directing Effects

The directing effect is governed entirely by the relative thermodynamic stability of the arenium intermediate for ortho, meta, and para electrophilic attack:

- **Ortho/Para Directors** ($+M > -I$, e.g., $-\text{OH}, -\text{OR}, -\text{NH}_2$; $+I/+H$, e.g., $-\text{CH}_3$): Attack at ortho/para places positive charge on the substituent carbon; a $+M$ donor forms a stable fourth resonance structure with complete octets.
- **Meta Directors** ($-M/-I$, e.g., $-\text{NO}_2, -\text{CN}, -\text{CHO}, -\text{COOH}$): Ortho/para attack places positive charge adjacent to positively polarized substituent carbon, causing severe repulsion. Meta attack avoids positive charge on the ipso carbon.

The Halogen Paradox in EAS: Deactivating yet Ortho/Para Directing

Halogens ($-\text{F}, -\text{Cl}, -\text{Br}, -\text{I}$) are unique in organic chemistry:

- **Reactivity (Deactivating):** The inductive effect ($-I$) of the highly electronegative halogen strongly withdraws σ-electron density from the entire π-system, reducing overall ring electron density. Hence, halobenzenes react **slower** than benzene in all EAS reactions.
- **Orientation (Ortho/Para):** When attack occurs at the ortho or para position, the halogen lone pair can be donated via resonance ($+M$) to form a fourth canonical contributor ($= \overset{+}{\text{X}}$) with complete valence octets. This stabilization is absent for meta attack.
- **Conclusion:** Inductive effect controls reactivity (deactivating), while resonance effect controls orientation (o/p-directing).

6.2 Nucleophilic Aliphatic Substitution (S_N1 vs S_N2)

6.2.1 The S_N1 Reaction Pathway

Two-Step Unimolecular Substitution (S_N1)

The S_N1 mechanism operates through a stepwise dissociation-addition mechanism:

1. **Step 1: Heterolysis to form Carbocation Intermediate [RDS]:**

$$\mathrm{R-X} \xrightarrow[\text{slow, RDS}]{k_1} \mathrm{R^+ + X^-} \tag{6.16}$$

   (Planar trigonal sp^2 carbocation intermediate). The rate depends solely on the concentration of the alkyl halide.
2. **Step 2: Nucleophilic Attack [Fast]:**

$$\mathrm{R^+ + Nu^-} \xrightarrow[\text{fast}]{k_2} \mathrm{R-Nu} \tag{6.17}$$

   The nucleophile can attack either the front face or back face of the unhindered planar carbocation.

S_N1 Governing Equations, Stereochemistry & Solvation

Rate Law: Rate $= k[\mathrm{R-X}]$ (First Order, Unimolecular).

Substrates: Benzyl $\approx$ Allyl $\approx 3^\circ > 2^\circ \gg 1^\circ > \mathrm{CH_3X}$.

Solvent Acceleration: Favored by polar protic solvents ($\mathrm{H_2O}$, MeOH, EtOH, AcOH) with high dielectric constant (ϵ), which solvate and stabilize both the carbocation and leaving group anion via hydrogen bonding.

Stereochemical Outcome (Racemization with Partial Inversion): The leaving group $\mathrm{X^-}$ forms an **intimate ion pair** ($\mathrm{R^+ \cdots X^-}$) shielding the front face. Backside attack occurs preferentially, leading to partial racemization with predominant inversion (50–70% inversion +30–50% retention).

Carbocation Rearrangements in S_N1

Because a free carbocation is formed in Step 1, S_N1 reactions are prone to Wagner-Meerwein rearrangements:

- **1,2-Hydride Shift:** 2-chloro-3-methylbutane solvolyzed in ethanol yields predominantly 2-ethoxy-2-methylbutane (3° product via hydride shift from 2° carbocation).

- **1,2-Methyl Shift:** Neopentyl bromide solvolyzed in water yields 2-methylbutan-2-ol (3° carbocation via methide shift).
- **Ring Expansion:** Cyclobutylmethyl halide yields cyclopentanol derivatives to relieve 90° ring angle strain.

6.2.2 The S_N2 Reaction Pathway

One-Step Bimolecular Concerted Substitution (S_N2)

The S_N2 mechanism occurs in a single, concerted step without any reactive intermediate:

1. The nucleophile approaches the electrophilic carbon from the **backside** (180° relative to the C − X leaving bond).
2. The reaction passes through a single, pentacoordinate, trigonal bipyramidal transition state:

$$\mathrm{Nu^- + R - CH_2 - X} \xrightarrow{\text{slow}} \left[\mathrm{Nu}^{\delta-} \cdots \mathrm{CH_2R} \cdots \mathrm{X}^{\delta-}\right]^{\ddagger} \tag{6.18}$$

$$\xrightarrow{\text{fast}} \mathrm{Nu - CH_2 - R + X^-} \tag{6.19}$$

The central carbon is approximately sp^2 hybridized with three equatorial bonds, while the nucleophile and leaving group share a collinear p-orbital axis at 180°.

S_N2 Rate Law, Steric Effects & Walden Inversion

Rate Law: $\mathrm{Rate} = k[\mathrm{R - X}][\mathrm{Nu^-}]$ (Order = 2)

Reactivity: $\mathrm{CH_3X} > 1^\circ > 2^\circ \ggg 3^\circ$

Relative Rates: $\mathrm{CH_3Br}$ (1200) > EtBr (40) > i-PrBr (1) > t-BuBr (10^{-3})

Stereochemical Outcome (100% Walden Inversion): Backside attack forces the three equatorial substituents to flip through a planar transition state, resembling an umbrella blown inside out by a gust of wind. This results in complete inversion of configuration at the stereocenter ($R \rightarrow S$ or $S \rightarrow R$, depending on Cahn-Ingold-Prelog priorities).

The Neopentyl Halide S_N2 Trap & Solvent Effects

- **Neopentyl Halide Inertness:** Neopentyl bromide, $(\mathrm{CH_3})_3\mathrm{C - CH_2 - Br}$, is a primary alkyl halide (1°). However, its S_N2 rate is 10^{-5} times slower than ethyl bromide! The three bulky methyl groups on the β-carbon completely

block backside access to the electrophilic α-carbon.

- **Polar Aprotic Solvents Accelerate S_N2:** Solvents such as DMSO (dimethyl sulfoxide), DMF (N,N-dimethylformamide), acetone, and acetonitrile coordinate strongly to cations via their electronegative oxygen atoms, but cannot hydrogen-bond to nucleophilic anions. The anions remain "naked" and desolvated, boosting their nucleophilicity by up to 10^5-fold.

6.2.3 Master Diagnostic: S_N1 vs S_N2 vs S_Ni

Parameter	S_N1 Mechanism	S_N2 Mechanism
Number of Steps	Two steps (carbocation intermediate).	One concerted step (pentacoordinate transition state).
Kinetic Rate Law	Rate $= k[\mathrm{R-X}]$ (1st order, unimolecular).	Rate $= k[\mathrm{R-X}][\mathrm{Nu^-}]$ (2nd order, bimolecular).
Substrate Preference	$3^\circ > 2^\circ \gg 1^\circ$; benzylic $\approx$ allylic $> 3^\circ$.	$\mathrm{CH_3X} > 1^\circ > 2^\circ \ggg 3^\circ$ (neopentyl extremely slow).
Nucleophile	Weak nucleophile is sufficient ($\mathrm{H_2O}, \mathrm{ROH}$).	Strong, unhindered nucleophile required ($\mathrm{OH^-}, \mathrm{RO^-}, \mathrm{CN^-}, \mathrm{I^-}$).
Solvent Requirement	Polar protic solvent ($\mathrm{H_2O}$, MeOH, EtOH) stabilizes ions.	Polar aprotic solvent (DMSO, DMF, acetone) desolvates $\mathrm{Nu^-}$.
Stereochemistry	Racemization with slight excess of inversion.	100% Walden Inversion of configuration.
Rearrangements	Readily observed (hydride/alkyl shifts, ring expansion).	Completely absent (no carbocation intermediate).
Leaving Group Order	$\mathrm{I^-} > \mathrm{Br^-} > \mathrm{Cl^-} \gg \mathrm{F^-}$ (governed by C – X bond energy).	$\mathrm{I^-} > \mathrm{Br^-} > \mathrm{Cl^-} \gg \mathrm{F^-}$ (governed by leaving group stability).

S_Ni Mechanism (Substitution Nucleophilic Internal): Thionyl Chloride Reaction

When an alcohol reacts with thionyl chloride ($\mathrm{SOCl_2}$):

- **In Ether (No Pyridine):** $\mathrm{R-OH + SOCl_2} \xrightarrow{\text{ether}} \mathrm{R-Cl + SO_2\uparrow + HCl\uparrow}$. Proceeds via an internal chlorosulfite intermediate ($\mathrm{R-O-SOCl}$) with intramolecular delivery of chlorine from the same face $\rightarrow$ **Complete Retention of Configuration (S_Ni).**
- **In Pyridine Medium:** Pyridine neutralizes HCl to form pyridinium chloride ($\mathrm{C_5H_5NH^+Cl^-}$), releasing free chloride ions ($\mathrm{Cl^-}$). These chloride ions attack the chlorosulfite from the backside $\rightarrow$ **Complete Inversion of Configuration (S_N2).**

6.3 Elimination Reactions ($E1$ vs $E2$) & Dehydration of Alcohols

6.3.1 $E1$ Elimination Mechanism

Unimolecular Elimination ($E1$): Steps & Thermodynamic Drive

$E1$ elimination proceeds through ionization to a planar carbocation, identical to Step 1 of S_N1:

1. **Step 1 (RDS):** $\mathrm{R-X} \xrightarrow{\text{slow, RDS}} \mathrm{R^+ + X^-}$.
2. **Step 2 (Fast):** A weak base ($\mathrm{H_2O}, \mathrm{EtOH}$) abstracts a β-hydrogen from the carbocation:

$$\mathrm{R-CH_2-\overset{+}{C}H-CH_3 + B: \xrightarrow{fast} R-CH=CH-CH_3 + BH^+} \tag{6.20}$$

3. **Thermodynamic Competition (S_N1 vs $E1$):** Because Step 1 is shared, S_N1 and $E1$ compete directly. Elimination yields 2 product molecules from 1 reactant ($\Delta S^\circ > 0$). In the Gibbs equation $\Delta G^\circ = \Delta H^\circ - T\Delta S^\circ$, higher temperatures increase the magnitude of $-T\Delta S^\circ$, favoring $E1$ **elimination over S_N1 substitution at elevated temperatures**.
4. **Regiochemistry:** Governed by **Saytzeff's Rule** (the more substituted, thermodynamically stable alkene with more hyperconjugative α-hydrogens is the major product; $trans > cis$).

6.3.2 $E2$ Elimination: Anti-Periplanar Geometry & Regioselectivity

Bimolecular Elimination ($E2$): Stereoelectronic Rigidity

$E2$ elimination occurs in a concerted manner upon treatment with a strong base (such as alcoholic KOH or $\mathrm{NaOCH_2CH_3}$):

$$\text{Rate} = k[\mathrm{R-X}][\text{Base}] \quad (\text{Order} = 2, \text{Molecularity} = 2) \tag{6.21}$$

Substrate Order: $3^\circ > 2^\circ > 1^\circ$ (developing double bond in transition state is stabilized by alkyl groups).

Stereoelectronic Requirement (Anti-Periplanar Geometry): The $\mathrm{C_\beta - H}$ bond and $\mathrm{C_\alpha - X}$ bond must lie in the same plane at a dihedral angle of 180° (**anti-coplanar**). This orientation allows the electron pair from the breaking $\mathrm{C-H}$ bond to smoothly overlap into the developing π-bond while minimizing electron repulsion between the incoming base and departing halide.

Saytzeff (Zaitsev) vs Hofmann Regioselectivity Matrix

1. Saytzeff Rule (Thermodynamic Major Product): With normal unhindered strong bases (OH^-, OEt^-) and good leaving groups (I^-, Br^-, Cl^-), the transition state resembles the product alkene. The more substituted, hyperconjugated alkene is formed as the major product:

$$CH_3CH_2CH(Br)CH_3 \xrightarrow{\text{alc. KOH},\Delta} \textbf{But-2-ene}\ (81\%, \text{Saytzeff}) + \text{But-1-ene}\ (19\%, \text{Hofmann})$$

2. Hofmann Rule (Kinetic Major Product): The less substituted terminal alkene is formed under two scenarios:

- **Scenario A: Poor Leaving Groups ($-F, -NR_3^+$):** Transition state develops carbanionic character ($1^\circ > 2^\circ > 3^\circ$), favoring terminal deprotonation:

$$CH_3CH_2CH(F)CH_3 \xrightarrow{EtO^-} \textbf{But-1-ene}\ (80\%) + \text{But-2-ene}\ (20\%)$$

- **Scenario B: Sterically Bulky Bases (*t*-BuOK):** Bulky base abstracts accessible terminal β-hydrogen:

$$CH_3CH_2CH(Br)CH_3 \xrightarrow{t\text{-}BuO^-} \textbf{But-1-ene}\ (73\%) + \text{But-2-ene}\ (27\%)$$

6.3.3 Acid-Catalyzed Dehydration of Alcohols

Dehydration of Alcohols: Step-by-Step Mechanism

Dehydration of alcohols to alkenes is an acid-catalyzed elimination ($E1$ for 2° and 3° alcohols; $E2$ for 1° alcohols):

1. **Step 1: Protonation of Hydroxyl Group (Fast Equilibrium):**

$$R - OH + H^+ \rightleftharpoons R - \overset{+}{O}H_2 \tag{6.22}$$

   (Converts poor leaving group $-OH$ into excellent leaving group H_2O).
2. **Step 2: Heterolysis to form Carbocation [RDS]:**

$$R - \overset{+}{O}H_2 \xrightarrow{\text{slow, RDS}} R^+ + H_2O \tag{6.23}$$

3. **Step 3: Carbocation Rearrangement (if applicable):** 1,2-hydride or 1,2-methyl shifts to yield the most stable carbocation.
4. **Step 4: Deprotonation (Fast):** Water or HSO_4^- abstracts β-H to form the Saytzeff alkene.

NCERT Class 12 Benchmark: Reaction Conditions & Temperature Trap

- **Reactivity Hierarchy:** 3° alcohol $>$ 2° alcohol $>$ 1° alcohol (parallels carbocation stability: $3^\circ > 2^\circ > 1^\circ$).
- **Ethanol Conditions:**
  - At 443 K(170°C) with conc. H_2SO_4: **Ethene** ($CH_2 = CH_2$) forms via intramolecular dehydration.
  - At 413 K(140°C) with excess ethanol: **Diethyl ether** ($C_2H_5OC_2H_5$) forms via bimolecular nucleophilic substitution (S_N2).
- 2° **Alcohol (Propan-2-ol):** Requires only 85% H_3PO_4 at 440 K.
- 3° **Alcohol (2-Methylpropan-2-ol):** Requires only 20% H_2SO_4 at mild temperature of 358 K.

6.4 Carbonyl Chemistry Mechanisms

6.4.1 Nucleophilic Addition to Aldehydes & Ketones (Ad_N)

Carbonyl Electrophilicity & Trajectory of Nucleophilic Attack

The carbonyl group ($>C=O$) is strongly polarized due to the high electronegativity of oxygen ($\chi_O = 3.5$ vs $\chi_C = 2.5$). The carbonyl carbon is planar sp^2 with an empty π^* LUMO.

1. **Burgi-Dunitz Angle:** The nucleophile attacks the electrophilic carbon at an angle of $\approx 107°$ relative to the $C=O$ bond axis, maximizing overlap with π^* while minimizing repulsion with oxygen lone pairs.
2. **Reactivity Sequence:**

$$HCHO > CH_3CHO > CH_3COCH_3 \tag{6.24}$$
$$> C_6H_5CHO > C_6H_5COCH_3 > C_6H_5COC_6H_5 \tag{6.25}$$

Governing Factors:

- **Electronic Effect:** Electron-donating alkyl groups ($+I$) disperse the partial positive charge on carbon ($\delta+$), reducing its electrophilicity. Conjugation with an aromatic ring (resonance) stabilizes the ground state carbonyl, diminishing reactivity.
- **Steric Effect:** As substituents on the carbonyl carbon increase in bulk, transition state crowding during conversion from trigonal planar (120°) to tetrahedral (109.5°) increases, raising the activation energy.

Key Nucleophilic Addition Mechanisms

1. Addition of HCN (Cyanohydrin Formation):

- Pure HCN is a weak acid ($pK_a = 9.2$) and reacts extremely slowly. The reaction is base-catalyzed:

$$HCN + OH^- \rightleftharpoons CN^- + H_2O \quad \text{(Generates strong nucleophile } CN^-\text{)} \tag{6.26}$$

- Step 1 (RDS): Attack of CN^- on carbonyl carbon: $R_2C=O + CN^- \xrightarrow{\text{slow}} R_2C(CN)O^-$.
- Step 2 (Fast): Proton transfer: $R_2C(CN)O^- + HCN \xrightarrow{\text{fast}} R_2C(CN)OH + CN^-$.

2. Addition of Sodium Bisulfite ($NaHSO_3$): Sulfur is more nucleophilic than oxygen. The bisulfite nucleophile attacks via sulfur to form a tetrahedral intermediate, followed by proton transfer to give a crystalline, water-soluble

bisulfite adduct:

$$\mathrm{RCHO} + \mathrm{NaHSO_3} \rightleftharpoons \mathrm{R-CH(OH)SO_3^-Na^+} \quad \text{(White crystalline precipitate)} \tag{6.27}$$

Used for the separation and purification of aldehydes and aliphatic methyl ketones (sterically hindered ketones do not react). **3. Addition of Ammonia Derivatives ($\mathrm{H_2N-Z}$):** Reagents: $\mathrm{Z} = -\mathrm{OH}$ (hydroxylamine $\rightarrow$ oxime), $-\mathrm{NH_2}$ (hydrazine $\rightarrow$ hydrazone), $-\mathrm{NHC_6H_5}$ (phenylhydrazine $\rightarrow$ phenylhydrazone), $-\mathrm{NH-C_6H_3(NO_2)_2}$ (2,4-DNP $\rightarrow$ orange Brady's precipitate), $-\mathrm{NHCONH_2}$ (semicarbazide $\rightarrow$ semicarbazone).

- **Optimum pH Control (pH ≈ 4.5):** If medium is too acidic (pH < 3), the amine reagent is completely protonated to $\mathrm{H_3\overset{+}{N}-Z}$, destroying its nucleophilicity. If medium is too basic (pH > 6), acid-catalyzed dehydration of the carbinolamine intermediate is too slow.

6.4.2 Aldol Condensation & Cross-Aldol Mechanism

Aldol Condensation: Step-by-Step Mechanism

Aldol condensation occurs in aldehydes and ketones possessing at least one α-hydrogen when treated with dilute alkali ($\mathrm{NaOH}, \mathrm{Ba(OH)_2}, \mathrm{Na_2CO_3}$):

1. **Step 1: Enolate Generation (Acid-Base Equilibrium):** Hydroxide abstracts an acidic α-proton ($pK_a \approx 19-20$, acidified by resonance stabilization and $-I$ of carbonyl):

$$\mathrm{CH_3CHO} + \mathrm{OH^-} \rightleftharpoons [\mathrm{^-CH_2CHO} \longleftrightarrow \mathrm{CH_2=CHO^-}] + \mathrm{H_2O} \tag{6.28}$$

(Resonance enolate carbanion) (6.29)

2. **Step 2: Nucleophilic Addition [RDS]:** The enolate carbanion attacks un-ionized aldehyde:

$$\mathrm{CH_3CHO} + \mathrm{CH_2=CHO^-} \xrightarrow{\text{slow, RDS}} \mathrm{CH_3-CH(O^-)-CH_2-CHO} \tag{6.30}$$

(Alkoxide intermediate) (6.31)

3. **Step 3: Protonation to Form β-Hydroxyaldehyde (Aldol):**

$$\mathrm{CH_3-CH(O^-)-CH_2-CHO} + \mathrm{H_2O} \rightleftharpoons \tag{6.32}$$

$$\mathrm{CH_3-CH(OH)-CH_2-CHO} + \mathrm{OH^-} \tag{6.33}$$

4. **Step 4: Condensation / Dehydration ($E1cB$ Mechanism):** Upon heating (Δ), the aldol eliminates water:

$$CH_3 - CH(OH) - CH_2 - CHO \xrightarrow[\Delta]{OH^-} CH_3 - CH = CH - CHO + H_2O \tag{6.34}$$

The resulting α,β-unsaturated enal/enone is thermodynamically stabilized by extended conjugation between the C = C double bond and the carbonyl group.

Cross-Aldol Condensation: Product Complexity & Directed Syntheses

- **Mixture of Two Aldehydes with α-H:** Reacting ethanal (CH_3CHO) with propanal (CH_3CH_2CHO) gives a complex mixture of **four** condensation products:
  1. Self-condensation of ethanal: But-2-enal.
  2. Self-condensation of propanal: 2-Methylpent-2-enal.
  3. Cross-condensation (ethanal enolate + propanal electrophile): Pent-2-enal.
  4. Cross-condensation (propanal enolate + ethanal electrophile): 2-Methylbut-2-enal.
- **Claisen-Schmidt Condensation (Directed Cross-Aldol):** To obtain a single clean product, one carbonyl component must lack α-hydrogens (e.g., benzaldehyde, C_6H_5CHO, or formaldehyde, HCHO). The partner with α-hydrogens acts exclusively as the nucleophile:

$$PhCHO + CH_3COCH_3 \xrightarrow[293\text{–}303\ K]{\text{dil. NaOH}} \tag{6.35}$$

$$Ph - CH = CH - COCH_3\ \text{(Benzalacetone)} + H_2O \tag{6.36}$$

6.4.3 Cannizzaro Reaction & Hydride Transfer Mechanism

Cannizzaro Disproportionation: Step-by-Step Hydride Shift

Aldehydes that lack α-hydrogens (e.g., $HCHO, C_6H_5CHO, (CH_3)_3CCHO$) cannot form an enolate. Upon heating with concentrated alkali (50% NaOH or KOH), they undergo self-oxidation-reduction (disproportionation):

1. **Step 1: Rapid Nucleophilic Addition of Hydroxide:**

$$Ar - CHO + OH^- \rightleftharpoons Ar - CH(OH)O^- \quad \text{(Monoanion hydrate)} \tag{6.37}$$

2. **Step 2: Second Deprotonation in Strongly Basic Media:**

$$Ar - CH(OH)O^- + OH^- \rightleftharpoons Ar - CH(O^-)_2 + H_2O \tag{6.38}$$

(**Dianion intermediate**, electron-rich hydride donor).

3. **Step 3: Intermolecular Hydride (H^-) Transfer [RDS]:** The electron-rich dianion expels a hydride ion (H^-) directly to the carbonyl carbon of a second aldehyde molecule. This direct hydride transfer is the slow, rate-determining step:

$$\mathrm{Ar-CH(O^-)_2 + Ar-CHO \xrightarrow{slow,\ RDS} Ar-COO^- + Ar-CH_2O^-} \quad (6.39)$$

4. **Step 4: Rapid Proton Exchange:** Alkoxide ($ArCH_2O^-$) rapidly abstracts a proton from water to yield alcohol ($ArCH_2OH$), leaving the carboxylate salt ($ArCOO^-Na^+$).

Kinetics & Crossed Cannizzaro Reaction

1. Kinetic Order:

$$\begin{aligned} \text{Moderate } [OH^-]: &\quad \text{Rate} = k[\mathrm{RCHO}]^2[OH^-] \text{ (Third order)} \\ \text{High } [OH^-]: &\quad \text{Rate} = k[\mathrm{RCHO}]^2[OH^-]^2 \text{ (Fourth order)} \end{aligned}$$

2. Crossed Cannizzaro (Regioselectivity of Oxidation): When formaldehyde (HCHO) is mixed with benzaldehyde (C_6H_5CHO), nucleophilic addition of OH^- occurs preferentially on formaldehyde:

$$\mathrm{C_6H_5CHO + HCHO \xrightarrow{conc.\ NaOH}} \textbf{Benzyl alcohol } (\mathrm{C_6H_5CH_2OH}) \quad (6.40)$$

$$+ \textbf{Sodium formate } (\mathrm{HCOO^-Na^+}) \quad (6.41)$$

Universal Rule: In crossed Cannizzaro involving formaldehyde, **formaldehyde is always oxidized to formate**, and the other aldehyde is reduced to the alcohol.

6.5 Carboxylic Acid Derivatives & Mechanisms

6.5.1 Fischer Esterification Mechanism

Acid-Catalyzed Esterification: Tetrahedral Intermediate Pathway

The reaction of a carboxylic acid with an alcohol in the presence of an inorganic acid catalyst (conc. H_2SO_4 or dry HCl) is an equilibrium addition-elimination process:

$$R - COOH + R' - OH \overset{H^+}{\rightleftharpoons} R - COOR' + H_2O \tag{6.42}$$

1. **Step 1: Protonation of Carbonyl Oxygen:** Protonation occurs selectively at the carbonyl oxygen (giving a resonance-stabilized cation $[R - C(OH)_2]^+$), increasing the electrophilicity of the carbonyl carbon:

$$R - COOH + H^+ \rightleftharpoons R - \overset{+}{C}(OH)_2 \tag{6.43}$$

2. **Step 2: Nucleophilic Attack of Alcohol [RDS]:** The alcohol oxygen lone pair attacks the activated carbonyl carbon:

$$R - \overset{+}{C}(OH)_2 + R'OH \xrightarrow{\text{slow, RDS}} R - C(OH)_2(\overset{+}{O}HR') \tag{6.44}$$

3. **Step 3: Proton Transfer (Tautomerism):** Proton shifts from the oxonium group to one of the hydroxyl groups:

$$R - C(OH)_2(\overset{+}{O}HR') \rightleftharpoons R - C(OH)(\overset{+}{O}H_2)(OR') \tag{6.45}$$

   This converts a poor leaving group ($-OH$) into a good leaving group (H_2O).
4. **Step 4: Elimination of H_2O & Rearomatization/Deprotonation:** The hydroxyl oxygen lone pair assists in expelling H_2O, followed by loss of H^+ to regenerate the acid catalyst and yield the ester ($RCOOR'$).

Isotopic Tracer Proof of Acyl-Oxygen Bond Cleavage

When benzoic acid reacts with ^{18}O-labeled methanol ($CH_3{}^{18}OH$), mass spectrometry confirms that the heavy isotope ^{18}O resides entirely in methyl benzoate, and **no** ^{18}O is found in the produced water:

$$C_6H_5 - CO - OH + H - {}^{18}OCH_3 \overset{H^+}{\rightleftharpoons} C_6H_5 - CO - {}^{18}OCH_3 + H_2O \tag{6.46}$$

Deduction: The reaction proceeds with **Acyl-Oxygen bond fission** ($R - CO$—OH cleavage) in the carboxylic acid, and **Alkyl-Oxygen bond retention** ($R'O$—H cleavage) in the alcohol!

Steric Hindrance Hierarchy in Esterification

Because the rate-determining step involves transition from planar sp^2 (120°) to a crowded tetrahedral sp^3 intermediate (109.5°), esterification rates are governed primarily by steric bulk:

$$\textbf{Acids:}\quad \mathrm{HCOOH} > \mathrm{CH_3COOH} > \mathrm{RCH_2COOH} > \mathrm{R_2CHCOOH}$$
$$\textbf{Alcohols:}\quad \mathrm{CH_3OH} > 1^\circ > 2^\circ \gg 3^\circ$$

Note on 3° *Alcohols:* Tertiary alcohols (e.g., tert-butanol) do **not** react via Fischer esterification. Instead, they dehydrate rapidly to alkenes or react via alkyl-oxygen cleavage through an S_N1 carbocation intermediate ($\mathrm{R'^+}$).

6.5.2 Decarboxylation of Carboxylic Acids

Soda-Lime Decarboxylation: Mechanism & The Carbanion Intermediate

Heating sodium or potassium salts of carboxylic acids with soda lime (a dry mixture of NaOH and CaO in a 3 : 1 mass ratio) eliminates carbon dioxide as carbonate, yielding an alkane with **one carbon atom less** than the parent carboxylate:

$$\mathrm{R-COONa + NaOH} \xrightarrow[\Delta]{\mathrm{CaO}} \mathrm{R-H + Na_2CO_3} \tag{6.47}$$

Step-by-Step Reaction Mechanism:

1. **Step 1: Dissociation & Attack by Base:** Under high thermal fusion (≈ 600 K), the carboxylate anion interacts with hydroxide ion:

$$\mathrm{R-COO^- + OH^-} \rightleftharpoons [\mathrm{R-COO\cdots H\cdots O}]^{2-} \tag{6.48}$$

2. **Step 2: Heterolytic Cleavage to Free Carbanion [RDS]:** The C – C bond connecting the alkyl group to the carboxylate carbon undergoes heterolytic cleavage, expelling carbon dioxide (which instantly neutralizes with base to form carbonate) and generating a reactive **carbanion intermediate** ($\mathrm{R^-}$):

$$\mathrm{R-COO^-} \xrightarrow{\text{slow, RDS}} \mathbf{R^-} + \mathrm{CO_2}$$
$$(\mathrm{CO_2 + 2OH^-} \rightarrow \mathrm{CO_3^{2-} + H_2O})$$

3. **Step 3: Rapid Proton Capture:** The carbanion abstracts a proton from water to form the alkane:

$$\mathbf{R^-} + \mathrm{H_2O} \xrightarrow{\text{fast}} \mathbf{R-H} + \mathrm{OH^-}$$

Rate-Determining Rule: Soda-lime decarboxylation rate is proportional to carbanion stability:

$$\text{Decarboxylation Rate} \propto \text{Carbanion Stability: } \mathrm{CH_3^-} > 1^\circ > 2^\circ > 3^\circ$$

Electron-withdrawing groups $(-I, -M)$ that disperse the negative charge tremendously accelerate the rate:

$$O_2N - CH_2 - COOH > NC - CH_2 - COOH > Cl_3C - COOH > CH_3COOH \tag{6.49}$$

Kolbe's Electrolytic Decarboxylation & Free Radical Coupling

When a concentrated aqueous solution of sodium or potassium carboxylate undergoes electrolysis, decarboxylative dimerization occurs at the anode:

$$2R - COO^-K^+ + 2H_2O \xrightarrow{\text{Electrolysis}} \mathbf{R - R} + 2CO_2 \uparrow + H_2 \uparrow + 2KOH \tag{6.50}$$

Electrochemical Mechanism at Electrodes:

- **Anode Oxidation (Decarboxylation & Radical Dimerization):**

$$2R - COO^- \xrightarrow{-2e^-} 2[R - COO^\bullet] \quad \text{(Acyloxy free radical)} \tag{6.51}$$

$$2[R - COO^\bullet] \xrightarrow{\text{fast, RDS}} 2R^\bullet + 2CO_2 \uparrow \quad \text{(Homolytic cleavage)} \tag{6.52}$$

$$2R^\bullet \xrightarrow{\text{fast}} \mathbf{R - R} \quad \text{(Radical-radical coupling)} \tag{6.53}$$

- **Cathode Reduction (Evolution of H_2 & Alkalinization):**

$$2H_2O + 2e^- \rightarrow \mathbf{H_2} \uparrow + 2OH^- \tag{6.54}$$

Diagnostic Electrolytic Phenomena:

1. **pH Drift:** As the reaction proceeds, H^+ is evolved as H_2 gas while OH^- accumulates in solution. Hence, the **pH of the electrolyte continuously rises** (pH > 7, alkaline).
2. **Gas Ratio at Electrodes:** At the anode, CO_2 and alkane vapor are evolved. At the cathode, pure H_2 gas is evolved. For sodium acetate, the molar ratio of gases is (Anodic gases : Cathodic gas) $= (2CO_2 + 1C_2H_6) : (1H_2) = 3 : 1$.

Four Cardinal NEET Traps in Decarboxylation Reactions

1. **The Methane Exclusion in Kolbe Electrolysis:** Methane (CH_4) **cannot** be prepared by Kolbe electrolysis because radical coupling requires at least two alkyl radicals ($2CH_3^\bullet \rightarrow CH_3 - CH_3$). Potassium formate (HCOOK) electrolyzed yields H_2 and CO_2, not alkane!
2. **Radical Disproportionation By-Products:** When sodium propanoate (CH_3CH_2COONa) is electrolyzed, the intermediate ethyl radicals ($CH_3CH_2^\bullet$) undergo both coupling (*n*-butane) and disproportionation:

$$CH_3CH_2^\bullet + CH_3CH_2^\bullet \rightarrow CH_2 = CH_2 \text{ (Ethene)} + CH_3 - CH_3 \text{ (Ethane)} \tag{6.55}$$

3. **Dicarboxylic Acid Electrolysis Shortcuts:**

- Potassium succinate ($KOOC - CH_2 - CH_2 - COOK$) yields **Ethene** ($CH_2 = CH_2$).
- Potassium maleate or fumarate ($KOOC - CH = CH - COOK$) yields **Ethyne** ($CH \equiv CH$).

4. **Role of CaO in Soda Lime:** CaO is not a reactant. It is used because:
   - It is intensely hygroscopic and keeps solid NaOH completely dry (deliquescence prevention).
   - It prevents melting/fusion of NaOH at high temperatures, avoiding corrosion and cracking of the glass tube.

Thermal Decarboxylation of β-Keto & Gem-Dicarboxylic Acids

Carboxylic acids possessing a carbonyl group at the β-carbon (β-keto acids like acetoacetic acid, and gem-dicarboxylic acids like malonic acid) undergo spontaneous decarboxylation upon gentle heating ($\approx 373 - 423$ K) **without any added base or catalyst**:

$$R - CO - CH_2 - COOH \xrightarrow{\Delta} R - CO - CH_3 + CO_2 \uparrow \tag{6.56}$$

Pericyclic Mechanism via 6-Membered Cyclic Transition State: The acidic carboxylic proton forms an intramolecular hydrogen bond with the β-carbonyl oxygen. Heating triggers a concerted 6-electron cyclic electron redistribution:

$$R - CO - CH_2 - COOH \xrightarrow[\text{6-ring T.S.}]{\Delta} R - C(OH) = CH_2 + CO_2 \uparrow \tag{6.57}$$

$$\xrightarrow{\text{tautomerism}} R - CO - CH_3 \text{ (Methyl Ketone)} \tag{6.58}$$

The enol intermediate undergoes instantaneous, virtually irreversible keto-enol tautomerism to yield the thermodynamically favored **methyl ketone**.

6.5.3 Hell-Volhard-Zelinsky (HVZ) Reaction & α-Halogenation

HVZ Mechanism: In Situ Enolization via Acyl Halide

Carboxylic acids undergo selective α-halogenation when treated with bromine or chlorine in the presence of a catalytic amount of red phosphorus:

$$R - CH_2 - COOH \xrightarrow[\text{(ii) } H_2O]{\text{(i) } X_2/\text{red P}} R - CH(X) - COOH \quad (X = Cl, Br) \tag{6.59}$$

Step-by-Step Mechanistic Progression:

1. **Formation of Phosphorus Trihalide:** $2P + 3X_2 \rightarrow 2PX_3$.

2. **Acyl Halide Generation:** Carboxylic acids themselves enolize too slowly for direct α-halogenation. The PX_3 converts a small catalytic fraction of carboxylic acid to the corresponding acyl halide:

$$3R - CH_2 - COOH + PX_3 \rightarrow 3R - CH_2 - COX + H_3PO_3 \tag{6.60}$$

3. **Rapid Enolization:** The acyl halide enolizes millions of times faster than the carboxylic acid because the electronegative halogen increases the acidity of the α-protons:

$$R - CH_2 - COX \rightleftharpoons R - CH = C(OH)X \quad \text{(Enol form)} \tag{6.61}$$

4. **Electrophilic Halogenation:** The enol attacks X_2 to produce the α-haloacyl halide:

$$R - CH = C(OH)X + X_2 \rightarrow R - CH(X) - COX + HX \tag{6.62}$$

5. **Transacylation Exchange:** The α-haloacyl halide reacts with unreacted parent carboxylic acid to yield the α-halocarboxylic acid and regenerate the acyl halide, sustaining the catalytic cycle:

$$R - CH(X) - COX + R - CH_2 - COOH \rightleftharpoons \tag{6.63}$$

$$R - CH(X) - COOH + R - CH_2 - COX \tag{6.64}$$

HVZ Substrate Scope & Synthetic Traps

- **Requirement for α-Hydrogen:** Carboxylic acids devoid of α-hydrogens (e.g., formic acid HCOOH, benzoic acid C_6H_5COOH, and trimethylacetic acid $(CH_3)_3CCOOH$) **never** undergo the HVZ reaction.
- **Halogen Limitations:** The reaction works efficiently with Br_2 and Cl_2. Direct fluorination causes extensive fragmentation; direct iodination fails because the reverse reaction with HI dominates.
- **Downstream Synthetic Utility:** The α-halogen is readily substituted:

$$R - CH(Br) - COOH \xrightarrow{\text{aq. KOH}} R - CH(OH) - COOH \quad (\alpha\text{-hydroxy acid})$$

$$R - CH(Br) - COOH \xrightarrow{\text{excess } NH_3} R - CH(NH_2) - COOH \quad (\alpha\text{-amino acid})$$

6.6 Aromatic & Nitrogen Mechanisms

6.6.1 Diazotization Mechanism of Primary Aromatic Amines

Diazotization Mechanism & Nitrosonium Ion Electrophile

Primary aromatic amines react with nitrous acid at ice-cold temperature (273 – 278 K) to form stable arenediazonium salts:

$$\mathrm{Ar-NH_2 + NaNO_2 + 2HCl \xrightarrow{273-278\ K} Ar-N_2^+Cl^- + NaCl + 2H_2O} \quad (6.65)$$

Step-by-Step Reaction Sequence:

1. **Generation of Nitrosonium Ion (NO^+):** Nitrous acid is unstable and prepared in situ:

$$\mathrm{NaNO_2 + HCl \rightarrow HNO_2 + NaCl}$$

$$\mathrm{HNO_2 + H^+ \rightleftharpoons H_2\overset{+}{O} - NO \rightleftharpoons NO^+ + H_2O} \quad \text{(Nitrosonium electrophile)}$$

2. **Nucleophilic Attack:** The aniline nitrogen lone pair attacks NO^+ to form *N*-nitrosaniline cation:

$$\mathrm{Ar-NH_2 + NO^+ \rightleftharpoons Ar-\overset{+}{N}H_2-NO} \quad (6.66)$$

$$\xrightarrow{-H^+} \mathrm{Ar-NH-NO} \quad (N\text{-nitrosamine}) \quad (6.67)$$

3. **Proton Tautomerism & Dehydration:**

$$\mathrm{Ar-NH-N=O \rightleftharpoons Ar-N=N-OH \xrightarrow{H^+} Ar-N=N-\overset{+}{O}H_2} \quad (6.68)$$

$$\xrightarrow{-H_2O} \mathrm{Ar-\overset{+}{N}\equiv N \longleftrightarrow Ar-\ddot{N}=\overset{+}{N}} \quad (6.69)$$

The diazonium ion is stabilized by resonance with the aromatic π-system.

Thermal Decomposition & Aliphatic vs Aromatic Diazonium Salts

- **Temperature Sensitivity (> 278 K/5°C):** If the temperature rises above 5°C, benzenediazonium chloride decomposes rapidly in aqueous solution:

$$\mathrm{Ar-N_2^+Cl^- + H_2O \xrightarrow{>278\ K} Ar-OH + N_2\uparrow + HCl} \quad (6.70)$$

(Forms phenol quantitatively) (6.71)

- **Aliphatic Primary Amines:** Aliphatic diazonium salts ($R-N_2^+Cl^-$) are exceptionally unstable even at 0°C because the alkyl group cannot disperse the positive charge by resonance. They instantaneously lose molecular N_2 to generate high-energy carbocations (R^+), yielding complex mixtures of alcohols, alkenes, and alkyl chlorides. The quantitative evolution of N_2 is utilized ana-

lytically for amino acid estimation (Van Slyke method).

6.6.2 Azo Dye Coupling Mechanism (Phenol vs Aniline)

Azo Coupling: Mild S_EAr & pH-Directed Chemoselectivity

Arenediazonium cations are weak electrophiles ($Ar-N_2^+$) that undergo electrophilic aromatic substitution only on highly activated aromatic rings:

1. **Coupling with Phenol (Orange Dye):**
   - **Optimum pH:** Alkaline medium (pH $\approx$ 9–10).
   - **Mechanistic Rationale:** Phenol deprotonates to phenoxide (PhO^-, superior $+M$ donor). Weak diazonium electrophile attacks para-position:

$$PhN_2^+Cl^- + PhOH \xrightarrow[\text{pH } 9-10]{OH^-} p\text{-}HO-C_6H_4-N=N-Ph\ (\textbf{Orange}) + Cl^- + H_2O$$

   - **High pH Hazard:**
     At pH $>$ 12, diazonium forms diazotate ($Ar-N=N-O^-$).
2. **Coupling with Aniline (Yellow Dye):**
   - **Optimum pH:** Mildly acidic (pH $\approx$ 4–5).
   - **Mechanistic Rationale:** At pH $\approx$ 4–5, free aniline exists to couple at para-position:

$$PhN_2^+Cl^- + PhNH_2 \xrightarrow[\text{pH } 4-5]{H^+} p\text{-}H_2N-C_6H_4-N=N-Ph\ (\textbf{Yellow}) + HCl$$

   - In basic medium, diazoaminobenzene forms ($Ph-N=N-NH-Ph$).

6.6.3 Reimer-Tiemann Reaction Mechanism

Dichlorocarbene Intermediate & Formylation of Phenol

The conversion of phenol to salicylaldehyde using chloroform and aqueous sodium hydroxide:

1. **Step 1: Generation of Dichlorocarbene (: CCl_2) [α-Elimination]:** Hydroxide ion deprotonates the relatively acidic proton of chloroform ($CHCl_3, pK_a \approx 15.5$ due to $-I$ of three chlorines):

$$CHCl_3 + OH^- \rightleftharpoons H_2O + {}^-:CCl_3 \quad \text{(Trichloromethyl carbanion)}$$

$$^{-}:\mathrm{CCl_3} \xrightarrow{\text{slow, } \alpha\text{-elimination}} :\mathrm{CCl_2}\ \text{(Singlet Dichlorocarbene)} + \mathrm{Cl^-}$$

Nature of $:\mathrm{CCl_2}$**:** Neutral reactive intermediate with 6 valence electrons and an empty unhybridized p-orbital → **powerful electrophile**.

2. **Step 2: Electrophilic Attack on Phenoxide:** Phenoxide attacks $:\mathrm{CCl_2}$ predominantly at the ortho position, forming a non-aromatic cyclohexadienone carbanion intermediate:

$$\mathrm{C_6H_5O^-} + :\mathrm{CCl_2} \rightarrow [o\text{-}\mathrm{C_6H_4(=O)} - \mathrm{CCl_2^-}] \tag{6.72}$$

3. **Step 3: Intramolecular Proton Shift & Rearomatization:** The aromatic proton from the ring shifts to the $-\mathrm{CCl_2^-}$ carbon, restoring aromatic sextet stability and forming the dichloromethylphenoxide intermediate:

$$[o\text{-}\mathrm{C_6H_4(=O)} - \mathrm{CCl_2^-}] \rightarrow o\text{-}\mathrm{O^-} - \mathrm{C_6H_4} - \mathrm{CHCl_2} \tag{6.73}$$

4. **Step 4: Alkaline Hydrolysis & Dehydration:** Hydroxide displaces both chlorine atoms via nucleophilic substitution, producing an unstable gem-diol intermediate that spontaneously loses water upon acidification to afford **salicylaldehyde** (o-hydroxybenzaldehyde).

Reimer-Tiemann Variant Trap: Chloroform vs Carbon Tetrachloride

- If **$\mathrm{CHCl_3}$** is employed: The intermediate is a gem-dichloride ($-\mathrm{CHCl_2}$), which hydrolyzes to give **Salicylaldehyde** (o-hydroxybenzaldehyde).
- If **$\mathrm{CCl_4}$** is employed: Phenoxide attacks $\mathrm{CCl_4}$ to generate a trichloromethyl intermediate ($-\mathrm{CCl_3}$). Alkaline hydrolysis of $-\mathrm{CCl_3}$ introduces three hydroxyl groups, which eliminate water to yield **Salicylic Acid** (o-hydroxybenzoic acid)!

6.6.4 Kolbe's (Kolbe-Schmitt) Reaction Mechanism

Carboxylation of Phenoxide via Carbon Dioxide Electrophile

Treatment of sodium phenoxide with carbon dioxide gas at 400 K and $4-7$ atm, followed by acidification, produces salicylic acid:

$$\mathrm{C_6H_5OH} \xrightarrow{\mathrm{NaOH}} \mathrm{C_6H_5O^-Na^+}$$
$$\mathrm{C_6H_5O^-Na^+} \xrightarrow[4\text{–}7\ \text{atm}]{\mathrm{CO_2,400\ K}} \text{Sodium Salicylate}$$
$$\xrightarrow{\mathrm{H^+}} \textbf{Salicylic Acid}$$

Mechanism & Ortho-Directing Sodium Chelate Effect:

1. **Electrophile:** Carbon dioxide ($\mathrm{O=C=O}$) is a relatively weak, neutral electrophile. Un-ionized phenol is too unreactive to react with $\mathrm{CO_2}$; conversion to sodium phenoxide ($\mathrm{Ph-O^-}$) provides the requisite nucleophilic activation.

2. **6-Membered Cyclic Transition State:** The Na^+ counter-ion coordinates simultaneously to the phenoxide oxygen and one of the carbonyl oxygens of CO_2. This cyclic coordination holds CO_2 directly above the ortho-carbon, directing electrophilic attack selectively to the **ortho position**.
3. **Rearomatization:** Proton transfer from the ring carbon to oxygen regenerates the aromatic ring, followed by acidification to yield salicylic acid.

Cation Size Effect: Sodium vs Potassium Phenoxide

- **Sodium Phenoxide (Na^+):** Forms a tight 6-membered cyclic chelate due to the small ionic radius of Na^+ (102 pm), giving **salicylic acid (ortho)** as the major product ($> 80\%$).
- **Potassium Phenoxide (K^+):** K^+ has a larger ionic radius (138 pm) and cannot form an effective cyclic chelate. At 473 K, attack shifts predominantly to the less hindered para position, yielding **p-hydroxybenzoic acid** as the major product.
- **Downstream Aspirin Synthesis:**

$$\text{Salicylic acid} + (CH_3CO)_2O \xrightarrow{H^+} \tag{6.74}$$

$$\textbf{Acetylsalicylic acid (Aspirin)} + CH_3COOH \tag{6.75}$$

6.6.5 Hofmann Bromamide Degradation Mechanism

Hofmann Bromamide Reaction: Concerted Intramolecular Rearrangement

Primary acid amides react with bromine in aqueous or ethanolic sodium hydroxide to yield primary amines containing **one carbon less** than the parent amide:

$$R - CONH_2 + Br_2 + 4NaOH \rightarrow R - NH_2 + Na_2CO_3 + 2NaBr + 2H_2O \tag{6.76}$$

Detailed Step-by-Step Mechanism:

1. **Step 1: Deprotonation & Formation of *N*-Bromoamide:** Hydroxide abstracts a proton from amide nitrogen ($pK_a \approx 15-16$), generating an anion that attacks Br_2:

$$R - CONH_2 + OH^- \rightleftharpoons R - CON^-H + H_2O$$

$$\xrightarrow{Br_2} R - CONHBr + Br^- \ (N\text{-bromoamide})$$

2. **Step 2: Second Deprotonation to Bromoamide Anion:** The remaining proton on nitrogen is more acidic due to the $-I$ effect of bromine:

$$R - CONHBr + OH^- \rightleftharpoons [R - CO - \bar{N} - Br] + H_2O$$

3. **Step 3: Concerted 1,2-Alkyl Shift [RDS] → Formation of Isocyanate:** Expulsion of bromide is concerted with intramolecular migration of R:

$$[\mathrm{R-CO-\bar{N}-Br}] \xrightarrow[\text{slow, RDS}]{-\mathrm{Br}^-} \mathbf{R-N=C=O}$$

(**Alkyl Isocyanate Intermediate**)

Stereochemical Consequence: The alkyl group migrates with **100% retention of stereochemical configuration** at the migrating chiral center!

4. **Step 4: Hydrolysis of Isocyanate to Carbamic Acid:** Hydroxide attacks the electrophilic carbonyl carbon of the isocyanate, followed by proton transfer to form an unstable carbamic acid:

$$\mathrm{R-N=C=O + OH^- \rightarrow R-NH-COO^-} \tag{6.77}$$

5. **Step 5: Spontaneous Decarboxylation:** The carbamate undergoes spontaneous decarboxylation, releasing CO_2 (which is trapped by alkali as Na_2CO_3) and yielding the **primary amine** ($\mathrm{R-NH_2}$).

Hofmann Bromamide Pitfalls, Stoichiometry & Reagent Trapping

- **Exact Stoichiometry:** For every 1 mol of amide, exactly **1** mol of **Br_2** and **4** mol of **NaOH** are consumed. A frequent NEET trap involves asking for the molar ratio: Amide : Br_2 : NaOH = 1 : 1 : 4.
- **Secondary and Tertiary Amides Fail:** Secondary amides ($\mathrm{RCONHR'}$) have only one hydrogen on nitrogen and cannot form the necessary *N*-bromoamide anion; tertiary amides have zero hydrogens. Thus, only 1° amides undergo this reaction.
- **Isocyanate Trapping Proof:** If the reaction is conducted in an anhydrous alcoholic medium ($\mathrm{R'OH}$) instead of water, the isocyanate intermediate is intercepted by alcohol to form a stable **urethane / carbamate** ($\mathrm{R-NH-COOR'}$), proving the existence of the isocyanate intermediate!
- **Phthalimide Reaction:** Phthalimide treated with Br_2/NaOH undergoes degradation to yield **anthranilic acid** (*o*-aminobenzoic acid).

6.7 Master Diagnostic Matrix: Reaction Mechanisms Overview

6.7.1 Reaction Matrix I: Hydrocarbons, Alkyl Halides & Alcohols

Reaction Name	Key Intermediate	Kinetic Order	Regio / Stereochemistry	Critical NEET Exam Pitfall
Nitration of Benzene	Arenium Ion (σ-complex)	2nd order $(1+1)$	$k_H/k_D \approx 1$ (no isotope effect)	HNO_3 acts as a base; aniline gives 47% meta without $-NH_2$ protection.
Friedel-Crafts Alkylation	$1^\circ, 2^\circ, 3^\circ$ Carbocation	2nd order	Thermodynamic rearrangement	Polyalkylation occurs; deactivated rings and aniline fail.
Friedel-Crafts Acylation	Acylium Ion $(R-C\equiv\overset{+}{O})$	2nd order	No rearrangement	Deactivates ring against polyacylation; requires >1 eq. $AlCl_3$.
Halogenation of Halobenzenes	Arenium Ion ($+M$ donor)	2nd order	Ortho/Para orientation	Halogens are deactivating overall ($-I > +M$) but direct ortho/para.
S_N1 **Substitution**	Planar Carbocation (sp^2)	1st order: k[RX]	Net partial racemization + excess inversion	Rearrangements occur; hindered substrate (3°) and polar protic solvent required.
S_N2 **Substitution**	Trigonal Bipyramid T.S.	2nd order: k[RX][Nu]	100% Walden Inversion	Neopentyl halides are extremely unreactive; polar aprotic solvents accelerate.
$E1$ **Elimination**	Planar Carbocation (sp^2)	1st order: k[RX]	Saytzeff major (more substituted)	High temperature favors $E1$ over S_N1 due to $+T\Delta S^\circ$.
$E2$ **Elimination**	Anti-periplanar T.S.	2nd order: k[RX][Base]	Anti-elimination; Saytzeff vs Hofmann	Hofmann product forms with poor leaving group (F^-) or bulky base (t-BuOK).
Dehydration of Alcohols	Alkyloxonium $\rightarrow$ Carbocation	1st order in alcohol	Saytzeff alkene	Ethanol at 443 K $\rightarrow$ ethene; at 413 K $\rightarrow$ diethyl ether (S_N2).

6.7.2 Reaction Matrix II: Carbonyls & Carboxylic Acid Derivatives

Reaction Name	Key Intermediate	Kinetic Order	Regio / Stereochemistry	Critical NEET Exam Pitfall
Aldol Condensation	Enolate Carbanion $\rightarrow$ Aldol	2nd order overall	α,β-unsaturated enal/enone	Must possess α-H; cross-aldol yields 4 products unless one partner lacks α-H.
Cannizzaro Reaction	Dianion hydrate $\rightarrow H^-$ shift	3rd / 4th order	Intermolecular hydride transfer	Requires NO α-H; in crossed Cannizzaro with HCHO, HCHO is always oxidized.
Fischer Esterification	Tetrahedral Intermediate	Equilibrium	Acyl-oxygen ($R-CO—OH$) cleavage	^{18}O tracer remains in ester; rate hindered by bulkiness on acid/alcohol.
Soda-Lime Decarboxylation	Free Carbanion (R^-)	1st order in carboxylate	Stereochemistry cage dependent	Rate $\propto$ carbanion stability ($CH_3^- > 1^\circ > 2^\circ > 3^\circ$); CaO prevents tube fusion and deliquescence.
Kolbe Electrolysis	Alkyl Free Radical ($R^\bullet$)	2nd order radical coupling	Symmetrical $R-R$ dimer	Methane cannot be synthesized; solution becomes strongly alkaline (pH $>$ 7) due to cathodic OH^-.

6.7.3 Reaction Matrix III: Phenols, Diazonium & Nitrogen Compounds

Reaction Name	Key Intermediate	Kinetic Order	Regio / Stereochemistry	Critical NEET Exam Pitfall
HVZ Halogenation	Enol of Acyl Halide	Catalytic in P	α-halogenation only	Must possess α-H; red P generates PX_3 which accelerates enolization.
Azo Dye Coupling	Diazonium Cation ($Ar - N_2^+$)	2nd order	Para-coupling exclusively	Phenol requires pH $\approx$ 9 – 10 (orange); Aniline requires pH $\approx 4-5$ (yellow).
Reimer-Tiemann	Dichlorocarbene (: CCl_2)	2nd order	Ortho-salicylaldehyde	Intermediate is : CCl_2 (6 valence e^-); using CCl_4 gives salicylic acid.
Kolbe Reaction	Sodium Phenoxide chelate	2nd order	Ortho-salicylic acid (Na^+)	Na^+ gives ortho (salicylic acid); K^+ at 473 K gives para-hydroxybenzoic acid.
Hofmann Degradation	Alkyl Isocyanate ($R - NCO$)	2nd order	Retention of chiral center	Carbon chain decreases by 1; consumes 1 mol Br_2 : 4 mol NaOH; 1° amides only.

PART VI

Practical Chemistry & Qualitative Analysis Lab Manual

CHAPTER 7

Practical Chemistry: Qualitative Salt Analysis & Volumetric Redox Titrations

With the NMC's renewed emphasis on practical laboratory skills, qualitative and quantitative analysis questions appear regularly. This chapter provides a complete laboratory manual covering systematic group separation of basic radicals (cations), confirmatory tests for acidic radicals (anions), and volumetric redox titrations with $KMnO_4$.

7.1 Systematic Qualitative Analysis: Acidic Radicals (Anions)

Class I: Dilute H_2SO_4 Soluble Anions ($CO_3^{2-}, S^{2-}, SO_3^{2-}, NO_2^-$)

Treated with dilute H_2SO_4 to observe volatile evolved gases:

Anion	Gas	Preliminary Observation	Confirmatory Test & Governing Equation
CO_3^{2-}	CO_2	Brisk effervescence; odorless, colorless gas.	Lime water turns milky ($CaCO_3\downarrow$); clears with excess CO_2: $CaCO_3 + H_2O + CO_2 \rightarrow Ca(HCO_3)_2$.
S^{2-}	H_2S	Colorless gas with rotten-egg smell.	Turns lead acetate paper black ($PbS\downarrow$); purple/violet with nitroprusside: $S^{2-} + [Fe(CN)_5NO]^{2-} \rightarrow [Fe(CN)_5NOS]^{4-}$.
SO_3^{2-}	SO_2	Colorless gas with suffocating odor of sulfur.	Acidified $K_2Cr_2O_7$ turns green: $K_2Cr_2O_7 + H_2SO_4 + 3SO_2 \rightarrow Cr_2(SO_4)_3 + K_2SO_4 + H_2O$.
NO_2^-	NO_2	Pungent reddish-brown fumes.	Starch-iodide turns blue: $2NO_2^- + 2I^- + 4H^+ \rightarrow 2NO + I_2 + 2H_2O$; brown ring with dil $H_2SO_4 + FeSO_4$.

Class II: Concentrated H_2SO_4 Anions ($Cl^-, Br^-, I^-, NO_3^-, CH_3COO^-$)

Treated with concentrated H_2SO_4 and gently heated:

Anion	Observation	Confirmatory Reactions & Identification
Cl^-	Colorless pungent fumes (HCl); white fumes with NH_4OH glass rod.	**Chromyl Chloride Test:** Salt $+K_2Cr_2O_7$ + conc. $H_2SO_4 \xrightarrow{\Delta} CrO_2Cl_2$ (red vapors) $\xrightarrow{NaOH} Na_2CrO_4 \xrightarrow{Pb^{2+}} PbCrO_4\downarrow$ (Yellow).
Br^-	Reddish brown vapors (Br_2) intensifying with MnO_2.	**Organic Layer Test:** Salt $+Cl_2$-water + $CHCl_3 \xrightarrow{shake}$ orange-brown layer ($2Br^- + Cl_2 \rightarrow 2Cl^- + Br_2$).
I^-	Deep violet vapors (I_2) turning starch paper blue.	**Layer Test:** Salt $+Cl_2$-water$+CHCl_3 \rightarrow$ violet layer. $AgNO_3$ gives bright yellow AgI ↓ insoluble in conc NH_4OH.
NO_3^-	Dense brown fumes of NO_2 with Cu turnings.	**Brown Ring Test:** Salt $+FeSO_4 \xrightarrow{\text{trickle conc. } H_2SO_4} [Fe(H_2O)_5(NO)]^{2+}$ (Brown Ring at junction).
CH_3COO^-	Characteristic vinegar odor.	**Ester Test:** Sweet fruity smell ($CH_3COOC_2H_5$). $FeCl_3$ **Test:** Blood-red coloration ($[Fe_3(OH)_2(CH_3COO)_6]^+$); boiling yields brown basic acetate.

Class III: Independent Group Anions (SO_4^{2-}, PO_4^{3-})

Do not evolve gases with dilute or concentrated acids; identified by direct precipitation:

$$\textbf{Sulfate } (SO_4^{2-}): \quad \text{Salt} + BaCl_2 \xrightarrow{\text{dil. HCl}} BaSO_4\downarrow \text{ (White ppt)}$$

$$\textbf{Phosphate } (PO_4^{3-}): \quad \text{Salt} + HNO_3 + (NH_4)_2MoO_4 \xrightarrow{\Delta} (NH_4)_{\mathbf{3}}[P(Mo_3O_{10})_4]\downarrow \textbf{ (Canary Yellow)}$$

7.2 Systematic Qualitative Analysis: Basic Radicals (Cations)

Group	Cations	Group Reagent	Precipitated As (Form & Color)
Group 0	NH_4^+	Dilute NaOH (warm)	Gas $NH_3\uparrow$ (pungent alkaline vapor)
Group I	Pb^{2+}	Dilute HCl	$PbCl_2\downarrow$ (White crystalline ppt)
Group II	$Pb^{2+}, Cu^{2+}, As^{3+}$	H_2S + dil. HCl	PbS, CuS (Black); As_2S_3 (Yellow)
Group III	Fe^{3+}, Al^{3+}	NH_4OH + solid NH_4Cl	$Fe(OH)_3\downarrow$ (Red-brown); $Al(OH)_3\downarrow$ (White gel)
Group IV	$Co^{2+}, Ni^{2+}, Mn^{2+}, Zn^{2+}$	$H_2S + NH_4OH + NH_4Cl$	CoS, NiS (Black); MnS (Buff); ZnS (Dirty white)
Group V	$Ba^{2+}, Sr^{2+}, Ca^{2+}$	$(NH_4)_2CO_3 + NH_4OH + NH_4Cl$	$BaCO_3, SrCO_3, CaCO_3\downarrow$ (White ppt)
Group VI	Mg^{2+}	$Na_2HPO_4 + NH_4OH$	$Mg(NH_4)PO_4\downarrow$ (White crystalline ppt)

Table 7.1: Systematic Overview: Analytical Group Separation of Basic Radicals.

Solubility Product (K_{sp}) Principles & Confirmatory Tests by Group

1. **Group 0 (NH_4^+):**
   - *Precipitation Principle:* Free NH_3 evolved due to base displacement: $NH_4^+ + OH^- \rightarrow NH_3 \uparrow + H_2O$.
   - *Confirmatory Test:* Nessler's reagent ($K_2[HgI_4]$ + KOH) yields brown precipitate of iodide of Millon's base: $H_2N\text{-}Hg\text{-}O\text{-}Hg\text{-}I \downarrow$.
2. **Group I (Pb^{2+}):**
   - *Precipitation Principle:* Lowest K_{sp} among metal chlorides ($K_{sp} \approx 1.7 \times 10^{-5}$).
   - *Confirmatory Test:* $PbCl_2$ dissolves in hot water and recrystallizes as golden spangles on cooling. With K_2CrO_4, forms bright yellow $PbCrO_4 \downarrow$, soluble in excess NaOH.
3. **Group II ($Pb^{2+}, Cu^{2+}, As^{3+}$):**
   - *Precipitation Principle (Common Ion Effect):* H^+ from HCl suppresses H_2S ionization ($H_2S \rightleftharpoons 2H^+ + S^{2-}$), keeping $[S^{2-}]$ extremely low ($\sim 10^{-21}$ M). Only sparingly soluble Group II sulfides exceed K_{sp}.
   - *Confirmatory Test:* Cu^{2+} +excess $NH_4OH \rightarrow [Cu(NH_3)_4]^{2+}$ (Deep Azure Blue); with $K_4[Fe(CN)_6] \rightarrow Cu_2[Fe(CN)_6] \downarrow$ (Chocolate Brown).
4. **Group III (Fe^{3+}, Al^{3+}):**
   - *Precipitation Principle (Common Ion Effect):* NH_4^+ from solid NH_4Cl suppresses NH_4OH ionization, keeping $[OH^-]$ sufficiently low so that only very low K_{sp} hydroxides of Group III precipitate, preventing Group IV/V/VI hydroxides from precipitating.
   - *Confirmatory Test:* $Fe^{3+} + K_4[Fe(CN)_6] \rightarrow Fe_4[Fe(CN)_6]_3 \downarrow$ (Prussian Blue). $Al(OH)_3$ dissolves in excess NaOH to form soluble $NaAlO_2$, reprecipitated with NH_4Cl.
5. **Group IV ($Co^{2+}, Ni^{2+}, Mn^{2+}, Zn^{2+}$):**
   - *Precipitation Principle:* Alkaline medium (NH_4OH) removes H^+, driving H_2S dissociation forward; high $[S^{2-}]$ precipitates the higher-K_{sp} sulfides.
   - *Confirmatory Test:* Ni^{2+} + DMG + $NH_4OH \rightarrow [Ni(DMG)_2] \downarrow$ (Rosy-red chelate). $Zn^{2+} + K_4[Fe(CN)_6] \rightarrow Zn_2[Fe(CN)_6] \downarrow$ (Bluish-white).
6. **Group V ($Ba^{2+}, Sr^{2+}, Ca^{2+}$):**
   - *Precipitation Principle:* Carbonate precipitation in ammoniacal medium; NH_4Cl keeps $[CO_3^{2-}]$ below $K_{sp}(MgCO_3)$.
   - *Confirmatory Test (Flame Test):* Ba^{2+} Apple Green, Sr^{2+} Crimson, Ca^{2+} Brick Red. With K_2CrO_4, Ba^{2+} yields yellow $BaCrO_4 \downarrow$.
7. **Group VI (Mg^{2+}):**
   - *Precipitation Principle:* Direct precipitation with Na_2HPO_4 in ammoniacal solution: $Mg^{2+} + NH_4^+ + PO_4^{3-} \rightarrow Mg(NH_4)PO_4 \downarrow$.
   - *Confirmatory Test:* White crystalline precipitate; scratching the inner test tube wall with a glass rod accelerates crystallization.

High-Yield Confirmatory Tests & Diagnostic Colorations

1. Nickel Ion (Ni^{2+}) with Dimethylglyoxime (DMG): Rosy-red precipitate stabilized by intramolecular hydrogen bonding:

$$Ni^{2+} + 2\,DMG + 2\,NH_4OH \longrightarrow [Ni(DMG)_2]\downarrow (\text{Rosy Red}) + 2\,NH_4^+ + 2\,H_2O \quad (7.1)$$

2. Ferric Ion (Fe^{3+}) Confirmatory Tests:

$$\text{Ferrocyanide: } 4Fe^{3+} + 3[Fe(CN)_6]^{4-} \longrightarrow Fe_4[Fe(CN)_6]_{\mathbf{3}}\downarrow (\textbf{Prussian Blue})$$
$$\text{Thiocyanate: } Fe^{3+} + SCN^- \longrightarrow [Fe(SCN)]^{\mathbf{2+}} (\textbf{Blood Red})$$

3. Copper Ion (Cu^{2+}) Confirmatory Tests:

$$\text{Excess } NH_4OH: \quad Cu^{2+} + 4NH_3 \longrightarrow [Cu(NH_3)_4]^{\mathbf{2+}} (\textbf{Azure Blue})$$
$$\text{Ferrocyanide: } 2Cu^{2+} + [Fe(CN)_6]^{4-} \longrightarrow Cu_2[Fe(CN)_6]\downarrow (\textbf{Chocolate Brown})$$

4. Flame Test Colors of Alkaline Earth Metals:

$$Ba^{2+} \implies \text{Apple Green}, \quad Sr^{2+} \implies \text{Crimson}, \quad Ca^{2+} \implies \text{Brick Red} \quad (7.2)$$

7.3 Volumetric Quantitative Redox Titrations

$KMnO_4$ vs Mohr's Salt Titration (Standardization)

Mohr's salt is a primary standard double salt: $(NH_4)_{\mathbf{2}}SO_4 \cdot FeSO_4 \cdot \mathbf{6}H_2O$ (M.W. $= 392\,g/mol$).

$$\text{Oxidation Half: } 5[Fe^{2+} \longrightarrow Fe^{3+} + e^-]$$
$$\text{Reduction Half: } MnO_4^- + 8H^+ + 5e^- \longrightarrow Mn^{2+} + 4H_2O$$
$$\textbf{Overall Net Redox: } MnO_4^- + \mathbf{5}Fe^{2+} + \mathbf{8}H^+ \longrightarrow Mn^{2+} + \mathbf{5}Fe^{3+} + \mathbf{4}H_2O$$

$$\frac{\mathbf{M_1V_1}}{\mathbf{1}}(KMnO_4) = \frac{\mathbf{M_2V_2}}{\mathbf{5}}(\text{Mohr's Salt}) \implies \mathbf{N_1V_1 = N_2V_2} \quad (7.3)$$

(Valence factor: $n = 5$ for $KMnO_4$, $n = 1$ for Fe^{2+})

Crucial Practical Precautions:

- **Acidifying Agent:** Strictly use **dilute** H_2SO_4.
  - **Why NOT** HCl**?** Permanganate oxidizes HCl to chlorine gas ($2MnO_4^- + 16HCl \rightarrow 2Mn^{2+} + 5Cl_2\uparrow + 8H_2O + 6Cl^-$), artificially consuming excess $KMnO_4$!

  - **Why NOT** HNO_3**?** Nitric acid is a strong oxidizing agent itself and competes with $KMnO_4$ in oxidizing Fe^{2+}.
- **Indicator:** $KMnO_4$ acts as a **self-indicator**. The endpoint is marked by the appearance of a **permanent faint pink color** (excess single drop of MnO_4^-).

$KMnO_4$ vs Oxalic Acid Titration

Oxalic acid is a primary standard dihydrate: $(COOH)_{\mathbf{2}} \cdot \mathbf{2}H_2O$ (M.W. = 126 g/mol, Equivalent Weight = 63 g/eq).

$$\begin{aligned} \text{Oxidation Half: } & 5[C_2O_4^{2-} \longrightarrow 2CO_2 \uparrow +2e^-] \\ \text{Red. Half: } & 2[MnO_4^- + 8H^+ + 5e^- \longrightarrow Mn^{2+} + 4H_2O] \\ \textbf{Net Redox: } & 2MnO_4^- + 5C_2O_4^{2-} + 16H^+ \longrightarrow \\ & 2Mn^{2+} + 10CO_2 \uparrow +8H_2O \end{aligned}$$

$$\frac{\mathbf{M_1V_1}}{\mathbf{2}}(KMnO_4) = \frac{\mathbf{M_2V_2}}{\mathbf{5}}(\text{Oxalic Acid}) \implies \mathbf{N_1V_1 = N_2V_2} \tag{7.4}$$

(Valence factor: $n = 5$ for $KMnO_4$, $n = 2$ for $C_2O_4^{2-}$)

Crucial Practical Precautions:

- **Pre-Heating Requirement:** Oxalic acid solution must be heated to 60 – 70°C before starting titration. At room temperature, the reaction is extremely slow.
- **Autocatalysis:** The initial disappearance of pink color is slow, but once the first few drops react, the reaction becomes rapid because product Mn^{2+} **acts as an autocatalyst**!
- **Do NOT boil (**> 70°C**):** High temperature causes thermal decomposition of oxalic acid into CO_2, CO, and H_2O.

Appendix: Chemistry Formula, Reaction & Reagent Matrices

APPENDIX A

Physical Chemistry Core Formulas & Thermodynamic Equations

This chapter compiles all mathematical formulations, equilibrium relations, colligative equations, electrochemical relations, and kinetic rate laws for rapid revision before major examinations.

A.1 Some Basic Concepts in Chemistry: Mole Concept (Unit 1)

A.1.1 Stoichiometric Mole Formulations

Fundamental Mole Formulations & Gas Volumes

$$n = \frac{w}{M} = \frac{N}{N_A} = \frac{V_{\text{STP}}}{22.4\,\text{L}} \tag{A.1}$$

$$N_A = 6.022 \times 10^{23}\,\text{mol}^{-1}, \quad 1\,\text{u} \approx 1.66 \times 10^{-27}\,\text{kg} \tag{A.2}$$

$$\text{VD} = \frac{\rho_{\text{gas}}}{\rho_{\text{H}_2}} \implies \text{Molar Mass} = 2 \times \text{VD} \tag{A.3}$$

$$\text{Mass \% } X = \frac{(\text{Atoms of } X) \times (\text{Atomic mass})}{\text{Molar mass}} \times 100 \tag{A.4}$$

Limiting Reagent Condition for Reaction $aA + bB \to cC + dD$:

$$\text{If } \frac{n_A}{a} < \frac{n_B}{b} \implies A \text{ is the limiting reagent (completely consumed).} \tag{A.5}$$

A.1.2 Solution Concentrations & Dilution Formulations

Molarity, Molality, Normality & Interconversions

$$M = \frac{n_{\text{solute}}}{V\ (\text{L})} = \frac{w_B \times 1000}{M_B \times V\ (\text{mL})} \quad [\text{Molarity; } T\text{-dependent}] \tag{A.6}$$

$$m = \frac{n_{\text{solute}}}{W_A\ (\text{kg})} = \frac{w_B \times 1000}{M_B \times W_A\ (\text{g})} \quad [\text{Molality; } T\text{-indep.}] \tag{A.7}$$

$$N = M \times n\text{-factor}, \qquad x_A + x_B = 1 \tag{A.8}$$

$$\text{ppm} = \frac{w_{\text{comp}}}{w_{\text{sol}}} \times 10^6, \qquad \%(w/w) = \frac{w_{\text{solute}}}{w_{\text{solution}}} \times 100 \tag{A.9}$$

High-Yield Interconversion Formulas (d = density in g mL^{-1}):

$$M = \frac{10 \times \%(w/w) \times d}{M_B}, \quad m = \frac{1000M}{1000d - MM_B} \tag{A.10}$$

$$m = \frac{x_B \times 1000}{(1 - x_B)M_A} \tag{A.11}$$

Dilution and Mixing of Solutions:

$$M_1V_1 = M_2V_2, \qquad M_{\text{mix}} = \frac{\sum M_iV_i}{\sum V_i} \tag{A.12}$$

$$N_1V_1 = N_2V_2 \quad [\text{at equivalence point}] \tag{A.13}$$

Mole Traps: STP Standards & Temperature Dependence

- **STP Standard Volume:** In IUPAC revised standard conditions ($T = 273.15\,\text{K}, P = 1\,\text{bar} = 10^5\,\text{Pa}$), molar volume is 22.7 L. Under traditional NCERT standard pressure ($1\,\text{atm} = 1.01325 \times 10^5\,\text{Pa}$), molar volume is 22.4 L. Unless specified as 1 bar, calculate using 22.4 L.
- **Molarity Changes with Temperature:** Because volume expands on heating ($V \propto T$), molarity M and normality N **decrease with increasing temperature**. Molality m, mole fraction x, and $\%(w/w)$ involve mass only and remain **strictly constant**.

A.2 Structure of Atom (Unit 2)

A.2.1 Quantum Mechanics, Wave-Particle Duality & Uncertainty

de Broglie, Heisenberg Uncertainty & Orbital Angular Momentum

$$\lambda = \frac{h}{p} = \frac{h}{mv} = \frac{h}{\sqrt{2mK}} = \frac{h}{\sqrt{2mqV}} \tag{A.14}$$

$$\Delta x \cdot \Delta p \geq \frac{h}{4\pi} \implies \Delta x \cdot \Delta v \geq \frac{h}{4\pi m} \quad \text{[Uncertainty]} \tag{A.15}$$

$$\Delta E \cdot \Delta t \geq \frac{h}{4\pi}, \qquad L_{\text{orb}} = \sqrt{l(l+1)}\hbar \tag{A.16}$$

$$\mu_{\text{spin}} = \sqrt{n(n+2)}\,\text{BM} \quad [n = \text{unpaired } e^-] \tag{A.17}$$

A.2.2 Quantum Numbers, Nodes & Electron Filling Principles

Quantum No.	Symbol	Allowed Values	Physical Significance
Principal	n	$1, 2, 3, \ldots$	Main energy level, shell size
Azimuthal	l	$0, \ldots, n-1$	Subshell shape $(0:s, 1:p, 2:d, 3:f)$
Magnetic	m_l	$-l, \ldots, +l$	Spatial orientation in field
Spin	m_s	$\pm 1/2$	Intrinsic spin orientation

Table A.1: The Four Quantum Numbers and Boundary Constraints.

Nodal Formulations for Atomic Orbitals

$$\text{Radial Nodes} = n - l - 1, \quad \text{Angular Nodes} = l \tag{A.18}$$

$$\text{Total Nodes} = (n - l - 1) + l = n - 1 \tag{A.19}$$

$$\text{Max } e^- \text{ in Shell } n = 2n^2, \quad \text{Orbitals in Shell } n = n^2 \tag{A.20}$$

$$\text{Max } e^- \text{ in Subshell } l = 2(2l+1) = 4l + 2 \tag{A.21}$$

Orbital Filling Rules & Exchange Energy Stability

- **Aufbau Principle ($(n+l)$ Rule):** Electrons enter lowest energy subshell first. Lower $(n+l)$ has lower energy. If $(n+l)$ identical, lower n has lower energy (e.g. $4s$ has $4+0=4$, $3d$ has $3+2=5 \implies 4s$ fills before $3d$).

- **Pauli Exclusion Principle:** No two electrons in an atom can have the identical set of all four quantum numbers. An orbital holds maximum 2 electrons with opposite spins.
- **Hund's Rule of Maximum Multiplicity:** Pairing does not occur in degenerate orbitals until each holds one electron with parallel spin.
- **Exchange Energy of Half/Fully Filled Subshells:** Number of possible electron exchanges among electrons of parallel spin:

$$K = \frac{n_p(n_p - 1)}{2} \implies \text{For } d^5 : K = \frac{5 \times 4}{2} = 10 \text{ exchanges}$$

A.3 Chemical Thermodynamics & Thermochemistry (Unit 4)

A.3.1 First Law, Work, Enthalpy & Heat Capacities

First Law, PV-Work & Enthalpy-Internal Energy Coupling

$$\Delta U = q + w \quad [\text{Work on system } > 0; \text{by system } < 0] \tag{A.22}$$

$$w_{\text{irr}} = -P_{\text{ext}}\Delta V \quad [\text{Irreversible against constant } P_{\text{ext}}] \tag{A.23}$$

$$w_{\text{rev, iso}} = -nRT\ln(V_2/V_1) = -2.303nRT\log_{10}(P_1/P_2) \tag{A.24}$$

$$w_{\text{free}} = 0\ (P_{\text{ext}} = 0) \implies \Delta U = 0,\ \Delta T = 0 \tag{A.25}$$

$$\Delta H = \Delta U + \Delta(PV) = \Delta U + \Delta n_g RT \tag{A.26}$$

$$C_p - C_v = R \implies \Delta H = q_p = nC_p\Delta T, \quad \Delta U = q_v = nC_v\Delta T \tag{A.27}$$

A.3.2 Thermochemistry & Hess's Law

Hess's Law, Reaction Enthalpy & Bond Dissociation Enthalpies

$$\Delta_r H^\circ = \sum \Delta_f H^\circ(\text{Products}) - \sum \Delta_f H^\circ(\text{Reactants}) \tag{A.28}$$

$$\Delta_r H^\circ = \sum BE(\text{Reactants}) - \sum BE(\text{Products}) \quad [\text{all gaseous}] \tag{A.29}$$

$$\Delta_r H = \sum \Delta_{\text{comb}} H^\circ(\text{Reactants}) - \sum \Delta_{\text{comb}} H^\circ(\text{Products}) \tag{A.30}$$

Kirchhoff's Law of Temperature Dependence of Enthalpy:

$$\frac{\Delta H_{T_2} - \Delta H_{T_1}}{T_2 - T_1} = \Delta C_p, \qquad \frac{\Delta U_{T_2} - \Delta U_{T_1}}{T_2 - T_1} = \Delta C_v \tag{A.31}$$

$$\Delta C_p = \sum C_{p,\text{prod}} - \sum C_{p,\text{react}} \tag{A.32}$$

A.3.3 Entropy, Second Law & Gibbs Free Energy Spontaneity

Entropy, Third Law & Spontaneity Criteria

$$\Delta S_{\text{total}} = \Delta S_{\text{system}} + \Delta S_{\text{surroundings}} \geq 0 \tag{A.33}$$

$$\Delta S_{\text{sys}} = nC_v \ln(T_2/T_1) + nR\ln(V_2/V_1) \tag{A.34}$$
$$= nC_p \ln(T_2/T_1) - nR\ln(P_2/P_1) \tag{A.35}$$
$$\Delta S_{\text{iso}} = 2.303nR\log_{10}(P_1/P_2), \quad \Delta S_{\text{isochoric}} = nC_v \ln(T_2/T_1) \tag{A.36}$$
$$\Delta G = \Delta H - T\Delta S, \quad \Delta G^\circ = -2.303RT\log_{10} K \tag{A.37}$$

ΔH	ΔS	$\Delta G = \Delta H - T\Delta S$	**Spontaneity Criterion**
$-$	$+$	Always Negative $(-)$	Spontaneous at ALL temperatures
$+$	$-$	Always Positive $(+)$	Non-spontaneous at ALL temperatures (reverse is spontaneous)
$-$	$-$	Negative at low T $(T < \frac{\Delta H}{\Delta S})$	Spontaneous at LOW temperature; non-spontaneous at high T
$+$	$+$	Negative at high T $(T > \frac{\Delta H}{\Delta S})$	Spontaneous at HIGH temperature; non-spontaneous at low T

Table A.2: Spontaneity Matrix Governed by Enthalpy and Entropy Interplay.

Thermodynamics Traps: Standard States & Bond Enthalpy Reversal

- **Bond Enthalpy Subtraction Order:** Enthalpy of reaction using formation enthalpies is [Products − Reactants], but using bond enthalpies it is [**Reactants** − **Products**]!
- **Reference Standard State Formation Enthalpy is Zero:** Only the most stable allotropic form at 298.15 K and 1 bar has $\Delta_f H^\circ = 0$: Carbon is **Graphite** (Diamond has $\Delta_f H^\circ = 1.9\,\text{kJ mol}^{-1}$), Sulfur is **Rhombic**, Phosphorus is **White**.

A.4 Equilibrium (Unit 6)

A.4.1 Chemical Equilibrium, $K_p - K_c$ Coupling & Le Chatelier

Equilibrium Constants, Reaction Quotient & Degree of Dissociation

$$K_p = K_c(RT)^{\Delta n_g} \quad [R = 0.0831\,\text{L bar mol}^{-1}\,\text{K}^{-1}] \tag{A.38}$$

$$Q < K \text{ (Forward)}, \quad Q = K \text{ (Equilibrium)}, \quad Q > K \text{ (Backward)} \tag{A.39}$$

$$\alpha = \frac{D - d}{(n-1)d} = \frac{M_{\text{theor}} - M_{\text{obs}}}{(n-1)M_{\text{obs}}} \tag{A.40}$$

van 't Hoff Equation:

$$\log_{10}\left(\frac{K_2}{K_1}\right) = \frac{\Delta H^\circ}{2.303R}\left[\frac{T_2 - T_1}{T_1 T_2}\right] = \frac{\Delta H^\circ}{2.303R}\left[\frac{1}{T_1} - \frac{1}{T_2}\right] \tag{A.41}$$

A.4.2 Ionic Equilibrium: Ostwald Dilution, Water Autoionization & pH

Ostwald's Dilution Law, Water Ion Product & Acid-Base pH

$$K_a \approx C\alpha^2 \implies \alpha = \sqrt{\frac{K_a}{C}}, \quad [\text{H}^+] = \sqrt{K_a C} \quad (\alpha \leq 0.05) \tag{A.42}$$

$$K_w = [\text{H}^+][\text{OH}^-] = 1.0 \times 10^{-14} \implies \text{pH} + \text{pOH} = 14 \tag{A.43}$$

$$\text{pH} = -\log_{10}[\text{H}^+], \qquad [\text{H}^+] = 10^{-\text{pH}} \tag{A.44}$$

A.4.3 Salt Hydrolysis Formulations

A.4.4 Buffer Solutions & Henderson-Hasselbalch Equations

Acidic and Basic Buffer Governing Equations

$$\text{pH} = \text{p}K_a + \log_{10}\left(\frac{[\text{Salt}]}{[\text{Acid}]}\right) \quad [\text{Acidic Buffer}] \tag{A.45}$$

Salt Type	K_h	h	pH at 25°C
SA + SB (NaCl)	None	0	7.00
WA + SB (CH_3COONa)	$\frac{K_w}{K_a}$	$\sqrt{\frac{K_w}{CK_a}}$	$7 + \frac{1}{2}pK_a + \frac{1}{2}\log C$
SA + WB (NH_4Cl)	$\frac{K_w}{K_b}$	$\sqrt{\frac{K_w}{CK_b}}$	$7 - \frac{1}{2}pK_b - \frac{1}{2}\log C$
WA + WB (CH_3COONH_4)	$\frac{K_w}{K_aK_b}$	$\sqrt{\frac{K_w}{K_aK_b}}$	$7 + \frac{1}{2}pK_a - \frac{1}{2}pK_b$

Table A.3: Exact Hydrolysis Parameters and pH Formulations for Canonical Salts.

$$\text{pOH} = pK_b + \log_{10}\left(\frac{[\text{Salt}]}{[\text{Base}]}\right), \quad \text{pH} = 14 - \text{pOH} \tag{A.46}$$

$$\text{Buffer Range} = pK_a \pm 1 \tag{A.47}$$

A.4.5 Solubility Product (K_{sp}) & Precipitation Criteria

Solubility Product Equations & Precipitation

$$A_xB_y \rightleftharpoons xA^{y+} + yB^{x-} \implies K_{sp} = (xS)^x(yS)^y = x^x y^y S^{x+y} \tag{A.48}$$

Canonical Stoichiometric Solubility Forms:

- 1 : 1 Salt ($AgCl$): $K_{sp} = S^2 \implies S = \sqrt{K_{sp}}$
- 1 : 2 Salt ($PbCl_2, Ag_2CrO_4$): $K_{sp} = 4S^3 \implies S = \sqrt[3]{K_{sp}/4}$
- 1 : 3 Salt ($Al(OH)_3$): $K_{sp} = 27S^4 \implies S = \sqrt[4]{K_{sp}/27}$
- 2 : 3 Salt ($Ca_3(PO_4)_2$): $K_{sp} = 108S^5 \implies S = \sqrt[5]{K_{sp}/108}$

Precipitation Criterion:

$$Q_{sp} < K_{sp} \text{ (Unsaturated)}; \quad Q_{sp} = K_{sp} \text{ (Saturated)} \tag{A.49}$$

$$Q_{sp} > K_{sp} \text{ (Precipitates)} \tag{A.50}$$

Ionic Equilibrium Traps: Autoionization of Water & pH Scaling

- **Low Concentration Acid pH (10^{-8} M HCl):** Never write pH = 8 for an acid! In extremely dilute solutions ($< 10^{-6}$ M), autoionization of water ($[H^+]_{\text{water}} = 10^{-7}$ M) cannot be neglected:

$$[H^+]_{\text{total}} = 10^{-8} + 10^{-7} = 1.1 \times 10^{-7}\,\text{M} \implies \text{pH} \approx 6.96 \quad (\text{strictly } < 7)$$

- **Temperature Dependence of K_w:** The autoionization of water ($H_2O \rightleftharpoons H^+ + OH^-$) is endothermic ($\Delta H > 0$). Heating water to 60°C increases $K_w \approx 10^{-13}$. Neutral water at 60°C has pH = $\frac{1}{2}pK_w = 6.5$, but remains strictly **neutral** because $[H^+] = [OH^-]$.

A.5 Solutions & Colligative Properties (Unit 5)

A.5.1 Gas Solubility & Raoult's Law

Henry's Law, Raoult's Law & Vapour Pressure

$$p = K_H \cdot x \implies x = \frac{p}{K_H} \quad \text{[Henry's Law]} \tag{A.51}$$

$$P_{\text{total}} = P_A + P_B = P_A^\circ x_A + P_B^\circ x_B = P_A^\circ + (P_B^\circ - P_A^\circ)x_B \tag{A.52}$$

$$y_A = \frac{P_A}{P_{\text{total}}}, \qquad \frac{1}{P_{\text{total}}} = \frac{y_A}{P_A^\circ} + \frac{y_B}{P_B^\circ} \tag{A.53}$$

A.5.2 Colligative Properties & van 't Hoff Factor

The Four Colligative Properties & van 't Hoff Factor (i)

$$\frac{P_A^\circ - P}{P_A^\circ} = i\, x_B \approx i \left(\frac{w_B M_A}{M_B w_A} \right) \quad \text{[RLVP]} \tag{A.54}$$

$$\Delta T_b = i\, K_b\, m = i \left(\frac{1000 K_b w_B}{M_B w_A} \right) \quad \text{[Boiling Point]} \tag{A.55}$$

$$\Delta T_f = i\, K_f\, m = i \left(\frac{1000 K_f w_B}{M_B w_A} \right) \quad \text{[Freezing Point]} \tag{A.56}$$

$$\Pi = i\, C\, R\, T = i \left(\frac{w_B R T}{M_B V} \right) \quad \text{[Osmotic Pressure]} \tag{A.57}$$

van 't Hoff Factor (i) Formulations:

$$i = \frac{\text{Observed Colligative Property}}{\text{Calculated Property}} = \frac{\text{Normal } M}{\text{Observed } M} \tag{A.58}$$

$$\alpha_{\text{dissoc}} = \frac{i - 1}{n - 1} \implies i = 1 + (n-1)\alpha \tag{A.59}$$

$$\alpha_{\text{assoc}} = \frac{1 - i}{1 - 1/n} \implies i = 1 - \left(1 - \frac{1}{n}\right)\alpha \tag{A.60}$$

Solutions Traps: Molality in RLVP & Abnormal Molar Mass

- **RLVP with Molality Shortcut:** For an aqueous solution, relative lowering

of vapour pressure can be directly expressed in terms of molality m:

$$\frac{P^\circ - P}{P} = \frac{m \times M_{\text{solvent}}}{1000} \implies \text{For water: } \frac{P^\circ - P}{P} = \frac{18m}{1000}$$

Notice the denominator is P (solution vapour pressure), NOT P°!

- **Elevation vs Depression Constant:** For water, $K_b = 0.52\,\text{K kg mol}^{-1}$, whereas $K_f = 1.86\,\text{K kg mol}^{-1}$. Freezing point depression is roughly 3.6 times more sensitive than boiling point elevation for the identical molal concentration.

A.6 Electrochemistry (Unit 7)

A.6.1 Galvanic Cells, Nernst Equation & Equilibrium

EMF, Nernst Equation & Free Energy Formulations

$$E^\circ_{\text{cell}} = E^\circ_{\text{cathode}} - E^\circ_{\text{anode}} = E^\circ_{\text{right}} - E^\circ_{\text{left}} \tag{A.61}$$

$$E_{\text{cell}} = E^\circ_{\text{cell}} - \frac{2.303RT}{nF}\log_{10} Q \tag{A.62}$$

$$E_{\text{cell}} = E^\circ_{\text{cell}} - \frac{0.0591}{n}\log_{10} Q \quad (\text{at } 298.15\,\text{K}) \tag{A.63}$$

$$\log_{10} K_c = \frac{nE^\circ_{\text{cell}}}{0.0591} \quad [\text{at equilibrium } E_{\text{cell}} = 0] \tag{A.64}$$

$$\Delta_r G = -nFE_{\text{cell}}, \quad \Delta_r G^\circ = -nFE^\circ_{\text{cell}} = -2.303RT\log_{10} K_c \tag{A.65}$$

Concentration Cells (Identical Electrodes, Different Concentrations $C_1 < C_2$):

$$E^\circ_{\text{cell}} = 0 \implies E_{\text{cell}} = \frac{0.0591}{n}\log_{10}\left(\frac{C_2}{C_1}\right) \quad (\text{Spontaneous if } C_2 > C_1) \tag{A.66}$$

A.6.2 Electrolytic Conductance & Kohlrausch's Law

Conductivity, Molar Conductivity & Kohlrausch Formulations

$$G = \frac{1}{R}, \quad \kappa = G\left(\frac{L}{A}\right) = \frac{G^*}{R} \quad [G^* = L/A] \tag{A.67}$$

$$\Lambda_m = \frac{1000\,\kappa}{M}, \quad \Lambda_{\text{eq}} = \frac{1000\,\kappa}{N} = \frac{\Lambda_m}{n\text{-factor}} \tag{A.68}$$

$$\Lambda_m = \Lambda^\circ_m - A\sqrt{C} \quad [\text{Debye-Hückel-Onsager}] \tag{A.69}$$

$$\Lambda^\circ_m(A_xB_y) = x\lambda^\circ_m(A^{y+}) + y\lambda^\circ_m(B^{x-}) \quad [\text{Kohlrausch}] \tag{A.70}$$

$$\alpha = \frac{\Lambda_m}{\Lambda^\circ_m}, \quad K_a = \frac{C\alpha^2}{1-\alpha}, \quad S = \frac{1000\,\kappa_{\text{salt}}}{\Lambda^\circ_m} \tag{A.71}$$

A.6.3 Faraday's Laws of Electrolysis

Faraday's First & Second Laws of Electrolysis

$$w = Z\,I\,t = \left(\frac{E}{96500}\right) I\,t = \left(\frac{M}{n \times 96500}\right) I\,t \tag{A.72}$$

$$\frac{w_1}{w_2} = \frac{E_1}{E_2} \quad \text{(for identical charge } Q\text{)} \tag{A.73}$$

Electrochemistry Traps: Intensive E° vs Extensive ΔG°

- **Electrode Potentials are Intensive Properties:** Never multiply E° by stoichiometric coefficients when balancing electrons! For half-reaction multiplied by 2: ΔG° doubles, but E° remains **unchanged**. When combining non-standard half-cells, always combine $\Delta G^\circ = -nFE^\circ$, never add potentials directly!
- **Conductivity vs Molar Conductivity on Dilution:** On dilution: Specific conductivity κ **decreases** (fewer ions per unit volume), but molar conductivity Λ_m **increases** (interionic attractions weaken for strong electrolytes; α increases for weak electrolytes).

A.7 Chemical Kinetics (Unit 8)

A.7.1 Rate Laws, Order & Molecularity

Reaction Rates & General Rate Expressions

For general reaction: $aA + bB \rightarrow cC + dD$:

$$\text{Rate} = -\frac{1}{a}\frac{d[A]}{dt} = -\frac{1}{b}\frac{d[B]}{dt} = +\frac{1}{c}\frac{d[C]}{dt} \tag{A.74}$$

$$\text{Rate Law} = k[A]^x[B]^y \implies \text{Order } n = x + y \tag{A.75}$$

$$\text{Units of } k = (\text{mol}\,\text{L}^{-1})^{1-n}\,\text{s}^{-1} \tag{A.76}$$

A.7.2 Integrated Rate Equations: Zero, First & n-th Order

Order	Rate Law	Integrated Form	Half-Life ($t_{1/2}$)	Completion
Zero ($n = 0$)	$-\frac{d[A]}{dt} = k$	$[A]_t = [A]_0 - kt$	$\frac{[A]_0}{2k} \propto [A]_0$	$t_{100\%} = \frac{[A]_0}{k}$
First ($n = 1$)	$-\frac{d[A]}{dt} = k[A]$	$k = \frac{2.303}{t}\log\frac{[A]_0}{[A]_t}$	$\frac{0.693}{k}$	$t_{100\%} \to \infty$
Second ($n = 2$)	$-\frac{d[A]}{dt} = k[A]^2$	$\frac{1}{[A]_t} - \frac{1}{[A]_0} = kt$	$\frac{1}{k[A]_0} \propto \frac{1}{[A]_0}$	$t_{100\%} \to \infty$
n-th	$-\frac{d[A]}{dt} = k[A]^n$	$\frac{1}{[A]_t^{n-1}} - \frac{1}{[A]_0^{n-1}} = (n-1)kt$	$t_{1/2} \propto \frac{1}{[A]_0^{n-1}}$	—

Table A.4: Systematic Summary of Integrated Rate Formulations and Half-Life Scaling.

First-Order Fractional Lifetimes Shortcuts

For any first-order process:

$$t_{50\%} = 1 \times t_{1/2} = \frac{0.693}{k}$$

$$t_{75\%} = 2 \times t_{1/2} \quad (75\% \text{ consumed, } 25\% \text{ remaining})$$

$$t_{87.5\%} = 3 \times t_{1/2} \quad (87.5\% \text{ consumed, } 12.5\% \text{ remaining})$$

$$t_{99\%} = 2 \times t_{90\%} \approx 6.64 \times t_{1/2} \approx \frac{4.606}{k}$$

$$t_{99.9\%} = 10 \times t_{1/2} \approx 3 \times t_{90\%} = \frac{6.909}{k}$$

A.7.3 Temperature Dependence: Arrhenius Equation & Catalysis

Arrhenius Equation, Activation Energy & Collision Theory

$$k = A\,e^{-E_a/RT} \implies \ln k = \ln A - \frac{E_a}{RT} \tag{A.77}$$

$$\log_{10} k = \log_{10} A - \frac{E_a}{2.303RT} \tag{A.78}$$

$$\log_{10}\left(\frac{k_2}{k_1}\right) = \frac{E_a}{2.303R}\left[\frac{T_2 - T_1}{T_1 T_2}\right] \tag{A.79}$$

$$\text{Slope} = -\frac{E_a}{2.303R}, \quad \text{Intercept} = \log_{10} A \tag{A.80}$$

$$\Delta_r H = E_{a,\text{forward}} - E_{a,\text{backward}}, \quad \text{Rate} = PZ_{AB}e^{-E_a/RT} \tag{A.81}$$

Kinetics Traps: Molecularity vs Order & Catalyst Action

- **Molecularity is NEVER Zero or Fractional:** Molecularity is the number of reacting species colliding simultaneously in an elementary step; it is strictly an integer $(1, 2, 3)$. Reactions with molecularity > 3 are statistically rare. Order is experimentally determined and can be 0, fractional, or negative.
- **Catalysts Do NOT Alter Equilibrium State:** A catalyst accelerates both forward and reverse reaction rates by the identical factor by providing an alternate reaction pathway with lower activation energy ($E_a' < E_a$). It does **not** alter $\Delta_r H$, $\Delta_r G^\circ$, or the equilibrium constant K_c.

APPENDIX B

Inorganic Trends & Organic Named Reactions Compendium

This quick-reference compendium brings together inorganic electronic configurations, crystal field splitting patterns, and all 35+ prescribed organic named reactions, complete with reagents, conditions, and functional distinction tests.

B.1 Classification of Elements & Periodicity (Unit 9)

B.1.1 Effective Nuclear Charge & Atomic Radii Trends

Slater's Rules, Screening Constant & Radius Scaling

$$Z_{\text{eff}} = Z - \sigma \quad [\sigma = \text{Shielding constant}] \tag{B.1}$$

$$\text{Shielding} : s > p > d > f \quad [d, f \text{ are very poor}] \tag{B.2}$$

$$r_{\text{cation}} < r_{\text{atom}} < r_{\text{anion}} \quad \left[\text{Size: } O^{2-} > F^{-} > Na^{+} > Mg^{2+} > Al^{3+}\right] \tag{B.3}$$

B.1.2 Ionization Enthalpy, Electron Gain Enthalpy & Electronegativity

Property	General Trend	High-Yield Exceptions
IE	Across: Inc. Down: Dec.	Be > B ($2s^2$ vs $2p^1$); N > O ($2p^3$ vs $2p^4$); Mg > Al; Tl > In.
$\Delta_{\mathbf{eg}}H$	Across: More Neg.	Cl > F and S > O (compact $2p$ repulsions in F, O).
χ	Across: Inc. Down: Dec.	F(4.0) > O(3.5) > N ≈ Cl(3.0) > Br(2.8) > I ≈ C ≈ S(2.5).

Table B.1: Systematic Summary of Periodic Trends and Anomaly Anchors.

Electronegativity Scales & Dipole Moment Formulations

$$\chi_{\text{Mulliken}} = \frac{\text{IE} + \text{EA}}{2}\ (\text{eV}), \quad \chi_{\text{Pauling}} = \frac{\text{IE} + \text{EA}}{5.6} \tag{B.4}$$

$$\mu = q \times d, \quad \%\,\text{Ionic} = \frac{\mu_{\text{obs}}}{\mu_{\text{calc}}} \times 100\% \tag{B.5}$$

$$\mu_{\text{net}} = \sqrt{\mu_1^2 + \mu_2^2 + 2\mu_1\mu_2\cos\theta} \implies \mu_{\text{ortho}} > \mu_{\text{meta}} > \mu_{\text{para}} = 0 \tag{B.6}$$

Periodic Traps: NH_3 vs NF_3 Dipole Moment & Electron Gain Enthalpy

- **Dipole Moment Comparison ($\mu_{\mathbf{NH_3}} > \mu_{\mathbf{NF_3}}$):** In NH_3, bond dipoles of three N − H bonds reinforce the lone-pair dipole ($\mu = 1.47\,\text{D}$). In NF_3, the highly electronegative fluorines pull electron density away, opposing the lone-pair dipole ($\mu = 0.23\,\text{D}$).
- **Most Negative $\Delta_{\mathbf{eg}}H$:** The element with the **most negative electron gain enthalpy** in the entire periodic table is **Chlorine (Cl)**, NOT Fluorine (Cl : $-349\,\text{kJ mol}^{-1}$ vs F : $-328\,\text{kJ mol}^{-1}$).

B.2 Chemical Bonding & Molecular Structure (Unit 3)

B.2.1 Formal Charge & VSEPR Theory Master Geometry Table

Formal Charge Equation & Steric Number Rule

$$\text{FC} = V - L - \frac{1}{2}S \quad [V = \text{Valence}, L = \text{Lone}, S = \text{Shared}] \tag{B.7}$$

$$SN = (\sigma\text{-bonds}) + (\text{Lone pairs on central atom}) \tag{B.8}$$

SN	Hyb.	BP	LP	Geometry	Examples
2	sp	2	0	Linear	$BeCl_2, CO_2, HCN$
3	sp^2	3	0	Trig. Planar	BF_3, BCl_3, SO_3
3	sp^2	2	1	Bent	SO_2, O_3, NO_2^-
4	sp^3	4	0	Tetrahedral	CH_4, NH_4^+, SO_4^{2-}
4	sp^3	3	1	Trig. Pyramidal	NH_3, PCl_3, H_3O^+
4	sp^3	2	2	Bent (104.5°)	H_2O, H_2S, OF_2
5	sp^3d	5	0	Trig. Bipyramidal	PCl_5, PF_5
5	sp^3d	4	1	Seesaw	SF_4, SeF_4
5	sp^3d	3	2	T-Shaped	ClF_3, BrF_3, ICl_3
5	sp^3d	2	3	Linear	XeF_2, I_3^-, ICl_2^-
6	sp^3d^2	6	0	Octahedral	$SF_6, [AlF_6]^{3-}$
6	sp^3d^2	5	1	Sq. Pyramidal	$BrF_5, IF_5, XeOF_4$
6	sp^3d^2	4	2	Square Planar	$XeF_4, [ICl_4]^-$
7	sp^3d^3	7	0	Pent. Bipyramidal	IF_7
7	sp^3d^3	6	1	Dist. Octahedral	XeF_6

Table B.2: Master VSEPR Geometries, Hybridization and Lone Pair Partitions.

Bent's Rule Formulations in Trigonal Bipyramidal (sp^3d)

- **Bent's Postulate:** More electronegative substituents prefer hybrid orbitals having **less s-character** (Axial bonds in TBP: pure pd character, longer and weaker).
- **Lone Pair Preference:** Lone pairs and electropositive substituents prefer hybrid orbitals having **more s-character** (Equatorial bonds in TBP: sp^2 character, shorter and stronger).
- **Application to PCl_5 and ClF_3:** In gaseous PCl_5, axial P – Cl bonds (214 pm) are longer than equatorial bonds (202 pm). In solid state, it dissociates into ionic $[PCl_4]^+[sp^3]$ and $[PCl_6]^-[sp^3d^2]$.

B.2.2 Molecular Orbital Theory (MOT)

Molecular Orbital Energy Levels, Bond Order & Magnetic State

$$\text{BO} = \frac{1}{2}(N_b - N_a), \quad \text{Stability} \propto \text{BO} \propto \frac{1}{\text{Bond Length}} \tag{B.9}$$

Case 1: Total Electrons ≤ 14 (B_2, C_2, N_2 with sp-mixing):

$$\sigma 1s < \sigma^* 1s < \sigma 2s < \sigma^* 2s \tag{B.10}$$

$$< (\pi 2p_x = \pi 2p_y) < \sigma 2p_z < (\pi^* 2p_x = \pi^* 2p_y) < \sigma^* 2p_z \tag{B.11}$$

Case 2: Total Electrons > 14 (O_2, F_2 without sp-mixing):

$$\sigma 1s < \sigma^* 1s < \sigma 2s < \sigma^* 2s \tag{B.12}$$

$$< \sigma 2p_z < (\pi 2p_x = \pi 2p_y) < (\pi^* 2p_x = \pi^* 2p_y) < \sigma^* 2p_z \tag{B.13}$$

Species	e^-	Configuration	BO	Magnetic Nature
N_2	14	$(\pi 2p)^4(\sigma 2p_z)^2$	3.0	Diamagnetic
N_2^+	13	$(\pi 2p)^4(\sigma 2p_z)^1$	2.5	Paramagnetic (1 e^-)
O_2^{2+}	14	$(\sigma 2p_z)^2(\pi 2p)^4$	3.0	Diamagnetic
O_2^+	15	$(\pi^* 2p)^1$	2.5	Paramagnetic (1 e^-)
O_2	16	$(\pi^* 2p_x)^1(\pi^* 2p_y)^1$	2.0	Paramagnetic (2 e^-)
O_2^-	17	$(\pi^* 2p)^3$	1.5	Paramagnetic (1 e^-)
O_2^{2-}	18	$(\pi^* 2p)^4$	1.0	Diamagnetic
C_2	12	$(\pi 2p)^4$	2.0	Diamagnetic (two π-bonds)

Table B.3: Molecular Orbital Configurations, Bond Orders and Paramagnetism for Homonuclear Diatomics.

B.3 Coordination Compounds (Unit 12)

B.3.1 Werner's Theory, Ligands & Chelate Effect

Primary vs Secondary Valency & Chelate Stability

$$\text{Primary Valency} = \text{Oxidation state; ionizable; non-directional} \tag{B.14}$$
$$\text{Secondary Valency} = \text{Coordination number; non-ionizable; directional} \tag{B.15}$$
$$\text{Chelate Effect : Rings formed} \implies \Delta S^\circ > 0 \implies \Delta G^\circ < 0 \tag{B.16}$$

B.3.2 Crystal Field Theory (CFT) & Splitting Formulations

Octahedral & Tetrahedral Crystal Field Splitting Parameters

$$\Delta_t = \frac{4}{9}\Delta_o < P \quad \text{(Tetrahedral: always high spin)} \tag{B.17}$$
$$\text{CFSE}_{\text{oct}} = \left(-0.4\, n_{t_{2g}} + 0.6\, n_{e_g}\right)\Delta_o + mP \tag{B.18}$$
$$\text{Spectrochemical : } \mathrm{I^-} < \mathrm{Br^-} < \mathrm{S^{2-}} < \mathrm{SCN^-} < \mathrm{Cl^-} < \mathrm{F^-} < \mathrm{OH^-} \tag{B.19}$$
$$< \mathrm{ox^{2-}} \approx \mathrm{H_2O} < \mathrm{NCS^-} < \mathrm{EDTA^{4-}} < \mathrm{NH_3} \tag{B.20}$$
$$< \mathrm{en} < \mathrm{bpy} < \mathrm{phen} < \mathrm{NO_2^-} < \mathrm{PPh_3} < \mathrm{CN^-} \approx \mathrm{CO} \tag{B.21}$$

High-Spin vs Low-Spin Splitting Conditions for Octahedral Complexes:

$$\Delta_o > P \implies \text{Strong field; } t_{2g} \text{ pairing} \implies \text{Low-Spin / Inner Orbital} \tag{B.22}$$
$$\Delta_o < P \implies \text{Weak field; no } t_{2g} \text{ pairing} \implies \text{High-Spin / Outer Orbital} \tag{B.23}$$

Config.	Weak Field ($\Delta_o < P$)	CFSE	Strong Field ($\Delta_o > P$)	CFSE
d^4 (Cr^{2+})	$t_{2g}^3 e_g^1$ ($n=4$)	$-0.6\,\Delta_o$	$t_{2g}^4 e_g^0$ ($n=2$)	$-1.6\,\Delta_o + P$
d^5 (Fe^{3+})	$t_{2g}^3 e_g^2$ ($n=5$)	$0\,\Delta_o$	$t_{2g}^5 e_g^0$ ($n=1$)	$-2.0\,\Delta_o + 2P$
d^6 (Fe^{2+})	$t_{2g}^4 e_g^2$ ($n=4$)	$-0.4\,\Delta_o$	$t_{2g}^6 e_g^0$ ($n=0$)	$-2.4\,\Delta_o + 2P$
d^7 (Co^{2+})	$t_{2g}^5 e_g^2$ ($n=3$)	$-0.8\,\Delta_o$	$t_{2g}^6 e_g^1$ ($n=1$)	$-1.8\,\Delta_o + P$

Table B.4: Crystal Field Electronic Configurations and CFSE Formulations for $3d^4 - 3d^7$ Octahedral Ions.

B.3.3 Stereoisomerism in Coordination Complexes

Geometrical and Optical Isomerism Rules

- **Square Planar Complexes ($CN = 4$):**
  - $[MA_2B_2]$: Exhibits 2 geometrical isomers (cis and trans).
  - $[MABCD]$: Exhibits 3 geometrical isomers (2 cis, 1 trans).
  - Square planar complexes **never exhibit optical isomerism** because the molecular plane is an inherent plane of symmetry (σ).
- **Octahedral Complexes ($CN = 6$):**
  - $[MA_4B_2]$: Exhibits 2 geometrical isomers (cis and trans). Both are optically inactive.
  - $[MA_3B_3]$: Exhibits 2 geometrical isomers (facial *fac* and meridional *mer*). Both are optically inactive.
  - $[M(AA)_3]$ (e.g. $[Co(en)_3]^{3+}$): Exhibits 0 geometrical, but exists as a pair of **optical enantiomers** (d and l).
  - $[M(AA)_2B_2]$ (e.g. $[Co(en)_2Cl_2]^{+}$): *cis*-isomer is chiral and optically active (d/l pair); *trans*-isomer has plane of symmetry and is **optically inactive**.

B.4 The d- and f-Block Elements (Unit 11)

B.4.1 Transition Metal Trends & Variable Oxidation States

Magnetic Moment, Enthalpy of Atomization & Electrode Potentials

$$\mu_{\text{spin}} = \sqrt{n(n+2)}\,\text{BM} \quad [n = \text{unpaired } d\text{-electrons}] \tag{B.24}$$

$$\text{Max O.S.} : \text{Sc } (+3),\ \text{Ti } (+4),\ \text{V } (+5),\ \text{Cr } (+6),\ \text{Mn } (+7),\ \text{Os/Ru } (+8) \tag{B.25}$$

B.4.2 Potassium Dichromate ($K_2Cr_2O_7$) & Permanganate ($KMnO_4$)

Reagent & Reaction Medium	Change in Oxidation State	n-Factor	Canonical Half-Reaction
$K_2Cr_2O_7$ in Acidic Medium	$Cr^{6+} \to Cr^{3+}$	6	$Cr_2O_7^{2-} + 14H^+ + 6e^- \to 2Cr^{3+} + 7H_2O$
$KMnO_4$ in Acidic Medium	$Mn^{7+} \to Mn^{2+}$	5	$MnO_4^- + 8H^+ + 5e^- \to Mn^{2+} + 4H_2O$
$KMnO_4$ in Neutral / Faintly Alkaline	$Mn^{7+} \to Mn^{4+}$	3	$MnO_4^- + 2H_2O + 3e^- \to MnO_2 + 4OH^-$
$KMnO_4$ in Strongly Alkaline Medium	$Mn^{7+} \to Mn^{6+}$	1	$MnO_4^- + e^- \to MnO_4^{2-}$

Table B.5: Stoichiometric Redox n-Factors for Dichromate and Permanganate Titrations.

High-Yield Oxidation Reactions of $KMnO_4$ and $K_2Cr_2O_7$

- **Acidic $KMnO_4$ converts:**

$$Fe^{2+} \to Fe^{3+}, \quad C_2O_4^{2-} \to CO_2, \quad S^{2-} \to S,$$
$$SO_3^{2-} \to SO_4^{2-}, \quad NO_2^- \to NO_3^-, \quad I^- \to I_2$$

- **Baeyer's Reagent converts:**

$$I^- \to IO_3^- \text{ (Iodate)}, \qquad S_2O_3^{2-} \to SO_4^{2-}$$

- **Chromyl Chloride Test (Cl^- Detection):**

$$K_2Cr_2O_7 + 4NaCl + 6H_2SO_4 \xrightarrow{\Delta} 2CrO_2Cl_2 \uparrow \text{ (Deep Red)} + 2KHSO_4 + 4NaHSO_4 + 3H_2O$$

B.4.3 Lanthanoid Contraction & Oxidation States

Lanthanoid Contraction Governing Principles

$$\text{Trend} : La^{3+}(106\,\text{pm}) \rightarrow Lu^{3+}(86\,\text{pm}) \quad [20\text{ pm contraction}] \tag{B.26}$$

$$\text{Radii} : Zr(160\,\text{pm}) \approx Hf(159\,\text{pm}), \quad Nb \approx Ta, \quad Mo \approx W \tag{B.27}$$

$$\text{Basicity} : La(OH)_3 > Ce(OH)_3 > \cdots > Lu(OH)_3 \tag{B.28}$$

High-Yield Oxidation States of Lanthanoids:

$$\text{Dominant: } +3; \quad Ce^{4+}\ (4f^0 \text{ oxidizer}), \quad Eu^{2+}\ (4f^7 \text{ reducer}), \tag{B.29}$$

$$Yb^{2+}\ (4f^{14} \text{ reducer}) \tag{B.30}$$

B.5 *p*-Block Elements (Unit 10)

B.5.1 Inert Pair Effect & Thermodynamic Stability Trends

Group	High OS	Low OS	Stability Order & Redox Consequence
Group 13	+3	+1	$Tl^{+} > Tl^{3+} \implies Tl^{3+}$ oxidizes; Al^{3+} is stable.
Group 14	+4	+2	$Pb^{2+} > Pb^{4+} \implies PbO_2$ oxidizes; Sn^{2+} reduces.
Group 15	+5	+3	$Bi^{3+} > Bi^{5+} \implies BiF_5$ only pentahalide; Bi^{5+} oxidizes.

Table B.6: Inert Pair Effect Manifestation Across Heavy *p*-Block Elements.

Group 15, 16 & 17 Hydrides High-Yield Periodic Trends

- **Group 15 Hydrides ($NH_3, PH_3, AsH_3, SbH_3, BiH_3$):**
  - Bond Angle: $NH_3(107.8°) > PH_3(93.6°) > AsH_3(91.8°) > SbH_3(91.3°)$ (Drago's rule).
  - Thermal Stability: $NH_3 > PH_3 > AsH_3 > SbH_3 > BiH_3$.
  - Reducing Character: $BiH_3 > SbH_3 > AsH_3 > PH_3 > NH_3$ (strongest reducing agent is BiH_3).
  - Boiling Point: $BiH_3 > SbH_3 > \mathbf{NH_3} > AsH_3 > PH_3$ (NH_3 elevated due to H-bonding!).
- **Group 16 Hydrides (H_2O, H_2S, H_2Se, H_2Te):**
  - Acidic Strength: $H_2O < H_2S < H_2Se < H_2Te$ ($E-H$ bond enthalpy decreases).
  - Boiling Point: $\mathbf{H_2O} > H_2Te > H_2Se > H_2S$ (water is highest due to extensive intermolecular H-bonding).
- **Group 17 Hydrides (HF, HCl, HBr, HI):**
  - Acidic Strength: HF < HCl < HBr < HI (HI is strongest acid).
  - Boiling Point: HF > HI > HBr > HCl (HF elevated due to Zig-Zag H-bonding).
  - Reducing Power: HI > HBr > HCl > HF.

p-Block Traps: Oxoacids of Phosphorus & Boric Acid Basicity

- **Basicity and Reducing Nature of Phosphorus Oxoacids:**
  - H_3PO_2 (Hypophosphorous / Phosphinic acid): Contains 1 P – OH and 2 P – H bonds. **Monobasic**; powerful reducing agent.
  - H_3PO_3 (Phosphorous / Phosphonic acid): Contains 2 P – OH and 1 P – H bonds. **Dibasic**; moderate reducing agent.
  - H_3PO_4 (Orthophosphoric acid): Contains 3 P – OH and 0 P – H bonds. **Tribasic**; non-reducing.
- **Orthoboric Acid (H_3BO_3):** Is **not a protonic acid**! It is a weak, **monoba-**

sic Lewis acid that accepts OH^- from water:

$$B(OH)_3 + 2H_2O \rightleftharpoons [B(OH)_4]^- + H_3O^+$$

B.6 General Organic Chemistry (GOC) & Electronic Effects

Fundamental Electronic Effects in Organic Molecules

Organic reaction mechanisms and intermediate stabilities are governed by four fundamental displacement effects:

1. **Inductive Effect (I-Effect):** Permanent polarization through σ-bonds due to electronegativity difference. Decreases rapidly with distance ($I \propto 1/d^3$, negligible beyond 3 carbons).
   - $-I$ **Order:** $-NF_3^+ > -NR_3^+ > -NO_2 > -CN > -COOH > -F > -Cl > -Br > -I > -OR > -OH > -C \equiv CH > -NH_2 > -C_6H_5 > -CH = CH_2 > -H$
   - $+I$ **Order:** $-O^- > -COO^- > -CR_3(3^\circ) > -CHR_2(2^\circ) > -CH_2R(1^\circ) > -CH_3 > -T > -D > -H$
2. **Electromeric Effect (E-Effect):** Temporary complete transfer of a shared pair of π-electrons to one of the atoms in the presence of an attacking reagent.
   - $+E$ **Effect:** π-electrons transferred to the atom where the electrophile attacks (e.g., addition of H^+ to alkenes).
   - $-E$ **Effect:** π-electrons transferred to the atom from which the nucleophile moves away (e.g., addition of CN^- to carbonyls).
3. **Resonance / Mesomeric Effect (R/M-Effect):** Permanent delocalization of conjugated π-electrons or lone pairs.
   - $+M$ **(Electron Donating via Lone Pair):** $-O^- > -NH_2 > -NHR > -OH > -OR > -NHCOR > -OCOR > -Ph > -X$ (F > Cl > Br > I).
   - $-M$ **(Electron Withdrawing via Multiple Bonds):** $-NO_2 > -CN > -SO_3H > -CHO >> C=O > -COOH > -COOR > -CONH_2$.
4. **Hyperconjugation (Baker-Nathan Effect):** Delocalization of σ_{C-H} electrons with an adjacent empty p-orbital or π^* orbital.

$$\text{Number of hyperconjugative structures} = \text{Number of } \alpha\text{-H atoms} \quad \text{(B.31)}$$

Stability Hierarchy of Reactive Intermediates

- **Carbocations** ($sp^2, 6e^-$)**:** Tropylium > CPM > Ph_3C^+ > Allyl ≈ Benzyl ≈ 3° > 2° > 1° > CH_3^+.
  Stabilized by: $+M$ > Hyperconj. > $+I$; destabilized by: $-M, -I$.
- **Carbanions** ($sp^3/sp, 8e^-$)**:** $HC \equiv C^-$ > $CH_2 = CH^-$ > CH_3^- > 1° > 2° > 3°.
  Stabilized by: $-M, -I$, high s-character, aromaticity.
- **Free Radicals** ($sp^2, 7e^-$)**:** $Ph_3C^\bullet$ > Allyl ≈ Benzyl > 3° > 2° > 1° > $CH_3^\bullet$.
  Governed by hyperconjugation and resonance.

Carbocation Rearrangements: Hydride, Alkyl Shift & Ring Expansion

Carbocations spontaneously undergo 1,2-shifts to achieve higher thermodynamic stability:

- **Migratory Aptitude in Carbocations:** H^- > Aryl (Ar) > Alkyl (3° > 2° > 1° > CH_3).
- **Ring Expansion:** 4-membered ring carbocation → 5-membered ring (relieves angle strain from 90° → 108°); 5-membered ring carbocation → 6-membered ring (relieves strain to stable chair conformation).
- **Ring Contraction:** 7-membered ring carbocation undergoes contraction to stable 6-membered ring.

Quantitative Acidity & Basicity Sequences

1. Acidity Orders (Stabilization of Conjugate Base A^-):

General:	$RSO_3H > RCOOH > H_2CO_3 > PhOH > H_2O > ROH > HC \equiv CH > NH_3 > RH$
Haloacids:	$CF_3COOH > CCl_3COOH > CHCl_2COOH > CH_2FCOOH > CH_2ClCOOH$
Phenols:	Picric acid > p-NO_2 > o-NO_2 > m-NO_2 > Phenol
Ortho-Effect:	Ortho-substituted benzoic acids > benzoic acid.

2. Basicity Orders of Amines (Availability of Lone Pair on N):

Gas Phase:	3° > 2° > 1° > NH_3 ($+I$ governed)
Aqueous (Me):	$(CH_3)_2NH > CH_3NH_2 > (CH_3)_3N > NH_3$ [**213**]
Aqueous (Et):	$(Et)_2NH > (Et)_3N > EtNH_2 > NH_3$ [**231**]
Aromatic:	Aliphatic ≫ NH_3 > p-toluidine > Aniline > p-NO_2.

NEET Trap: Ortho-Effect, H-Bonding & Volatility

- ***o*-Nitrophenol vs *p*-Nitrophenol:** *o*-nitrophenol possesses **intramolecular H-bonding** (chelation), making it steam volatile with lower boiling point and less water solubility. *p*-nitrophenol possesses **intermolecular H-bonding**, giving it a higher boiling point and higher acidity ($pK_a = 7.15$ vs 7.23 for ortho due to absence of steric repulsion).
- **Picric Acid ($pK_a = 0.38$):** Stronger than carbonic acid and even many carboxylic acids; readily gives effervescence with aqueous $NaHCO_3$.
- **Phenol with $NaHCO_3$:** Phenol does NOT evolve CO_2 with $NaHCO_3$ (pK_a phenol $\approx 10 > pK_a$ $H_2CO_3 \approx 6.4$). Nitrophenols with ≥ 2 nitro groups DO dissolve in $NaHCO_3$.

B.7 Hydrocarbons: Alkanes, Alkenes, Alkynes & Arenes

Alkane Synthesis & Free Radical Transformations

1. Wurtz Reaction:

$$2R - X + 2Na \xrightarrow{\text{dry ether}} R - R + 2NaX$$

(Free radical; even-carbon symmetrical alkanes)

2. Decarboxylation (Soda Lime):

$$R - COONa + NaOH \xrightarrow[\Delta]{\text{CaO (3:1)}} R - H + Na_2CO_3$$

(Product contains 1 fewer carbon atom). **3. Kolbe Electrolytic Synthesis:**

$$\text{At Anode: } 2R - COO^- \xrightarrow{-2e^-} 2R^\bullet \xrightarrow{-2CO_2} R - R + 2CO_2$$

$$\text{At Cathode: } 2H_2O + 2e^- \rightarrow H_2 \uparrow + 2OH^- \quad \text{(solution pH rises)}$$

4. Free Radical Halogenation Reactivity & Selectivity:

$$\text{Reactivity: } F_2 > Cl_2 > Br_2 > I_2 \qquad \text{Selectivity: } 3^\circ\text{-H} > 2^\circ\text{-H} > 1^\circ\text{-H} \quad \text{(B.32)}$$

Relative rates of abstraction at 300°C:

- Chlorination ($1^\circ : 2^\circ : 3^\circ = 1 : 3.8 : 5.0$) → Low selectivity, governed by probability factor.
- Bromination ($1^\circ : 2^\circ : 3^\circ = 1 : 82 : 1600$) → Highly selective for 3° carbon.

Alkene Addition Reactions & Regiochemical Rules

1. Electrophilic Addition of HX (Markownikoff's Rule): The electrophile (H^+) adds to the double bond to form the more stable carbocation intermediate (allows rearrangement!).

$$CH_3 - CH = CH_2 + HCl \longrightarrow CH_3 - CH(Cl) - CH_3 \quad \text{(2-chloropropane)} \tag{B.33}$$

2. Peroxide Effect / Kharasch Effect (Anti-Markownikoff Addition): Occurs **only with** HBr in the presence of organic peroxides ($R - O - O - R$) via free radical mechanism.

$$CH_3CH = CH_2 + HBr \xrightarrow{\text{peroxide}} CH_3CH_2CH_2Br \text{ (1-bromopropane)}$$

Why only HBr*?* Addition of $Cl^\bullet$ is endothermic in step 2; abstraction of H from HI is endothermic in step 1. Only with HBr are both propagation steps exothermic.

3. Hydroboration-Oxidation (Syn-Addition, Anti-Markownikoff):

$$\begin{aligned} 6RCH = CH_2 &\xrightarrow{B_2H_6} 2(RCH_2CH_2)_3B \\ &\xrightarrow{H_2O_2/OH^-} 3RCH_2CH_2OH + B(OH)_3 \end{aligned}$$

Regioselectivity: Anti-Markownikoff; Stereochemistry: **Syn-addition**; strictly no rearrangement.

4. Oxymercuration-Demercuration (Markownikoff, No Rearrangement):

$$\begin{aligned} RCH = CH_2 &\xrightarrow[H_2O]{Hg(OAc)_2} RCH(OH)CH_2HgOAc \\ &\xrightarrow{NaBH_4/OH^-} RCH(OH)CH_3 \end{aligned}$$

Proceeds via cyclic mercurinium ion $\implies$ strictly NO rearrangement.

5. Ozonolysis (Reductive vs Oxidative Cleavage):

$$R_2C = CHR' \xrightarrow{O_3/CH_2Cl_2} \text{Molar Ozonide} \xrightarrow[Zn/H_2O]{\text{Reductive}} R_2C = O + R'CHO + ZnO \tag{B.34}$$

If oxidative (H_2O_2 or Ag_2O), any aldehyde formed is oxidized to carboxylic acid ($R'COOH$).

Alkynes: Acidity, Hydration & Stereoselective Reduction

1. Terminal Alkyne Acidity: *sp*-hybridized carbon has 50% *s*-character, stabilizing conjugate carbanion:

$$RC \equiv CH + NaNH_2 \longrightarrow RC \equiv C^-Na^+ + NH_3 \uparrow$$

$$\mathrm{RC \equiv CH + [Ag(NH_3)_2]^+ \longrightarrow RC \equiv C\text{-}Ag \downarrow \ (White\ ppt)}$$
$$\mathrm{RC \equiv CH + Cu_2Cl_2/NH_4OH \longrightarrow RC \equiv C\text{-}Cu \downarrow \ (Red\ ppt)}$$

2. Kucherov Reaction (Hydration of Alkynes):

$$\mathrm{HC \equiv CH + H_2O \xrightarrow[333\,K]{HgSO_4/H_2SO_4} [CH_2 = CHOH] \xrightarrow{tautomerism} CH_3CHO\ (Acetaldehyde)}$$

All other terminal and non-terminal alkynes yield ketones:

$$\mathrm{CH_3 - C \equiv CH + H_2O \xrightarrow{Hg^{2+}/H^+} CH_3 - CO - CH_3 \quad (Acetone)} \qquad \text{(B.35)}$$

3. Stereospecific Reduction of Alkynes:

- **Lindlar's Catalyst** ($\mathrm{H_2/Pd - CaCO_3/quinoline}$)**:** Syn-addition → ***cis*-Alkene**.
- **Birch Reduction** ($\mathrm{Na/liq.\ NH_3}$ **with EtOH):** Anti-addition → ***trans*-Alkene**.

Aromatic Hydrocarbons: EAS Directing Groups & Activating Power

Electrophilic Aromatic Substitution (EAS) involves arenium ion (σ-complex / Wheland intermediate):

Group	Substituents	Dir.	Reactivity
Strong Act.	$-OH, -O^-, -NH_2, -NR_2$	o/p	Greatly increased
Mod. Act.	$-OR, -NHCOR, -OCOR$	o/p	Moderately increased
Weak Act.	$-CH_3, -C_2H_5, -Ar$	o/p	Slightly increased
Halogens	$-F, -Cl, -Br, -I$	**o/p**	**Decreased** ($-I > +M$)
Mod. Deact.	$-CHO, -COR, -COOH, -SO_3H$	m	Moderately decreased
Strong Deact.	$-NO_2, -NR_3^+, -CF_3, -CN$	m	Greatly decreased

NEET Trap: Friedel-Crafts Reactions Limitations

1. **Aniline does NOT undergo Friedel-Crafts reaction:** The Lewis acid catalyst ($AlCl_3$) complexes with the lone pair on the $-NH_2$ group: $Ph - NH_2 + AlCl_3 \rightarrow Ph - \overset{+}{N}H_2 - \bar{A}lCl_3$. The resulting positive charge strongly deactivates the aromatic ring.
2. **Strongly deactivated rings do not react:** Nitrobenzene, benzoic acid, and benzonitrile do not undergo alkylation or acylation (used as solvents).
3. **Polyalkylation vs Monoacylation:** Alkylation introduces an activating alkyl group, leading to polyalkylation. Acylation introduces a deactivating acyl group ($-COR$), so it strictly stops at monoacylation.
4. **Rearrangement in Alkylation:** 1-chloropropane + $Ph - H \xrightarrow{AlCl_3}$ isopropylbenzene (Cumene), due to $1^\circ \rightarrow 2^\circ$ carbocation hydride shift!

B.8 Haloalkanes & Haloarenes: S_N1, S_N2, $E1$, $E2$

Nucleophilic Substitution Mechanisms: S_N1 vs S_N2 Core Matrix

Parameter	S_N1 (Unimolecular)	S_N2 (Bimolecular)
Kinetics	First Order: Rate $= k[\mathrm{R-X}]$	Second Order: Rate $= k[\mathrm{R-X}][\mathrm{Nu}^-]$
Mechanism	Two steps (carbocation intermediate)	One step concerted (transition state)
Reactivity of RX	$3^\circ > 2^\circ > 1^\circ > \mathrm{CH_3X}$ (carbocation)	$\mathrm{CH_3X} > 1^\circ > 2^\circ > 3^\circ$ (sterics)
Stereochemistry	**Racemization** (partial inversion)	**100% Walden Inversion**
Nucleophile	Weak ($\mathrm{H_2O}$, ROH); low conc.	Strong ($\mathrm{OH^-}$, $\mathrm{RO^-}$, $\mathrm{CN^-}$); high conc.
Solvent	**Polar Protic** ($\mathrm{H_2O}$, EtOH, AcOH)	**Polar Aprotic** (DMSO, DMF, Acetone)
Rearrangement	**Possible** (hydride/alkyl shift)	**Strictly Impossible**
Substrate Limits	Allylic & Benzylic react rapidly	Vinyl ($\mathrm{CH_2=CHX}$) and Aryl unreactive

Ambident Nucleophiles in Haloalkane Reactions

Ambident nucleophiles possess two nucleophilic centers with differing electronegativities:

$$\mathrm{R-X + KCN} \xrightarrow{\text{aq. EtOH}} \mathrm{R-CN} \quad \text{(Ionic; C-attack)}$$

$$\mathrm{R-X + AgCN} \xrightarrow{\text{aq. EtOH}} \mathrm{R-NC} \quad \text{(Covalent; N-attack)}$$

$$\mathrm{R-X + KNO_2} \xrightarrow{\text{aq. EtOH}} \mathrm{R-ONO} \quad \text{(Ionic; O-attack)}$$

$$\mathrm{R-X + AgNO_2} \xrightarrow{\text{aq. EtOH}} \mathrm{R-NO_2} \quad \text{(Covalent; N-attack)}$$

Elimination Regiochemistry: Saytzeff vs Hofmann Product

- **Saytzeff (Zaitsev) Rule:** Base removes β-hydrogen from the β-carbon having fewer hydrogens, yielding the more substituted, thermodynamically more stable alkene (major product with small bases: $\mathrm{OH^-}, \mathrm{EtO^-}, \mathrm{MeO^-}$).
- **Hofmann Rule:** Base removes β-hydrogen from the least hindered β-carbon, yielding the less substituted alkene as major product. Favored when:
  1. Attacking base is bulky (e.g., Potassium *tert*-butoxide: $t\text{-BuO}^-\mathrm{K}^+$).
  2. Leaving group is poor (e.g., Fluoride $-\mathrm{F}$, or quarternary ammonium $-\mathrm{NR_3^+}$).

NEET Trap: Nucleophilic Inactivity of Haloarenes

Chlorobenzene ($\mathrm{C_6H_5Cl}$) does not undergo nucleophilic substitution under normal conditions due to:

1. **Resonance:** Delocalization of chlorine lone pair gives partial double bond

character to the C − Cl bond (169 pm vs 177 pm in R − Cl).
2. **Hybridization of Carbon:** Chlorine is attached to an sp^2 hybridized carbon (33.3% s-character), which is more electronegative and holds electrons tighter than sp^3 carbon.
3. **Instability of Phenyl Cation:** Self-ionization to phenyl cation is energetically forbidden.
4. **Electrostatic Repulsion:** Electron-rich π-cloud of benzene repels incoming nucleophile.

Remedy (S_NAr Mechanism): Introduction of strong electron-withdrawing groups ($-NO_2$) at ***ortho* and *para* positions** stabilizes the Meisenheimer intermediate anionic complex, drastically decreasing required temperature:

- Chlorobenzene: NaOH, 623 K, 300 atm (Dow Process).
- p-Nitrochlorobenzene: NaOH, 443 K, H^+.
- 2,4-Dinitrochlorobenzene: NaOH, 368 K, H^+.
- 2,4,6-Trinitrochlorobenzene: Warm with H_2O alone at 323 K!

B.9 Oxygen Compounds: Alcohols, Phenols, Ethers, Carbonyls & Acids

Alcohols & Phenols Preparation & Conversions

1. Grignard Reagent (RMgX) Carbonyl Additions:

$$\begin{aligned}
\mathrm{HCHO} + \mathrm{RMgX} &\xrightarrow{\mathrm{H_3O^+}} 1^\circ \text{ Alcohol} \\
\mathrm{R'CHO} + \mathrm{RMgX} &\xrightarrow{\mathrm{H_3O^+}} 2^\circ \text{ Alcohol} \\
\mathrm{R'COR''} + \mathrm{RMgX} &\xrightarrow{\mathrm{H_3O^+}} 3^\circ \text{ Alcohol} \\
\mathrm{CO_2} + \mathrm{RMgX} &\xrightarrow{\mathrm{H_3O^+}} \mathrm{RCOOH} \text{ (Carboxylic acid)}
\end{aligned}$$

2. Cumene Hydroperoxide Process (Industrial Phenol):

$$\begin{aligned}
\mathrm{Benzene} + \mathrm{CH_3CH{=}CH_2} &\xrightarrow{\mathrm{H_3PO_4}} \mathrm{Cumene} \\
&\xrightarrow{\mathrm{O_2}} \mathrm{Cumene\ Hydroperoxide} \\
&\xrightarrow{\text{dil. } \mathrm{H_2SO_4}} \mathrm{Phenol} + \mathrm{CH_3COCH_3}
\end{aligned}$$

3. Kolbe-Schmitt Reaction (Synthesis of Salicylic Acid):

$$\begin{aligned}
\mathrm{Sodium\ phenoxide} + \mathrm{CO_2} &\xrightarrow[4\text{–}7\ \mathrm{atm}]{400\ \mathrm{K}} \mathrm{Sodium\ salicylate} \\
&\xrightarrow{\mathrm{H^+}} \mathrm{Salicylic\ acid}
\end{aligned}$$

4. Reimer-Tiemann Reaction (Synthesis of Salicylaldehyde):

$$\text{Phenol} + CHCl_3 + 3NaOH \xrightarrow{340\,K} \text{Salicylaldehyde}$$

(**Electrophile: Dichlorocarbene** : CCl_2). If CCl_4 is used, salicylic acid is main product.

5. Esterification (Fischer):

$$RCOOH + R'OH \underset{}{\overset{\text{conc. } H_2SO_4}{\rightleftharpoons}} RCOOR' + H_2O$$

(Reactivity of ROH: $CH_3OH > 1^\circ > 2^\circ > 3^\circ$).

Ethers: Williamson Synthesis & Cleavage with HI

1. Williamson Ether Synthesis:

$$R - X + R' - O^-Na^+ \longrightarrow R - O - R' + NaX \quad (\textbf{Mechanism: Strictly } S_N2) \tag{B.36}$$

- Alkyl halide must be 1° (or CH_3X).
- If 3° alkyl halide is used with alkoxide, **elimination ($E2$) dominates exclusively**:

$$(CH_3)_3C - Br + CH_3O^- \longrightarrow CH_3 - C(CH_3) = CH_2 + CH_3OH + Br^- \tag{B.37}$$

(Isobutylene) (B.38)

2. Cleavage of Ethers by Halogen Acid (HI):

- **Alkyl Alkyl Ethers ($1^\circ/2^\circ$ groups):** Follows S_N2. Iodine ion attacks the less sterically hindered alkyl group:

$$CH_3 - O - C_2H_5 + HI \xrightarrow{\text{cold}} CH_3 - I + C_2H_5 - OH \tag{B.39}$$

- **One group is 3° (or Benzyl):** Follows S_N1 via stable carbocation; iodine attacks the 3° carbon:

$$(CH_3)_3C - O - CH_3 + HI \xrightarrow{\text{cold}} (CH_3)_3C - I + CH_3 - OH \tag{B.40}$$

- **Alkyl Aryl Ethers (Anisole):** The $O -$ Aryl bond has partial double bond character and never cleaves:

$$C_6H_5 - O - CH_3 + HI \xrightarrow{\Delta} C_6H_5 - OH\ (\text{Phenol}) + CH_3 - I \tag{B.41}$$

Carbonyl Compounds: Nucleophilic Addition & Carbonyl Condensations

1. Relative Reactivity Towards Nucleophilic Addition:

$$\text{HCHO} > \text{CH}_3\text{CHO} > \text{CH}_3\text{COCH}_3 > \text{PhCHO} > \text{PhCOCH}_3 > \text{PhCOPh} \tag{B.42}$$

Governed by: Steric hindrance at carbonyl carbon and $+I/+M$ electron donation reducing electrophilicity of carbonyl carbon.

2. Aldol Condensation (α-H Required): Carbonyl compounds with at least one α-hydrogen react in dilute alkali (10% NaOH or Ba(OH)_2):

$$\begin{aligned} 2\text{CH}_3\text{CHO} &\xrightarrow{\text{dil. NaOH}} \text{CH}_3 - \text{CH(OH)} - \text{CH}_2 - \text{CHO} \\ &\xrightarrow[\Delta]{-\text{H}_2\text{O}} \text{CH}_3 - \text{CH} = \text{CH} - \text{CHO} \quad \text{(Crotonaldehyde)} \end{aligned} \tag{B.43}$$

3. Cannizzaro Reaction (No α-H Present): Self-oxidation and self-reduction in concentrated alkali (50% NaOH):

$$2\text{HCHO} + \text{conc. NaOH} \longrightarrow \text{CH}_3\text{OH (Methanol)} + \text{HCOONa (Sodium formate)} \tag{B.44}$$

$$\begin{aligned} 2\text{C}_6\text{H}_5\text{CHO} + \text{conc. NaOH} \longrightarrow{}& \text{C}_6\text{H}_5\text{CH}_2\text{OH (Benzyl alcohol)} \\ &+ \text{C}_6\text{H}_5\text{COONa (Sodium benzoate)} \end{aligned} \tag{B.45}$$

Cross Cannizzaro with Formaldehyde: Formaldehyde undergoes oxidation exclusively (HCOO^-) because hydride transfer occurs from the more nucleophilically attacked formaldehyde.

Named Carbonyl Reductions, Oxidations & Acid Reactions

1. **Clemmensen Reduction:** $\text{Zn} - \text{Hg} + \text{conc. HCl} \quad (> \text{C} = \text{O} \rightarrow > \text{CH}_2)$; *Avoid if acid-sensitive groups present.*
2. **Wolff-Kishner Reduction:** $\text{NH}_2 - \text{NH}_2 + \text{KOH/glycol}, \ \Delta \quad (> \text{C} = \text{O} \rightarrow > \text{CH}_2)$; *Avoid if base-sensitive groups present.*
3. **Rosenmund Reduction:** $\text{R} - \text{COCl} + \text{H}_2 \xrightarrow{\text{Pd/BaSO}_4,\text{quinoline}} \text{R} - \text{CHO} + \text{HCl}$.
4. **Stephen Reaction:** $\text{R} - \text{CN} + \text{SnCl}_2/\text{HCl} \rightarrow \text{R} - \text{CH} = \text{NH} \cdot \text{HCl} \xrightarrow{\text{H}_3\text{O}^+} \text{R} - \text{CHO} + \text{NH}_4\text{Cl}$.
5. **Etard Reaction:** $\text{C}_6\text{H}_5 - \text{CH}_3 + \text{CrO}_2\text{Cl}_2/\text{CS}_2 \rightarrow \text{Complex} \xrightarrow{\text{H}_3\text{O}^+} \text{C}_6\text{H}_5 - \text{CHO}$.
6. **Gattermann-Koch:** $\text{C}_6\text{H}_6 + \text{CO} + \text{HCl} \xrightarrow{\text{anhyd. AlCl}_3/\text{CuCl}} \text{C}_6\text{H}_5 - \text{CHO} + \text{HCl}$.
7. **Hell-Volhard-Zelinsky (HVZ) Reaction:** Carboxylic acids with α-hydrogen react with halogen in presence of red phosphorus:

$$\text{R} - \text{CH}_2 - \text{COOH} + \text{X}_2 \xrightarrow[\text{(ii) H}_2\text{O}]{\text{(i) Red P}} \text{R} - \text{CH(X)} - \text{COOH}$$

(α-halocarboxylic acid; X = Cl, Br) (B.46)

B.10 Nitrogen Compounds: Amines & Diazonium Salts

Synthesis & Reactions of Amines

1. Gabriel Phthalimide Synthesis (Pure 1° Aliphatic Amines):

$$\text{Phthalimide} \xrightarrow{\text{KOH}} \text{K-phthalimide} \xrightarrow{\text{R}-\text{X}}$$
$$N\text{-Alkylphthalimide} \xrightarrow{\text{NaOH(aq)}} \text{R} - \text{NH}_2 + \text{Phthalate}$$

Limitation: Aromatic primary amines cannot be prepared because aryl halides do not undergo S_N2 displacement.

2. Hoffmann Bromamide Degradation: Conversion of an amide to a 1° amine containing **one carbon less**:

$$\text{R} - \text{CONH}_2 + \text{Br}_2 + 4\text{NaOH} \longrightarrow \text{R} - \text{NH}_2 + \text{Na}_2\text{CO}_3 + 2\text{NaBr} + 2\text{H}_2\text{O} \quad \text{(B.47)}$$

Intermediate is alkyl isocyanate ($\text{R} - \text{N} = \text{C} = \text{O}$); migration of alkyl group occurs with **retention of configuration**.

3. Carbylamine Test (Isocyanide Test - Strictly 1° Amines): Both aliphatic and aromatic 1° amines react with chloroform and alc. KOH to form foul-smelling isocyanides:

$$\text{R} - \text{NH}_2 + \text{CHCl}_3 + 3\text{KOH} \xrightarrow{\Delta} \text{R} - \text{NC (Alkyl isocyanide)} + 3\text{KCl} + 3\text{H}_2\text{O} \quad \text{(B.48)}$$

2° and 3° amines do NOT show this test.

4. Hinsberg Test (Distinction of 1°, 2°, 3° Amines): Reagent: Benzenesulfonyl chloride ($\text{C}_6\text{H}_5\text{SO}_2\text{Cl}$).

- 1° Amine → N-alkylbenzenesulfonamide → possesses acidic H on N → **Soluble in alkali** (KOH).
- 2° Amine → N,N-dialkylbenzenesulfonamide → no acidic H → **Insoluble in alkali**.
- 3° Amine → No replaceable H on N → **Does not react with Hinsberg reagent**.

Diazonium Chemistry: Arenediazonium Salt Reactions

Diazotization: $\text{Ar} - \text{NH}_2 + \text{NaNO}_2 + 2\text{HCl} \xrightarrow{273-278\,\text{K}} \text{Ar} - \text{N}_2^+\text{Cl}^- + \text{NaCl} + 2\text{H}_2\text{O}$.

$$\text{Sandmeyer: } \text{Ar} - \text{N}_2^+\text{Cl}^- \xrightarrow{\text{Cu}_2\text{X}_2/\text{HX}} \text{Ar} - \text{X} \quad (\text{X} = \text{Cl}, \text{Br}, \text{CN})$$

$$\text{Gattermann: } \mathrm{Ar-N_2^+Cl^-} \xrightarrow{\mathrm{Cu/HX}} \mathrm{Ar-X} \quad (\mathrm{X=Cl, Br})$$

$$\text{Balz-Schiemann: } \mathrm{Ar-N_2^+Cl^-} \xrightarrow{\mathrm{HBF_4}} \mathrm{Ar-N_2^+BF_4^-} \xrightarrow{\Delta} \mathrm{Ar-F + BF_3 + N_2\uparrow}$$

$$\text{Iodination: } \mathrm{Ar-N_2^+Cl^- + KI} \xrightarrow{\text{warm}} \mathrm{Ar-I + KCl + N_2\uparrow}$$

$$\text{Hydrolysis: } \mathrm{Ar-N_2^+Cl^- + H_2O} \xrightarrow{373\,\mathrm{K}} \mathrm{Ar-OH + HCl + N_2\uparrow}$$

Reduction to Arene ($\mathrm{Ar-H}$):

$$\mathrm{Ar-N_2^+Cl^- + H_3PO_2 + H_2O} \xrightarrow{\mathrm{Cu^+}} \mathrm{Ar-H + H_3PO_3 + HCl + N_2\uparrow}$$

$$\mathrm{Ar-N_2^+Cl^- + CH_3CH_2OH} \longrightarrow \mathrm{Ar-H + CH_3CHO + HCl + N_2\uparrow}$$

Azo Coupling Reactions (Formation of Dyes):

$$\text{With Phenol: } \mathrm{Ph-N_2^+Cl^- + C_6H_5-OH} \xrightarrow{\mathrm{pH\,9\text{–}10}} \text{p-hydroxyazobenzene (\textbf{Orange dye})}$$

$$\text{With Aniline: } \mathrm{Ph-N_2^+Cl^- + C_6H_5-NH_2} \xrightarrow{\mathrm{pH\,4\text{–}5}} \text{p-aminoazobenzene (\textbf{Yellow dye})}$$

B.11 Biomolecules: Carbohydrates, Amino Acids & Nucleic Acids

Carbohydrates Classification & Reactions of D-Glucose

1. Classification & Reducing Nature:

- **Reducing Sugars:** All monosaccharides (aldoses & ketoses) and disaccharides with a free hemiacetal/hemiketal -OH at anomeric carbon (Maltose, Lactose). Reduce Tollens and Fehling solutions.
- **Non-Reducing Sugars:** Glycosidic linkage involves anomeric carbons of both units (e.g., **Sucrose** = α-D-glucopyranosyl-(1→2)-β-D-fructofuranoside).
- **Invert Sugar:** Sucrose ($[\alpha]_D = +66.5^\circ$) on hydrolysis gives D-glucose ($+52.5^\circ$) and D-fructose (-92.4°). The mixture is levorotatory ($-39.9^\circ/2 = -19.9^\circ$).

2. Chemical Reactions of D-Glucose ($\mathrm{C_6H_{12}O_6}$):

$$\mathrm{Glucose + HI},\ \Delta \longrightarrow n\text{-hexane} \quad (\text{Proves open chain of 6 carbons})$$

$$\mathrm{Glucose + NH_2OH} \longrightarrow \text{Glucose oxime} \quad (\text{Confirms carbonyl } > \mathrm{C=O})$$

$$\mathrm{Glucose + HCN} \longrightarrow \text{Glucose cyanohydrin}$$

$$\mathrm{Glucose + Br_2/H_2O} \longrightarrow \text{Gluconic acid} \quad (\text{Mild oxidation of } -\mathrm{CHO})$$

$$\text{Glucose} + \text{conc. } HNO_3 \longrightarrow \text{Saccharic acid} \quad (\text{Oxidizes } -\text{CHO and } 1^\circ\text{-OH})$$
$$\text{Glucose} + 5(CH_3CO)_2O \longrightarrow \text{Glucose pentaacetate}$$
$$(\text{Confirms 5 independent } -\text{OH groups})$$

3. Why the Cyclic Structure of Glucose? D-Glucose does not give Schiff's test, 2,4-DNP test, and does not form hydrogen sulphite addition product with $NaHSO_3$. The pentaacetate does not react with hydroxylamine (absence of free $-$CHO).

$$\begin{aligned} &\alpha\text{-D-(+)-glucose } ([\alpha]_D = +112^\circ) \\ &\quad \underset{\text{in solution}}{\overset{\text{Mutarotation}}{\rightleftharpoons}} \text{Equilibrium } (+52.7^\circ) \\ &\quad \overset{\beta}{\rightleftharpoons} \beta\text{-D-(+)-glucose } (+19^\circ) \end{aligned} \tag{B.49}$$

α and β isomers differ only at $C-1$ (anomeric carbon) $\implies$ **Anomers**.

Amino Acids, Peptide Bonds & Protein Structure

1. Zwitterion & Isoelectric Point (pI): In aqueous solution, the carboxyl group loses a proton and amino group accepts it:

$$R - CH(NH_2) - COOH \rightleftharpoons R - CH(NH_3^+) - COO^- \quad (\text{Zwitterion}) \tag{B.50}$$

$$\text{At pH} < \text{pI} : \text{cation } R - CH(NH_3^+) - COOH \implies \text{cathode} \tag{B.51}$$
$$\text{At pH} > \text{pI} : \text{anion } R - CH(NH_2) - COO^- \implies \text{anode} \tag{B.52}$$

For neutral amino acid: $\text{pI} = \frac{pK_{a1}+pK_{a2}}{2}$.

- **Chirality:** All naturally occurring amino acids are optically active and possess **L-configuration**, **except Glycine** ($H_2N - CH_2 - COOH$, achiral).
- **Essential Amino Acids (10):** Valine, Leucine, Isoleucine, Phenylalanine, Methionine, Tryptophan, Threonine, Lysine, Histidine, Arginine. [**Mnemonic: PVT TIM HALL**].

2. Protein Structural Levels:

- **Primary:** Specific sequence of amino acids linked by covalent peptide bonds ($-CO-NH-$).
- **Secondary:** Local conformation of polypeptide chain stabilized by **hydrogen bonds** between $>C=O$ and $-NH-$ groups: α-helix (intramolecular H-bonds) and β-pleated sheets (intermolecular H-bonds).
- **Tertiary:** Overall 3D folding of polypeptide chains stabilized by H-bonds, disulphide linkages ($-S-S-$), van der Waals forces, and electrostatic attractions.

- **Quaternary:** Spatial arrangement of two or more subunits (e.g., Hemoglobin with 2α and 2β chains).
- **Denaturation:** Heat, pH change breaks $2°, 3°, 4°$ structures; **Primary structure remains completely intact**! (e.g., coagulation of egg white, curdling of milk).

Nucleic Acids: Nucleotides, Chargaff's Rules & Double Helix

1. Structural Components:

- **Pentose Sugar:** β-D-ribose (RNA) vs β-D-2-deoxyribose (DNA).
- **Nitrogenous Bases:**
  - **Purines (Bicyclic):** Adenine (A), Guanine (G).
  - **Pyrimidines (Monocyclic):** Cytosine (C), Thymine (T, DNA only), Uracil (U, RNA only).
- **Nucleoside:** Sugar + Base (attached at $C-1'$ of sugar via β-glycosidic linkage).
- **Nucleotide:** Nucleoside + Phosphate (esterified at $C-5'$ of sugar).
- **Phosphodiester Linkage:** Connects $C-3'$ of one nucleotide to $C-5'$ of adjacent nucleotide.

2. Chargaff's Rules (for Double-Stranded DNA):

$$\%A = \%T \quad \text{and} \quad \%G = \%C \implies \frac{A+G}{T+C} = 1 \qquad (\text{Purines} = \text{Pyrimidines}) \tag{B.53}$$

Base pairing: A pairs with T via **2 hydrogen bonds** (A = T); G pairs with C via **3 hydrogen bonds** (G ≡ C).

B.12 Master Identification Tests & Distinction Matrix

Comprehensive Tabular Matrix for Distinguishing Organic Functional Pairs

Compound Pair	Reagent / Test	Observed Visual Distinction
$1^\circ, 2^\circ, 3^\circ$ Alcohols	Lucas Reagent (conc. HCl + anhyd. $ZnCl_2$)	3°: immediate cloudiness; 2°: turbidity in 5 min; 1°: no turbidity at room temp.
Aliphatic vs Aromatic Aldehydes	Fehling's Solution ($A + B$)	Aliphatic: Red ppt (Cu_2O); Benzaldehyde: No reaction (does not reduce Fehling).
Aldehyde vs Ketone	Tollens' Reagent ($[Ag(NH_3)_2]^+$)	Aldehydes: Bright silver mirror (Ag); Ketones: Negative.
$CH_3 - C = O$ or $CH_3 - CH(OH) -$	Iodoform Test (I_2 + NaOH)	Yellow crystalline $CHI_3 \downarrow$ (m.p. $119^\circ C$) with antiseptic odor. Positive for ethanol, acetaldehyde, 2-propanol.
Carboxylic Acid vs Phenol	$NaHCO_3$ (aqueous)	Carboxylic acid: Brisk effervescence of CO_2; Phenol: No effervescence.
Phenol vs Alcohol	Neutral $FeCl_3$ solution	Phenol: Violet coloration ($[Fe(OC_6H_5)_6]^{3-}$); Alcohols: Negative.
1° Amine vs $2^\circ/3^\circ$	Carbylamine Test ($CHCl_3$ + alc. KOH)	Only 1° amine gives extremely foul smell of isocyanide ($R - NC$).
$1^\circ, 2^\circ, 3^\circ$ Amines	Hinsberg Reagent ($C_6H_5SO_2Cl$)	1°: Soluble in KOH; 2°: Insoluble in KOH; 3°: No reaction.
Terminal vs Internal Alkyne	Ammoniacal Cu_2Cl_2	Terminal ($R - C \equiv CH$): Red ppt of copper acetylide; Internal: Negative.
Alkene/Alkyne vs Alkane	Bromine water (Br_2/CCl_4)	Unsaturated: Decolorization of reddish-brown bromine color without HBr evolution.

High-Yield Reagent Conversion Master Reference

- PCC **(Pyridinium Chlorochromate) / Collins Reagent:** Oxidizes 1° alcohols strictly to aldehydes (RCHO) without over-oxidation to carboxylic acid. Oxidizes 2° alcohols to ketones.
- **DIBAL-H (Diisobutylaluminium Hydride):** Selectively reduces esters ($RCOOR'$) and nitriles (RCN) at low temperature ($-78^\circ C$) to **aldehydes** (RCHO).
- $LiAlH_4$ **vs** $NaBH_4$**:** $LiAlH_4$ reduces acids, esters, amides, nitriles, aldehydes, and ketones. $NaBH_4$ is weaker and selectively reduces **only aldehydes, ketones, and acid halides** (does not touch esters, acids, or amides).
- CrO_3 **in glacial acetic acid vs Jones Reagent (**CrO_3/H_2SO_4**):** Mild vs strong chromic acid oxidation.
- **Fenton's Reagent (**$FeSO_4 + H_2O_2$**):** Generates hydroxyl free radicals ($OH^\bullet$) for selective oxidation of alcohols to carbonyls.

About the Author & Errata Submissions

Satya Prakash is an educator, curriculum researcher, and author dedicated to creating rigorous, student-first learning resources for competitive examinations in India. His focus is on eliminating cognitive overload, dismantling obsolete legacy syllabi, and restoring conceptual rigor and mathematical transparency to competitive pre-medical preparation.

Official Project Repository & Continuous Improvement

Science education thrives on continuous scrutiny and collective refinement. If you identify a typo, numerical discrepancy, or pedagogical ambiguity, or if you wish to suggest an enhancement for future editions, please submit your feedback:

- **Direct Author Email:** spdedsec@hotmail.com
- **GitHub Open-Source Repository:** `https://github.com/spdedsec/neet-study-materials`
- **Errata Tracker:** Verified corrections and updates are posted directly to the project repository.

Authored by Satya Prakash · All Rights Reserved