JACOBI COEFFICIENTS OF MEASURES ON RECIPROCAL SPECTRA: A SPECTRAL DICTIONARY AND THE NATURAL INDEX

GORDON SPEAGLE

ABSTRACT. For an increasing sequence $\Lambda = \{\lambda_m\}$ with $\lambda_m \sim m^p$ we study the recurrence coefficients a_n, b_n of the discrete measure $\sum_m \lambda_m^{-s/p}\delta_{1/\lambda_m}$; the lattice $\lambda_m = m^p$ gives the measures $\sum_k k^{-s}\delta_{k^{-p}}$ of the earlier papers of this series, for which $n^p b_n \to c(p)$. For the lattice and every $p \neq 1$ we prove an expansion of $n^p b_n/c$ to order $n^{-7} + n^{-2-3p}$. It is a double series in $1/n$ and in $\zeta(-jp)n^{-jp}$, computed by a finite residue recursion and exact computer algebra. In the natural index $N = n + \frac{\alpha}{2} + \frac{1}{2}$, $\alpha = (s-1-p)/p$, the integer layer is even in $1/N$ to all orders: separating the roles of α as Bessel order and as Szegő exponent, we prove a reflection symmetry and, by steepest descent in a second frame of non-integer index, that the Szegő exponent enters only through N. The first-order zeta terms are $\varepsilon_j N^{-1-jp}(1 + g_j N^{-2})$ with $\varepsilon_j = \binom{2j}{j} p\, c^j \zeta(-jp)$ and g_j quadratic in α, for all j, and, formally, the zeta layers are Toda flows of the integer layer. For Hurwitz lattices $\lambda_m = (m+\delta)^p$, $-1 < \delta \leq 0$, the offset only shifts the Szegő exponent, so the lattice expansion holds with N shifted by δ and $\zeta(-jp)$ replaced by $\zeta(-jp, 1+\delta)$; for $\lambda_m = (m+\delta)^p + \kappa$, $p > 1$, we prove the first pole term. We write the zeta layer as a Mellin–Barnes integral with kernel $\pi[\cot(\pi w p/2) - \cot(\pi w)]$ and solve the constrained equilibrium problem for a general decreasing node density explicitly; its soft edge carries every pole of the node zeta function with the weight $\pi\cot(\pi q)$. For general spectra these statements become a conjectural dictionary between the expansion of b_n and the spectral zeta function of Λ, which we test numerically on perturbed lattices, Bessel zeros and spectra with multiplicities (with predictions fixed before the measures were computed), and on planar drums and anharmonic oscillators; for the oscillators $-\frac{\mathrm{d}^2}{\mathrm{d}q^2} + q^N$ the predicted n^{-2} shift is the rational number $(N-1)/(6(N+2))$.

1. INTRODUCTION

1.1. **Setting.** Let $p > 0$, $s > 1$, and let $\Lambda = \{\lambda_m\}_{m\geq 1}$ be an increasing sequence of positive numbers with $\lambda_m \sim m^p$. We call Λ a *node set* and study the discrete measure on its reciprocals

$$\sigma_\Lambda := \sum_{m\geq 1} \lambda_m^{-s/p}\, \delta_{1/\lambda_m}. \tag{1.1}$$

For the *lattice* $\lambda_m = m^p$ this is the measure $\sigma_{p,s} = \sum_k k^{-s}\delta_{k^{-p}}$ of the earlier papers of this series [10, 11, 12]. Let P_n be the monic orthogonal polynomials of σ_Λ; they satisfy

$$xP_n = P_{n+1} + a_n P_n + b_n^2 P_{n-1} \qquad (n \geq 0,\ b_0 := 0,\ b_n > 0), \tag{1.2}$$

and a_n, b_n are the entries of the Jacobi matrix whose spectrum is $\{1/\lambda_m\} \cup \{0\}$. With

$$I(p) := \frac{\sqrt{\pi}}{2}\,\frac{\Gamma(\frac{1}{p}+\frac{1}{2})}{\Gamma(\frac{1}{p}+1)}, \qquad c(p) := \frac{1}{4}\Big(\frac{\pi}{2I(p)}\Big)^p, \tag{1.3}$$

Date: October 2026.

2020 *Mathematics Subject Classification.* Primary 42C05; Secondary 47B36, 31A15, 30E15, 11M35.

Key words and phrases. Orthogonal polynomials, Jacobi matrices, recurrence coefficients, discrete measures, constrained equilibrium, Riemann–Hilbert problems, spectral zeta functions, Hurwitz zeta function, Mellin transforms, Toda hierarchy.

the lattice satisfies the pointwise law $n^p b_n \to c(p)$, $n^p a_n \to 2c(p)$, with rate $O(1/n)$ and an explicit $1/n$ term for every $p > 0$ [11, Theorems 1.2 and 1.4]. We write $c = c(p)$,

$$\alpha := \frac{s-1-p}{p}, \qquad y_n := \frac{n^p b_n}{c} - 1, \tag{1.4}$$

so that $\alpha > -1$ is the Bessel order of the hard edge at 0 in [11]. The questions of this paper are: how does the expansion of y_n continue, and how does it depend on Λ? The organising object is the *node zeta function*

$$Z_\Lambda(w) := \sum_{m \ge 1} \lambda_m^{-w}, \tag{1.5}$$

continued meromorphically from $\operatorname{Re} w > 1/p$; for the lattice $Z_\Lambda(w) = \zeta(pw)$. Two quantities recur: the numbers

$$W(q) := \binom{2q}{q} c^q = \frac{(4c)^q\,\Gamma(q+\frac12)}{\sqrt{\pi}\,\Gamma(q+1)} \qquad (q \in \mathbb{C} \setminus \{-\tfrac12, -\tfrac32, \dots\}), \tag{1.6}$$

which satisfy $W(1/p) = 1$ for every p (this is the definition (1.3) of c), and the coefficients

$$\varepsilon_j := W(j)\, p\, \zeta(-jp) = \binom{2j}{j} p\, c^j \zeta(-jp) \qquad (j \ge 1), \tag{1.7}$$

so that $\varepsilon_1 = \mathsf{k}_p := 2pc\,\zeta(-p)$ is the constant of [11, Theorem 1.4(ii)].

1.2. **Results.** The first result is an expansion of the lattice coefficients far beyond the $1/n$ term. Its form is simplest in the shifted index

$$N := n + \frac{\alpha}{2} + \frac12. \tag{1.8}$$

The rational functions f_4, f_6 of (p, α) and $g_{1,4}$ are given in (5.3)–(5.5).

Theorem A (the lattice to order $n^{-7} + n^{-2-3p}$)**.** *Let $p > 0$, $p \ne 1$, $s > 1$. The recurrence coefficients of $\sigma_{p,s}$ satisfy*

$$\begin{aligned} \frac{n^p b_n}{c} = \Big(\frac{n}{N}\Big)^p \Big[1 + \frac{f_2}{N^2} + \frac{f_4}{N^4} + \frac{f_6}{N^6} + \frac{\varepsilon_1}{N^{1+p}}\Big(1 + \frac{g_1}{N^2} + \frac{g_{1,4}}{N^4}\Big) \\ + \sum_{j \ge 2} \frac{\varepsilon_j}{N^{1+jp}}\Big(1 + \frac{g_j}{N^2}\Big) + \frac{3p+1}{2p}\,\frac{\varepsilon_1^2}{N^{2+2p}}\Big] + O\big(n^{-7} + n^{-2-3p}\big), \end{aligned} \tag{1.9}$$

where the sum is over the finitely many $j \ge 2$ with $1 + jp < \min(7, 2+3p)$,

$$f_2 = \frac{p+1}{12} - \frac{(p+2)\alpha^2}{8}, \qquad g_j = A_j + B_j\alpha^2, \qquad B_j = \frac{(p+2)\big(j^2p^2 + (j+1)p + 2\big)}{8(2j-1)p},$$

$$A_j = \frac{j^2(j^2+2j-2)p^3 + 2(j+1)(2j^2-1)p^2 + (5j^2+2j-6)p + 2(j-2)}{24(2j-1)p}.$$

Theorem A is the case $K = 6$, $B = 2$ of Theorem 5.7, which gives a double expansion of $n^p b_n/c$ in $1/n$ and in the parameters $\mu_{n,j} = 2\zeta(-jp)/(jn^{jp})$ of the band factor, to every order, through a finite residue recursion. Its proof combines the third-order Riemann–Hilbert analysis of [11], extended to order K, with exact computer algebra over $\mathbb{Q}(p, \alpha)$ and, for the zeta terms, over $\mathbb{Q}(p, \alpha, j)$ with j a symbol. Three structural facts are visible in (1.9): the shift $n \mapsto N$ absorbs the whole $1/n$ term and the $(s-1)^2$ part of the n^{-2} term; every layer computed is even in $1/N$; and the first-order zeta terms are dressed by an even factor $1 + g_j N^{-2} + \cdots$ with

g_j quadratic in α. Theorem D below explains the first two to all orders for the integer layer. Not every displayed term is a coefficient of b_n for every p: Table 5.1 lists which are.

The second result explains the zeta coefficients. The band factor of [11] enters the first-order correction through one linear functional e_n (Section 4), whose asymptotic series is $\sum_j \varepsilon_j n^{-jp}$.

Theorem B (the zeta layer as a Mellin–Barnes integral)**.** *Let $p > 0$, $p \neq 1$, and $0 < c_0 < 1$. The band-factor functional e_n of the lattice satisfies*

$$e_n = \frac{1}{2\pi i} \int_{c_0 - i\infty}^{c_0 + i\infty} p\, W(w)\, \zeta(-wp)\, n^{-wp}\, \pi \Big[\cot \frac{\pi w p}{2} - \cot(\pi w) \Big]\, \mathrm{d}w,$$

and, for every $J \geq 1$, $e_n = \sum_{j \leq J} \varepsilon_j n^{-jp} + O(n^{-(J+1)p})$. The kernel $-\pi \cot(\pi w) W(w) (4c)^{-w}$ is the Mellin transform of the truncated per-node law $(1-u)^{-1/2} \mathbf{1}_{u<1}$ (Lemma 4.1).

The third result is on the side of potential theory. In the variable $x = n^p \cdot (\text{node})$, the n nodes that carry the polynomial P_n have the continuum density $\sigma(y) = \frac{1}{p} y^{-1/p-1}$, and the zero distribution of P_n is the equilibrium measure of unit mass constrained by σ [9, 6, 10]. We solve this problem for every constraint with $y\sigma(y)$ decreasing.

Theorem C (constrained equilibrium)**.** *Let σ be a positive, locally Hölder continuous density on $(0, Y]$, $0 < Y \leq \infty$, with $y\sigma(y)$ strictly decreasing and, if $Y = \infty$, $\int_1^\infty \sigma(y) \log y \, \mathrm{d}y < \infty$. Then the soft-edge condition $\int_a^Y \sigma(y)(1 - a/y)^{-1/2}\, \mathrm{d}y = 1$ has exactly one root $a \in (0, Y)$, and the constrained equilibrium measure is $\nu\, \mathrm{d}x|_{[0,a]} + \sigma\, \mathrm{d}y|_{[a,Y]}$ with the explicit density ν of (3.2), $0 < \nu < \sigma$. For the lattice $a = 4c(p)$. For a counting function $\mathcal{N}(\lambda) \sim \sum_\rho c_\rho \lambda^\rho$, the root $a_n = 4c(1 + \eta_n) n^{-p}$ of $\int (1 - a_n \lambda)^{-1/2} \mathbf{1}_{a_n \lambda < 1}\, \mathrm{d}\mathcal{N}(\lambda) = n$ satisfies*

$$n\eta_n = p \sum_{j \geq 0} W(j) Z_{\mathcal{N}}(-j) n^{-jp} + p \sum_q R_q W(q)\, \pi \cot(\pi q)\, n^{-pq} + O(n\eta_n^2),$$

the second sum over the poles $q \neq -1/p$ of $Z_{\mathcal{N}}(-w) = \int \lambda^w\, \mathrm{d}\mathcal{N}(\lambda)$, with residues R_q (Proposition 3.4).

The proof of the equilibrium statement uses no input from outside this series: the positivity of the logarithmic energy of signed measures of mass zero, without compact support, is [10, Lemma 3.3].

The fourth result explains the natural index and the evenness. In the residue recursion the exponent α plays two roles, the Bessel order ν of the hard edge and the Szegő exponent β of the weight; separating them gives formal series $B(n; \beta, \nu) = \sum_k d_k(p, \beta, \nu) n^{-k}$, the integer layer of a decoupled model (Section 6), with $B(n; \alpha, \alpha)$ the integer layer of the lattice.

Theorem D (shift and evenness)**.** *Identically in (p, β, ν), $B(n; \beta, \nu) = (n/N)^p F_p(N; \nu)$ with $N = n + \frac{\beta+1}{2}$ and a formal series F_p that does not depend on β; moreover $F_p(N; \nu) = 1 + \sum_{k \geq 1} f_{2k}(p, \nu^2) N^{-2k}$ is even in N and in ν. The zeta layers of weight at most two (μ_1, μ_2, μ_1^2) obey the same shift. In particular, for the lattice and every $p \neq 1$, the integer layer of $n^p b_n / c$ is $(n/N)^p (1 + f_2 N^{-2} + f_4 N^{-4} + \cdots)$ to every order, with no odd powers of $1/N$.*

This is Theorem 6.3 with Corollary 6.4. The evenness comes from an exact reflection $d_k(p, -\beta - 2, \nu) = (-1)^k d_k(p, \beta, \nu)$ (Theorem 6.1); the shift is proved by steepest descent of the same lattice problem in a second frame, of non-integer index $n + \gamma$, in which the Szegő exponent is $\alpha - 2\gamma$ and the Bessel order is still α (Lemma 6.5), and by comparing the two expansions of b_n. In the same model the zeta layers are Toda flows of the integer layer, with the times $-2\zeta(-jp)/j$ (Proposition 6.6).

The fifth result treats the first node sets other than the lattice. For the Hurwitz lattice $\lambda_k = (k+\delta)^p$ the offset δ changes only the Szegő exponent, $\beta' = \alpha + 2\delta$, at fixed Bessel order, and Theorem D turns this into a shifted natural index.

Theorem E (Hurwitz lattices)**.** *Let $p \neq 1$, $-1 < \delta \leq 0$ ($\delta < 0$ if $p < 1$), $a = 1+\delta$ and $1 < s < 1+2p$. For σ_Λ with $\lambda_k = (k+\delta)^p$, the double expansion of $n^p b_n/c$ is that of the decoupled model with $(\beta,\nu) = (\alpha+2\delta,\alpha)$ and band-factor parameters $2\zeta(-jp,a)/(jn^{jp})$. Consequently expansion (1.9) holds with N replaced by $N_\delta = n + \frac{\alpha}{2} + \frac{1}{2} + \delta$ and $\zeta(-jp)$ by $\zeta(-jp,a)$, with the same f_{2k}, g_j; the integer layer and the zeta layers of weight at most two are those of the lattice in N_δ to all orders.*

For the shifted lattices $\lambda_k = (k+\delta)^p + \kappa$, $p > 1$, whose node zeta function has a pole at $q = 1 - \frac{1}{p}$, we prove that $n^p b_n/c$ is the Hurwitz expansion plus the pole term $-2(2-p)c\kappa n^{-p} = pR_q W(q)\pi\cot(\pi q) n^{-p}$ of Theorem C, up to $O(n^{-1-p})$ (Theorem 7.2).

1.3. **The spectral dictionary.** For the lattice, Theorems A and B identify the coefficients of b_n with values of ζ, and Theorem E does the same for Hurwitz lattices with values of $\zeta(\cdot,a)$. For a general node set the Riemann–Hilbert analysis has not been carried out, and we formulate what the theorems and the numerical evidence suggest (Section 9). In the natural index $N_\Lambda := n + \frac{\alpha}{2} - Z_\Lambda(0)$ (which is (1.8) for the lattice, $\zeta(0) = -\frac{1}{2}$), the expansion of $n^p b_n/c$ should read, at leading order in each spectral datum,

$$\Big(\frac{n}{N_\Lambda}\Big)^p \Big[1 + \frac{f_2}{N_\Lambda^2} + \cdots + \sum_{j\geq 1} \frac{W(j)\, p\, Z_\Lambda(-j)}{N_\Lambda^{1+jp}} + \sum_q \frac{p\, R_q\, W(q)\, \pi\cot(\pi q)}{N_\Lambda^{1+pq}} + \cdots\Big],$$

the second sum running over the poles q of $Z_\Lambda(-w) - \zeta(-pw)$, with w-residues R_q. The integer residues of $\pi\cot(\pi w) W(w) p Z_\Lambda(-w)$ are the zeta terms of Theorem B; its residues at the poles of Z_Λ are the pole terms of Theorem C. In a refined form (Conjecture 9.5) the expansion is the decoupled model with Szegő exponent $\alpha - 2Z_\Lambda(0) - 1$, flowed along the Toda hierarchy with times read off from $Z_\Lambda(-j)$ and from the density and mass of the nodes, plus the pole terms; it is proved for Hurwitz lattices and, at first order in the poles, for their constant shifts, and was tested on nine further measures with predictions fixed before the measures were computed (§8.5); one coefficient outside its claimed precision failed and led to a refinement (Remark 9.7). The dictionary is exact for the measures on reciprocal squared Bessel zeros (Proposition 5.2), it reads the index offset of the spectrum from $Z_\Lambda(0)$, and it predicts, for the eigenvalues of $-\frac{\mathrm{d}^2}{\mathrm{d}q^2} + q^N$ with N even, an n^{-2} shift equal to $(N-1)/(6(N+2))$ (Section 8.7). For drums at $p = 1$, outside the range of the theorems, the perimeter appears at order $n^{-1/2}$ and the corners at order n^{-1}, as the kernel predicts (Section 8.6). The dictionary has a limit: subleading terms of each pole, one order lower, are not given by it (Remark 9.2), and for spectra whose nodes deviate from a smooth node function by a fraction of the spacing the expansion is replaced by noise of relative size $n^{-1/2}$, whose variance appears numerically to be set by the spectral form factor (Section 8.8).

1.4. **Organisation and notation.** Section 2 restates the results of the earlier papers that are used, together with two results of the same project that extend them by one order. Section 3 proves Theorem C, Section 4 proves Theorem B, and Section 5 proves Theorem A and its general form. Section 6 introduces the decoupled model and proves Theorem D, and states the Toda structure of the zeta layers. Section 7 proves Theorem E and the theorem on shifted Hurwitz lattices; the analytic estimates of these two sections are transfers of those of [11], which we outline and do not reproduce. Section 8 collects the examples and the numerical

evidence, including the pre-registered tests of Conjecture 9.5, and Section 9 the conjectures. Section 10 relates the results to the literature. Appendix A lists the scripts.

Throughout, $p > 0$ and $c = c(p)$; α, y_n, W, ε_j, N are as in (1.4)–(1.8). The Bessel order α is that of [11], and $\mathsf{k}_p = \varepsilon_1$; the constant κ_p of [10] does not occur. We write $\binom{2q}{q} = \Gamma(2q+1)/\Gamma(q+1)^2$ for complex q. The letter N denotes the natural index, except in Section 8.7, where q^N is a potential. Constants C may change from line to line. Numbers obtained by fitting floating-point data are quoted with the spread of the fit over fit windows and nuisance exponents in parentheses; such a spread is not an error bound.

2. Setting and prior results

This section fixes the Riemann–Hilbert frame of [11] for the lattice $\sigma_{p,s}$ and restates the expansions that the later sections refine. Theorem 2.1 is proved in [11]. Theorem 2.5 and Lemmas 2.2–2.4 extend it by one order and to all band-factor parameters; their proofs are third-order versions of [11, Sections 8–10] and are not reproduced here.

2.1. The first two orders.

Theorem 2.1 ([11, Theorem 1.4])**.** *Let $s > 1$. If $p \geq 1$, then $n^p b_n/c = 1 - \frac{s-1}{2n} + O(n^{-2})$ and $n^p a_n/(2c) = 1 - \frac{s-1+p}{2n} + O(n^{-2})$. If $0 < p < 1$, both brackets carry the further term $\mathsf{k}_p n^{-1-p}$ and the remainder is $O(n^{-1-2p} + n^{-2})$.*

The $1/n$ coefficient of $n^p b_n/c$ is $d_1 := -(s-1)/2 = -p(\alpha+1)/2$. Since $(1+\delta/n)^{-p} = 1 - p\delta/n + O(n^{-2})$, it is the first term of the index shift $(n/N)^p$ with $N = n + \frac{\alpha+1}{2}$; this is the origin of (1.8).

2.2. The frame.

We use the notation of [11, Sections 4, 8 and 10]. After the rescaling $z = n^p x$ the polynomials oscillate on the band $[0, 4c]$, with a hard edge at 0 and a soft edge at $4c$. The weight on the band is $W_0 = \mathsf{w}_n z^\alpha (4c - z) H_n(z)$, where $\mathsf{w}_n > 0$ is a constant and H_n is the *band factor*, which collects the discreteness of the nodes. With $\mathcal{R}(z) := (z(z-4c))^{1/2}$ (branch $\sim z$ at ∞), $\mathsf{T} := \frac{1}{\sqrt{2}}\left(\begin{smallmatrix} 1 & 1 \\ i & -i \end{smallmatrix}\right)$ and $P_0^\infty = \mathsf{T}((z-4c)/z)^{\sigma_3/4}\mathsf{T}^{-1}$, the error matrix R of the steepest-descent analysis has jumps $I + O(1/n)$ on two small circles around 0 and $4c$ and exponentially small jumps elsewhere, and

$$n^{2p} b_n^2 = (Q_1)_{12}(Q_1)_{21}, \qquad Q_1 = P_1 + \mathsf{R}_1, \tag{2.1}$$

where P_1 and R_1 are the $1/z$ coefficients at infinity of the global parametrix and of R [11, Lemmas 4.1, 4.2 and Proposition 5.6]. The subscript "mod" denotes quantities computed for the *model weight* $z^\alpha(4c - z)$, i.e. with H_n replaced by a constant.

The band factor is $H_n = e^{-2G}$ with an explicit function G [11, Lemmas 8.1 and 10.1]. For $p > 1$, G has a jump $i\mathfrak{e}$ on $(-\infty, 0)$, with $|\mathfrak{e}(y)| \leq \pi \exp(-2\pi \sin(\pi/p) n |y|^{-1/p})$, and

$$G(z) - G(1) = \frac{1}{2\pi} \int_{-\infty}^{0} \mathfrak{e}(y)\, \frac{z-1}{(y-z)(y-1)}\, \mathrm{d}y \qquad (p > 1) \tag{2.2}$$

[11, Lemma 8.1]. For $0 < p < 1$, G jumps across the two seams $\arg z = \pm\vartheta$, $\vartheta = \pi p/2$, by $\mathfrak{j}(|z|)$, where $\mathfrak{j}(r) := -\log(1 - e^{-2\pi n r^{-1/p}})$, and

$$G(z) - G(1) = \frac{1}{2\pi i} \int_0^\infty \mathfrak{j}(r) \left[e^{i\vartheta} \mathfrak{K}(re^{i\vartheta}, z) - e^{-i\vartheta} \mathfrak{K}(re^{-i\vartheta}, z) \right] \mathrm{d}r, \qquad \mathfrak{K}(\zeta, z) := \frac{z-1}{(\zeta - z)(\zeta - 1)} \tag{2.3}$$

[11, Lemma 10.1(ii)]. Expanding $\mathfrak{K}$ in powers of z/ζ gives the moments of the jump densities:

Lemma 2.2. *Let $p \neq 1$ and $k \geq 1$. The k-th moment of the jump density, $m_k := \frac{1}{2\pi}\int_{-\infty}^{0} \mathfrak{e}(y) y^{-k-1}\,\mathrm{d}y$ for $p > 1$ and the corresponding seam integral for $p < 1$, equals $\zeta(-kp)/(kn^{kp})$. For every $K \geq 1$,*

$$\log H_n(z) - \log H_n(1) = -\sum_{k \leq K} \mu_{n,k}(z^k - 1) + O(n^{-(K+1)p}), \qquad \mu_{n,k} := \frac{2\zeta(-kp)}{kn^{kp}},$$

uniformly for $|z| \leq 4c + 2$, boundary values included, with one derivative.

For $p < 1$ this is [11, Lemma 10.1(iii)–(iv)]. For $p > 1$ the proof is the same computation on the negative axis: with $q(r) = \exp(-a_n r^{-1/p})$, $a_n = 2\pi i n e^{-i\pi/p}$, $\operatorname{Re} a_n = 2\pi n \sin(\pi/p) > 0$, one expands $\mathfrak{e} = -2\operatorname{Im}\sum_m q^m/m$, integrates termwise by [5, (5.9.1)], sums by [5, (25.2.1)] and applies the reflection formula [5, (25.4.1)]. At $p = 2$ all m_k vanish and H_n is constant, in agreement with [11, Lemma 8.1].

2.3. **Tilts and the gauge.** For $\mu = (\mu_1, \dots, \mu_K) \in \mathbb{C}^K$ put $q_\mu(x) := \exp(-\sum_k \mu_k x^k)$ and

$$Q_k(z) := \sum_{j=0}^{k-1} \binom{2j}{j} c^j z^{k-1-j}, \qquad M_\mu(z) := \sum_k \mu_k Q_k(z), \tag{2.4}$$

so that $Q_k(4c) = c^{k-1}k\binom{2k}{k}/2$. The arcsine moments $\frac{1}{\pi}\int_0^{4c} x^j (x(4c-x))^{-1/2}\,\mathrm{d}x = \binom{2j}{j}c^j$ give the Szegő function of the tilt explicitly.

Lemma 2.3 (gauge)**.** *For the weight $z^\alpha(4c - z)q_\mu(z)$, the jump data on the two circles are $\Delta_k^\mu = G_\mu \Delta_k^{\mathrm{mod}} G_\mu^{-1}$, where, with $h := \mathcal{R}M_\mu/2$ and $S := \sinh(h)/\mathcal{R}$,*

$$G_\mu(z) = P_0^\infty e^{-\mathcal{R}M_\mu \sigma_3/2}(P_0^\infty)^{-1} = \mathsf{T}\begin{pmatrix} \cosh h & -(z-4c)S \\ -zS & \cosh h \end{pmatrix}\mathsf{T}^{-1}$$

is entire in (z, μ) with $\det G_\mu = 1$. In particular Δ_k^μ is meromorphic in each disc, with Laurent coefficients entire in μ.

By Lemma 2.2, the band factor is the tilt q_{μ_n} with $\mu_n = (\mu_{n,1}, \dots, \mu_{n,K})$, up to $O(n^{-(K+1)p})$. The first-order effect of a tilt on b_n is exactly linear:

Lemma 2.4 (linear response)**.** *Let $e(\mu)$ be the $1/n$ coefficient of $\big((P_1 + \mathsf{R}_1^{(1)}(\mu)/n)_{12}(P_1 + \mathsf{R}_1^{(1)}(\mu)/n)_{21}\big)^{1/2}/c$ minus d_1, where $\mathsf{R}_1^{(1)}(\mu) = \operatorname{Res}_0 \Delta_1^\mu + \operatorname{Res}_{4c} \Delta_1^\mu$. Then, for every $\mu \in \mathbb{C}^K$,*

$$e(\mu) = cp\, M_\mu(4c) = cp \sum_k \mu_k Q_k(4c).$$

The proof expands G_μ at $4c$ and at 0. With $M_4 := M_\mu(4c)$, $G_\mu(0)$ commutes with the residue of Δ_1^{mod} at 0, $M_\mu'(4c)$ drops out, and the residue at $4c$ moves by

$$M_4 c^2 p \begin{pmatrix} M_4 c - \alpha - 1 & -i(M_4 c - \alpha - 2) \\ -i(M_4 c - \alpha) & -(M_4 c - \alpha - 1) \end{pmatrix},$$

which is quadratic in M_4 but changes $i(A_{21} - A_{12})/(2c)$, the combination that enters the $1/n$ coefficient, by exactly cpM_4. The identity $Q_k(4c) = c^{k-1}k\binom{2k}{k}/2$ follows from the recursion $a_{k+1} = 4a_k + \binom{2k}{k}$, $a_1 = 1$, satisfied by both $\sum_{j<k} 4^{k-1-j}\binom{2j}{j}$ and $k\binom{2k}{k}/2$.

2.4. **The second order.**

Theorem 2.5. *Let $p > 0$, $p \neq 1$, $s > 1$ and $K \geq 1$. Then*

$$\frac{n^p b_n}{c} = 1 + \frac{d_1}{n} + \frac{d_2}{n^2} + \sum_{j=1}^{K} \frac{\varepsilon_j}{n^{1+jp}} + O\big(n^{-3} + n^{-2-p} + n^{-1-(K+1)p}\big),$$

$$d_2 = d_2(p,s) = \frac{(p+1)(s-1)^2}{8p} - \frac{(p+2)(s-1-p)^2}{8p^2} + \frac{p+1}{12}.$$

In particular, for $0 < p < 1$ with $1/p \notin \mathbb{N}$ the n^{-2} coefficient is d_2; for $p = 1/j_0$, $j_0 \geq 2$, it is $d_2 + \varepsilon_{j_0} = d_2 - p/12$, because $W(j_0) = \binom{2j_0}{j_0} c (1/j_0)^{j_0} = 1$.

For $p > 1$ the terms with $j \geq 2$ are $O(n^{-3})$, and the statement is $n^p b_n/c = 1 + d_1/n + d_2/n^2 + \mathsf{k}_p n^{-1-p} + O(n^{-3})$. For $p = 2$, $d_2 = (s^2 + 6s - 11)/16$ and $\mathsf{k}_p = 0$. The proof of Theorem 2.5 is the second Neumann iterate of the small-norm problem, with the second asymptotic coefficients of the Bessel and Airy functions on the two circles and the band factor entering through G_{μ_n}; the $1/n^2$ term at $\mu = 0$ is an exact Laurent computation, and the zeta terms are Lemma 2.4 with $\mu_{n,j} \cdot cpQ_j(4c) = \varepsilon_j n^{-jp}$. Section 5 carries this out at every order.

3. The constrained equilibrium

At leading order the zeros of P_n, rescaled by n^p, follow the equilibrium measure of unit mass constrained by the continuum density of the nodes. For the lattice this problem was solved in [10, Theorem 1.3] in the index variable. Here we solve it for a general constraint density in the edge variable, by a Cauchy-transform argument that gives the density explicitly, and we show that the soft edge of the discrete analogue carries the poles of the node zeta function with the weight $\pi \cot(\pi q)$. Nothing in this section concerns b_n directly; the link is discussed in Section 9.

3.1. **Theorem C.** Let $0 < Y \leq \infty$ and let σ be a density on $(0, Y]$ such that

- (H1) σ is positive and locally Hölder continuous on $(0, Y)$ (the Hölder condition is used only in (iii));
- (H2) $y \mapsto y\sigma(y)$ is strictly decreasing on $(0, Y)$;
- (H3) if $Y = \infty$, then $\int_1^\infty \sigma(y) \log y \, \mathrm{d}y < \infty$.

By (H2), $g(y) := \sqrt{y}\,\sigma(y) = y\sigma(y)/\sqrt{y}$ is strictly decreasing, and $\sigma(y) \geq y_0\sigma(y_0)/y$ for $y < y_0$, so $\int_0 \sigma = \infty$; if $Y = \infty$, (H3) gives $\int^\infty \sigma < \infty$ and hence $y\sigma(y) \to 0$. For a positive measure μ on $[0, Y]$ let $U^\mu(z) := \int \log \frac{1}{|z-x|} \, \mathrm{d}\mu(x)$ and $I(\mu) := \int U^\mu \, \mathrm{d}\mu$. The *constrained equilibrium problem* is to minimise $I(\mu)$ over positive μ on $[0, Y]$ with $\mu([0, Y]) = 1$ and $\mu \leq \sigma$. The *soft-edge condition* is

$$F(a) := \int_a^Y \sigma(y) \Big(1 - \frac{a}{y}\Big)^{-1/2} \mathrm{d}y = 1. \tag{3.1}$$

Theorem 3.1. *Assume* (H1)–(H3).

- (i) *Equation* (3.1) *has exactly one root $a \in (0, Y)$.*
- (ii) *Put $s(y) := \sigma(y)\sqrt{y/(y-a)}$ on (a, Y) and*

$$\nu(x) := \frac{1}{\pi}\sqrt{\frac{a-x}{x}} \int_a^Y \frac{s(y)}{y-x} \, \mathrm{d}y \qquad (0 < x < a). \tag{3.2}$$

Then $\mu := \nu \, \mathrm{d}x|_{[0,a]} + \sigma \, \mathrm{d}y|_{[a,Y]}$ has unit mass, $0 < \nu < \sigma$ on $(0, a)$, and $\pi\sqrt{x(a-x)}\,\nu(x) = 1 - \int_a^Y \sigma(y)\sqrt{y(y-a)}\,(y-x)^{-1} \, \mathrm{d}y$.

(iii) *There is a constant ℓ with $U^\mu = \ell$ on $[0,a]$ and $U^\mu < \ell$ on $(a,Y]$.*
(iv) *μ is the unique solution of the constrained equilibrium problem.*

Proof. (i) Substituting $y = au$, $F(a) = \int_1^{Y/a} a\sigma(au)(1-1/u)^{-1/2}\,\mathrm{d}u$. For fixed u, $a\sigma(au) = (au\,\sigma(au))/u$ is strictly decreasing in a by (H2), and the upper limit decreases, so F is strictly decreasing. On $[a_0,a_1] \subset (0,Y)$ we have $a\sigma(au) \le a_0\sigma(a_0u)$ by (H2), so F is continuous by dominated convergence. As $a \to 0$, $F(a) \ge \int_a^Y \sigma \to \infty$. If $Y < \infty$, $F(a) \to 0$ as $a \to Y$. If $Y = \infty$, split at $2a$: $\int_{2a}^\infty \le \sqrt{2}\int_{2a}^\infty \sigma \to 0$, and, since $\sigma(y) \le a\sigma(a)/y$ on $[a,2a]$, $\int_a^{2a} \le a\sigma(a)\int_1^2 (1-1/u)^{-1/2}\,\mathrm{d}u \to 0$.

(ii) Let $S(z) := \int_a^Y s(y)(z-y)^{-1}\,\mathrm{d}y$ and $\tilde G(z) := ((z-a)/z)^{1/2}S(z)$ (principal branch), analytic off $[0,Y]$, and let $G(z) := \int (z-x)^{-1}\,\mathrm{d}\mu(x)$. For fixed $y \in (a,Y)$ put

$$k_y(z) := \frac{((z-a)/z)^{1/2} - \sqrt{(y-a)/y}}{z-y}, \qquad \rho_y(x) := \frac{1}{\pi}\sqrt{\frac{a-x}{x}}\,\frac{1}{y-x} > 0 \quad (0<x<a).$$

The singularity of k_y at $z=y$ is removable, so k_y is analytic off $[0,a]$; it is $O(|z|^{-1/2})$ at 0, bounded at a, and $\sim (1-\sqrt{(y-a)/y})/z$ at infinity; since $((z-a)/z)^{1/2}$ takes the values $\pm i\sqrt{(a-x)/x}$ on $(0,a)$ from above and below, k_y jumps by $-2\pi i\rho_y$ there. As $[0,a]$ is compact, Liouville's theorem gives $k_y(z) = \int_0^a \rho_y(x)(z-x)^{-1}\,\mathrm{d}x$, and ρ_y has mass $1-\sqrt{(y-a)/y}$. Since $\sigma = s\sqrt{(y-a)/y}$,

$$\tilde G(z) = \int_a^Y s(y)\Big[\frac{\sqrt{(y-a)/y}}{z-y} + k_y(z)\Big]\,\mathrm{d}y = \int_a^Y \frac{\sigma(y)\,\mathrm{d}y}{z-y} + \int_0^a \frac{\nu(x)\,\mathrm{d}x}{z-x} = G(z),$$

by Fubini's theorem (everything is positive, $|z-x|^{-1} \le |\operatorname{Im} z|^{-1}$, and $\int s(1-\sqrt{(y-a)/y})\,\mathrm{d}y \le F(a)$), because $\int s(y)\rho_y(x)\,\mathrm{d}y = \nu(x)$. Hence μ has mass $\int_a^Y \sigma + \int s(1-\sqrt{(y-a)/y}) = \int s = F(a) = 1$; $\operatorname{Re} G(x \pm i0) = 0$ on $(0,a)$, where S is real-analytic; and $\nu > 0$ because $s > 0$. With $R(y) := \sqrt{y(y-a)}$ one has $R\sigma = s\,(y-a)$, which gives $\tilde G = R(z)^{-1}\big[F(a) + \int \sigma R\,(z-y)^{-1}\,\mathrm{d}y\big]$; the difference of the two expressions for ν is $(F(a)-1)/(\pi\sqrt{x(a-x)})$, which vanishes by (3.1). For the upper bound write $s(y) = g(y)(y-a)^{-1/2}$ and use $\int_a^\infty (y-a)^{-1/2}(y-x)^{-1}\,\mathrm{d}y = \pi(a-x)^{-1/2}$:

$$\nu(x) < \frac{1}{\pi}\sqrt{\frac{a-x}{x}}\,g(a)\,\frac{\pi}{\sqrt{a-x}} = \frac{g(a)}{\sqrt{x}} \le \frac{g(x)}{\sqrt{x}} = \sigma(x) \qquad (0<x<a).$$

(iii) On $(0,a)$, U^μ is constant because $\operatorname{Re} G = 0$ there. For $z \in (a,Y)$, $\varphi := U^\mu - \ell$ satisfies $\varphi(a) = 0$ and $\varphi'(z) = -\operatorname{Re} G(z \pm i0) = -((z-a)/z)^{1/2}H(z)$ with $H(z) := \mathrm{PV}\int_a^Y s(y)(z-y)^{-1}\,\mathrm{d}y$, which exists since σ is locally Hölder. Writing $s = g\cdot(y-a)^{-1/2}$,

$$H(z) = \int_a^Y \frac{(g(y)-g(z))(y-a)^{-1/2}}{z-y}\,\mathrm{d}y + g(z)\,\mathrm{PV}\int_a^Y \frac{(y-a)^{-1/2}}{z-y}\,\mathrm{d}y.$$

The first integrand is positive since g is decreasing. The principal value is 0 if $Y=\infty$, and $(z-a)^{-1/2}\log\frac{\sqrt{Y-a}+\sqrt{z-a}}{\sqrt{Y-a}-\sqrt{z-a}} > 0$ if $Y < \infty$. Hence $H > 0$, $\varphi' < 0$ and $\varphi < 0$ on $(a,Y]$.

(iv) Since $\log^+|x-y| \le \log(1+x) + \log(1+y)$ and (H3) bounds the logarithmic moment of every $\mu' \le \sigma$, $I(\mu') > -\infty$ for every competitor; let $\mu' \le \sigma$ have unit mass and $I(\mu') < \infty$, and $\tau := \mu' - \mu$. Then $I(\mu') - I(\mu) = I(\tau) + 2\int (U^\mu - \ell)\,\mathrm{d}\tau$. The integrand vanishes on $[0,a]$ (at every point, since the density of μ is at most $g(a)x^{-1/2}$ on $(0,a)$ and bounded on $[a,Y]$, so U^μ is continuous); on $(a,Y]$, $\tau \le 0$ and $U^\mu - \ell < 0$, so the integral is ≥ 0. For $I(\tau)$: τ is a finite signed measure with $\tau(\mathbb{R}) = 0$, and $\iint |\log|x-y||\,\mathrm{d}|\tau|\,\mathrm{d}|\tau| < \infty$ (the positive part of

the logarithm by (H3), the negative part because μ has density at most $g(a)\max(x^{-1/2}, a^{-1/2})$ and $I(\mu') < \infty$). So $I(\tau) \geq 0$, with equality only if $\tau = 0$, by [10, Lemma 3.3]. □

Remark 3.2. (a) The proof of (ii)–(iii) uses only that g is decreasing; (H2) enters through the monotonicity of F in (i). So (H2) may be replaced by "$\sqrt{y}\,\sigma(y)$ strictly decreasing and $\int_0^Y \sigma > 1$"; uniqueness of the root then follows from (iv), since a is the left end of the saturated set of the unique minimiser. Neither condition is necessary, but they cannot simply be dropped: in a discretised energy minimisation, a constraint density with a dip below the edge gives three roots of (3.1) and a minimiser with two bands, and a shallower dip gives one root but a saturated interval inside the band. (b) Without (3.1), ν acquires a multiple of $(a-x)^{-1/2}$, which violates $\nu \geq 0$ or $\nu \leq \sigma$. Under (3.1), $\nu(a^-) = g(a)/\sqrt{a} = \sigma(a)$, so μ is continuous at a. If σ is Hölder of order $> \frac{1}{2}$ near a (for instance C^1), then $\sigma - \nu = C\sqrt{a-x}\,(1+o(1))$ with $C = -h(a)/(\pi\sqrt{a}) > 0$, where $h(a) := \int_a^Y (g(y)-g(a))(y-a)^{-3/2}\,\mathrm{d}y - 2g(a)(Y-a)^{-1/2}\mathbf{1}_{Y<\infty}$. (c) In node language (3.1) is the statement that $\frac{1}{n}\sum_{u_k<1}(1-u_k)^{-1/2} \approx 1$, with $u_k = a\lambda_k/n^p$: the edge is fixed by a sum of the per-node quantity $(1-u)^{-1/2}$ over the saturated nodes.

Corollary 3.3 (the lattice)**.** *For $\sigma(y) = \frac{1}{p}y^{-1/p-1}$ on $(0,\infty)$ the root of (3.1) is $a = 4c(p)$.*

Proof. $y\sigma(y) = \frac{1}{p}y^{-1/p}$ is strictly decreasing and (H3) holds. Equation (3.1) reads $a^{-1/p}\frac{1}{p}B(\frac{1}{p},\frac{1}{2}) = 1$, i.e. $a^{1/p} = \sqrt{\pi}\,\Gamma(\frac{1}{p}+1)/\Gamma(\frac{1}{p}+\frac{1}{2}) = \pi/(2I(p))$, and $a = 4c(p)$ by (1.3). □

The density $\sigma(y) = \frac{1}{p}y^{-1/p-1}$ is the image of Lebesgue measure on the index variable u under $y = u^{-p}$, so Corollary 3.3 concerns the problem of [10, Theorem 1.3]; its edge $a = 4c$ is the right end of the interval on which the potential of the measure μ^* there is constant [10, Theorem 1.3(ii)]. The derivation above does not use the Riemann–Hilbert analysis of [11].

3.2. **The soft edge carries the poles.** Let $L(u) := (1-u)^{-1/2}\mathbf{1}_{u<1}$; its Mellin transform $\widetilde{L}(w) := \int_0^\infty u^{-w-1}L(u)\,\mathrm{d}u$ is computed in Lemma 4.1: $\widetilde{L}(w) = B(-w,\frac{1}{2}) = -\pi\cot(\pi w)\,\Gamma(w+\frac{1}{2})/(\sqrt{\pi}\,\Gamma(w+1))$, so that

$$-(4c)^q\,\widetilde{L}(q) = W(q)\,\pi\cot(\pi q). \tag{3.3}$$

At $q \in -\frac{1}{2} - \mathbb{N}_0$ the right side is a limit of the form $0\cdot\infty$ and is defined by the left side; for instance $\widetilde{L}(-\frac{1}{2}) = \pi$.

Let $\mathcal{N}$ be a C^1 counting function on $[\lambda_0,\infty)$, $\lambda_0 > 0$, with $\mathcal{N}' > 0$, and suppose that for exponents $\rho_0 = 1/p > \rho_1 > \rho_2 > \cdots$ and constants c_ρ with $c_{1/p} = 1$,

$$\mathcal{N}'(\lambda) = \sum_{k<K} \rho_k c_{\rho_k}\lambda^{\rho_k-1} + O(\lambda^{\rho_K-1}) \qquad (\lambda\to\infty) \tag{3.4}$$

for every K (if the list is finite, the remainder is $O(\lambda^{-M})$ for every M). Then $Z_{\mathcal{N}}(s) := \int_{\lambda_0}^\infty \lambda^{-s}\,\mathrm{d}\mathcal{N}(\lambda)$ continues meromorphically to $\mathbb{C}$, with simple poles at $s = \rho$ of residue ρc_ρ. In the variable $w = -s$, $w \mapsto Z_{\mathcal{N}}(-w)$ has poles at $q = -\rho$ with residues $R_q = -\rho c_\rho$; the *Weyl pole* is $q = -1/p$, $R_{-1/p} = -1/p$. We assume that $\rho c_\rho = 0$ whenever $\rho \in \{-1,-2,\dots\}$, i.e. no pole of $Z_{\mathcal{N}}(-w)$ sits at a positive integer.

Proposition 3.4. *Under these assumptions, as $A \to 0^+$,*

$$\int_{\lambda_0}^\infty L(A\lambda)\,\mathrm{d}\mathcal{N}(\lambda) \sim -\sum_q R_q\widetilde{L}(q)A^q + \sum_{j\geq 0}\binom{2j}{j}4^{-j}Z_{\mathcal{N}}(-j)A^j, \tag{3.5}$$

with remainder $O(A^{-\rho_K})$ after the terms with $q < -\rho_K$ and $j < -\rho_K$. Consequently the root of $\int L(A\lambda)\,\mathrm{d}\mathcal{N}(\lambda) = n$, written $A = 4c(1+\eta)n^{-p}$, satisfies

$$\eta = \frac{p}{n}\Big[\sum_{j\ge 0} W(j)Z_{\mathcal{N}}(-j)n^{-jp} + \sum_{q\neq -1/p} R_q W(q)\,\pi\cot(\pi q)\,n^{-pq}\Big] + O(\eta^2), \tag{3.6}$$

and the higher orders follow by reversion of the series.

Proof. For one power, $\mathrm{d}\mathcal{N} = \rho c_\rho \lambda^{\rho-1}\,\mathrm{d}\lambda$, the substitution $u = A\lambda$ and the Taylor expansion of $(1-u)^{-1/2}$ on $[0, A\lambda_0]$ give

$$\int_{\lambda_0}^{1/A} (1-A\lambda)^{-1/2}\rho c_\rho \lambda^{\rho-1}\,\mathrm{d}\lambda = \rho c_\rho\Big[A^{-\rho}B(\rho,\tfrac12) - \sum_{j\ge 0}\binom{2j}{j}4^{-j}A^j\frac{\lambda_0^{\rho+j}}{\rho+j}\Big],$$

first for $\rho > 0$ and then for all $\rho \notin -\mathbb{N}$ by analytic continuation in ρ. The first term is $-R_q\widetilde{L}(q)A^q$ with $q = -\rho$. For the remainder r_K of (3.4), split at $\lambda = 1/(2A)$: on $[\lambda_0, 1/(2A)]$ expand $(1-A\lambda)^{-1/2}$ in powers of $A\lambda$, and on $[1/(2A), 1/A]$ use $|r_K| \le C\lambda^{\rho_K-1}$; both give the integer terms up to $O(A^{-\rho_K})$. The coefficient of A^j collects $-\rho c_\rho \lambda_0^{\rho+j}/(\rho+j)$ over the powers and $\int \lambda^j r_K\,\mathrm{d}\lambda$, i.e. the continued value $Z_{\mathcal{N}}(-j)$. This proves (3.5). The Weyl pole contributes $\frac{1}{p}B(\frac1p,\frac12)A^{-1/p} = n(1+\eta)^{-1/p}$ by Corollary 3.3, so the equation is $n(1+\eta)^{-1/p} + (\text{rest}) = n$; by (3.3) and $\binom{2j}{j}4^{-j}(4c)^j = W(j)$, the rest at $\eta = 0$ is the bracket of (3.6), and the implicit function theorem gives (3.6). □

The termwise proof avoids shifting the contour of the Mellin–Barnes form

$$\frac{1}{2\pi i}\int \widetilde{L}(w)Z_{\mathcal{N}}(-w)A^w\,\mathrm{d}w$$

of (3.5), which would need bounds on $Z_{\mathcal{N}}$ on vertical lines. The numbers $Z_{\mathcal{N}}(-j)$ depend on the cutoff λ_0; they are not the values $Z_\Lambda(-j)$ of a node set, which enter through the band factor (Section 4). What (3.6) contributes for a node set is the pole part: the weight $W(q)\pi\cot(\pi q)$ of each pole of $Z(-w)$ other than the Weyl pole.

Remark 3.5 (exact root for a drum-type count). For $p = 1$ ($c = \frac12$) and $\mathrm{d}\mathcal{N} = (1 - \frac{\beta}{2}\lambda^{-1/2})\,\mathrm{d}\lambda$ on $(0,\infty)$, the condition is $2/A - \frac{\pi\beta}{2}A^{-1/2} = n$. With $A = 2(1+\eta)/n$ and $v := (1+\eta)^{-1/2}$ it reads $v^2 - \epsilon v - 1 = 0$, $\epsilon := \pi\beta/(2\sqrt{2n})$, so $1+\eta = v^{-2} = 1 - \epsilon + \epsilon^2/2 + O(\epsilon^3)$ and

$$n\eta = -\frac{\pi\beta}{2\sqrt{2}}\,n^{1/2} + \frac{\pi^2}{16}\beta^2 + O(n^{-1/2}).$$

The first term is the pole term of (3.6) at $q = -\frac12$, $R_q = \beta/2$, with $W(q)\pi\cot(\pi q) = -\pi(4c)^{-1/2}$; the second is quadratic in the pole residue and is produced by the reversion. Both are compared with b_n in Section 8.6.

Remark 3.6 (numerical checks). For the lattice at $p = 0.6$ and $p = 1.5$, for the drum density of Remark 3.5 cut at the first node, and for a fourth density, the root, the two forms of ν, the mass, $0 < \nu/\sigma < 1$ and the outer inequality of Theorem 3.1 were evaluated by quadrature, the outer inequality both through H and by direct evaluation of U^μ, and $-((z-a)/z)^{1/2}H(z)$ was compared with a difference quotient of U^μ; the root agrees with $4c$ to 10^{-12} in the lattice cases, and the two forms of ν to 10^{-21}, which tests the root and the quadrature rather than (ii). The first-order pole term of (3.6) was fitted from the root over $n = 100$–6400 for $\lambda_k = (k+\gamma k^{-m})^p$ at $(p,m) = (1.5,\frac12), (0.75,\frac14), (1.25,\frac12), (1.8,\frac34)$; it agrees with $pR_qW(q)\pi\cot(\pi q)$ to between 10^{-8} (the printed digits) and 10^{-6} (scripts `m1_theoremQ`, `m1_edge`).

4. The kernel

By Lemma 2.4 and Lemma 2.2, the zeta terms of Theorem 2.5 are the values at $\mu = \mu_n$ of the linear functional $cpM_\mu(4c)$, truncated to K terms. This section evaluates the same functional on the band factor itself, without truncation, and writes it as one Mellin–Barnes integral (Theorem 4.2). Its kernel contains $-\pi\cot(\pi w)$, which is also the Mellin transform of the truncated per-node law of Section 3 (Lemma 4.1). The proofs are for the lattice; Section 9 discusses other node sets.

4.1. The truncated per-node law.

Lemma 4.1. *Let* $L(u) := (1-u)^{-1/2}\mathbf{1}_{u<1}$. *For* $\operatorname{Re} w < 0$,

$$\widetilde{L}(w) := \int_0^\infty u^{-w-1}L(u)\,\mathrm{d}u = B\big(-w,\tfrac12\big) = -\pi\cot(\pi w)\,\frac{\Gamma(w+\frac12)}{\sqrt{\pi}\,\Gamma(w+1)}.$$

$\widetilde{L}$ *continues meromorphically to* $\mathbb{C}$; *its poles in* $\operatorname{Re} w \ge 0$ *are simple, at* $w = j \in \mathbb{N}_0$, *with* $-\operatorname{Res}_{w=j}\widetilde{L} = \binom{2j}{j}4^{-j}$, *the Taylor coefficients of* L *at* 0. *For* $a = 4c$, $(a/n^p)^w\Gamma(w+\frac12)/(\sqrt{\pi}\Gamma(w+1)) = W(w)n^{-pw}$.

Proof. The integral is $B(-w,\frac12) = \Gamma(-w)\sqrt{\pi}/\Gamma(\frac12 - w)$ by [5, (5.12.1)]. By the reflection formula [5, (5.5.3)], $\Gamma(\frac12 - w)\Gamma(\frac12 + w) = \pi/\cos(\pi w)$ and $\Gamma(-w)\Gamma(1+w) = -\pi/\sin(\pi w)$, so $\sqrt{\pi}\,\Gamma(-w)/\Gamma(\frac12 - w) = \Gamma(-w)\Gamma(w+\frac12)\cos(\pi w)/\sqrt{\pi} = -\pi\cot(\pi w)\Gamma(w+\frac12)/(\sqrt{\pi}\,\Gamma(w+1))$. The residue of $-\pi\cot(\pi w)$ at j is -1, and $\Gamma(j+\frac12)/(\sqrt{\pi}\,j!) = \binom{2j}{j}4^{-j}$. □

The two analytic continuations of $(1-u)^{-1/2}$ to $u > 1$ are $\mp i|1-u|^{-1/2}$, and their Mellin transforms carry $\Gamma(-w)\Gamma(1+w)e^{\pm i\pi w} = -\pi e^{\pm i\pi w}/\sin(\pi w)$. Their average is L, and the average of the two transforms is $-\pi\cot(\pi w)$: the cotangent is the signature of truncating the per-node law at the edge $u = 1$.

4.2. The band-factor functional.

For a function f analytic near the jump contour of G put, with the notation of (2.2)–(2.3),

$$\mathcal{M}_n[f] := \frac{1}{2\pi}\int_{-\infty}^{0} \mathfrak{e}(y)f(y)\,\mathrm{d}y \qquad (p>1),$$

$$\mathcal{M}_n[f] := \frac{1}{\pi}\int_0^\infty \mathfrak{j}(r)\,\operatorname{Im}\big[e^{i\vartheta}f(re^{i\vartheta})\big]\,\mathrm{d}r \qquad (0<p<1),$$

so that $\mathcal{M}_n[y^{-k-1}] = m_k = \zeta(-kp)/(kn^{kp})$ by Lemma 2.2 (for $0<p<1$ the integrand is real and $\mathcal{M}_n$ is the seam integral of [11, Lemma 10.1(iii)]). Since $\sum_{j\ge0}\binom{2j}{j}c^jt^j = (1-4ct)^{-1/2}$ and $\sum_l (4c)^l t^l = (1-4ct)^{-1}$,

$$\sum_{k\ge1} Q_k(4c)\,y^{-k-1} = k_c(y) := y^{-2}\Big(1-\frac{4c}{y}\Big)^{-3/2} \qquad (|y|>4c). \tag{4.1}$$

The *band-factor functional* is

$$e_n := 2cp\,\mathcal{M}_n[k_c], \tag{4.2}$$

that is, $e_n = \frac{cp}{\pi}\int_0^\infty \mathfrak{e}(-r)\,r^{-1/2}(r+4c)^{-3/2}\,\mathrm{d}r$ for $p>1$ and $e_n = \frac{2cp}{\pi}\int_0^\infty \mathfrak{j}(r)\operatorname{Im}[e^{i\vartheta}k_c(re^{i\vartheta})]\,\mathrm{d}r$ for $0<p<1$. On the seams $\operatorname{Im}(1-4c/\zeta) = (4c/r)\sin\vartheta > 0$, so the principal branch in k_c is never crossed. By Lemma 2.4 and Lemma 2.2, $e(\mu_n) = cp\sum_{k\le K}\mu_{n,k}Q_k(4c) = 2cp\sum_{k\le K}m_kQ_k(4c)$ is the truncation of e_n obtained by expanding k_c in (4.1).

Theorem 4.2. *Let $p > 0$, $p \neq 1$, $s > 1$, and $0 < c_0 < 1$.*

(i) $e_n = 2cp \sum_{k \le K} m_k Q_k(4c) + O(n^{-(K+1)p})$ *for every* K.

(ii) $e_n = \dfrac{1}{2\pi i} \displaystyle\int_{c_0 - i\infty}^{c_0 + i\infty} p\, W(w)\, \zeta(-wp)\, n^{-wp}\, \pi \Big[\cot \frac{\pi w p}{2} - \cot(\pi w) \Big]\, \mathrm{d}w.$

(iii) *For every* $J \ge 1$, $e_n = \sum_{j \le J} \varepsilon_j n^{-jp} + O(n^{-(J+1)p})$.

Part (iii) follows from (i) and $2cpm_jQ_j(4c) = \varepsilon_j n^{-jp}$; the content of the theorem is the exact representation (ii) and its kernel.

Proof. (i) Split the integral at $|y| = 2(4c+2)$. On the far part, expand k_c by (4.1) to K terms with remainder $O(|y|^{-K-2})$; the near part, and the near parts of the moments, are $O(e^{-n/C})$ because the jump densities are [11, Lemmas 8.1(i) and 10.1]. The remainder is bounded as in Lemma 2.2.

(ii), $p > 1$. For complex w with $\operatorname{Re} w > 0$, the computation of Lemma 2.2 with k replaced by w (the Fubini bound depends only on $\operatorname{Re} w$) gives

$$I(w) := \int_0^\infty \mathfrak{e}(-r) r^{-w-1}\, \mathrm{d}r = -2p\, \Gamma(wp) \zeta(1+wp) (2\pi n)^{-wp} \sin\Big(\pi w - \frac{\pi w p}{2} \Big).$$

The strip is set by $\mathfrak{e}(-r) \to \pi(1 - 2/p)$ as $r \to \infty$ and by the exponential decay of $\mathfrak{e}(-r)$ as $r \to 0$. By [5, (25.4.1)] at $1 + wp$, $p\Gamma(wp)\zeta(1+wp)(2\pi)^{-wp} = -\pi\zeta(-wp)/(w \sin(\pi w p/2))$, so

$$I(w) = \frac{2\pi}{w}\, \zeta(-wp)\, n^{-wp} \Big[\sin(\pi w) \cot \frac{\pi w p}{2} - \cos(\pi w) \Big].$$

Next, $r^{-1/2}(r + 4c)^{-3/2} = r^{-2}(1 + 4c/r)^{-3/2}$, and the Mellin–Barnes representation

$$(1+x)^{-3/2} = \frac{1}{2\pi i} \int_{(c_1)} \frac{\Gamma(-s)\Gamma(\frac{3}{2} + s)}{\Gamma(\frac{3}{2})}\, x^s\, \mathrm{d}s \qquad (x > 0,\ -\tfrac{3}{2} < c_1 < 0) \tag{4.3}$$

is the Mellin inversion of the Beta integral [5, (5.12.3)], whose transform is absolutely integrable on the line. Put $w = s + 1$, so that $\operatorname{Re} w = c_0 \in (0,1)$, and exchange the integrals (absolute convergence). Using $\Gamma(1-w)/w = \pi/(\sin(\pi w)\Gamma(1+w))$ and $\Gamma(\frac{3}{2}) = \sqrt{\pi}/2$, the factors collect into $pW(w)\zeta(-wp)n^{-wp}\pi[\sin(\pi w)\cot(\pi wp/2) - \cos(\pi w)]/\sin(\pi w)$, which is the stated integrand.

(ii), $0 < p < 1$. The Mellin transform of the seam density is $\int_0^\infty \mathfrak{j}(r) r^{-w-1}\, \mathrm{d}r = p(2\pi n)^{-wp}\Gamma(wp)\zeta(1+wp)$ for $\operatorname{Re} w > 0$, by the computation of [11, Lemma 10.1(iii)] with complex w; the strip comes from $\mathfrak{j}(r) \sim \frac{1}{p} \log r$ at infinity and the super-exponential decay at 0. Expand $(1+v)^{-3/2}$ with $v = -4ce^{-i\vartheta}/r$ by (4.3) (now $\arg v = \pi - \vartheta \in (0,\pi)$, where the representation continues). With $w = s + 1$ the angular factor is $\operatorname{Im}[e^{i\vartheta} e^{-2i\vartheta} e^{-i\vartheta s} e^{i\pi s}] = -\sin((\pi - \vartheta)w)$, and with $\vartheta = \pi p/2$ and the same two identities the integrand becomes $pW(w)\zeta(-wp)n^{-wp}\pi \sin(\pi w - \frac{\pi w p}{2})/(\sin(\pi w)\sin\frac{\pi w p}{2})$. Finally $\sin(\pi w - \frac{\pi w p}{2})/(\sin(\pi w)\sin\frac{\pi w p}{2}) = \cot\frac{\pi w p}{2} - \cot(\pi w)$.

(iii) On vertical lines $|\cot(\pi wp/2) - \cot(\pi w)| \le Ce^{-\min(p,2)\pi|\operatorname{Im} w|}$, while $\zeta(-wp)$ and $W(w)$ grow at most polynomially, so the contour can be moved to the right. The poles of $-\pi\cot(\pi w)$ at $w = j \ge 1$ give $-\operatorname{Res} = pW(j)\zeta(-jp)n^{-jp} = \varepsilon_j n^{-jp}$. The poles of $\cot(\pi wp/2)$ at $w = 2l/p$ lie on the zeros $\zeta(-2l) = 0$ and are removable; where $2l/p = j$ is an integer the product is regular and $\varepsilon_j = 0$. □

Remark 4.3 (reading of Theorem 4.2). (a) The second kernel $\pi\cot(\pi wp/2)$ comes only from writing $I(w)$ with $\zeta(-wp)$: in the form $-2p\Gamma(wp)\zeta(1+wp)(2\pi n)^{-wp}\sin(\pi w - \pi wp/2)$ there are no poles at $w = 2l/p$. So the band factor contributes exactly the integer residues of

$-\pi\cot(\pi w)$. (b) The pole of $\zeta(-wp)$ at $w=-1/p$ (the Weyl pole) and the point $w=0$ lie to the left of the strip. The $j=0$ term $pW(0)Z(0)=p\zeta(0)$, which is part of d_1, does not come from e_n but from the index normalisation of the frame. (c) e_n is a smooth function of n with an expansion in powers of n^{-p}. The literal sum $\frac{p}{n}\sum_{u_m<1}(1-u_m)^{-1/2}$, $u_m=4cm^p/n^p$, differs from its Mellin pole part by an oscillation of order $n^{-1/2}$ set by the fractional part of $(n^p/4c)^{1/p}$, which is not present in b_n. Theorem 4.2 says that the band factor produces the pole part only. (d) The negative-axis geometry ($p>1$) and the seam geometry ($p<1$) give the same kernel. At $p=1$ the two cotangents coincide and their poles merge, which is where the expansions of [11] change form. (e) The residue sum in (iii) is asymptotic only: $\zeta(-jp)$ grows factorially in j.

Numerically, the quadrature of e_n and the Mellin–Barnes integral agree to all 16 printed digits at $(p,n)=(1.5,10),(1.25,20),(3,10),(0.6,10),(0.3,8),(0.75,20)$, and the partial residue sums approach them (scripts `spec_kernel`, `m2_thmK_plt1`).

4.3. **Perturbed index functions.** For a node set other than the lattice, the analysis of [11] has not been carried out. The following computation shows what the band factor would contribute if the construction carried over with the lattice's index function $t(z)=nz^{-1/p}$ replaced by $t(z)=\kappa(n^p/z)$, κ the inverse of $k\mapsto\lambda_k$.

Proposition 4.4. *Let $p>1$, $m>0$, and for $\lambda_k=(k+\gamma k^{-m})^p$ let $\delta t_+(r):=-\gamma n^{-m}(re^{i\pi})^{m/p}$ be the first-order change of $t_+=nr^{-1/p}e^{-i\pi/p}$ on the negative axis, and $\delta\mathfrak{e}:=2\,\mathrm{Im}[\mathcal{F}'(t_+)\delta t_+]$ with $\mathcal{F}'(t)=2\pi i/(e^{2\pi it}-1)$. Then, for $\mathrm{Re}\,w>(m+1)/p$,*

$$\int_0^\infty \delta\mathfrak{e}(-r)r^{-w-1}\,\mathrm{d}r=-2\pi\gamma p\,n^{-pw}\zeta(1+m-pw)\Big[\cos(\pi w)+\sin(\pi w)\tan\frac{\pi(pw-m)}{2}\Big].$$

The proof is the computation of Lemma 2.2: expand $\mathcal{F}'$ in powers of $e^{-2\pi it_+}$, integrate termwise by [5, (5.9.1)], sum $\sum_{m'}m'^{m-pw}=\zeta(pw-m)$ by [5, (25.2.1)] and apply [5, (25.4.1)]; a quadrature check agrees to 10^{-13} at $w=2.3$ and 3.1 ($p=1.5$, $m=1$). With $\delta Z(-w)=pw\gamma\zeta(1+m-pw)$, the first-order change of $Z_\Lambda(-w)$, the change of e_n is $\frac{1}{2\pi i}\int pW(w)\delta Z(-w)n^{-pw}\pi[-\cot(\pi w)-\tan(\pi(pw-m)/2)]\,\mathrm{d}w$ on a line $\mathrm{Re}\,w>(m+1)/p$.

Remark 4.5. Under that hypothetical construction, the integer residues give $pW(j)\delta Z(-j)n^{-jp}$, the zeta terms of the perturbed node set, exactly as for the lattice. The pole of $\delta Z(-w)$ at $q=m/p$ lies to the left of the line, so the band factor does not carry the pole term $pR_qW(q)\pi\cot(\pi q)n^{-1-pq}$ of Section 9; in the dictionary that term comes from the equilibrium, Proposition 3.4. The tan-pole at $w=(m+1)/p$ and the change $\log|t'|$ of the node density sit at order n^{-2-m}, the order of the subleading pole terms of Remark 9.2. None of this is a statement about b_n.

5. The natural index and the expansion of the lattice

This section proves Theorem A. The $1/n$ term and the $(s-1)^2$ part of the n^{-2} term of the lattice are one index shift (§5.1); for Bessel zeros the shift is exact to all orders (§5.2). We then extend the analysis of [11] and Theorem 2.5 to every order K in $1/n$ and every order B in the band-factor parameters (§§5.3–5.4). The analytic part is the order-K version of estimates already proved at order 2; the algebraic part is a finite residue recursion, which we evaluate in exact arithmetic. Strategy: the band factor is a tilt (Lemma 2.2), tilts act on the jump data by an entire gauge (Lemma 2.3), so every coefficient of the expansion is an entire function of the tilt parameters μ with n-independent Taylor coefficients, and the Taylor coefficients are computed by the recursion.

5.1. **The index shift.** Put $\delta_N := \frac{\alpha+1}{2}$, so that $N = n + \delta_N$, and

$$f_2 = f_2(p,\alpha) := \frac{p+1}{12} - \frac{(p+2)\alpha^2}{8}. \tag{5.1}$$

Proposition 5.1. *For all p, s, $d_1 = -p\delta_N$ and $d_2(p,s) = \frac{p(p+1)}{2}\delta_N^2 + f_2$. Equivalently, $1 + d_1/n + d_2/n^2 = (n/N)^p(1 + f_2N^{-2}) + O(n^{-3})$.*

Proof. With $s - 1 = p(\alpha + 1)$ and $s - 1 - p = p\alpha$, the three terms of d_2 in Theorem 2.5 are $p(p+1)(\alpha+1)^2/8 = \frac{p(p+1)}{2}\delta_N^2$, $-(p+2)\alpha^2/8$ and $(p+1)/12$. Since $(1+\delta/n)^{-p} = 1 - p\delta/n + \frac{p(p+1)}{2}\delta^2 n^{-2} + O(n^{-3})$, the claim follows. □

So the lattice expansion to order n^{-2} is an index shift by δ_N followed by a bare function with no $1/N$ term. The shift has two parts: $\frac{\alpha}{2}$, attached to the hard edge of Bessel order α, and $\frac{1}{2} = -\zeta(0)$, the regularised count of the nodes (Section 9).

5.2. **Bessel zeros.** For $\mu > -1$ let $j_{\mu,1} < j_{\mu,2} < \cdots$ be the positive zeros of the Bessel function J_μ and $\lambda_k := (j_{\mu,k}/\pi)^2$. For $p = s = 2$ the measure (1.1) is $\sigma_\Lambda = \pi^2 \sum_k j_{\mu,k}^{-2} \delta_{\pi^2/j_{\mu,k}^2}$, and $\mu = \frac{1}{2}$ gives the lattice $\lambda_k = k^2$.

Proposition 5.2. *The recurrence coefficients of σ_Λ satisfy, for $n \geq 1$, with $N := n + \mu/2$,*

$$\frac{n^2 b_n}{c(2)} = \frac{4n^2}{(2n+\mu)\sqrt{(2n+\mu)^2 - 1}} = \Big(\frac{n}{N}\Big)^2 \Big(1 - \frac{1}{4N^2}\Big)^{-1/2}.$$

The bare function $(1 - \frac{1}{4N^2})^{-1/2} = 1 + \frac{1}{8N^2} + \frac{3}{128N^4} + \frac{5}{1024N^6} + \cdots$ is even in $1/N$, and its coefficients are $f_2(2, -\frac{1}{2})$, $f_4(2, -\frac{1}{2})$, $f_6(2, -\frac{1}{2})$ of (5.1), (5.3), (5.4).

Proof. The Lommel polynomials of order $\nu = \mu + 1$ satisfy the monic recurrence $xh_n = h_{n+1} + \beta_n^2 h_{n-1}$ with $\beta_n^2 = 1/(4(n+\nu-1)(n+\nu))$, and their Jacobi operator has spectrum the closure of $\{\pm 1/j_{\nu-1,k}\}$ [14, §3]; the orthogonality measure is symmetric with masses proportional to $j_{\nu-1,k}^{-2}$. By the quadratic transformation of symmetric measures (as in [11, Remark 9.6]), the image measure in $y = x^2$ has $b_n^{(y)} = \beta_{2n-1}\beta_{2n}$, and σ_Λ is its image under $y \mapsto \pi^2 y$. Hence $b_n = \pi^2/(4(2n+\mu)\sqrt{(2n+\mu)^2-1})$, and $c(2) = \pi^2/16$ gives the first identity. The second is $(2N)\sqrt{4N^2-1} = 4N^2(1 - \frac{1}{4N^2})^{1/2}$. □

The first identity agrees with the Jacobi coefficients computed from the zeros (mpmath zeros for $k \leq 120$, McMahon's expansion beyond, a continuous tail) to $1.6 \cdot 10^{-15}$ at $\mu = 0$ and $\mu = 1.5$, $n \leq 40$ (script `spec_lommel`). Here $\alpha = (s - 1 - p)/p = -\frac{1}{2}$, and the large-$k$ form $j_{\mu,k} = \pi(k + \frac{\mu}{2} - \frac{1}{4}) + O(1/k)$ gives $Z_\Lambda(0) = \zeta(0, \frac{\mu}{2} + \frac{3}{4}) = -\frac{\mu}{2} - \frac{1}{4}$, so $N = n + \frac{\alpha}{2} - Z_\Lambda(0)$: Proposition 5.2 is the natural-index law of Section 9, exactly.

5.3. **The residue recursion.** We use the frame of §2.2 for $p \neq 1$: the clockwise circles ∂U_0, ∂U_r around 0 and $4c$, the local maps f (hard edge) and $\tilde{\varphi}$ (soft edge), and the Szegő quotients ρ, ω of [11, Sections 8–10]. Let $M := \frac{1}{\sqrt{2}}\left(\begin{smallmatrix} 1 & i \\ i & 1 \end{smallmatrix}\right)$, let $\circ$ be the entrywise product, let $a_k = a_k(\alpha)$, $b_k = b_k(\alpha)$ be the coefficients of the Bessel expansions [5, (10.17.1), (10.17.8)] and u_k, v_k those of the Airy expansions [5, (9.7.2)], and put

$$\Xi_k := M^{-1}(M \circ C_k),\ C_k := \begin{pmatrix} (-1)^k a_k & a_k \\ (-1)^k b_k & b_k \end{pmatrix}, \qquad X_k := M^{-1}(M \circ C_k'),\ C_k' := \begin{pmatrix} u_k & (-1)^k u_k \\ v_k & (-1)^k v_k \end{pmatrix}.$$

Lemma 5.3 (the jumps to every order)**.** *Let $p \neq 1$ and $K, J \geq 1$. Uniformly on $\partial U_0 \cup \partial U_r$, boundary values included, the jump of the error matrix on the circles is $I + \sum_{k \leq K} \Delta_k n^{-k} + O(n^{-K-1})$, where Δ_k equals*

$$\Delta_k^0 = (2f^{1/2})^{-k} P_0^\infty \rho^{\sigma_3} (\sigma_3 \Xi_k \sigma_3) \rho^{-\sigma_3} (P_0^\infty)^{-1} \qquad \text{on } \partial U_0,$$
$$\Delta_k^r = (-1/\tilde{\varphi})^k P_0^\infty (-\omega)^{-\sigma_3/2} X_k (-\omega)^{\sigma_3/2} (P_0^\infty)^{-1} \qquad \text{on } \partial U_r,$$

and $\Delta_k = G_{\mu_n} \Delta_k^{\mathrm{mod}} G_{\mu_n}^{-1} + O(n^{-(J+1)p})$.

Proof sketch. For $k = 1$ this is [11, Lemmas 9.1, 9.2 and 10.6], and for $k = 2$ the third-order version cited in §2.4. The proof is the same with $K + 1$ terms of the expansions [5, (10.40.1)–(10.40.4), (10.17.5), (10.17.6), (10.17.11), (10.17.12)] of the Bessel model [8, §6] and [5, (9.7.5)–(9.7.6)] of the Airy model, whose Poincaré remainders are $O(n^{-K-1})$ uniformly because the arguments are of size n on the circles. The sign pattern $(-1)^k$ is the same in every sector, for every k: each column of the Airy model is a constant times $\mathrm{Ai}(Z)$ or $\mathrm{Ai}'(Z)$ with $Z = e^{2\pi i m/3}\zeta$, $|\mathrm{ph}\, Z| \leq \pi - \delta$, and $Z^{3/2} = \pm\zeta^{3/2}$, so $(-1)^k$ sits on the column that carries $e^{-\xi}$ whichever branch occurs; the column carrying $e^{\pm\xi}$ is fixed at $k = 0$. The Bessel model is the same with $Y = \pm w$. The n-dependence of the weight is Lemma 2.2 at order J, and the conjugation by G_{μ_n} is Lemma 2.3, whose proof commutes diagonal factors and does not depend on k. □

Lemma 5.4 (meromorphy)**.** *For every k, Δ_k^{mod} is meromorphic in U_0 and in U_r, and $\Delta_k^\mu := G_\mu \Delta_k^{\mathrm{mod}} G_\mu^{-1}$ is meromorphic with Laurent coefficients that are entire functions of μ.*

Proof sketch. For $p > 1$: $\hat{P}^{\mathrm{mod}} (P_0^\infty)^{-1}$ is analytic in $U_e \setminus \{e\}$ and has the expansion of Lemma 5.3 uniformly on compact subsets, so $\Delta_k = \lim n^k(\cdots)$ is analytic and single-valued there; its explicit formula grows at most polynomially at e, so e is a pole. For $0 < p < 1$, where the glued hard-edge parametrix carries seam jumps inside U_0, one checks directly that Δ_k has no jump: from the definitions, $X_k = \frac{1}{2}(u_k + v_k)I - \frac{1}{2}(u_k - v_k)\sigma_2$ for even k and $X_k = \frac{1}{2}(u_k + v_k)\sigma_3 + \frac{1}{2}(v_k - u_k)i\sigma_1$ for odd k, so $JX_kJ^{-1} = (-1)^k X_k$ for $J = i\sigma_2$; on $(4c - \epsilon, 4c)$, $\tilde{\varphi}_+ = -\tilde{\varphi}_-$ contributes the same $(-1)^k$, and $P_{0,+}^\infty = P_{0,-}^\infty J$, $(-\omega)_+^{1/2}(-\omega)_-^{1/2} = 1$. The hard edge is the same with $\sigma_3 \Xi_k \sigma_3$, $f_+^{1/2} = -f_-^{1/2}$ and the relations for ρ of [11, Section 10]. The gauge G_μ is entire in (z, μ) and satisfies the same relations. □

For $\mu \in \mathbb{C}^J$ define, near each $e \in \{0, 4c\}$, $\mathsf{R}_{\mathrm{in}}^{(0)} := I$ and for $m \geq 1$

$$\mathsf{F}_m(\mu) := \sum_{j=1}^{m} \mathsf{R}_{\mathrm{in}}^{(m-j)}(\mu) \Delta_j^\mu, \qquad \mathsf{R}_{\mathrm{out}}^{(m)}(\mu) := \sum_{e \in \{0,4c\}} \big(\text{principal part of } \mathsf{F}_m(\mu) \text{ at } e\big), \tag{5.2}$$
$$\mathsf{R}_{\mathrm{in}}^{(m)}(\mu) := \mathsf{R}_{\mathrm{out}}^{(m)}(\mu) - \mathsf{F}_m(\mu), \qquad \mathsf{R}_1^{(m)}(\mu) := \mathrm{Res}_0\, \mathsf{F}_m(\mu) + \mathrm{Res}_{4c}\, \mathsf{F}_m(\mu).$$

By Lemma 5.4 these are well defined, $\mathsf{R}_{\mathrm{in}}^{(m)}$ is analytic in each disc, and all entries are entire in μ. For $m = 1, 2$ and $\mu = 0$ they are the objects of [11, Proposition 9.5] and its third-order version.

Proposition 5.5 (small norm to order K)**.** *Let $p \neq 1$ and $K, J \geq 1$. Then $\mathsf{R}_1 = \sum_{m \leq K} \mathsf{R}_1^{(m)}(\mu_n) n^{-m} + O(n^{-K-1} + n^{-1-(J+1)p})$, and the same expansion holds for R uniformly at positive distance from its jump contour.*

Proof sketch. Write $\eta_n := n^{-K-1} + n^{-1-(J+1)p}$. By Lemma 5.3 the jump is $I + \sum_{k \leq K} \Delta_k^{\mu_n} n^{-k}$ on the circles up to $O(\eta_n)$, and off the circles it is I up to errors that are $O(\eta_n)$ in $L^1 \cap L^2 \cap L^\infty$,

as in [11, Propositions 9.5 and 10.8]; the Cauchy operator is bounded uniformly in n. One proves by induction on $m = 1, \dots, K+1$ that the boundary value R_- equals $\sum_{k<m} \mathsf{R}_-^{(k)} n^{-k} + O_{L^2}(n^{-m} + n^{-1-(J+1)p})$, where $\mathsf{R}^{(k)}$ is $\mathsf{R}_{\mathrm{in}}^{(k)}(\mu_n)$ inside the discs and $\mathsf{R}_{\mathrm{out}}^{(k)}(\mu_n)$ outside. In the step, insert the claim into $\mathsf{R} = I + \mathcal{C}(\mathsf{R}_-(J-I))$: the new error is $O(n^{-m-1} + n^{-1-(J+1)p})$, and the term of order n^{-m} is the Cauchy integral over the clockwise circles of $\mathsf{F}_m(\mu_n)$, which by the residue theorem is $\mathsf{R}_{\mathrm{out}}^{(m)}$ outside the discs and $\mathsf{R}_{\mathrm{out}}^{(m)} - \mathsf{F}_m = \mathsf{R}_{\mathrm{in}}^{(m)}$ inside. The coefficient R_1 is $-\frac{1}{2\pi i}\int \mathsf{R}_-(J-I)$, and $-\frac{1}{2\pi i}\oint_{\mathrm{cw}} \mathsf{F}_m = \operatorname{Res}_0 \mathsf{F}_m + \operatorname{Res}_{4c} \mathsf{F}_m$. □

Define $\mathfrak{D}_K(n;\mu) := ((Q_1)_{12}(Q_1)_{21})^{1/2}/c$, expanded to order n^{-K}, where $Q_1 = P_1 + \sum_{m\le K} \mathsf{R}_1^{(m)}(\mu) n^{-m}$, and write $\mathfrak{D}_K(n;\mu) = \sum_{\mathbf{r}} \mu^{\mathbf{r}} \mathfrak{D}_{\mathbf{r},K}(n)$ for its Taylor series in μ. Each $\mathfrak{D}_{\mathbf{r},K}$ is a polynomial in $1/n$ of degree at most K. A polynomial in $1/n$ is *written in natural form* when it is re-expanded in $1/N$ by $1/n = N^{-1}/(1-\delta_N N^{-1})$; a layer $\mu^{\mathbf{r}}$ carries $n^{-|\mathbf{r}|_{wp}} = N^{-|\mathbf{r}|_{wp}}(N/n)^{|\mathbf{r}|_{wp}}$, with $|\mathbf{r}|_w := \sum_j j r_j$.

Lemma 5.6 (exact algebra)**.** *The recursion* (5.2) *and the extraction* $\mathfrak{D}_K$ *give, identically in the symbols:*

(a) $\mu = 0$, $K = 6$: $\mathfrak{D}_{0,6}(n) = (n/N)^p[1 + f_2 N^{-2} + f_4 N^{-4} + f_6 N^{-6} + O(N^{-7})]$, *with no* N^{-1}, N^{-3}, N^{-5} *terms,* f_2 *as in* (5.1) *and* f_4, f_6 *as in* (5.3), (5.4)*; in particular* $d_1 = -p(\alpha+1)/2$ *and* d_2 *is that of Theorem 2.5;*

(b) *first order in* μ_j*, for every integer* $j \ge 1$ *(*j *a symbol),* $K = 4$*: the coefficient of* n^{-1} *per unit* $\binom{2j}{j}\mu_j$ *is* $pj/2$ *(with* $c = 1$*), and in natural form the layer is a multiple of* $N^{-1-jp}[1 + g_j N^{-2} + O(N^{-4})]$*, with no* N^{-1}*,* N^{-3} *term in the bracket and* $g_j = A_j + B_j\alpha^2$ *as in Theorem A; the same holds at* $K = 5$ *for the integers* $j = 1, \dots, 5$*, and for* $j = 1$ *the* N^{-4} *coefficient of the bracket is* $g_{1,4}$ *of* (5.5)*;*

(c) *second order in* μ_1*,* $K = 5$*: per unit* μ_1^2 *the natural form has no* N^{-1-2p} *term, the coefficient* $p(3p+1)/2$ *at* N^{-2-2p}*, no* N^{-3-2p} *and no* N^{-5-2p} *term, and* $p^2 h_2$ *at* N^{-4-2p}*, with* h_2 *as in* (5.6)*.*

Proof. With $c = 1$ every input is a rational function of (p, α): the local maps have convergent series with rational coefficients, π cancels in f, the square roots are of series whose leading coefficient is 1 or a perfect square, and the factors $2^{-1/2}$ in T and M occur in conjugate pairs. The recursion is carried out on truncated Laurent series over $\mathbb{Q}(p,\alpha)[i]$; every series carries the index of its highest exactly known coefficient, and every principal part checks that its coefficients down to the residue are known, otherwise the run stops. For (b), μ_j is normalised by $\binom{2j}{j}$; the gauge to first order is $\partial_{\mu_j}\Delta_k^{\mu}|_{\mu=0} = [\mathcal{G}_j, \Delta_k^{\mathrm{mod}}]$ with $\mathcal{G}_j = \mathsf{T}\left(\begin{smallmatrix} 0 & -(z-4c)Q_j/2 \\ -zQ_j/2 & 0 \end{smallmatrix}\right)\mathsf{T}^{-1}$, and it needs only the Taylor data of $Q_j/\binom{2j}{j}$ at 0 and at $4c = 4$, which are rational in j: $\prod_{i\le m}(j-i)/(2(2j-2i-1))$ at 0 and $j(j-1)\cdots(j-m)/(4^{m+1}(\frac{1}{2})_{m+1})$ at 4, from $\sum_j Q_j(z)t^j = t/((1-zt)\sqrt{1-4t})$. So (b) is a computation over $\mathbb{Q}(p,\alpha,j)[i]$. For (c) the gauge is taken to second order, $\cosh h = 1 + z(z-4)\mu^2 Q_1^2/8 + O(\mu^4)$, $S = \mu Q_1/2 + O(\mu^3)$. The scripts and their outputs are listed in Appendix A; each output begins with the SHA-256 hashes of the scripts that produced it. Since $(z-4c)Q_j$ and zQ_j scale as c^j under $z = c\zeta$, μ_j enters only as $c^j\mu_j$; the integer layer and the ratios g_j, h_2 are independent of c. □

The closed forms are

$$f_4 = \frac{(p+1)(6p^3+46p^2+55p+3)}{1440\,p(3p+2)} - \frac{5(p+1)(p+2)\alpha^2}{96(3p+2)} - \frac{p(p+2)^2\alpha^4}{128(3p+2)}, \tag{5.3}$$

$$f_6 = \frac{\phi_0 + \phi_2\alpha^2 + \phi_4\alpha^4 + \phi_6\alpha^6}{(3p+2)^2(5p+2)}, \tag{5.4}$$

$$\begin{aligned}
\phi_0 &= -\frac{p+1}{362880\,p^2}\big(450p^8 + 1590p^7 - 316p^6 - 23761p^5 - 66864p^4 - 65507p^3 \\
&\qquad - 18220p^2 - 1632p - 40\big), \\
\phi_2 &= \frac{(p+1)(p+2)(90p^6 + 270p^5 + 664p^4 - 165p^3 - 1649p^2 - 288p + 28)}{11520\,p}, \\
\phi_4 &= -\frac{(p+1)(p+2)^2(20p^4 + 45p^3 + 162p^2 + 72p + 16)}{1536}, \\
\phi_6 &= \frac{p(p+2)^3(16p^3 + 7p^2 - 4p - 4)}{3072},
\end{aligned}$$

$$\begin{aligned}
g_{1,4} = \frac{p+1}{5760\,p^2(3p+2)}\big(&9p^6 + 117p^5 + 161p^4 - 569p^3 - 1550p^2 - 976p + 48\big) \\
&+ \frac{(p+1)(p+2)(3p^4 + 33p^3 + 124p^2 + 182p + 108)\alpha^2}{192\,p(3p+2)} \\
&+ \frac{(p+2)^3(p^3 - 4p - 4)\alpha^4}{128\,p(3p+2)},
\end{aligned} \tag{5.5}$$

$$h_2 = \frac{(p+1)(p+2)(3p^2 + 2p + 1)}{8p^2} + \frac{(p+2)(2p^3 - p^2 - 7p - 6)\alpha^2}{16p^2}. \tag{5.6}$$

For $j = 1$, $g_1 = (p+1)(p^2+3p-2)/(24p) + (p+2)(p^2+2p+2)\alpha^2/(8p)$. At the Bessel point $(p, \alpha) = (2, -\frac{1}{2})$, $f_4 = \frac{3}{128}$ and $f_6 = \frac{5}{1024}$, in agreement with Proposition 5.2.

5.4. **The theorem.**

Theorem 5.7. *Let $p > 0$, $p \neq 1$, $s > 1$, and $K, J, B \geq 1$.*

(i) Double expansion. $\dfrac{n^p b_n}{c} = \displaystyle\sum_{|\mathbf{r}| \leq B} \mu_n^{\mathbf{r}} \mathfrak{D}_{\mathbf{r},K}(n) + O\big(n^{-K-1} + n^{-2-(B+1)p} + n^{-1-(J+1)p}\big)$.

(ii) Natural forms of the layers. $\mathfrak{D}_{0,6}$, $\mu_{n,j}\mathfrak{D}_{e_j,4}$ *and* $\mu_{n,1}^2\mathfrak{D}_{2e_1,5}$ *satisfy*

$$\begin{aligned}
\mathfrak{D}_{0,6}(n) &= \Big(\frac{n}{N}\Big)^p\Big[1 + \frac{f_2}{N^2} + \frac{f_4}{N^4} + \frac{f_6}{N^6} + O(N^{-7})\Big], \\
\mu_{n,j}\mathfrak{D}_{e_j,4}(n) &= \Big(\frac{n}{N}\Big)^p \frac{\varepsilon_j}{N^{1+jp}}\Big[1 + \frac{g_j}{N^2} + O(N^{-4})\Big] \qquad (j \geq 1), \\
\mu_{n,1}^2\mathfrak{D}_{2e_1,5}(n) &= \Big(\frac{n}{N}\Big)^p \frac{\varepsilon_1^2}{N^{2+2p}}\Big[\frac{3p+1}{2p} + \frac{h_2}{N^2} + O(N^{-4})\Big],
\end{aligned}$$

with the brackets free of odd powers of $1/N$ to the orders shown; for $j = 1$ the second bracket is $1 + g_1N^{-2} + g_{1,4}N^{-4} + O(N^{-5})$.

(iii) The coefficients of b_n. *With $K = 6$, $B = 2$ and $J \geq 6/p - 1$, expansion (1.9) holds.*

Proof. (i) By (2.1) and Proposition 5.5, $n^{2p}b_n^2 = (Q_1)_{12}(Q_1)_{21}$ with $Q_1 = P_1 + \sum_{m \leq K} \mathsf{R}_1^{(m)}(\mu_n)n^{-m} + O(n^{-K-1} + n^{-1-(J+1)p})$. The n^0 term of $(Q_1)_{12}(Q_1)_{21}$ is c^2, since P_1 does not depend on μ, so each coefficient of $\mathfrak{D}_K(n;\mu)$ is a polynomial in the entries of the $\mathsf{R}_1^{(i)}(\mu)$, $i \leq K$, which are entire in μ and independent of n. Taylor's theorem in μ on $|\mu| \leq 1$ applies coefficientwise. The n^0 coefficient is 1 for all μ, and the n^{-1} coefficient is

$d_1 + cpM_\mu(4c)$, exactly linear in μ by Lemma 2.4; hence $\mathfrak{D}_{\mathbf{r},K} = O(n^{-2})$ for $|\mathbf{r}| \geq 2$, and the Taylor remainder after order B is $O(|\mu|^{B+1}n^{-2})$. Since $|\mu_n| = O(n^{-p})$, this is $O(n^{-2-(B+1)p})$.

(ii) This is Lemma 5.6 in natural form, with the normalisations $\mu_{n,j}\binom{2j}{j}pc^j j/2 = \varepsilon_j n^{-jp}$ and $\mu_{n,1}^2 p(3p+1)/2 \cdot c^2 = \varepsilon_1^2 n^{-2p}(3p+1)/(2p)$.

(iii) In (i) keep $\mathbf{r} = 0$ at $K = 6$; $\mathbf{r} = e_j$ for every j with $1 + jp < \min(7, 2+3p)$, at $K = 5$ ($j \leq 5$) or $K = 4$; and $\mathbf{r} = 2e_1$ at $K = 5$. The other terms with $|\mathbf{r}| = 2$ ($\mu_1\mu_2, \mu_2^2, \dots$) are $O(n^{-2-3p})$ because $\mathfrak{D}_{\mathbf{r}} = O(n^{-2})$ and $|\mathbf{r}|_w \geq 3$, and those with $|\mathbf{r}| = 3$ are $O(n^{-2-3p})$. The truncation errors are bounded as follows. The $j = 1$ bracket is known to relative order N^{-5}, an error $O(n^{-6-p})$, which is $O(n^{-2-3p})$ for $p < 2$ and $O(n^{-8})$ for $p \geq 2$. For $j \geq 2$ the bracket error is $O(n^{-5-jp})$, and $5 + 2p > 2 + 3p$ for $p < 3$, $5 + 2p \geq 11$ for $p \geq 3$. The h_2 term of the ε_1^2 layer is $O(n^{-4-2p})$, which is $O(n^{-2-3p})$ for $p < 2$ and $O(n^{-7})$ for $p \geq \frac{3}{2}$. Finally $n^{-1-(J+1)p} = O(n^{-7})$. □

A term CN^{-e} of (1.9) is a coefficient of the expansion of $n^p b_n/c$ (in powers of N, or equivalently, after re-expansion, of n) exactly when $e < \min(7, 2+3p)$. Table 5.1 lists the resulting ranges. The terms outside their range are true identities of the layers in Theorem 5.7(ii) but are not determined as coefficients of b_n, because the uncomputed layers $\mu_1\mu_2$, μ_1^3 enter at n^{-2-3p}.

term	exponent e	a coefficient of $n^p b_n/c$ for
$f_2 N^{-2}$	2	all $p \neq 1$
$f_4 N^{-4}$	4	$p > \frac{2}{3}$
$f_6 N^{-6}$	6	$p > \frac{4}{3}$
$\varepsilon_j N^{-1-jp}$	$1 + jp$	$(j-3)p < 1$ and $1 + jp < 7$
$\varepsilon_1 g_1 N^{-3-p}$	$3 + p$	$\frac{1}{2} < p < 4$
$\varepsilon_2 g_2 N^{-3-2p}$	$3 + 2p$	$1 < p < 2$
$\varepsilon_1 g_{1,4} N^{-5-p}$	$5 + p$	$\frac{3}{2} < p < 2$
$\frac{3p+1}{2p}\varepsilon_1^2 N^{-2-2p}$	$2 + 2p$	$p < \frac{5}{2}$
$\varepsilon_j g_j N^{-3-jp}$ ($j \geq 3$), $\varepsilon_1^2 h_2 N^{-4-2p}$		never at $B = 2$

TABLE 5.1. Which terms of (1.9) are coefficients of b_n ($p \neq 1$). Coincident exponents merge: at $p = 1/j_0$, $\varepsilon_{j_0} N^{-2}$ merges with $f_2 N^{-2}$, and $f_2 + \varepsilon_{j_0} = f_2 - p/12$, as in Theorem 2.5.

Remark 5.8. (a) Theorem 5.7 does not prove evenness in $1/N$ to all orders, nor any closed form of the bare function; the computed layers are listed in (ii), and the layers $\mu_1\mu_2$, μ_2^2, μ_1^3 and higher are not computed. Evenness of the integer layer to all orders is Corollary 6.4 below. Nothing is claimed at $p = 1$, where the hard-edge and band-factor constructions of [11] change. (b) The same recursion evaluated numerically at 40 digits, with truncation depths 40–64, gives $f_1 = f_3 = f_5 = f_7 = 0$ to 10^{-35} at every sampled (p, α) with $p \in [\frac{1}{3}, 12]$, each f_{2k} even in α, and $f_8(3,0) = 1311857/632311515$, $f_8(2,0) = 3899/2^{21}$ after rational recognition. The denominators of f_4, f_6 are $3p+2$ and $(3p+2)^2(5p+2)$; they come from the factors $1/(a'+k)$, $a' = \frac{1}{p} + \frac{1}{2}$, of the hard-edge local map, $a' + k = ((2k+1)p + 2)/(2p)$, and $632311515 = 3^5 \cdot 5 \cdot 11^3 \cdot 17 \cdot 23$ is $3^5 \cdot 5\,(3p+2)^3(5p+2)(7p+2)$ at $p = 3$ (scripts `m4_rh4`, `m4_f8_denominator`). These statements are numerical, except $f_7 = 0$ and the evenness in α, which hold to all orders by Corollary 6.4. (c) Heuristically, evenness in α after the shift reflects the fact that the Bessel coefficients $a_k(\alpha)$, $b_k(\alpha)$ depend on α only through α^2, while elsewhere α enters only through the power of z at the hard edge, which the shift by $\frac{\alpha}{2}$ absorbs. Section 6 makes this precise: the two roles of α separate, the Szegő exponent is absorbed by the

natural index (Theorem 6.3), and the evenness in $1/N$ is a reflection symmetry (Theorem 6.1, Corollary 6.4). (d) The factor $1/(2j-1)$ in g_j, read as a function of a continuous Mellin variable w, is a simple pole at $w=\frac{1}{2}$, the Mellin signature of $(1-u)^{+1/2}$, against $(1-u)^{-1/2}$ for the leading kernel (Lemma 4.1). Partial fractions give $B_j=\frac{p+2}{8p}\big[\frac{p^2 j}{2}+\frac{p(p+2)}{4}+\frac{(p+2)(p+4)}{4(2j-1)}\big]$. This suggests a second-order edge kernel (Problem 9.9).

6. Reflection, shift and the Toda structure

Theorem 5.7 computes the layers of the lattice expansion to fixed orders and observes that they are even in $1/N$. This section explains the evenness and the natural index to all orders. The exponent α plays two roles in the residue recursion, and separating them gives a two-parameter family of formal problems (§6.1). On this family there are two exact symmetries: a reflection, which is algebra (Theorem 6.1), and a shift, which we prove by steepest descent of the same lattice problem in a second, non-integer frame (Theorem 6.3). Together they give the evenness of the integer layer to all orders (Corollary 6.4). The zeta layers are Toda flows of the integer layer (§6.4).

6.1. **The decoupled model.** The exponent α enters the jump data of Lemma 5.3 in two ways: as the Bessel order ν in the coefficients $a_k(\nu), b_k(\nu)$, which are polynomials in ν^2, and as the Szegő exponent β in $\rho^2=\phi^{2(\beta+1)}$ at the hard edge and in the factor $\phi^{-2(\beta+1)}$ at the soft edge, where $\phi(z)=(z^{1/2}+(z-4c)^{1/2})/(2c^{1/2})$ is the exterior conformal map of $[0,4c]$. For the lattice $\nu=\beta=\alpha$. Keeping ν and β independent, the residue recursion (5.2) and the extraction of §5.3, with the gauge of Lemma 2.3 for the band-factor parameters μ, define formal series

$$\frac{n^p\, b(n;\beta,\nu;\mu)}{c}=\sum_{\mathbf{r}}\mu^{\mathbf{r}}\,\mathfrak{D}_{\mathbf{r}}(n;\beta,\nu),\qquad B(n;\beta,\nu):=\mathfrak{D}_0(n;\beta,\nu)=\sum_{k\ge 0}d_k(p,\beta,\nu)\,n^{-k},\tag{6.1}$$

whose truncations at order n^{-K} are the polynomials $\mathfrak{D}_{\mathbf{r},K}$ of §5.3 when $\beta=\nu=\alpha$. We call this the *decoupled model* $M(\beta,\nu)$ and B its *integer layer*. Each d_k is a rational function of (p,β,ν), $d_0=1$: with $c=1$ the recursion and the extraction are field operations on Laurent data that are rational in p (Lemma 5.6), polynomial in ν through $a_k(\nu), b_k(\nu)$, and polynomial in β, since β enters only through binomial series of $(1+\text{series without constant term})^{2(\beta+1)}$.

Theorem 6.1 (reflection)**.** *As formal series, $b(n;\beta,\nu;0)=b(-n;-\beta-2,\nu;0)$; that is, $d_k(p,-\beta-2,\nu)=(-1)^k d_k(p,\beta,\nu)$ for all k.*

Proof. With $T=\frac{1}{\sqrt{2}}\left(\begin{smallmatrix}1 & 1\\ i & -i\end{smallmatrix}\right)$ we have $\sigma_2 T=T\sigma_3$, so $P_0^\infty=T\mathfrak{a}_\infty^{\sigma_3}T^{-1}$ and $P_1=-c\sigma_2$ commute with σ_2. The core matrices X_k and $\sigma_3\Xi_k\sigma_3$ lie in the span of I,σ_2 for even k and of $\sigma_3, i\sigma_1$ for odd k, so conjugation by σ_2 multiplies them by $(-1)^k$, while $\sigma_2 d^{\sigma_3}\sigma_2=d^{-\sigma_3}$ for scalars d. Hence $\sigma_2\Delta_k(\beta)n^{-k}\sigma_2=\Delta_k(-\beta-2)(-n)^{-k}$, the recursion is covariant under conjugation by a constant matrix, and $(Q_1)_{12}(Q_1)_{21}$ is invariant under conjugation by σ_2. □

Lemma 6.2 (self-similar Buyarov–Rakhmanov identity)**.** *Let Φ denote the phase of the lattice equilibrium in the scaled frame, with hard-edge branch $2f^{1/2}$ and soft-edge branch $\tilde{\varphi}$. Then $(1+pz\partial_z)\Phi=-2\log\phi$ up to the additive constants of the branches; in particular $(1+pz\partial_z)(2f^{1/2})=2\,\mathrm{arsinh}(v/2)$ at $z=-v^2$ ($c=1$).*

Proof. At the hard edge $2f^{1/2}=\pi\sqrt{-z}\,\mathfrak{a}(z)$, and the series of $\mathfrak{a}$ has the factors $1/(a'+k)$ with $a'=\frac{1}{p}+\frac{1}{2}$. Since $(1+pz\partial_z)[\sqrt{-z}\,z^k]=p(a'+k)\sqrt{-z}\,z^k$, these factors cancel term by term and leave the arcsine series. The soft-edge branch is the analytic continuation of the same phase. □

Under the self-similar scaling of the lattice problem, $1+pz\partial_z$ is the derivative in n of $n\Phi(xn^p)$; so Lemma 6.2 says that the n-flow of the phase is generated by $\log\phi$, which also generates the β-flow ($\partial_\beta\rho^2 = 2\log\phi\cdot\rho^2$). This suggested the following theorem; its proof does not use the lemma.

6.2. **The shift theorem.**

Theorem 6.3 (shift)**.** *Identically in (p,β,ν), as formal series:*

(i) *$B(n;\beta,\nu) = (n/N)^p F_p(N;\nu)$ with $N := n + \frac{\beta+1}{2}$, where $F_p(N;\nu)$ is a formal power series in $1/N$ that does not depend on β; equivalently, for every real γ (the* frame identity*)*

$$B(n+\gamma;\beta-2\gamma,\nu) = \Big(1+\frac{\gamma}{n}\Big)^p B(n;\beta,\nu), \tag{6.2}$$

the left side re-expanded in $1/n$;

(ii) *for $\mathbf{r}\in\{e_1,e_2,2e_1\}$ the same holds with the exponent p replaced by $(1+|\mathbf{r}|_w)p$: the series $(N/n)^{(1+|\mathbf{r}|_w)p}\mathfrak{D}_{\mathbf{r}}(n;\beta,\nu)$, re-expanded in $1/N$, does not depend on β.*

The infinitesimal form of (6.2) is the flow identity $\partial_n B - 2\partial_\beta B = (p/n)B$. The equivalence of (i) and (6.2) is algebra: if (i) holds, then $(n+\gamma)+\frac{\beta-2\gamma+1}{2} = N$ and $B(n+\gamma;\beta-2\gamma,\nu) = ((n+\gamma)/N)^p F_p(N) = (1+\gamma/n)^p (n/N)^p F_p(N)$. Conversely, given (6.2), fix β_0 and put $\gamma := (\beta_0-\beta)/2$; then $(N/n)^p B(n;\beta,\nu) = (N/(n-\gamma))^p B(n-\gamma;\beta_0,\nu)$, and $n-\gamma = N - \frac{\beta_0+1}{2}$ does not involve β. All re-expansions are between formal series in $1/n$, $1/(n-\gamma)$, $1/N$, which differ by constants.

Corollary 6.4 (evenness)**.** *$F_p(N;\nu) = F_p(-N;\nu)$, and F_p depends on ν only through ν^2. Hence $F_p(N;\nu) = 1+\sum_{k\ge1} f_{2k}N^{-2k}$ with each f_{2k} rational in p and polynomial in ν^2. For the lattice and every $p\neq1$, the integer layer of Theorem 5.7 is, for every K,*

$$\mathfrak{D}_{0,K}(n) = \Big(\frac{n}{N}\Big)^p\Big[1+\sum_{2k\le K}\frac{f_{2k}(p,\alpha^2)}{N^{2k}}\Big] + O(N^{-K-1}), \qquad N = n+\frac{\alpha}{2}+\frac{1}{2};$$

in particular all odd coefficients f_{2k+1} vanish, and the natural index of the lattice is the Szegő shift $\frac{\beta+1}{2}$ at $\beta=\alpha$.

Proof. By Theorem 6.1, $B(n;\beta,\nu) = B(-n;-\beta-2,\nu)$. The natural index of the right side is $-n+\frac{-\beta-1}{2} = -N$, so Theorem 6.3(i) gives $(n/N)^pF_p(N) = ((-n)/(-N))^pF_p(-N)$. The coefficients $a_k(\nu)$, $b_k(\nu)$ depend on ν only through ν^2, and β no longer occurs in F_p. Each d_k is polynomial in ν, hence so is each f_{2k}. □

Corollary 6.4 settles the integer-layer part of Remark 5.8(a)–(c): $f_7=0$ and the evenness in α hold to all orders, while the values $f_8, f_{10},\dots$ and the observed degree k of f_{2k} in α^2 remain numerical. The α-even part has the heuristic explanation of Remark 5.8(c): α enters the integer layer through ν^2 and through β, and the shift absorbs β.

6.3. **Proof of the shift theorem: a second frame.** The decoupled model is not related to itself by any transformation of the formal recursion that we know. In particular, the Christoffel shift $x\sigma_{p,s} = \sigma_{p,s+p}$ of [11] moves the Bessel order with the Szegő exponent, $(\beta,\nu)\mapsto(\beta+1,\nu+1)$, and any polynomial modification of the measure that is regular and nonzero at 0 moves neither. The proof instead analyses the same exact object twice: the degree-n Riemann–Hilbert problem Y of the lattice is steepest-descended in the frame of an arbitrary real index $m = n+\gamma$, and the two resulting expansions of the same number b_n are compared.

Let $U_m(z) := m^{pn\sigma_3} Y(zm^{-p})$, the rescaling of [11] with m^p in place of n^p; the degree stays n. Put $t_m := mz^{-1/p}$, so that the nodes $X_k = (m/k)^p$ satisfy $t_m(X_k) = k$, and let g, ϕ, $\tilde\varphi$ be the lattice g-function and phases of [11], which do not depend on the index. With the principal branch of z^γ (cut $(-\infty, 0]$) and the constant $C = m^{p\gamma\sigma_3}$, consider

$$\tilde U := C\, U_m z^{\gamma\sigma_3} C^{-1}, \qquad \tilde U z^{-m\sigma_3} = C(U_m z^{-n\sigma_3})C^{-1} \to I \quad (z \to \infty).$$

The conjugation by a constant diagonal matrix leaves the product of the off-diagonal entries of the $1/z$ coefficient unchanged, so $\tilde U$ still encodes $m^{2p} b_n^2$.

Lemma 6.5 (the m-frame)**.** *(i) The residue conditions of $\tilde U$ are those of the lattice measure $\sigma_{p,\tilde s}$, $\tilde s = s - 2p\gamma$, in the frame of index m, so that its band weight has the exponent $\tilde\alpha := \alpha - 2\gamma$. After Steps 1–4 of* [11, Section 5] *with n replaced by m in every place except the degree, the jumps are those of* [11] *with weight exponent $\tilde\alpha$, there is no jump on $(-\infty, 0)$, and the $(2,1)$ entry of the lip jumps in the hard-edge disc is $\omega^\natural e^{-4\zeta^{1/2}}$, $\zeta = m^2 f$, with $\omega^\natural = \omega^{(\tilde\alpha)} e^{\mp 2\pi i\gamma}$ in $\mathbb{C}_\pm$, whose boundary values on the band are $e^{\mp i\pi\alpha}$. That is, the m-frame problem has Bessel order α and Szegő exponent $\alpha - 2\gamma$.*

(ii) Let $p > 1$, $|\alpha| < 1$ and $0 \le \gamma < (1 - |\alpha|)/2$. Then for all $K, J, B \ge 1$, with $\mu_{m,j} = 2\zeta(-jp)/(jm^{jp})$,

$$\frac{m^p b_n}{c} = \sum_{|\mathbf{r}| \le B} \mu_m^{\mathbf{r}}\, \mathfrak{D}_{\mathbf{r},K}(m; \alpha - 2\gamma, \alpha) + O\big(m^{-K-1} + m^{-2-(B+1)p} + m^{-1-(J+1)p}\big).$$

Proof sketch. (i) The residue of $\tilde U_{j2}$ at X_k acquires the factor $(c_1/c_2) X_k^{-2\gamma} = k^{2p\gamma}$, which turns the mass k^{-s} into $k^{-\tilde s}$. The computation of the jumps in [11, Section 5] uses the integrality of the index only on $(-\infty, 0)$, where $e^{m\phi\sigma_3}$ jumps by $e^{-2\pi i m\sigma_3}$ and the cut of $z^{\gamma\sigma_3}$ contributes $e^{2\pi i\gamma\sigma_3}$; the product is $e^{-2\pi i n\sigma_3} = I$ because the degree n is an integer. In the hard-edge disc $\phi = \mp i\pi - 2f^{1/2}$ in $\mathbb{C}_\pm$ [11], so $e^{2m\phi} = e^{\mp 2\pi i\gamma} e^{-4mf^{1/2}}$; the Szegő quotient $\omega^{(\tilde\alpha)}$ tends to $e^{\mp i\pi\tilde\alpha}$ on the band, and $e^{\mp i\pi(\tilde\alpha + 2\gamma)} = e^{\mp i\pi\alpha}$. At the soft edge every jump is written with $\tilde\varphi$, which does not depend on the index. The normalisation at infinity is that of [11] with m, since $g - \operatorname{Log} z$ has no jump on $(-\infty, 0)$ for every real m.

(ii) These are the jumps of the Hurwitz problem of Section 7 under the dictionary $(n, \delta, \alpha + 2\delta) \mapsto (m, -\gamma, \alpha - 2\gamma)$, with the lattice band factor instead of the Hurwitz one and without the moved nodes. The analytic estimates of that problem (the outer scalar, the a-priori bound, the comparison at the hard edge with Bessel order α and arbitrary real Szegő exponent, the band-factor gauge and the order-K small-norm expansion, steps (H-b)–(H-e) of §7.3) are statements for a fixed index in which the index enters only as a real scale, through t_m, the Bessel variable $\zeta = m^2 f$ and the moments $\mu_{m,j}$; none of them uses its integrality. The range of γ is the range $-1 < \delta \le 0$ of those estimates, with $|\tilde\alpha| < 1$. The band factor transforms homogeneously: $\mu_{n,j}(n/m)^{jp} z^j = \mu_{m,j} z^j$. The expansion then follows as in the proof of Theorem 5.7(i), for the decoupled model at index m. □

The proof of Lemma 6.5(ii) is a transfer of estimates, written out step by step against [11] and the Hurwitz analysis in the notes of the repository named in the data availability statement; we do not repeat it. There is no circularity: Section 7 uses Theorem 6.3 only for the natural form (Theorem 7.1(ii)), not for the estimates.

Proof of Theorem 6.3. Two expansions of one number. Fix $p \in (1,2) \setminus \mathbb{Q}$, $\alpha \in (-\frac12, \frac12)$, $\gamma \in (0, \frac14)$. As $n \to \infty$ through the integers, Theorem 5.7(i) and Lemma 6.5(ii), the latter multiplied by $(n/m)^p$, both express $n^p b_n/c$ as a finite sum $\sum_e c_e n^{-e}$ plus $o(n^{-E})$, for any given E (choose K, B, J large); in the m-frame the factors $(n/m)^{(1+|\mathbf{r}|_w)p}$ and m^{-k} are convergent series in

$1/n$. In both, the exponents are $e = k + |\mathbf{r}|_w p$, $k \in \mathbb{N}_0$, and since p is irrational, the exponents $k+jp$ are pairwise distinct. Two finite sums that agree up to $o(n^{-E})$ have the same coefficients below E; at $e = k \in \mathbb{N}_0$ only $\mathbf{r} = 0$ contributes. Hence, coefficientwise, $B(n+\gamma; \alpha - 2\gamma, \alpha) = (1+\gamma/n)^p B(n; \alpha, \alpha)$.

From an open set to an identity. The coefficient of n^{-k} in the difference of the two sides is a rational function $R_k(p, \alpha, \gamma)$, which vanishes on $((1,2) \setminus \mathbb{Q}) \times (-\frac{1}{2}, \frac{1}{2}) \times (0, \frac{1}{4})$. Writing $R_k = N_k/D_k$ with polynomials, $N_k(p, \cdot, \cdot)$ vanishes on an open set of (α, γ) for infinitely many p, so each of its coefficients, a polynomial in p, has infinitely many zeros; thus $R_k \equiv 0$. Renaming $\alpha \mapsto \nu$ and putting $\gamma = (\nu - \beta)/2$ gives $B(n_0 + \frac{\nu-\beta}{2}; \beta, \nu) = ((n_0 + \frac{\nu-\beta}{2})/n_0)^p B(n_0; \nu, \nu)$ for every β. With $F_p(N; \nu) := (N/n_0)^p B(n_0; \nu, \nu)$, $n_0 := N - \frac{\nu+1}{2}$, this is (i).

The zeta layers. The same comparison identifies the coefficient of n^{-k-Jp} for each weight $J = |\mathbf{r}|_w$: $\sum_{|\mathbf{r}|_w = J} \hat{\mu}^{\mathbf{r}} [X_{\mathbf{r}}]_k = 0$, where $\hat{\mu}_j = 2\zeta(-jp)/j$ and $X_{\mathbf{r}} := (n/m)^{(1+J)p} \mathfrak{D}_{\mathbf{r}}(m; \alpha - 2\gamma, \alpha) - \mathfrak{D}_{\mathbf{r}}(n; \alpha, \alpha)$. For $J = 1$ only $\mathbf{r} = e_1$ occurs, and $\zeta(-p) \neq 0$ for $p \in (1,2)$, so $X_{e_1} = 0$. For $J = 2$ the relation reads $\zeta(-2p)[X_{e_2}]_k + 4\zeta(-p)^2 [X_{2e_1}]_k = 0$; the brackets are rational in p (after removing a common factor c^2), and if $[X_{e_2}]_k \not\equiv 0$, then $\zeta(-2p)/\zeta(-p)^2$ would be a rational function of p. It is not, since it vanishes at every odd integer $p \geq 3$ ($\zeta(-2p) = 0$, $\zeta(-p) \neq 0$). So $X_{e_2} = X_{2e_1} = 0$, and the identity argument extends both to all (p, α, γ); (ii) follows as (i) did. □

For weights $J \geq 3$ the separation would need the linear independence of the monomials $\prod_j \zeta(-jp)^{r_j}$, $|\mathbf{r}|_w = J$, over $\mathbb{Q}(p)$, which we do not prove; the shift of the layers of weight ≥ 3 is open. Exact computer algebra confirms the frame identity (6.2) identically in $(p, \alpha, \beta, \gamma)$ to order n^{-4}, and 40-digit evaluation of the recursion to order n^{-8} at 32 points, including points with $p < 1$ and with γ outside the range of Lemma 6.5; controls with a half shift $\beta = \alpha - \gamma$, the wrong sign, no frame factor, no index shift, or the Bessel order moved with β, all fail (scripts `m4_shift_exact`, `sigma_frame_exact`, `sigma_frame_num`). These checks confirm the algebraic consequence; they do not test the analysis of Lemma 6.5(ii).

6.4. **The zeta layer as a Toda flow.** In the unscaled variable the band factor $\exp(-\sum_k \mu_{n,k} z^k)$, $z = n^p x$, is $\exp(\sum_k t_k x^k)$ with

$$t_k := -\frac{2\zeta(-kp)}{k}, \tag{6.3}$$

a weight factor independent of n; these are the times of the Toda hierarchy. For a weight $w_t = e^{\sum_j t_j x^j} w$ with monic orthogonal polynomials π_n, $h_n = \int \pi_n^2 w_t$ and Jacobi matrix $\mathcal{J}$ (orthonormal basis), $\partial_{t_j} \pi_n$ has degree $< n$, so $\partial_{t_j} h_n = (\mathcal{J}^j)_{nn} h_n$ and, since $b_n^2 = h_n / h_{n-1}$,

$$\partial_{t_j} \log b_n^2 = (\mathcal{J}^j)_{nn} - (\mathcal{J}^j)_{n-1,n-1}, \qquad \partial_{t_1} a_n = b_{n+1}^2 - b_n^2. \tag{6.4}$$

Proposition 6.6 (Toda structure)**.** *Let $p \neq 1$. The μ-Taylor coefficients $\mathfrak{D}_{\mathbf{r}}$ of Theorem 5.7(i), and their analogues for a_n, are the Toda-hierarchy derivatives of the integer layers $b_n^{(0)} = cn^{-p} B(n)$ and $a_n^{(0)} = 2cn^{-p} A(n)$, with $\partial_{\mu_j} = -n^{jp} \partial_{t_j}$. For example, as identities of formal series in $1/n$ evaluated at $\mu = 0$,*

$$\partial_{\mu_1}(n^p b_n) = -\tfrac{1}{2} n^p (a_n - a_{n-1})\, n^p b_n, \qquad \tfrac{1}{2} \partial_{\mu_1}^2 (n^p b_n) = \tfrac{1}{2} n^{2p} \partial_{t_1}^2 (n^p b_n),$$
$$\partial_{\mu_2}(n^p b_n) = -\tfrac{1}{2} n^{2p} \big[(\mathcal{J}^2)_{nn} - (\mathcal{J}^2)_{n-1,n-1}\big] n^p b_n, \qquad (\mathcal{J}^2)_{nn} = a_n^2 + b_n^2 + b_{n+1}^2,$$

with $\partial_{t_1}^2 \log b_n^2 = b_{n+1}^2 - 2b_n^2 + b_{n-1}^2$.

Proof sketch. Consider the tilted lattice measures $e^{\sum_j \tau_j x^j} \sigma_{p,s}$ with small real τ. In the scaled variable the tilt is the entire factor $\exp(\sum_j \tau_j n^{-jp} z^j)$, bounded on the contours of the frame;

combined with the band factor it is again of the gauge form of Lemma 2.3, with $\mu_{n,j}$ replaced by $\mu_{n,j} - \tau_j n^{-jp}$. So Theorem 5.7(i) applies to b_n of the tilted measures uniformly for τ in a compact set, with layers polynomial in τ, and the same holds for a_n through the extraction $n^p a_n = (Q_2)_{12}/(Q_1)_{12} - (Q_1)_{22}$ of [11]. The hierarchy (6.4) holds exactly for the tilted measures at every n; inserting the expansions and taking finite differences in τ isolates each Taylor coefficient. □

The proof is only sketched: the a_n-extraction to every order and the uniformity in τ are not written out in detail. The three displayed identities hold numerically to the central-difference error (10^{-20} to 10^{-27}) at three values of (p, α) (script `m4_toda`). In this reading the zeta layer is not new data: it is a fixed, n-independent Toda flow of the integer layer, with times (6.3) given by the zeta values. The diagonal integer layer behaves like B:

Conjecture 6.7 (diagonal shift)**.** *For the decoupled model, the diagonal coefficient satisfies* $n^p a(n; \beta, \nu)/(2c) = (n/M)^p G_p(M; \nu^2)$ *with* $M := n + \frac{\beta}{2} + 1 = N + \frac{1}{2}$ *and* G_p *even in* $1/M$ *and independent of* β*, with*

$$G_p(M; \nu^2) = 1 + \frac{a_2}{M^2} + O(M^{-4}), \qquad a_2 = -\frac{p+1}{24} + \frac{(p+1)(p+2)\nu^2}{8}.$$

At $(p, \nu) = (2, -\frac{1}{2})$, $G_2 = (1 - 1/(4M^2))^{-1}$.

The evidence is numerical: the recursion at 40 digits to order 8, with β and ν decoupled (19 checks), and a_2 reconstructed from $p \in \{\frac{1}{2}, \dots, 6\}$ and asserted at three further points (scripts `m4_toda`, `m4_diag`). The value at the Bessel point is the diagonal of the Lommel measures. So b_n is centred at N and a_n at $N + \frac{1}{2}$, the arrangement of [3, Theorem 5.2] for varying weights, and Theorem 6.1 corresponds to the reversal symmetry $b_n \leftrightarrow b_{-n}$, $a_n \leftrightarrow a_{-n-1}$ of the Toda lattice. The proof of Theorem 6.3 should apply to a_n through the same invariance of the extraction once the order-K expansion of a_n is written for $M(\beta, \nu)$.

Remark 6.8 (the parity of the zeta layers). Given Proposition 6.6, the parity statements of Conjecture 9.4 reduce to the integer layers. If B is even in N and A is even in $M = N + \frac{1}{2}$, then $a_n - a_{n-1} = A(N + \frac{1}{2}) - A(N - \frac{1}{2})$ (in natural form) is odd in N, so each first-order zeta term is $\varepsilon_j N^{-1-jp}$ times an even series; second differences $b_{n+1}^2 - 2b_n^2 + b_{n-1}^2$ are even, so the ε_1^2 layer is N^{-2-2p} times an even series, as found in Theorem 5.7(ii). Higher Toda derivatives are built from such differences and products, and we expect the full monomial structure to follow; we have not written this out. By Corollary 6.4 the evenness of B is proved; that of A is Conjecture 6.7. The $1/(2j-1)$ in g_j has a heuristic reading in the same picture: for a slowly varying Jacobi matrix, $(\mathcal{J}^j)_{nn}$ is approximately $\frac{1}{2\pi}\int_0^{2\pi}(a + 2b\cos\theta)^j \, \mathrm{d}\theta$, whose first variation off the point $a = 2b$ involves $\binom{2j-2}{j-1}/\binom{2j}{j} = j/(2(2j-1))$.

7. Hurwitz lattices

For the Hurwitz lattice $\lambda_k = (k+\delta)^p$ the node zeta function is $Z_\Lambda(w) = \zeta(pw, a)$, $a := 1 + \delta$, so the dictionary of §1.3 predicts the natural index $n + \frac{\alpha}{2} - \zeta(0, a) = n + \frac{\alpha}{2} + \frac{1}{2} + \delta$ and the zeta terms with $\zeta(-jp, a)$ in place of $\zeta(-jp)$. This section proves it. The mechanism is an index transfer: the offset δ does not change the Bessel order of the hard edge, it changes only the Szegő exponent, $\alpha \mapsto \beta' = \alpha + 2\delta$; the Hurwitz problem is the decoupled model $M(\alpha + 2\delta, \alpha)$ of §6.1, and Theorem 6.3 converts the shifted Szegő exponent into the shifted natural index.

7.1. **Statement.**

Theorem 7.1 (Hurwitz lattices)**.** *Let $p > 0$, $p \neq 1$, $-1 < \delta \leq 0$ (with $\delta < 0$ if $p < 1$), $a := 1+\delta$, $1 < s < 1+2p$, and*

$$\sigma^{\delta} := \sum_{k\geq 1} (k+\delta)^{-s}\, \delta_{(k+\delta)^{-p}},$$

the measure σ_Λ of (1.1) for $\lambda_k = (k+\delta)^p$. Put $\beta' := \alpha + 2\delta$, $\mu^{\delta}_{n,j} := 2\zeta(-jp,a)/(jn^{jp})$, $N_\delta := n + \frac{\alpha}{2} + \frac{1}{2} + \delta$ and $\varepsilon_j(\delta) := \binom{2j}{j} p\, c^j \zeta(-jp,a)$.

(i) Reduction. *For all $K, J, B \geq 1$, the recurrence coefficients of σ^{δ} satisfy*

$$\frac{n^p b_n}{c} = \sum_{|\mathbf{r}|\leq B} (\mu^{\delta}_n)^{\mathbf{r}}\, \mathfrak{D}_{\mathbf{r},K}(n;\beta',\alpha) + O\big(n^{-K-1} + n^{-2-(B+1)p} + n^{-1-(J+1)p}\big),$$

with the layers $\mathfrak{D}_{\mathbf{r},K}(n;\beta,\nu)$ of the decoupled model (6.1), truncated at order n^{-K}.

(ii) Natural form. *In N_δ, the integer layer is that of the lattice to all orders, $\mathfrak{D}_0(n;\beta',\alpha) = (n/N_\delta)^p F_p(N_\delta;\alpha)$, and the layers e_1, e_2, $2e_1$ are those of the lattice to all orders, with $\zeta(-jp)$ replaced by $\zeta(-jp,a)$. For every $j \geq 1$ the layer e_j is $(n/N_\delta)^p \varepsilon_j(\delta) N_\delta^{-1-jp}(1 + g_j N_\delta^{-2} + O(N_\delta^{-4}))$. In particular, expansion (1.9) of Theorem A holds for σ^{δ} with N replaced by N_δ and ε_j by $\varepsilon_j(\delta)$, with the same $f_2, f_4, f_6, g_j, g_{1,4}$ and the same remainder.*

At $\delta = 0$ this is Theorem 5.7. The $1/n$ coefficient is $-p(\alpha+1)/2 - p\delta = -\frac{s-1}{2} - p\delta$, the first prediction tested in Table 8.3, and the n^{-1-p} coefficient is $\varepsilon_1(\delta) = 2pc\,\zeta(-p,a)$. The layers of weight ≥ 3 (μ_3, $\mu_1\mu_2$, μ_1^3) are given by (i), but their natural form beyond the orders of Lemma 5.6 is open, as for the lattice. Nothing is claimed for $\delta > 0$: then $t = \delta$, in the index variable $t = nz^{-1/p}$ of [11], is a zero of the interpolating function that carries no mass (a "phantom node"), and the pole removal of [11, Section 5] fails there. Nothing is claimed at $p = 1$ with $\delta \neq 0$.

The constants. The proof produces the band-factor moments as Fourier sums (step (H-c) below),

$$m_k(\delta) = -\frac{p}{\pi}\Gamma(kp)(2\pi n)^{-kp} S_k(\delta), \qquad S_k(\delta) := \sum_{m\geq 1} \frac{\sin(\pi kp/2 - 2\pi m\delta)}{m^{1+kp}}, \tag{7.1}$$

and $\mu^{\delta}_{n,k} = 2m_k(\delta)$. Their identification with Hurwitz zeta values is Hurwitz's formula, which we use in the form stated in [4, Theorem 1.1] (where it is quoted from Titchmarsh): *for $0 < a \leq 1$ and $\operatorname{Re} s < 0$,*

$$\zeta(s,a) = \frac{2\Gamma(1-s)}{(2\pi)^{1-s}}\Big\{\sin\frac{\pi s}{2}\sum_{m\geq 1}\frac{\cos(2\pi ma)}{m^{1-s}} + \cos\frac{\pi s}{2}\sum_{m\geq 1}\frac{\sin(2\pi ma)}{m^{1-s}}\Big\}.$$

We apply it at $s = -kp$, where $\operatorname{Re} s = -kp < 0$, and $a = 1+\delta \in (0,1]$. Since $\cos(2\pi ma) = \cos(2\pi m\delta)$, $\sin(2\pi ma) = \sin(2\pi m\delta)$ and $\sin(-\pi kp/2) = -\sin(\pi kp/2)$, the bracket is $-S_k(\delta)$, and $2\Gamma(1+kp)(2\pi)^{-1-kp} = \frac{kp}{\pi}\Gamma(kp)(2\pi)^{-kp}$. Hence $\zeta(-kp,a) = -\frac{kp}{\pi}\Gamma(kp)(2\pi)^{-kp}S_k(\delta) = kn^{kp}m_k(\delta)$, which is the normalisation of Theorem 7.1. (The formula also agrees with direct evaluation of $\zeta(-kp,a)$ to $5\cdot 10^{-42}$ at 48 parameter points, script `m3a_hurwitz_formula`.)

7.2. **The index transfer and the α-frame.** In the variable $t = nz^{-1/p}$ the Hurwitz nodes sit at $t \in a + \mathbb{N}_0$, and the mass at a node is still $n^p t^{-s}$ up to the constant of [11]; so the weight on the band, its power z^α and the Bessel order are unchanged, and only the phase changes,

$\theta = \pi t \mapsto \theta_\delta = \pi(t-\delta)$. Interpolation is done with $\sin \pi(t-\delta)$ and $1/\Gamma(t-\delta)$, so the outer scalar of [11] is built from

$$j_\delta(t) := \operatorname{Ln}\Gamma(t-\delta) - (t-\delta-\tfrac{1}{2})\log t + t - \tfrac{1}{2}\log 2\pi = j(t-\delta) + (t-\delta-\tfrac{1}{2})\operatorname{Log}(1-\delta/t) + \delta,$$

where $j = j_0$ is the Binet function of [11]. Nothing changes at the hard edge ($t \to \infty$), but at $z = \infty$ ($t \to 0$) the outer scalar F_δ now behaves like $C_\delta(-z)^\delta$ with $C_\delta \neq 0$. Normalising by the gauge

$$F^\natural := F_\delta\,(-z)^{-\delta}$$

restores the normalisation at infinity, at fixed degree, scale and g-function, and moves the power $(-z)^{-2\delta}$ into the band weight: the Szegő exponent becomes $\beta' = \alpha + 2\delta = \alpha - 2Z_\Lambda(0) - 1$. On the lips in the hard-edge disc the gauge leaves the phase $e^{\pm 2\pi i\delta}$, so the $(2,1)$ entry of the lip jump is $\omega^\natural e^{-4\zeta^{1/2}}$ with $\omega^\natural = \omega^{(\beta')}e^{\pm 2\pi i\delta}$, whose boundary values on the band are $e^{\mp i\pi\beta'}e^{\pm 2\pi i\delta} = e^{\mp i\pi\alpha}$. This is the data of a Bessel model of order α glued to the Szegő function of $x^{\beta'}$: the decoupled model $M(\beta', \alpha)$.

The hard-edge comparison must therefore be made in the *α-frame*: the local parametrix is the Bessel model of order α, and β' enters only through the quotient $\rho^\natural = \rho^{(\alpha)}/G$, with $G := (-z)^{-\delta} D_{x^{2\delta}}$, D the Szegő function on $[0, 4c]$. By the formula $D_{x^b}(z)^2 = (4cz/\Upsilon(z))^b$ of [11], $|G| = |4c/\Upsilon|^\delta$ with $\Upsilon = (z^{1/2} + (z-4c)^{1/2})^2$, $\Upsilon(0) = -4c$, so G is bounded above and below near 0 uniformly in n, for every real β'. A comparison in a β'-frame (Bessel order β') would fail: the a-priori bound $|z|^{-|\beta'|/2}$ is not square integrable at 0 once $|\beta'| \geq 1$, which happens for δ near -1.

7.3. **Outline of the proof.** The proof follows [11, Sections 5–10] and Section 5, in six steps.

(H-a) *Interpolation.* The lemmas of [11, Section 5] with $t \mapsto t - \delta$: node masses as residues, and the top node $X_1 = (n/a)^p > n^p$, so that the expansion at infinity holds for $|z| > X_1$ and the contour and the far-arc bounds are taken with constants that depend also on a.

(H-b) *Outer scalar and a-priori bound.* By the shift identity for j_δ, every bound on j_δ reduces to the Binet bound of [11] on j; this gives the sector bounds of the outer scalar ($p > 1$: Binet cut; $p < 1$: seam version) and its behaviour $C_\delta(-z)^\delta$ at infinity. The a-priori bound $|Q| \leq C_{n,\eta}|z|^{-|\alpha|/2-\eta}$ near 0 and the L^2 boundary values follow by the column factorisation of [11, Appendix C], with the exponent of the lattice, $|\alpha|/2$, because G is bounded above and below.

(H-c) *Band factor.* The jump densities of the Binet cut ($p > 1$) or of the seams ($p < 1$) are $2\operatorname{Im}\operatorname{Log}(1 - e^{-2\pi i(t_+ - \delta)})$; their moments are (7.1), by [5, (5.9.1), (5.2.1), (25.2.1)] and absolute convergence, with the Fubini bounds of Lemma 2.2, uniformly in δ. The gauge of Lemma 2.3 then holds with μ_n^δ on both circles.

(H-d) *Index transfer.* With $F^\natural$ in place of the outer scalar, the jumps are those of $M(\beta', \alpha)$ with the Hurwitz band factor, as in §7.2. This is exact algebra on the jumps of [11, Section 5].

(H-e) *Hard edge and small norm.* In the α-frame the comparison of [11, Section 7] holds verbatim, and the order-K small-norm expansion of Proposition 5.5 holds with the circle jumps $\Delta_k(\beta', \alpha)$; Lemmas 5.3 and 5.4 hold for $M(\beta, \nu)$, since β enters only through $\rho^2 = \phi^{2(\beta+1)}$ and $\phi^{-2(\beta+1)}$, bounded above and below on the circles. For $p < 1$ the glued hard-edge parametrix of [11, Section 10] is rebuilt with the gauged weight, using a Cauchy-difference bound on the seam band factor that is uniform in δ. The step to (i) also needs the linear response of Lemma 2.4 for $M(\beta, \nu)$; it holds with $d_1 = -p(\beta+1)/2$, by the same computation with the residue at 0 depending

on ν only and the shift at $4c$ on β only (verified by exact computer algebra, script `m3a_e37_decoupled`).

(H-f) *Natural form.* Exact computer algebra for $M(\beta,\nu)$, identically in (p,ν,β): the integer layer is β-free in $N = n + \frac{\beta+1}{2}$ to order N^{-4}, and the first-order zeta layer, for $j = 1, 2, 3$ and for symbolic j, is $\varepsilon_j N^{-1-jp}(1+g_j N^{-2}+O(N^{-4}))$ with β-free $g_j = A_j + B_j\nu^2$ (scripts `m4_shift_exact`, `m3a_g5_exact`).

Part (i) is (H-a)–(H-e) followed by the proof of Theorem 5.7(i). Part (ii) is (i) with (H-f) and Theorem 6.3 at $(\beta,\nu) = (\alpha+2\delta,\alpha)$: by Theorem 6.3(i) the integer layer in $N_\delta = n + \frac{\beta'+1}{2}$ is the β-free series $F_p(N_\delta;\alpha)$, the same as for the lattice in its own natural index, and by Theorem 6.3(ii) the same holds for the layers $e_1, e_2, 2e_1$. The estimates of steps (H-a)–(H-e) are written out in the notes of the repository named in the data availability statement; here we only list them. Numerically, the model $M(\alpha+2\delta,\alpha)$ with the Hurwitz μ^δ reproduces all 36 stored fits of Hurwitz lattices of §8.2, and excludes the alternatives (Bessel order moved with β', no Szegő shift, lattice μ) (script `m3a_hd_model`).

7.4. **Shifted Hurwitz lattices.** The first node sets with poles of Z_Λ that we can treat are the Hurwitz lattices shifted by a constant, $\lambda_k = (k+\delta)^p + \kappa$. For $p > 1$, $Z_\Lambda(0) = -\frac{1}{2} - \delta$ does not depend on κ, and $w \mapsto Z_\Lambda(-w)$ has the moved Weyl pole $q = 1 - \frac{1}{p}$, with residue $R_q = -\kappa q/p$, and further poles at $q_m = m - \frac{1}{p}$, $m \geq 2$, whose terms are $O(n^{-2p})$. The pole law of Conjecture 9.1(c) predicts the term $pR_qW(q)\pi\cot(\pi q)n^{-p} = -2(2-p)c\kappa\, n^{-p}$ (§8.3).

Theorem 7.2 (shifted Hurwitz lattices)**.** *Let $p > 1$, $-1 < \delta \leq 0$, $a = 1+\delta$, $\kappa \neq 0$ with $\kappa > -a^p$, $1 < s < 1+2p$, and $\sigma^{\delta,\kappa} := \sum_k \lambda_k^{-s/p}\delta_{1/\lambda_k}$ with $\lambda_k = (k+\delta)^p + \kappa$. Then, with $\beta' = \alpha + 2\delta$,*

$$\frac{n^p b_n}{c} = \sum_{0\leq k<1+p} d_k(p,\beta',\alpha)\, n^{-k} - 2(2-p)\, c\,\kappa\, n^{-p} + O(n^{-1-p}).$$

Equivalently, the n^{-k} coefficients with $k < p$ do not depend on κ and are those of Theorem 7.1, and the coefficient of n^{-p} is that of the Hurwitz lattice plus the pole term $pR_qW(q)\pi\cot(\pi q)$ of Conjecture 9.1(c).[1]

By Theorem 6.3, $\sum_k d_k(p,\beta',\alpha)n^{-k} = (n/N_\delta)^p(1 + f_2N_\delta^{-2} + f_4N_\delta^{-4} + \cdots)$, so the integer part is the lattice's natural form in N_δ, and the pole term sits at the same natural index. For $1 < p < 2$ the pole term lies between d_1/n and d_2/n^2; at $p = 2$ its coefficient vanishes.

Outline. The index map is a Möbius change of variable: κ-nodes satisfy $t_\kappa(z) = k+\delta$ with $t_\kappa = t\circ\zeta_\kappa$, $\zeta_\kappa(z) = z/(1-\kappa z/n^p)$. Composing the Hurwitz outer scalar with ζ_κ gives a scalar with the right jumps, which, unlike the Hurwitz one, behaves like $C(-z)^{-1/2}$ at infinity: the exponent at infinity of an unnormalised outer scalar is not an invariant. The invariant is the exponent of the normalising gauge at the hard edge. We take the gauge $G_\kappa = (-z)^{-\delta}\exp\mathcal{C}[\rho_1]$, where $\rho_1 = t - t_\kappa$ is minus the index offset and $\mathcal{C}$ a Cauchy transform on $(-\infty,0)$; then the normalised scalar tends to a nonzero constant at infinity, and at the hard edge $G_\kappa = (-z)^{-\delta}E_\kappa(0)(1+O(n^{1-p}|z|^{1-1/p}))$, so the Szegő shift is $\beta' = \alpha + 2\delta$, independent of κ. The jumps are those of $M(\beta',\alpha)$ with a band factor H_n^κ and an extra lip factor $e^{\pm 2\pi i\rho_1}$. The integer powers of $\log H_n^\kappa$ give the Toda

[1] The last step of the proof, the first-order response of the model to the branched tilt (the splitting of the gauge of a tilt $e^{-\mu z^\rho}$ into an analytic part with the Taylor data of the symbolic-j layer at $j = \rho$ and a remainder $\pm i\tan(\pi\rho)z^\rho$), and the proposition that uses it, were re-read independently just before this draft was assembled; no mathematical error was found, and the minor corrections from that reading were still being incorporated into the underlying notes at the time of writing.

parameters $2Z_\Lambda(-j)/j$ minus the density coefficients (the gauge supplies the term $\kappa^j\zeta(0,a)$), and the branched part is the tilt $\sum_m 2\pi\cot(\pi q_m)a_m(z/n^p)^{q_m}$, $a_m=-R_{q_m}/q_m$. The branched tilt and the lip factor come from one function, analytic across $(-\delta_0,0)$, so the circle jumps stay meromorphic (Lemma 5.4) and no further hard-edge term arises. The a-priori bound, the α-frame comparison and the small-norm expansion are transferred from steps (H-b)–(H-e); for $1<p<2$ the Hölder exponent of the band factor at 0 drops below $\frac12$, but every use of it survives because the lip coefficient carries the gauge only through a factor with exponent $\frac12$. The first-order response of the model to the tilt z^q is the symbolic-j layer of Lemma 5.6(b) continued to $j=q$, so that a tilt $e^{\tau x^q}$ of the weight contributes $-\frac{pqW(q)}{2}\tau(n/N_\delta)^pN_\delta^{-1-pq}$ at leading order; for $\tau=2\pi\cot(\pi q)a_1$ this is exactly $pR_qW(q)\pi\cot(\pi q)(n/N_\delta)^pN_\delta^{-1-pq}$. □

The proof gives more: up to $O(n^{-2p})$, n^pb_n/c is the full double series of $M(\beta',\alpha)$ with the Toda parameters of Λ, plus for each pole the first-order layer of Lemma 5.6(b) continued to $j=q_m$, evaluated at the tilt $\tau_m=2\pi\cot(\pi q_m)a_m$. It says nothing about products of pole terms. Fits at $(p,\delta,\kappa)=(2.5,-0.3,\pm0.3)$ and $(1.5,-0.4,0.4)$ confirm d_1, d_2 of $M(\alpha+2\delta,\alpha)$ to 10^{-10} and $5\cdot10^{-8}$, and the n^{-p} term to $4\cdot10^{-7}$; the Szegő exponent read from the behaviour of the unnormalised scalar at infinity is excluded (script `w_outer_e2e`). For $p<1$ the route fails: the gauge density ρ_1 is of size $|\kappa|n^{(1-p)(2+p)/p}$ on the Bessel scale, so the lattice g-function cannot carry it; the rows $p=0.6$ and $p=0.3$ of §8.3 are therefore evidence, not instances of Theorem 7.2.

8. Examples and numerical evidence

This section tests Theorem 5.7 on the lattice and collects the evidence for the conjectures of Section 9 on other node sets. No proof uses it. The computations are floating-point; the scripts and their stored outputs are listed in Appendix A.

Method. The coefficients b_n, $n\le1600$, are computed in quadruple precision by adding the atoms $1/\lambda_m$, $m\le K$, one at a time and restoring tridiagonal form by plane rotations, with a Gauss–Jacobi rule for the continuous tail beyond K; K is 100 to 200 times the largest n. Expansion coefficients are obtained by least-squares fits of n-windows ($n\ge200$, 400, 800 or mid-range windows) after subtracting the terms that are known exactly, with a pool of nuisance exponents; the number in parentheses is the spread of the fitted value over windows and pools. When a prediction was formed after looking at some of the data, we say which rows were used to form it; those rows are not evidence for it.

8.1. **The lattice in natural form.** Table 8.1 maps the lattice coefficients to $F(N):=(N/n)^pn^pb_n/c$, subtracts $1+f_2N^{-2}+\sum_j\varepsilon_jN^{-1-jp}$ and fits the remainder. The N^{-1} and N^{-2} residuals re-confirm d_1 and d_2. The N^{-3} residual is zero to the precision of the tail windows, against lattice n^{-3} coefficients of size 0.1 to 1 (for example -0.267 at $p=1.5$, $s=2$); its sign follows α at $p=1.5$, but the large-n end of the data is noise-dominated at this scale, and mid-range windows $n\in[10,400]$ give N^{-3} coefficients below $4\cdot10^{-8}$ in all cases of Table 8.2. Theorem 5.7 proves the absence of the N^{-3} term for $p>1$.

For the zeta layer, after subtracting the integer layer and the bare terms ε_jN^{-1-jp}, mid-range fits at $p\in\{1.5,2.5\}$ give an N^{-2-p} coefficient below $4\cdot10^{-7}$ and an N^{-3-p} coefficient equal to ε_1g_1 within $4\cdot10^{-6}$ to $6\cdot10^{-5}$, with spreads $8\cdot10^{-4}$ to 10^{-2} (script `m4_zeta_lattice`); at $p=1.5$ the values $g_1=95/288$ ($\alpha=0$) and $22/9$ ($\alpha=1$) were first seen in these fits.

8.2. **Hurwitz lattices.** For $-1<\delta\le0$ (and $\delta<0$ if $p<1$) and $1<s<1+2p$ the statements tested here are Theorem 7.1; the rows with $a\ge\frac32$ ($\delta>0$) or $s\ge1+2p$ are outside it, and all rows predate the theorem. For $\lambda_m=(m+\delta)^p$, $\delta>-1$, one has $Z_\Lambda(w)=\zeta(pw,a)$ with

p	s	α	N^{-1} residual	N^{-2} residual	N^{-3} residual
1.5	2	$-\frac{1}{3}$	$-4\cdot10^{-11}$	$+1.5\cdot10^{-7}$	$-3.0\cdot10^{-4}$ $(2\cdot10^{-4})$
1.5	3	$+\frac{1}{3}$	$+5\cdot10^{-11}$	$-1.6\cdot10^{-7}$	$+3.2\cdot10^{-4}$ $(2\cdot10^{-4})$
1.25	2	-0.2	$+4\cdot10^{-11}$	$-1.5\cdot10^{-7}$	$+5.2\cdot10^{-4}$ $(4\cdot10^{-4})$
2.5	2	-0.6	$-2\cdot10^{-11}$	$+5\cdot10^{-8}$	$-6\cdot10^{-5}$ $(6\cdot10^{-5})$
3	2	$-\frac{2}{3}$	$+2\cdot10^{-12}$	$-4\cdot10^{-9}$	$+2.3\cdot10^{-6}$ $(5\cdot10^{-7})$
1.5	2.5	0	$+2\cdot10^{-12}$	$-7\cdot10^{-9}$	$+1.3\cdot10^{-5}$ $(2\cdot10^{-5})$
2	3	0	$-2\cdot10^{-13}$	$+4\cdot10^{-10}$	$-3\cdot10^{-7}$ $(3\cdot10^{-7})$

TABLE 8.1. Residual coefficients of $F(N)-1-f_2N^{-2}-\sum_j\varepsilon_jN^{-1-jp}$ for the lattice, tail windows $n\geq200,400$ (script `m4_natural_index`).

p	α	f_4 fitted	f_4, (5.3)	difference
2	0	0.044921952 $(1\cdot10^{-6})$	0.044921875	$7.7\cdot10^{-8}$
2	1	-0.064452710 $(2\cdot10^{-6})$	-0.064453125	$4.2\cdot10^{-7}$
4	0	0.083271330 $(3\cdot10^{-7})$	0.083271329	$2.9\cdot10^{-10}$
4	1	-0.108692879 $(9\cdot10^{-7})$	-0.108692956	$7.7\cdot10^{-8}$
2.5	0.5	0.029254295 $(1\cdot10^{-4})$	0.029206272	$4.8\cdot10^{-5}$

TABLE 8.2. The N^{-4} coefficient of the lattice, mid-range windows (script `m4_f4_lattice`). The polynomial $f_4(2,\alpha)=(23-40\alpha^2-16\alpha^4)/512$ was first read off fits of this kind; the exact computation of Lemma 5.6 was built independently.

$a=1+\delta$, so $Z_\Lambda(-j)=\zeta(-jp,a)$ and $Z_\Lambda(0)=\frac{1}{2}-a$. The natural-index law of Section 9 predicts the $1/n$ coefficient $d_1=-\frac{s-1}{2}-p\delta$ and the n^{-1-p} coefficient $2pc\,\zeta(-p,a)$ in place of k_p. Table 8.3 compares both with fits; the lattice value k_p is excluded in every row.

p	s	a	d_1 fitted	d_1 predicted	n^{-1-p} fitted	$2pc\,\zeta(-p,a)$
0.6	2	$\frac{1}{2}$	-0.19999996	-0.2	$+0.0306156$ $(8\cdot10^{-5})$	$+0.0306142$
0.6	2	$\frac{3}{2}$	-0.80000024	-0.8	-0.309374 $(1.4\cdot10^{-4})$	-0.309386
0.6	2	2	-1.10000058	-1.1	-0.605309 $(3\cdot10^{-4})$	-0.605320
0.75	3	2	-1.75000033	-1.75	-0.779282 $(1.3\cdot10^{-3})$	-0.779293
1.5	2	$\frac{3}{2}$	-1.25000000	-1.25	-0.57244 $(5.5\cdot10^{-3})$	-0.57263
1.5	2	2	-2.00000000	-2	-1.74157 $(5.6\cdot10^{-3})$	-1.74211

TABLE 8.3. Hurwitz lattices (script `spec_hur`). The lattice values of the n^{-1-p} coefficient are -0.0899767 $(p=0.6)$, -0.0918692 $(p=0.75)$ and -0.0432947 $(p=1.5)$.

The n^{-2} coefficients of Hurwitz lattices at $p=0.6$ $(a=\frac{1}{2},\frac{3}{2},2)$, 0.75 and 1.5 agree with the index-shifted lattice value $d_2+\frac{p(p+1)}{2}\delta^2-(p+1)\delta d_1$ to between $2\cdot10^{-6}$ and $3\cdot10^{-5}$ (spreads $3\cdot10^{-5}$ to $9\cdot10^{-4}$), for shifts of size 0.3 to 3 (script `spec_hur_d2`). At $p=1.5$ the rule "the lattice expansion as a function of $n+\delta$, multiplied by $(n/(n+\delta))^p$, with every $\zeta(-jp)$ replaced by $\zeta(-jp,a)$" also predicts the n^{-3} coefficients ($\delta=-\frac{1}{2},\frac{1}{2},1$; agreement within the fit ranges, $1\cdot10^{-4}$ to $1.3\cdot10^{-2}$, relative $3\cdot10^{-4}$ to $2\cdot10^{-3}$ of the shifts; this uses the fitted lattice d_3) and the $n^{-3.5}$ coefficients, where the alternative "zeta terms unshifted" is excluded at $\delta=\frac{1}{2},1$ (script

`spec_shift_hi`). At $p = \frac{1}{3}$ and $\frac{1}{4}$ ($\delta = -\frac{1}{2}$) the first two, resp. three, coefficients $2pc\,\zeta(-p,a)$, $6pc^2\zeta(-2p,a)$, $20pc^3\zeta(-3p,a)$ agree to between $4\cdot10^{-7}$ and $3\cdot10^{-4}$, and at $p=\frac{1}{3}$ the merged n^{-2} coefficient requires the term $\frac{1}{3}\zeta(-1,a)$: without it the fit is off by seven spreads (script `spec_hur_small`).

8.3. **The pole law on synthetic lattices.** For $\lambda_k = (k+\gamma k^{-m})^p$ the counting function is unchanged at order k^0, and to first order in γ, $Z_\Lambda(-w) = \zeta(-pw) + pw\gamma\zeta(m+1-pw) + O(\gamma^2)$ has a new pole at $q = m/p$ with residue $R = -m\gamma/p$. The pole law (Conjecture 9.1(c)) predicts the term $pRW(q)\pi\cot(\pi q)n^{-1-pq} = -m\gamma W(m/p)\pi\cot(\pi m/p)n^{-1-m}$ in $n^p b_n/c$. Table 8.4 compares it with fits of $n(y_n - y_n^{\mathrm{lat}})$ after subtraction of the first-order zeta terms $pW(j)\delta Z(-j)n^{-jp}$. For $m=1$, $W(1/p)=1$, so the rows test the cotangent; the rows $m=\frac{1}{2}$ test W as well.

p	m	$\cot(\pi m/p)$	predicted	fitted, $n \ge 200$ / 400 / 800	role
0.6, 1.5	1	-0.5774	$+0.906900$	$+0.906900$	identification
1.25	1	-1.3764	$+2.162016$	$+2.162016$ / $+2.162017$ / $+2.162022$	prediction
1.8	1	-0.1763	$+0.276974$	$+0.276974$ / $+0.276974$ / $+0.276973$	prediction
0.8	1	$+1$	-1.570796	-1.570783 / -1.570792 / -1.570813	prediction
0.75	1	$+0.5774$	-0.906900	-0.906866 / -0.906885 / -0.906954	prediction
1.5	$\frac{1}{2}$	$+0.5774$	-0.424704	-0.424703 / -0.424697 / -0.424630	prediction
1.25	$\frac{1}{2}$	$+0.3249$	-0.235183	-0.235177 / -0.235112 / -0.232695	prediction

TABLE 8.4. The pole law, $s=2$, $\gamma=\frac{1}{2}$ (scripts `spec_pole`, `spec_gamma`). The kernel $\pi\cot(\pi w)$ was identified from the first row; the predictions of the other rows were fixed before the corresponding measures were computed. For $m=\frac{1}{2}$, $W(m/p) = 0.9366$ ($p=1.5$) and 0.9216 ($p=1.25$). The $n\ge800$ windows drift where few points remain.

A constant shift. For $\lambda_k = k^p + \kappa$ the index offset grows like k^{1-p}, and the expansion changes at relative order n^{-p}, not n^{-1-p}. To first order, $Z_\Lambda(-w) = \zeta(-pw) + \kappa w\zeta(p(1-w))$ has its Weyl pole moved to $q = 1-1/p$, with residue $-\kappa q/p$, and the pole law gives $-2(2-p)c\kappa\, n^{-p}$ in $n^p b_n/c$, using $W(-1/p) = p\tan(\pi/p)/\pi$. Fits of the n^{-p} coefficient give -1.20342 (0.13), -0.60154 (0.04), -1.22311 (0.74) at $(p,\kappa) = (0.6,1), (0.6,\frac{1}{2}), (0.3,1)$, against -1.20247, -0.60123, -1.22262 (script `spec_kappa_fit`); the first row was the identification point, and at $p=0.3$ the spread is 60% of the value, so that row is not evidence. For $p>1$ the law is now Theorem 7.2 (with $\delta=0$); the rows here have $p<1$, where it is not proved. The Hankel determinants of the shifted measure are those of the old nodes with an m-dependent weight: since $\varphi(y) = y/(1+\kappa y)$ satisfies $\varphi(a)-\varphi(b) = (a-b)/((1+\kappa a)(1+\kappa b))$, Heine's formula gives $D_m(\sigma_\kappa) = D_m((1+\kappa y)^{-2(m-1)}\sigma')$ with $\sigma' = (1+\kappa y)^{-s/p}\sigma$, and a first-order expansion of $\log D_m$ in κ reproduces the law, with a relative gap that decays like κn^{-p} (script `spec_mobius_pred`).

8.4. **The subleading pole term.** The pole law gives the leading power n^{-1-pq} of a pole. The same pole contributes at n^{-2-pq}, and this term is not removed when $\cot(\pi q)=0$ (Remark 9.2). For the family of §8.3, the part linear in γ of the n^{-2-m} coefficient of $n^p b_n/c$, isolated by antisymmetrising in $\gamma \in \{\pm\frac{1}{4},\pm\frac{1}{2}\}$ and Richardson extrapolation, is compared in Table 8.5 with

$$S = -\frac{mpq_1}{2}W(q_1) + \mathcal{L}\,\frac{(1+m+p)\,d_1}{p}, \qquad q_1 := \frac{m+1}{p}, \qquad \mathcal{L} := -mW(q)\pi\cot(\pi q), \tag{8.1}$$

per unit γ, with $d_1 = -(s-1)/2$. The first term is the response to the change of node density $\log|t'|$, a tilt $\propto z^{q_1}$ branched at the hard edge; the second is the leading pole term evaluated at the shifted index $n + (s-1)/(2p) = n + \delta_N$.

p	s	m	q_1	S fitted	(8.1)	role
2	2	1	1	$-1.233702\ (1\cdot10^{-6})$	-1.233701	identification
2.5	2	1	0.8	$-0.258693\ (2\cdot10^{-5})$	-0.258696	identification
3	2	1	0.667	$+0.371552\ (1\cdot10^{-6})$	$+0.371551$	identification
2.2	2	1	0.909	$-0.777016\ (7\cdot10^{-5})$	-0.776976	prediction
4	2	1	0.5	$+1.261975\ (4\cdot10^{-7})$	$+1.261975$	prediction
3	2	2	1	$-8.270517\ (3\cdot10^{-5})$	-8.270552	prediction
3	3	1	0.667	$+1.883052\ (1\cdot10^{-6})$	$+1.883050$	prediction
2.5	3	1	0.8	$+0.659993\ (2\cdot10^{-5})$	$+0.659993$	prediction
1.8	2	1	1.111	$-1.847995\ (3\cdot10^{-4})$	-1.849477	see text
1.5	2	1	1.333	$-3.39874\ (1.4\cdot10^{-2})$	-3.44204	see text
1.25	2	1	1.6	$-6.54\ (0.26)$	-7.02	see text

TABLE 8.5. The subleading pole coefficient (script `spec_subpole`). At $p = 2$, $\mathcal{L} = 0$ and $S = -W(1) = -\pi^2/8$.

For $q_1 \le 1$ the formula holds to the fit precision in the eight rows; three of them were used to form it. For $q_1 > 1$ the exponents crowd near $p = 1$ and the fits depend on the nuisance pool: refits of the stored data over twenty pools (windows $n \ge 150$ to 600, three to seven nuisance exponents) give medians minus prediction of $-2\cdot10^{-3}$ ($p = 1.5$), $+3\cdot10^{-3}$ ($p = 1.8$) and $-9\cdot10^{-2}$ ($p = 2.5$, $m = 2$), with signs depending on the pool. The same decomposition, at $p = 2$, $m = 1$, gives a nonzero n^{-3} coefficient -3.70γ for this family; for Bessel zeros (§5.2), whose index correction at $p = 2$ is tied to the offset, the corresponding terms cancel, and the expansion is the pure index shift of Proposition 5.2.

8.5. **Conjecture 9.5: predictions before fits.** To test Conjecture 9.5 on node sets with poles, with $Z(0) \ne -\frac{1}{2}$, and with masses that are not pure powers, every coefficient was predicted from the spectral data alone (regularised $Z(0)$ and $Z(-j)$, residues, density and mass factors) and the recursion of §5.3 at 40 digits (Toda responses by central differences), and the predictions were stored before any of the nine measures was computed; the test script asserts the time stamps. The conjecture has no fitted constant. The alternatives are the conjecture with one piece switched off: no Szegő shift ($\beta = \alpha$), no pole term, no $Z(-j)$, no density or mass tilt, no natural-index shift of the pole terms. A coefficient passes if $|\text{fit} - \text{prediction}| \le \max(3\cdot\text{spread}, 2\cdot10^{-6}(1 + |\text{prediction}|))$. Of the 41 coefficients tested, 40 pass; the exception is the sphere at n^{-4}, outside the claimed precision (Remark 9.7). Table 8.6 shows a selection.

The families are: (A) $\lambda_k = k^p + ak^{p-1}$, $s = 2$, which is a Hurwitz offset $\delta = a/p$ plus decaying corrections, so $Z(0) = -\frac{1}{2} - \frac{a}{p}$, with poles at $q_i = (i-1)/p$; (B) Bessel zeros $\lambda_k = (j_{\nu',k}/\pi)^P$, $s = 2$, with $Z(0) = -\frac{\nu'}{2} - \frac{1}{4}$ and poles at $q = (2i-1)/P$ (the residues use McMahon's expansion, checked against computed zeros to $5\cdot10^{-26}$); (M2) the nodes $l^{3/2}$ with masses $(2l+1)l^{-s}$, $s = 3$, a mass factor with the fractional time $\rho = \frac{2}{3}$; (M1) the sphere, nodes $l(l+1)$ with multiplicity $2l+1$, $s = 4$.

The n^{-1} coefficients test the Szegő shift $\beta = \alpha - 2Z(0) - 1$ at $Z(0) \ne -\frac{1}{2}$: they agree to 10^{-8} or better, and $\beta = \alpha$ is excluded by 10^3–10^5 tolerances. The n^{-3} rows of the families A and B test, beyond Conjecture 9.1, the natural-index shift of the pole terms with $Z(0) \ne -\frac{1}{2}$ and the density tilt as a fractional Toda time. At $q_1 = \frac{4}{3}$ (family A, $p = 1.5$) the agreement

measure	coeff.	prediction	fit (spread)	difference	next alternative
A: $p=1.5$, $a=-\frac{3}{4}$	n^{-2}	$+0.0984431$	$+0.0984431$ $(3\cdot10^{-5})$	$-3\cdot10^{-9}$	no pole: 1392
	$n^{-2.5}$	$+0.2215758$	$+0.2215749$ $(4\cdot10^{-5})$	$-9\cdot10^{-7}$	no $Z(-j)$: 1912
	n^{-3}	$+0.0699377$	$+0.0699393$ $(1.1\cdot10^{-3})$	$+1.6\cdot10^{-6}$	no tilt: 25
A: $p=1.5$, $a=0.6$	n^{-2}	$+1.0955036$	$+1.0955031$ $(1.6\cdot10^{-4})$	$-5\cdot10^{-7}$	no pole: 150
A: $p=3$, $a=-0.6$	n^{-2}	$+0.1347742$	$+0.1347742$ $(5\cdot10^{-9})$	$+5\cdot10^{-10}$	no pole: 31968
	n^{-3}	$+0.0452648$	$+0.0452649$ $(2.7\cdot10^{-6})$	$+1.0\cdot10^{-7}$	no shift: 1466
A: $p=0.6$, $a=-0.3$	$n^{-1.6}$	$+0.0854664$	$+0.0854637$ $(3.7\cdot10^{-4})$	$-3\cdot10^{-6}$	no $Z(-j)$: 77
	n^{-2}	$+0.1329122$	$+0.1327116$ $(1.1\cdot10^{-3})$	$-2.0\cdot10^{-4}$	no pole: 27
B: $\nu'=0$, $P=1.5$	n^{-2}	$+0.1957151$	$+0.1957150$ $(1.4\cdot10^{-5})$	$-1.3\cdot10^{-7}$	no pole: 562
B: $\nu'=1$, $P=1.5$	$n^{-2.5}$	-0.0664936	-0.0664957 $(5\cdot10^{-4})$	$-2\cdot10^{-6}$	no $Z(-j)$: 42
B: $\nu'=0$, $P=3$	n^{-2}	$+0.0742502$	$+0.0742502$ $(3\cdot10^{-9})$	$+5\cdot10^{-10}$	no pole: 10692
	n^{-3}	$+0.0049259$	$+0.0049264$ $(1.2\cdot10^{-5})$	$+4.8\cdot10^{-7}$	no shift: 276
M2	n^{-2}	$+0.1180556$	$+0.1180558$ $(3.8\cdot10^{-6})$	$+2.7\cdot10^{-7}$	no mass tilt: 21830
	n^{-3}	$+0.1900476$	$+0.1900481$ $(1.1\cdot10^{-3})$	$+4.8\cdot10^{-7}$	no shift: 90
M1 (sphere)	n^{-3}	-5	-4.99999992 $(3\cdot10^{-6})$	$8\cdot10^{-8}$	one tilt only: 12851
	n^{-4}	$+7.5449219$	$+7.776241$ $(3.3\cdot10^{-4})$	$+0.23132$	not claimed

TABLE 8.6. Conjecture 9.5 against fits (scripts `w_predict`, `w_test`). The last column gives the miss of the closest alternative in units of the tolerance. The row $a=0.6$ has $\delta=0.4>0$, outside Theorem 7.1; the rows $p=0.6$ are the weakest tests. Fit-free residuals $n^e(y_n-\sum_{e'<e}W_{e'}n^{-e'})-W_e$ at $n=200,400,800,1600$ decay in the rows shown except the sphere's n^{-4} (script `w_resid`).

is $2\cdot10^{-6}$–$3\cdot10^{-5}$ against spreads of 10^{-3}: consistent, not precise. For the Bessel anchor at $P=2$ (Proposition 5.2) the prediction of Conjecture 9.5, the bare model $M(\nu'-1,-\frac{1}{2})$, reproduces the coefficients $n^{-1},\dots,n^{-4}$ of the exact closed form to $2\cdot10^{-21}$. The probes of Remark 9.7 were designed after the sphere's n^{-4} failure (script `w_probe`); the two measures $k^2+\frac{1}{4}$ at $s=2$ and $s=3.5$ were then computed after the predictions of the refinement had been stored (script `w_probe2`).

8.6. **Drums.** Let λ run over the Dirichlet eigenvalues of a planar domain of area $\mathcal{A}$ and perimeter $\mathcal{P}$, rescaled to $\tilde\lambda=\mathcal{A}\lambda/4\pi$, so that the counting function is $\tilde\lambda-\beta\tilde\lambda^{1/2}+c_0+$(fluctuations) with $\beta=\mathcal{P}/\sqrt{4\pi\mathcal{A}}$; the constant $c_0=Z(0)$ carries corners and curvature ($\frac{1}{4}$ for rectangles, $\frac{1}{3}$ for the equilateral triangle, $\frac{1}{6}$ for smooth simply connected domains). We take $p=1$, $s=2$, which is outside the range of the theorems. A *smoothed drum* is the node set defined by the counting function without fluctuations, $\mathcal{N}(\lambda_k)=k-\frac{1}{2}-c_0$. The perimeter is a pole of $Z(-w)$ at $q=-\frac{1}{2}$ with residue $\beta/2$, where $\cot(\pi q)=0$ and W has a pole; the limit $W(q)\pi\cot(\pi q)\to-\pi(4c)^{-1/2}$ of (3.3) gives the prediction $n(y_n-y_n^{\mathrm{lat}})\supset-\frac{\pi\beta}{2\sqrt{2}}n^{1/2}$, which is the first term of Remark 3.5.

The corner term is isolated by comparing two smoothed drums with the same β (the square's) and $c_0=\frac{1}{4},\frac{1}{3}$: the n^0 coefficient of the difference is 0.0833487 $(2\cdot10^{-5})$, against $\Delta c_0=\frac{1}{12}$, as the term $pZ(0)$ of the dictionary requires. The absolute n^0 coefficient has median $c_0+\frac{1}{2}+\frac{\pi^2}{16}\beta^2$ over fits in all smoothed drums, to $2\cdot10^{-4}$, but single fits of it are unstable; the β^2 part is the second term of Remark 3.5, i.e. it is produced by solving the continuum soft-edge condition exactly rather than to first order. For the actual eigenvalues of the golden rectangle, the deviation from the smoothed drum has mean near 0 and a standard deviation growing like $n^{0.55}$: the counting fluctuations of the spectrum dominate everything below the perimeter term (§8.8).

smoothed drum	β	fitted $n^{1/2}$ coefficient	$-\pi\beta/(2\sqrt{2})$
square	1.128379	-1.253317 $(1\cdot10^{-5})$	-1.253314
equilateral triangle	1.286074	-1.428473 $(1\cdot10^{-5})$	-1.428469
disk	1	-1.110722 $(7\cdot10^{-6})$	-1.110721
golden rectangle	1.161199	-1.289770 $(1\cdot10^{-5})$	-1.289767

TABLE 8.7. The perimeter term at $p=1$ (script `spec_corners`). The prediction, made before the run, was that no $n^{1/2}$ term occurs, which forgot the pole of W; the value in the last column is the $0\cdot\infty$ limit of the same formula, with no free parameter.

8.7. **Anharmonic oscillators.** For the eigenvalues E_K ($K\ge1$) of $-\frac{\mathrm{d}^2}{\mathrm{d}q^2}+q^N$ on $\mathbb{R}$, $N\ge4$ even, the semiclassical quantisation reads $K-\frac12=AE^a+BE^{-a}+O(E^{-3a})$, $a=\frac12+\frac1N$, with $A=\frac1\pi\int_{-1}^{1}\sqrt{1-t^N}\,\mathrm{d}t=\frac{2}{\pi N}B(\frac1N,\frac32)$. With $p:=1/a=2N/(N+2)$ and $\lambda_K:=A^pE_K$,

$$\lambda_K^{1/p}=k'+\gamma/k'+O(k'^{-3}),\qquad k'=K-\tfrac12,\qquad \gamma=-AB,$$

a Hurwitz lattice with $a=\frac12$ plus an index correction with $m=1$, whose pole $q=1/p$ has $W(q)=1$ and $\cot(\pi/p)=-\tan(\pi/N)\neq0$. (For Bessel zeros, and for linear potentials, the poles of $Z(-w)$ sit at half-integers, where the cotangent vanishes; here it does not.) The predictions are $d_1=-\frac{s-1}{2}+\frac{p}{2}$ ($Z(0)=\zeta(0,\frac12)=0$), no n^{-1-p} term (the heat-trace exponents of q^N contain no integer, so $Z(-1)=0$), and an n^{-2} coefficient equal to the Hurwitz-shifted lattice value plus the pole term $-\gamma\pi\cot(\pi/p)=AB\pi\cot(\pi/p)$.

The pole term is rational. If the second-order quantisation condition

$$\frac{1}{\pi}\int_{-q_+}^{q_+}\sqrt{E-V}\,\mathrm{d}q-\frac{1}{24\pi}\,\partial_E^2\int_{-q_+}^{q_+}V'(q)^2(E-V)^{-1/2}\,\mathrm{d}q=K-\tfrac12+O(E^{-3a})$$

holds, then for $V=q^N$ the substitutions $q=E^{1/N}t$, $u=t^N$ give $B=-\frac{N}{12\pi}e(e-1)B(2-\frac1N,\frac12)$ with $e=\frac32-\frac1N$, and the reflection formulas collapse the product to

$$AB=-\frac{(N-1)\cot(\pi/N)}{6\pi(N+2)},\qquad\text{hence}\qquad AB\,\pi\cot\frac{\pi}{p}=\frac{N-1}{6(N+2)},$$

since $\pi/p=\pi/2+\pi/N$. The cotangent of the kernel cancels the $\cot(\pi/N)$ of the semiclassical coefficient. The values are $\frac{1}{12},\frac{5}{48},\frac{7}{60},\frac18,\frac{11}{84}$ for $N=4,\dots,12$; with B fitted from exact eigenvalues they are reproduced to between 10^{-10} and $7\cdot10^{-8}$ (script `spec_homog`); at $N=4$ the fitted $B=-0.0476724854$ equals $-1/(12\pi A)$ to $4\cdot10^{-10}$.

N	d_1 fitted (predicted)	n^{-1-p} fitted	n^{-2} fitted	predicted	without pole term
4	$+0.16666667$ $(\frac16)$	$1\cdot10^{-5}$ $(3\cdot10^{-3})$	$+0.276010$ $(1\cdot10^{-5})$	$+0.276042$	$+0.192708$
6	$+0.25000000$ $(\frac14)$	$5\cdot10^{-5}$ $(2\cdot10^{-3})$	$+0.315918$ $(3\cdot10^{-5})$	$+0.315972$	$+0.211806$

TABLE 8.8. Oscillators $-\frac{\mathrm{d}^2}{\mathrm{d}q^2}+q^N$, $s=2$, eigenvalues from a banded harmonic-oscillator basis converged to $3\cdot10^{-16}$ for $K\le1200$, with the semiclassical series beyond (scripts `spec_quartic`, `spec_qN`). With the sign of the pole term reversed the predictions would be 0.109375 and 0.107639.

8.8. **Irregular spectra.** If the nodes deviate from a smooth node function by a fixed fraction of the spacing, the expansion is replaced by an irregular component. For iid jitter $\lambda_m = (m+\xi_m)^p$, ξ_m uniform on $(-0.3, 0.3)$, the residual of y_n after a smooth fit has root mean square of order $n^{-1/2}$ (for example $1.3\cdot10^{-2}$ on $[100,200)$ and $4.2\cdot10^{-3}$ on $[800,1600)$ at $p=0.6$), and the fitted d_1 is meaningless. Jitter confined to nodes deep inside the band for all n considered is invisible (root mean square 10^{-9} to $3\cdot10^{-7}$), and jitter confined to the first 300 nodes stops producing noise once the soft-edge threshold $\lambda \approx n^p/(4c)$ has passed them (scripts `spec_jitter`, `spec_window`).

Single-node kernel. Moving node $k_0 \in \{100, 200, 400, 600\}$ by ±0.05 and following $F(n;k_0) := [D(+h) - D(-h)]/(2h)$, $D = n(y_n - y_n^{\mathrm{lat}})$, in $\tau = (n-n_*)/n_*^{1/3}$, where n_* is the index at which the node crosses the edge, gives one curve for all k_0: outside the band ($\tau \geq 2$) it is the derivative of the per-node law $-p(1-u)^{-1/2}$ to 1–3%; near $\tau = 0$ it stays finite; inside the band it oscillates with a period of about ten nodes, and its sums over n nearly cancel (script `spec_edgekernel`).

Two ensembles. Replacing the first 3000 nodes of the lattice ($p = 0.6$, $s = 2$) by unfolded eigenphases of random unitary matrices or by a Poisson bridge gives the root mean square of D in Table 8.9. Both grow like $n^{1/2}$ (exponents 0.50 and 0.52), and the Poisson amplitude is larger by the stable factor 3.3–4.0. The noise is not a linear statistic realisation by realisation (correlations between -0.04 and 0.47 between D and its linearisation), but the linearisation has the same root mean square in every window (script `spec_noiselin`). The power spectrum of the linear kernel, measured with sinusoidal displacements $\epsilon\sin(2\pi k/\Lambda_0)$, scales as $n\kappa(\omega)$ with $\kappa(\omega) \propto \omega$ for small ω (script `spec_kernelfft`). The formula $\operatorname{Var} D \approx n\int_0^\pi \kappa(\omega) S_h(\omega)\,\mathrm{d}\omega/\pi$, with displacement spectrum $S_h = K(\omega)/(4\sin^2(\omega/2))$, $K = 1$ (Poisson) and $K = \min(\omega/2\pi, 1)$ (unitary ensemble), predicts a ratio 3.95 and root mean squares 5.67, 8.02, 11.36 (unitary) and 22.5, 31.7, 44.9 (Poisson) on the last three windows, against the measured 5.38, 7.46, 11.56 and 20.8, 28.3, 41.8 for one seed.

n-window	unitary eigenphases	Poisson bridge	ratio
$[50, 100)$	2.67 ± 0.36	8.7 ± 2.5	3.3
$[100, 200)$	3.76 ± 0.34	15.0 ± 4.8	4.0
$[200, 400)$	5.38 ± 0.18	19.3 ± 3.4	3.6
$[400, 800)$	7.40 ± 0.33	26.3 ± 2.1	3.6
$[800, 1600)$	10.68 ± 0.50	39.8 ± 5.1	3.7

TABLE 8.9. Root mean square of $n(y_n - y_n^{\mathrm{lat}})$, mean and standard deviation over five seeds (scripts `spec_stats`, `spec_stats_summary`).

9. Conjectures and open problems

The theorems of Sections 3–7 concern the lattice, Hurwitz lattices and shifted Hurwitz lattices, apart from the potential theory of Section 3, which holds for general densities. This section states what they and the evidence of Section 8 suggest for other node sets, and what is missing for proofs.

9.1. **The natural-index law.** Call Λ *regular* if $\lambda_m^{1/p}$ has an asymptotic expansion $m + a - 1 + \sum_{i\geq1} \gamma_i m^{-r_i}$ with exponents $0 < r_1 < r_2 < \cdots \to \infty$, or if Λ differs from such a set in finitely many elements. Then Z_Λ is meromorphic, $Z_\Lambda(0)$ and $Z_\Lambda(-j)$ are defined by continuation, and

$w \mapsto Z_\Lambda(-w) - \zeta(-pw, a)$ has poles only at points q produced by the corrections $\gamma_i m^{-r_i}$ (at $q = r_i/p$ to first order).

Conjecture 9.1 (natural-index law)**.** *Let Λ be regular and $N_\Lambda := n + \frac{\alpha}{2} - Z_\Lambda(0)$. Then $n^p b_n/c = (n/N_\Lambda)^p F$, where F has an asymptotic expansion in powers of $1/N_\Lambda$ and of the spectral monomials below, with*

(a) *no term N_Λ^{-1}, and the term $f_2 N_\Lambda^{-2}$, with f_2 as in (5.1);*
(b) *for $j \geq 1$ the term $pW(j)Z_\Lambda(-j)N_\Lambda^{-1-jp}$;*
(c) *for every pole $q \neq -1/p$ of $Z_\Lambda(-w) - \zeta(-pw)$, with residue R_q, the term $pR_qW(q)\pi\cot(\pi q)N_\Lambda^{-1-pq}$, with the value $-(4c)^q\widetilde{L}(q)$ at $q \in -\frac{1}{2} - \mathbb{N}_0$.*

Coefficients of coincident exponents add. The statement concerns the leading power of each datum in (b) and (c); it says nothing about the terms of order N_Λ^{-2-pq} (Remark 9.2).

For the lattice, (a) and (b) are Theorem 5.7 (with $Z_\Lambda(0) = -\frac{1}{2}$, $W(j)p\zeta(-jp) = \varepsilon_j$), and (c) is empty. For Bessel zeros the whole expansion is Proposition 5.2. For Hurwitz lattices, synthetic corrections, smoothed drums and the oscillators q^N the evidence is in Section 8. In the form (b)–(c) the two kinds of spectral data are the residues of one integrand,

$$\frac{p}{n} \cdot \frac{1}{2\pi i} \int_{\uparrow} -W(w)\,\pi\cot(\pi w)\,Z_\Lambda(-w)\,n^{-wp}\,\mathrm{d}w,$$

at the integers (Theorem 4.2) and at the poles of Z_Λ (Proposition 3.4). The Weyl pole of Z_Λ is excluded: it is the leading term n of the soft-edge condition, not a correction. A moved Weyl pole, as for $\lambda_k = k^p + \kappa$, is included (§8.3). If a pole meets an integer, the two residues diverge separately and their sum produces a term $n^{-1-jp}\log n$; one weak test at $p = m = \frac{1}{2}$ is consistent with the predicted coefficient.

Two consequences make the law a spectral dictionary. Through $Z_\Lambda(0)$, the $1/n$ coefficient $d_1 = -p\alpha/2 + pZ_\Lambda(0)$ reads off the index offset of the spectrum, such as the constant a of a Hurwitz lattice (§8.2) or the corner and curvature constant of a drum (§8.6). Through $Z_\Lambda(-j)$ and the poles, it reads off the coefficients of the heat trace of the operator whose spectrum is Λ, weighted by $W(q)\pi\cot(\pi q)$.

Remark 9.2 (subleading pole terms). Each pole also contributes at order n^{-2-pq}, one order below its leading term, and this contribution is not removed by $\cot(\pi q) = 0$. For Bessel zeros at $p = 2$ the McMahon correction has $q = \frac{1}{2}$; its leading term vanishes, and its subleading term at n^{-3} cancels the zeta term $2pcZ_\Lambda(-1)$ exactly, which is why Proposition 5.2 is a pure index shift; a pure Hurwitz lattice with the same offset does carry the zeta term at n^{-3}. For the synthetic family the subleading term is (8.1) when $q_1 \leq 1$:

Conjecture 9.3. *For $\lambda_k = (k + \gamma k^{-m})^p$, $m > 0$, $p \neq 1$, $q_1 = (m+1)/p \leq 1$, the part linear in γ of the n^{-2-m} coefficient of $n^p b_n/c$ is γS with S as in (8.1).*

Both terms of (8.1) are of the type that Conjecture 9.1 does not describe: a change of node density on the band, and an index shift of a first-order term. In Theorem 5.7 the same shift is visible as the factor $(n/N)^p$ in front of every layer. This suggests that every first-order term is to be evaluated at the natural index, plus a density term.

9.2. **The natural form to all orders.**

Conjecture 9.4. *For the lattice and $p \neq 1$, $F := (N/n)^p n^p b_n/c$ depends on n only through N^{-2} and the monomials $X_j := \varepsilon_j N^{-1-jp}$: $F = \sum_{\mathbf{r}} c_{\mathbf{r}}(N^{-2}) \prod_j X_j^{r_j}$, with each $c_{\mathbf{r}}$ a power series.*

Theorem 5.7 proves the instances $\mathbf{r} = e_j$ to relative order N^{-3} for every j (and N^{-4} for $j = 1$) and $\mathbf{r} = 2e_1$ to relative order N^{-3}, with $c_{2e_1}(0) = (3p+1)/(2p)$, and Corollary 6.4 proves the instance $\mathbf{r} = 0$ to all orders. Theorem 6.3(ii) shows that the layers e_1, e_2, $2e_1$ are natural in N to all orders (their coefficients in N do not depend on the Szegő exponent); their evenness beyond the orders of Theorem 5.7(ii), and the layers of weight ≥ 3, are open. By Remark 6.8 the parity part of the conjecture would follow from Conjecture 6.7. The numbers $(3p+1)/(2p) = 1+(p+1)/(2p)$, like ε_j, do not depend on α. A closed form for $c_0 = F_p$ is known only at $(p,\alpha) = (2,-\frac{1}{2})$, where it is $(1-x/4)^{-1/2}$ in $x = N^{-2}$. The hard-edge denominators $\prod_k((2k+1)p+2)$ of Remark 5.8(b) suggested a hypergeometric structure in $a' = \frac{1}{p}+\frac{1}{2}$, but the series F_p appears to be divergent: evaluating the recursion to order 40 at 120–170 digits, at $(p,\nu) = (3, 0.3)$, the coefficients are small up to $f_{16} \approx -0.031$ and then grow factorially with oscillating sign ($f_{30} \approx -9.0\cdot 10^4$, $f_{40} \approx -1.9\cdot 10^{12}$), and Borel–Padé transforms at six values of (p,ν) show a complex-conjugate pair of singularities (scripts in `checks/large_order`). These statements are numerical; they do not affect Theorem 5.7, which concerns finite truncations.

9.3. **A general form: weighted Weyl-type spectra.** The theorems of Sections 6 and 7 suggest a single statement for general node sets, in which the Szegő-shifted model, the Toda flow and the pole terms of Conjecture 9.1 appear together. Let $\sigma := \sum_k w_k \delta_{1/\lambda_k}$ with:

- (W1) distinct nodes $0 < \lambda_1 < \lambda_2 < \cdots \to \infty$ (eigenvalues of an operator with multiplicity are merged into one node, and the multiplicity goes into the mass);
- (W2) an index function $\mathfrak{n}$ with $\mathfrak{n}(\lambda_k) = k$ and an expansion, differentiable termwise, $\mathfrak{n}(\lambda) = \lambda^{1/p} + \sum_{r\in\mathcal{R}} A_r \lambda^r + \mathfrak{n}_0 + O(\lambda^{-M})$, with $\mathcal{R} \subset (-\infty, \frac{1}{p}) \setminus \{0\}$ finite for each M (this excludes fluctuating counting functions, §8.8);
- (W3) masses $w_k = \lambda_k^{-s/p} v(\lambda_k)$ with $v(\lambda) = v_0\lambda^\tau(1+\sum_\rho b_\rho \lambda^{-\rho})$, $\rho > 0$; we put $s_{\mathrm{eff}} := s - p\tau > 1$ and, in this subsection, $\alpha := (s_{\mathrm{eff}} - 1 - p)/p$;
- (W4) $p \neq 1$, and (W5) no pole q below is an integer (otherwise a term $n^{-1-pq}\log n$ appears and nothing is claimed from that order on).

The spectral data are read from (Λ, v): the node zeta function $Z(w) = \sum_k \lambda_k^{-w}$, continued by (W2), with $Z(0) = \mathfrak{n}_0 - \frac{1}{2}$; the poles $q = -r$ of $Z(-w)$, $r \in \mathcal{R}$, with residues $R_q = qA_{-q}$ (the Weyl pole $q = -\frac{1}{p}$ is excluded); the *density factor* $D(x) := (\mathrm{d}\mathfrak{n}/\mathrm{d}\lambda)/(\frac{1}{p}\lambda^{1/p-1})$ at $x = 1/\lambda$, which is exactly $1 - p\sum_q R_q x^{q+1/p}$ up to the error term of (W2); the *mass factor* $V(x) := v(1/x)x^\tau/v_0$; the *band tilt* $\log V + \log D =: \sum_\rho B_\rho x^\rho$; and the *Toda times*

$$t_j := -\frac{2Z(-j)}{j} + B_j \quad (j \in \mathbb{N}), \qquad t_\rho := B_\rho \quad (\rho \notin \mathbb{N}).$$

For the lattice, $Z(0) = -\frac{1}{2}$, $Z(-j) = \zeta(-jp)$, there are no poles, $D = V = 1$, and t_j are the times (6.3).

Conjecture 9.5. *Under (W1)–(W5), with $\beta := \alpha - 2Z(0) - 1$, $\nu := \alpha$ and $N := n + \frac{\beta+1}{2} = n + \frac{\alpha}{2} - Z(0)$,*

$$\frac{n^p b_n}{c} \simeq \sum_{\mathbf{r}} \mu_n^{\mathbf{r}} \mathfrak{D}_{\mathbf{r}}(n;\beta,\nu) + \sum_{q\notin\mathbb{N}} pR_q W(q)\pi\cot(\pi q)\Big(\frac{n}{N}\Big)^p N^{-1-pq} - \sum_{\rho\notin\mathbb{N}} \frac{p\rho W(\rho)}{2}\, t_\rho \Big(\frac{n}{N}\Big)^p N^{-1-p\rho},$$

with $\mu_{n,j} := -t_j n^{-jp}$ and the layers (6.1) of the decoupled model $M(\beta,\nu)$. The claim is: the integer and Toda layers to every order, each pole and fractional term to relative order $O(N^{-2})$; products of two pole or fractional terms, and of such a term with a Toda time, are not claimed.

The first sum is the decoupled model flowed along the Toda hierarchy (Proposition 6.6); for $\rho = j \in \mathbb{N}$ the first-order Toda response is $\varepsilon_j = pW(j)Z(-j) = -\frac{pjW(j)}{2}t_j$ at $B_j = 0$, so there is one linear response $-\frac{p\rho W(\rho)}{2}t_\rho$ for every time, integer or not. The lattice is Theorem 5.7; Hurwitz lattices are Theorem 7.1 ($\beta = \alpha + 2\delta$); the shifted Hurwitz lattices are Theorem 7.2, at first order in the poles; and Conjecture 9.1 is the first order of Conjecture 9.5 in each datum. For the Bessel zeros $\lambda_k = (j_{\nu',k}/\pi)^2$, $s = 2$, the poles are half-integers, where $\cot(\pi q) = 0$, and the Toda times computed from the spectral data vanish ($|t_1| < 10^{-21}$, $|t_2| < 10^{-16}$ at $\nu' = 0, 1$, the precision of the regularised zero sums); so Conjecture 9.5 predicts that the measure is the bare model $M(\nu' - 1, -\frac{1}{2})$, which is what Proposition 5.2 says. In this reading the cancellation at n^{-3} of Remark 9.2 is the statement $t_1 = 0$. The tests of Conjecture 9.5, with predictions fixed before the measures were computed, are in §8.5.

Remark 9.6 (multiplicities). For an operator spectrum with multiplicities the zeta function in Conjecture 9.5 is that of the *distinct* nodes, and the multiplicity is a mass factor. With multiplicities counted, already the Weyl exponent is wrong: for the eigenvalues $l(l+1)$ of the sphere, with multiplicity $2l+1$, the distinct nodes have $p = 2$, and the fits give $n^{-2}b_n \to c(2)$ with $d_1 = -2.00000000$ at $s = 4$, while the spectral zeta function with multiplicities has Weyl exponent 1. For these nodes $Z(-j) = 0$ for every $j \geq 1$: with $t = l + \frac{1}{2} \in \frac{3}{2} + \mathbb{N}_0$, $(t^2 - \frac{1}{4})^j = \sum_m c_m t^{2m}$, and $\zeta(-2m, \frac{3}{2}) = \zeta(-2m, \frac{1}{2}) - 2^{-2m} = -2^{-2m}$ for $m \geq 0$ (since $\zeta(s, \frac{1}{2}) = (2^s - 1)\zeta(s)$ and $\zeta(-2m) = 0$ for $m \geq 1$), so $Z(-j) = -(\frac{1}{4} - \frac{1}{4})^j = 0$. The mass and density factors cancel, so Conjecture 9.5 predicts the bare model $M(1, 0)$ at $p = 2$; the fits agree at n^{-1}, n^{-2}, n^{-3} to $8 \cdot 10^{-8}$, and the alternatives "mass tilt only" and "density tilt only" are excluded.

Remark 9.7 (the dressing of pole terms). At n^{-4} the sphere misses $M(1, 0)$ by $+0.23132$ (spread $3 \cdot 10^{-4}$), outside the precision claimed in Conjecture 9.5. The probes $k^2 \pm \frac{1}{4}$ (moved Weyl pole $q = \frac{1}{2}$, residue $R = \mp\frac{1}{16}$) miss by ∓ 0.23131: the term is odd in the residue, so it is of first order in the pole, one order below its leading power n^{-1-pq}. It is explained by the following refinement, found after this failure: each pole term carries the dressing of the zeta layer continued in the index,

$$pR_qW(q)\pi\cot(\pi q)\Big(\frac{n}{N}\Big)^p N^{-1-pq}\big[1 + g(q)N^{-2} + O(N^{-4})\big], \qquad g(w) := A(w) + B(w)\nu^2,$$

with A_j, B_j of Theorem A read as functions of w (and the same for the fractional times). By Remark 5.8(d), g has a simple pole at $w = \frac{1}{2}$, where $\cot(\pi w)$ vanishes, with $\operatorname{Res}_{w=1/2} g = -(p+2)^2(p+4)(1-4\nu^2)/(256p)$. So a pole at $q = \frac{1}{2}$, whose leading term vanishes, contributes the finite term

$$R_{1/2}\, W(\tfrac{1}{2})\, \frac{\pi^2(p+2)^2(p+4)(1-4\nu^2)}{256}\, n^{-3-p/2},$$

which is $\frac{3\pi^2}{8}R_{1/2} = \mp 0.2313188$ for the probes at $p = 2$, $\nu = 0$ (fits -0.2313059 with spread $7{\cdot}10^{-4}$, and 7.7762408 against 7.7762407 for the sphere's n^{-4} coefficient). It vanishes at $\nu = \pm\frac{1}{2}$ for every p, which is why the Lommel measures ($\nu = -\frac{1}{2}$) and the earlier tests with $q = \frac{1}{2}$ poles (all at $\nu = \pm\frac{1}{2}$) show nothing. The refinement has one test made after it was formulated and before the measures were computed: for $k^2 + \frac{1}{4}$ at $s = 3.5$ ($\nu = \frac{1}{4}$) it predicts $+2.9296023$ at n^{-4}, the fit gives $+2.9296226$ (spread $3.6 \cdot 10^{-3}$), and the unrefined value $+3.1030915$ is excluded; at $s = 2$ ($\nu = -\frac{1}{2}$) it predicts no extra term, as found. The dressing at poles $q \neq \frac{1}{2}$ is untested. Its premise, that pole terms are the analytically continued zeta layer and nothing else, holds in the decoupled model with the pole structure of §7.4 (the tilt and the lip factor come from one function), and Theorem 7.2 proves it at leading order.

Remark 9.8 (a hard-edge companion for branched tilts). For a branched tilt $e^{-\mu z^\rho}$, $\rho \notin \mathbb{Z}$, that does not come from a pole (for example a mass factor), the gauge of Lemma 2.3 splits near 0 into an analytic part, whose Taylor data are those of the symbolic-j layer of Lemma 5.6(b) at $j = \rho$, and a remainder $\pm i \tan(\pi\rho) z^\rho$ in $\mathbb{C}_\pm$. In the formal model the remainder acts only through the lip jumps at the Bessel scale, and a Mellin computation with the products of the Bessel functions K_ν, K'_ν gives, besides the symbolic-j layer at $j = \rho$, a first-order *companion* term proportional to

$$(p+2)^{2\rho+1}\Phi_\nu(\rho)\Big(\frac{n}{N}\Big)^{(\rho+1)p} N^{-2\rho-2}\sum_{k\ge 0} c_k N^{-k}, \qquad \Phi_\nu(\rho) = \frac{\Gamma(-1-2\rho)\Gamma(\nu+\rho+1)}{\Gamma(-\rho)^2\Gamma(\nu-\rho)},$$

with $c_0 = (p\rho + 2\rho + 1)/2$, $c_1 = c_3 = 0$, and each c_k given by a finite exact computation; it is β-free in N. At $p = -1$, where $M(\nu-1,\nu)$ is the Laguerre weight, this reproduces the exact branched Laguerre series through N^{-4} identically in (ν,ρ), and pre-registered residues at order N^{-7} at $p = \frac{3}{2}$ and 3 hold. The companion is of relative order $N^{-2\rho-1}$; it is beyond the precision claimed in Conjecture 9.5 for $\rho > \frac{1}{2}$, which covers all the tests of §8.5, and would have to be added for $\rho < \frac{1}{2}$. That these formal layers are asymptotic expansions for actual measures with branched mass factors is not proved. Pole terms have no companion: for them the tilt and the lip factor combine to an analytic gauge.

Problem 9.9. *Prove that the N^{-2} dressing of a pole term at q is $g(q)$ with $g(w) = A(w) + B(w)\nu^2$ the continuation of g_j (Remark 9.7), or find the correct second-order edge kernel. By Remark 5.8(d) it should combine a polynomial in $\theta = u\frac{\mathrm{d}}{\mathrm{d}u}$ acting on $(1-u)^{-1/2}$ with a multiple of $(1-u)^{1/2}$. The data of Table 8.5 and of drums are test cases.*

9.4. **What a proof needs.** The derivation of Conjecture 9.1(c) from the analysis of [11] needs that analysis for a node set other than the lattice, with three ingredients: the g-function of the constrained equilibrium of the n-dependent continuum node density (Theorem 3.1); the band factor given by the Cauchy integral of the jump of $\log(1-e^{-2\pi i t_\kappa})$ for the index function $t_\kappa(z) = \kappa(n^p/z)$ (Proposition 4.4); and parametrix corrections whose first-order dependence on the perturbation is $O(n^{-2-m})$. Under these, the first-order change of $n^p b_n/c$ at order n^{-1-m} is the edge shift of (3.1) for the perturbed density, and Proposition 3.4 gives its value. For Hurwitz lattices the index normalisation of the outer function is the gauge $(-z)^{-\delta}$ of §7.2, and the shift $-p\delta$ of d_1 arises as the Szegő shift $\beta' = \alpha + 2\delta$ (Theorem 7.1); for $\lambda_k = (k+\delta)^p + \kappa$, $p > 1$, the lattice g-function still suffices, and the pole term comes from a branched band tilt rather than from the equilibrium (Theorem 7.2). For a general Weyl-type node set the template of Theorem 7.1 needs four pieces: an outer scalar built from a regularised determinant of Λ composed with an analytic index map, whose normalising gauge has the exponent $-Z_\Lambda(0) - \frac{1}{2}$ at the hard edge; band-factor moments equal to $Z_\Lambda(-j)/(jn^{jp})$; the n-dependent equilibrium above; and the linear response of Lemma 2.4 for branched tilts, including the companion of Remark 9.8. For $p < 1$ the lattice g-function cannot carry the index offset of a shifted lattice (§7.4), and the n-dependent equilibrium is needed already there.

Remark 9.10 (a proved instance of the per-node law). Conjecture 9.1, read at leading order per node, says that one node with $u = 4c\lambda n^{-p} < 1$ outside the band contributes $-p(1-u)^{-1/2}/n$ to y_n when it is removed: this is the derivative of the soft-edge condition of Remark 3.2(c). For one atom k_0^{-p} of $\sigma_{p,s}$ this is proved, uniformly for $u_n = 4c(k_0/n)^p \le 1-\eta$, with error $O(n^{-1-\epsilon})$, $\epsilon = \min(2p, 1)$ for $p < 1$ and $\epsilon = 1$ for $p \ge 1$, from the rank-one update of the Hankel matrices and a contraction argument for the minimal solution of the recurrence at the deleted node; the proof uses only the first-order expansions of a_n and b_n.

9.5. **Further problems.** (a) *The case $p = 1$.* Drums and the measures $\sum_k k^{-s}\delta_{1/k}$ have $p = 1$, where the two cotangents of Theorem 4.2 merge and [11] uses a different hard-edge analysis; Section 8.6 indicates that the dictionary persists. (b) *Higher layers.* The layers $\mu_1\mu_2$, μ_2^2, μ_1^3 control the coefficients below n^{-2-3p} and are not computed, and their shift (Theorem 6.3(ii) for weight ≥ 3) is open; it would follow from the linear independence of the products $\prod_j \zeta(-jp)^{r_j}$ over $\mathbb{Q}(p)$, or from a version of Lemma 6.5 for the tilted measures of Proposition 6.6, in which the μ_j are free parameters. (c) *Irregular spectra.* For spectra with displacements of the order of the spacing there is no deterministic expansion; Section 8.8 suggests that the variance of the irregular part is $n\int \kappa(\omega)S_h(\omega)\,\mathrm{d}\omega/\pi$ with a universal kernel spectrum κ. A closed form for κ, presumably from the Airy kernel of the soft edge, is open, as is a proof that rigidity at wavelengths of two to forty spacings sets the noise.

10. Relation to the literature

Constrained equilibrium. Existence, uniqueness and the structure (band, saturated region, balayage) of equilibrium problems with an upper constraint are classical [9, 6]; these results assume a finite constraint (Dragnev–Saff) or a compact interval (Rakhmanov) and are not used in the proof of Theorem 3.1, which establishes existence and uniqueness directly. For conjugate-gradient convergence, according to [2], Beckermann and Kuijlaars derived endpoint equations of the same Cauchy-integral form as (3.1), with an external field and compactly supported constraints; a formula for the case without external field is attributed to [1], which we have not been able to consult. We have not found the explicit density formula of Theorem 3.1, the criterion that $y\sigma(y)$ be decreasing, or the treatment of a constraint of infinite mass with a hard edge at 0. The fully saturated equilibrium of [15] ($y\sigma$ constant on a q-lattice) is the boundary case of hypothesis (H2).

Spectral data and Jacobi parameters. Orthogonal polynomials on reciprocal spectra, with asymptotics at fixed argument expressed through a regularised characteristic function, are studied in [13]. We know of no expansion of the recurrence coefficients themselves in terms of spectral zeta values at negative arguments, as in Theorem 4.2, and no reading of geometric spectral invariants (perimeter, corners) from Jacobi parameters. The Lommel family [14] is the classical exactly solvable case (Proposition 5.2).

Expansions in a shifted index. For varying weights e^{-nV} the recurrence coefficients have expansions in even powers of $1/n$ after a half-index shift; the odd terms are removed by a reindexing symmetry of the string equations [3], and for even weights the coefficients are generating functions of maps [7]. For Jacobi-type weights on $[-1, 1]$ the first correction vanishes [8]. The present setting is a fixed discrete measure without a string equation: the shift involves the spectral constant $Z(0)$, and the evenness comes with the zeta layer and its own even dressing (Theorem 5.7, Conjecture 9.4). Theorem 6.1 plays the role of the reindexing symmetry of [3], and the evenness to all orders (Corollary 6.4) is obtained without a string equation, from a second steepest-descent frame at a non-integer index (Lemma 6.5). We have not searched the literature systematically for this device, which is a gauge $z^{\gamma\sigma_3}$ of the exact Riemann–Hilbert problem combined with a change of frame. The centring of b_n at N and of a_n at $N + \frac{1}{2}$ (Conjecture 6.7) is the arrangement of [3, Theorem 5.2]. The identities of the Toda hierarchy used in Proposition 6.6 are elementary and proved in the text.

Hurwitz zeta values. The band-factor moments of a Hurwitz lattice come out of the analysis as Fourier series, and their identification with $\zeta(-jp, a)$ is Hurwitz's formula, which we quote from [4, Theorem 1.1]; there it is attributed to Hurwitz and quoted from Titchmarsh's monograph, which we have not consulted. The index transfer of Section 7, a change of the Szegő exponent at fixed Bessel order, and the statements of Theorems 7.1 and 7.2, have not been

checked against the literature on discrete orthogonal polynomials beyond the works cited in this section.

Appendix A. Reproducibility

All scripts are in the directory `research/G1/RH/spectral/checks` of the repository named in the data availability statement, next to their stored outputs (`.out`). They use the project's toolkit (quadruple-precision recurrence coefficients, cached under names that include a hash of the node array and of (p, s, K), so that a changed measure never reuses a cache), `mpmath` and `sympy`. The scripts that test a prediction end with assertions on it and print a line of the form "ALL n CHECKS PASSED" or "ALL IDENTITIES HOLD".

Exact computer algebra (Lemma 5.6). The outputs begin with the SHA-256 hashes of the scripts that produced them:

script	SHA-256
`m4_rh_exact.py`	`1e2904089fa3053cd6a7249dd4cd6149 2be3bb9c90b46c0932ff2507c9d9ffcc`
`m4_zeta_exact.py`	`3df62305bba0a4801a767c252eeee000 31e499f7c80a36946542fea6cd834041`
`m4_rh_exact2.py`	`143a308b8744bfccf3110c3a3f4a60a8 f192268b0f65f78b64bf8bd76c116521`
`m4_zeta_exact2.py`	`badda6a534fbf621c2bbef3eca6c7822 b26d13493a655a1655860449a8bab9f4`

The runs and what they assert, identically in the symbols:

output	configuration	content
`m4_rh_exact_k4.out`	$K = 4$	Lemma 5.6(a) to f_4
`m4_rh_exact_k6.out`	$K = 6$	Lemma 5.6(a): d_1, d_2, $f_1 = f_3 = f_5 = 0$, f_2, f_4, f_6
`m4_zeta_exact.out`	$K = 5$, $j = 1, \dots, 5$	Lemma 5.6(b) for integer j; $g_{1,4}$
`m4_zeta_exact2_A.out`	$K = 4$, j symbolic	Lemma 5.6(b) for all j
`m4_zeta_exact2_B.out`	$K = 5$, second order in μ_1	Lemma 5.6(c)

The first two runs also assert the coefficient identities $b_k = a_k + (k - \frac{1}{2})a_{k-1}$ (the termwise derivative of [5, (10.40.2)]) and the recursion for u_k, v_k of [5, (9.7.2)] for $k \le 8$; the symbolic run `m4_zeta_exact2_A` asserts the Taylor data of $Q_j/\binom{2j}{j}$ against the integer polynomials for $j \le 8$.

Floating-point scripts.

used in	scripts
Remark 3.6	`m1_theoremQ`, `m1_edge`
§4	`spec_kernel` (parts K0, K, P, Q), `m2_thmK_plt1`
§5.2	`spec_lommel`
Remark 5.8(b)	`m4_rh4`, `m4_f8_denominator`
§8.1	`m4_natural_index`, `m4_f4_lattice`, `m4_zeta_lattice`
§8.2	`spec_hur`, `spec_hur_d2`, `spec_shift_hi`, `spec_hur_small`
§8.3	`spec_pole`, `spec_gamma`, `spec_kappa`, `spec_kappa_fit`, `spec_mobius`, `spec_mobius_pred`, `spec_resonance`
§8.4	`spec_subpole`
§8.6	`spec_corners`
§8.7	`spec_quartic`, `spec_qN`, `spec_homog`
§6.3	`m4_shift_exact`, `sigma_frame_exact` (exact), `m4_shift_num`, `sigma_frame_num`
§6.4	`m4_toda`, `m4_diag`
§9.2	`large_order/large_order`, `large_order/borel`
Theorem 7.1	`m3a_hurwitz_formula`, `m3a_e37_decoupled`, `m3a_g5_exact` (exact), `m3a_hd_model`
Theorem 7.2	`w_outer_e2e`
§8.5, §9.3	`w_predict`, `w_test`, `w_resid`, `w_probe`, `w_probe2`
§8.8	`spec_jitter`, `spec_window`, `spec_edgekernel`, `spec_stats`, `spec_stats_summary`, `spec_noiselin`, `spec_kernelfft`

The shell script `run_all.sh` in the same directory regenerates the outputs from scratch in dependency order (readers of other scripts' caches last), including the five exact runs (about 35 minutes) and the coefficient computations (one to two and a half minutes per measure with $n \leq 1600$ on two cores). The scripts `spec_quartic`, `spec_qN`, `spec_corners`, `spec_edgekernel`, `spec_noiselin` and `spec_kernelfft` are not in `run_all.sh` and are run individually. The scripts listed for Sections 6 and 7, §9.2 and §8.5 are not in `run_all.sh` either and are run individually; the measures of §8.5 are cached outside the repository.

Acknowledgements

The research and the exposition were developed with the assistance of an AI system (Claude, Anthropic); the author takes responsibility for the content.

Data and code availability

The scripts that produced the exact identities of Section 5 and the numbers of Section 8, with their outputs, are available in the repository
https://github.com/speagleg/jacobi-asymptotics; Appendix A lists them.

References

1. Bernhard Beckermann and Arno B. J. Kuijlaars, *Superlinear convergence of conjugate gradients*, SIAM J. Numer. Anal. **39** (2001).
2. ———, *Superlinear CG convergence for special right-hand sides*, Electron. Trans. Numer. Anal. **14** (2002), 1–19.
3. Pavel Bleher and Alexander Its, *Asymptotics of the partition function of a random matrix model*, Ann. Inst. Fourier (Grenoble) **55** (2005), 1943–2000.
4. Atul Dixit and Rahul Kumar, *On Hurwitz zeta function and Lommel functions*, 2019, arXiv:1912.01199.
5. *NIST Digital Library of Mathematical Functions*, https://dlmf.nist.gov, release of 15 September 2026.
6. P. D. Dragnev and E. B. Saff, *Constrained energy problems with applications to orthogonal polynomials of a discrete variable*, J. Anal. Math. **72** (1997), 223–259.

7. Nicholas M. Ercolani, Kenneth D. T.-R. McLaughlin, and Virgil U. Pierce, *Random matrices, graphical enumeration and the continuum limit of Toda lattices*, 2006, arXiv:math-ph/0606010.
8. A. B. J. Kuijlaars, K. T.-R. McLaughlin, W. Van Assche, and M. Vanlessen, *The Riemann–Hilbert approach to strong asymptotics for orthogonal polynomials on* $[-1, 1]$, Adv. Math. **188** (2004), 337–398, arXiv:math/0111252v2.
9. E. A. Rakhmanov, *Equilibrium measure and the distribution of zeros of the extremal polynomials of a discrete variable*, Sb. Math. **187** (1996), no. 8, 1213–1228.
10. G. Speagle, *Hankel determinants of discrete measures with power-law atoms, and of zeta values*, 2026, Preprint.
11. ———, *Recurrence coefficients of discrete measures with power-law atoms: pointwise asymptotics and the* $1/n$ *term*, 2026, Preprint.
12. ———, *Stability and instability of recurrence asymptotics for discrete measures with power-law atoms*, 2026, Preprint.
13. Pavel Šťovíček and Grzegorz Świderski, *On positive Jacobi matrices with compact inverses*, 2024, arXiv:2409.19644.
14. W. Van Assche, *Compact Jacobi matrices: from Stieltjes to Krein and* $M(a, b)$, Ann. Fac. Sci. Toulouse Math. (6) (1996), 195–215, Numéro spécial Stieltjes; arXiv:math/9510214.
15. Walter Van Assche and Thomas Van Baelen, *Zero distribution of orthogonal polynomials on a q-lattice*, 2019, arXiv:1910.12518.