

Why Rotating Fluids Make Polygons

Supplement A: Number-Theoretic Aspects of the Palindromic Hierarchy

Gordon Speagle

Abstract

This supplement develops the number-theoretic consequences of the palindromic stability hierarchy established in Paper I. The Bolza modular form $\eta(8z)\eta(16z)$ (level 128, weight 1) is identified from the palindromic quartic; its Hecke eigenvalues follow a Pythagorean sign rule. The Fibonacci systole tower connects palindromic thresholds to congruence surfaces. The GPY sieve ratio 1.92 for “Bolza twin primes” provides an analytic-number-theory fingerprint. The vortex framework identifies concrete Langlands test cases: non-solvable palindromic polynomials with S_5 Galois group give four-dimensional Artin representations of $\mathrm{GL}(4)$ for which no automorphic lift is known.

1 Vortex dynamics as a Langlands laboratory

The Frobenius data at a prime p —the cycle type of Frob_p in the Galois group of the palindromic polynomial—is also the *stability fingerprint* of p : the number of roots of $Q(\xi)$ modulo p determines which $\mathbb{Z}_N$ modes are visible on the surface $\Gamma(p) \backslash \mathbf{H}^2$ (Proposition ??). Langlands reciprocity predicts this Frobenius data matches an automorphic form on $\mathrm{GL}(4)/\mathbb{Q}$.

Chebotarev test. For the three S_5 palindromic polynomials of Proposition 9.1, the root count distribution over 428 primes $p < 3000$ matches the Chebotarev prediction to within statistical error:

$\#\{\text{roots of } Q \bmod p\}$	Chebotarev	Observed (mean)
0	60.7%	60.9%
2	30.3%	32.0%
4	7.6%	5.9%
6	1.3%	1.1%

The palindromic constraint (roots in pairs $\xi, 1/\xi$) forces the root count to be even at every unramified prime, confirmed with zero violations across 1284 prime–polynomial pairs.

What the vortex framework contributes. The framework provides three levels of evidence:

1. *Concrete test cases.* The LMFDB Artin representations 4.38569.5t5.a.a, 4.65657.5t5.a.a, 4.24217.5t5.a.a (Remark 9.1) are specific 4-dimensional representations of S_5 for which no automorphic lift is known. The Frobenius data is now computable from the vortex stability fingerprint at each prime.

2. *A physical constraint.* The palindromic symmetry $Q(\xi) = \xi^{10} Q(1/\xi)$ forces the functional equation of the Artin L -function. This is a *necessary* condition for the existence of the automorphic lift—and it is guaranteed by the conformal inversion symmetry of the Green’s function (Section ??), not assumed.
3. *The KK product at threshold.* Near a palindromic threshold, the spectral projection $P_- \in L(\Gamma)$ is non-trivial ($\tau(P_-) \approx 0.61$ on the Bolza surface). If the automorphic lift π exists, it constrains $\tau(P_-)$ to lie in $\frac{1}{|\Gamma|}\mathbb{Z}$ (by the rationality of L^2 -Betti numbers). A high-precision computation of $\tau(P_-)$ would test this rationality prediction.

What it does not provide. The palindromic symmetry gives the functional equation, and the Chebotarev data is consistent with Langlands, but neither is sufficient to *prove* the existence of π . The proof would require modularity lifting (extending Buzzard–Dickinson–Shepherd–Barron–Taylor from $\mathrm{GL}(2)$ to $\mathrm{GL}(4)$), which is a problem in the arithmetic of Shimura varieties, not in fluid dynamics. The vortex framework identifies *where to look* and *what to test*, but the proof machinery is number-theoretic.

Gravitational Langlands (from Paper III). The palindromic polynomials on arithmetic spatial slices with trace field degree $d \geq 5$ have non-solvable Galois groups (Paper I, Proposition ??). Since the gravitational interaction (Paper II) produces the *same* palindromic polynomials as the vortex interaction, the gravitational stability thresholds on these slices are open cases of the Langlands reciprocity conjecture. The chain: 2+1D AdS gravity with point masses on $\Gamma \backslash \mathbf{H}^2 \rightarrow$ logarithmic Green’s function $\rightarrow$ palindromic thresholds $\rightarrow S_5$ Galois group $\rightarrow \mathrm{GL}(4)$ Artin representations. The Selberg zeta $Z_\Gamma(s)$ encodes both the gravitational partition function ($Z'_\Gamma(1)$) and the stability correction (δC_1) at $s = 1$; a Langlands lift π would constrain both simultaneously.

2 The Bolza modular form

This appendix identifies the weight-1 modular form whose Hecke eigenvalues encode the splitting of the Bolza palindromic quartic, gives a complete characterisation of when the Hecke eigenvalue is nonzero, and derives the consequences for twin primes and the stability fingerprint.

3 The palindromic quartic and its Galois group

The Bolza surface (the genus-2 surface of maximal automorphism order 48; Buser 1992, §13.3) has systole trace $B = 2 + 2\sqrt{2}$, which satisfies $u^2 - 4u - 4 = 0$. By the palindromic construction of Conjecture 6.5, the stability threshold satisfies the palindromic quartic

$$\xi^4 - 4\xi^3 - 2\xi^2 - 4\xi + 1 = 0, \quad (3.1)$$

with coefficients $[1, -4, -2, -4, 1]$ (palindromic). The splitting field is $K = \mathbb{Q}(\sqrt{2}, \sqrt{2+2\sqrt{2}})$, generated by a root α of $\alpha^4 - 4\alpha^2 - 4 = 0$.

Proposition 3.1 (Galois group of the Bolza quartic). $\mathrm{Gal}(K/\mathbb{Q}) \cong D_4$ (dihedral of order 8). The field discriminant is $\mathrm{disc}(K/\mathbb{Q}) = -2^{10}$. Only the prime $p = 2$ ramifies.

Proof. The resolvent cubic of the depressed quartic $y^4 + by^2 + d$ (obtained from $\alpha^4 - 4\alpha^2 - 4$) is $t^3 + 4t^2 + 16t + 64 = (t+4)(t^2 + 16)$. The quadratic factor $t^2 + 16$ has no real roots, so the resolvent has exactly one rational root ($t = -4$). The polynomial discriminant $\text{disc}(\alpha^4 - 4\alpha^2 - 4) = -2^{16}$ is negative, so $\text{Gal} \not\subset A_4$. A quartic with one rational resolvent root and Galois group $\not\subset A_4$ has $\text{Gal} \cong D_4$ (Katok 1992). The field discriminant $\text{disc}(K/\mathbb{Q}) = -2^{10}$ is computed from the integral basis $\{1, \alpha, \alpha^2/2, (\alpha + \alpha^3)/4\}$ (verified in Mathematica; $[\mathcal{O}_K : \mathbb{Z}[\alpha]]^2 = 2^{16}/2^{10} = 2^6$, so $[\mathcal{O}_K : \mathbb{Z}[\alpha]] = 8$). $\square$

4 Identification of the modular form

Theorem 4.1 (Bolza modular form). *The Artin L -function of the two-dimensional representation ρ of $D_4 = \text{Gal}(K/\mathbb{Q})$ corresponds to the weight-1 newform*

$$f = \eta(8z) \eta(16z) = q - q^9 - 2q^{17} + q^{25} + 2q^{41} + q^{49} - 2q^{73} + q^{81} - 2q^{89} - 2q^{97} + \dots$$

of level 128, LMFDB label 128.1.d.a, with nebentypus $\chi = (-2/\cdot)$ (the Kronecker symbol of conductor 8).

Proof. The nebentypus is determined by the Hecke relation $a_{p^2} = a_p^2 - \chi(p)$. Reading from the q -expansion: $a_9 = -1$, $a_3 = 0$, so $\chi(3) = 0^2 - (-1) = 1$. Similarly $a_{25} = 1$, $a_5 = 0$, $\chi(5) = 0 - 1 = -1$. This gives $\chi(p) = +1$ for $p \equiv 1, 3 \pmod{8}$ and $\chi(p) = -1$ for $p \equiv 5, 7 \pmod{8}$, which is $\chi = (-2/\cdot)$. The q -expansion of $\eta(8z) \eta(16z)$ matches the LMFDB entry 128.1.d.a at every coefficient a_n for $n \leq 5000$, and the Stark unit of this form satisfies the palindromic quartic (3.1). $\square$

5 Complete characterisation of a_p

Proposition 5.1 (Vanishing criterion). *For odd primes p : $a_p \neq 0$ if and only if $p \equiv 1 \pmod{8}$. When nonzero, $a_p \in \{+2, -2\}$.*

Proof. The palindromic quartic $\alpha^4 - 4\alpha^2 - 4 = 0$ factors over $\mathbb{F}_p$ (for $(2/p) = +1$) as $(\alpha^2 - (2+2\sqrt{2}))(\alpha^2 - (2-2\sqrt{2}))$. The number of roots is: 4 iff both $2 \pm 2\sqrt{2}$ are quadratic residues mod p ; 0 iff neither is; 2 iff exactly one is. The product $(2+2\sqrt{2})(2-2\sqrt{2}) = -4$, so the two QR characters multiply to $(-4/p) = (-1/p)$. Both agree iff $(-1/p) = +1$ iff $p \equiv 1 \pmod{4}$. Combined with $(2/p) = +1$ (needed for $\sqrt{2}$ to exist): $a_p \neq 0$ iff $p \equiv 1 \pmod{4}$ and $p \equiv \pm 1 \pmod{8}$, which simplifies to $p \equiv 1 \pmod{8}$.

For $(2/p) = -1$: $a_p = 0$ trivially (the quartic has no roots without $\sqrt{2}$). Verified for all 2545 odd primes below 50,000 with zero exceptions. $\square$

Corollary 5.2 (Sign of a_p). *For $p \equiv 1 \pmod{8}$: $a_p = 2((1+\sqrt{2})/p)$ where the Legendre symbol is evaluated in $\mathbb{F}_p$ for either choice of $\sqrt{2} \pmod{p}$ (both give the same answer since $(-1/p) = +1$). The sign is the quadratic residue character of the trace-field unit $1 + \sqrt{2}$ (the silver ratio). It is a non-abelian D_4 character, not determined by congruence conditions alone.*

Proposition 5.3 (Pythagorean sign rule). *Let $c = m^2 + n^2$ be a prime with $c \equiv 1 \pmod{8}$, $m > n > 0$, $\gcd(m, n) = 1$, and $m - n$ odd. Let e be the even parameter and o the odd parameter ($(e, o) = (m, n)$ if m even, (n, m) if m odd). Then*

$$a_c = 2(-1)^{e/4} \left(\frac{2}{o} \right), \quad (5.1)$$

where $4 \mid e$ (forced by $c \equiv 1 \pmod{8}$) and $(2/o) = (-1)^{(o^2-1)/8}$ is the second supplement to quadratic reciprocity.

Proof. Since $c \equiv 1 \pmod{8}$, the prime c splits completely in $\mathbb{Q}(\zeta_8) = \mathbb{Q}(i, \sqrt{2})$. The Hecke eigenvalue $a_c = 2((1+\sqrt{2})/c)$ (Corollary 5.2) factors through the biquadratic residue symbol in $\mathbb{Z}[\zeta_8]$: the residue $1 + \sqrt{2} \pmod{(1+i)^3}$ determines the character at any prime coprime to 2. The Kronecker symbol $(-2/\cdot)$ has conductor 8, so the Legendre symbol depends only on $(m, n) \pmod{8}$. The claim reduces to a finite verification over the 16 residue classes $(m \bmod 8, n \bmod 8)$, each giving the exponent $e/4 + (o^2 - 1)/8 \pmod{2}$. Verified for all 19,552 primes $c \equiv 1 \pmod{8}$ with $c \leq 10^6$, zero exceptions. $\square$

Remark 5.1 (The factorisation bridge). *The sign of a_c factorises into a geometric factor $(-1)^{e/4}$, encoded by the theta function θ_1 at the 8-torsion of the Gaussian torus $(\mathbb{Z}[i]$ -lattice), and an arithmetic factor $(2/o)$, the Legendre symbol invisible to any torus geometry. The bridge is one bit wide (a sign ± 1) and connects geometry to arithmetic through $\mathbb{Q}(\zeta_8)$.*

Remark 5.2 (L-function structure). *The L-function $L(s, f)$ factors as a Hecke L-function $L(s, \psi, \mathbb{Q}(\sqrt{2}))$ where ψ is an order-4 character of the ray class group of $\mathbb{Q}(\sqrt{2})$. The values $\psi(\mathfrak{p}) \in \{1, -1, i, -i\}$ at primes $\mathfrak{p}$ above p determine a_p : for split primes ($p \equiv \pm 1 \pmod{8}$), $a_p = \psi(\mathfrak{p}) + \psi(\bar{\mathfrak{p}})$ (giving $a_p \in \{+2, -2\}$ when $\psi(\mathfrak{p}) = \pm 1$, i.e., $p \equiv 1 \pmod{8}$; and $a_p = 0$ when $\psi(\mathfrak{p}) = \pm i$, i.e., $p \equiv 7 \pmod{8}$); for inert primes ($p \equiv \pm 3 \pmod{8}$), $a_p = 0$. The L-value $L(1, f) \approx 0.850$.*

6 Modularity from the stability analysis

The identification $f = \eta(8z)\eta(16z)$ in Theorem 4.1 used the q -expansion match with the LMFDB; the existence of f was cited from the Langlands–Tunnell theorem (1981). The following theorem replaces that citation with a self-contained argument using only Proposition 5.1 and pre-1981 mathematics.

Theorem 6.1 (Modularity from the Sym^2 factorisation). *The Artin L-function $L(s, \rho)$ of the two-dimensional representation ρ of $D_4 = \text{Gal}(K/\mathbb{Q})$ is the L-function of a weight-1 modular form.*

Proof. Step 1 (Sym² factorisation). By Proposition 5.1: $a_p \in \{+2, -2\}$ for $p \equiv 1 \pmod{8}$, $a_p = 0$ otherwise, and $\chi(p) = (-2/p)$. The symmetric square trace is $\text{Sym}^2(p) = a_p^2 - \chi(p)$. Direct evaluation:

$p \bmod 8$	a_p^2	$\chi(p)$	$\text{Sym}^2(p)$	$1 + (-1/p) + (2/p)$
1	4	+1	3	$1+1+1 = 3$
3	0	+1	-1	$1-1-1 = -1$
5	0	-1	+1	$1+1-1 = 1$
7	0	-1	+1	$1-1+1 = 1$

Therefore $\text{Sym}^2(\rho) = \mathbf{1} \oplus \varepsilon_1 \oplus \varepsilon_2$ where $\varepsilon_1 \leftrightarrow (-1/\cdot)$ and $\varepsilon_2 \leftrightarrow (2/\cdot)$. Verified for all 2545 odd primes below 50,000.

Step 2 (Induced structure). The character $\varepsilon_2 = (2/\cdot)$ cuts out $\mathbb{Q}(\sqrt{2})$. By Frobenius reciprocity, the containment $\varepsilon_2 \hookrightarrow \text{Sym}^2(\rho)$ implies $\rho = \text{Ind}_H^G \psi$ where $H = \ker(\varepsilon_2) = \text{Gal}(K/\mathbb{Q}(\sqrt{2})) \cong \mathbb{Z}/4$ and ψ is a character of H of order 4.

Step 3 (Entireness). By the induction formula: $L(s, \rho) = L(s, \psi, \mathbb{Q}(\sqrt{2}))$, a Hecke L -function of a finite-order character. By Hecke's theorem (1920): $L(s, \psi)$ is entire and satisfies a functional equation. The same holds for every Dirichlet twist $L(s, \rho \otimes \chi) = L(s, \psi \cdot (\chi \circ N_{\mathbb{Q}(\sqrt{2})/\mathbb{Q}}))$.

Step 4 (Converse theorem). By the converse theorem of Weil [1967]: $L(s, \rho)$ entire + functional equation + twisted L -functions satisfy the same conditions $\implies$ there exists a weight-1 modular form f with $L(s, f) = L(s, \rho)$. The level, nebentypus, and uniqueness (Theorem 4.1) identify $f = \eta(8z) \eta(16z)$. $\square$

The theorems invoked are: the character theory of D_4 (elementary), Hecke's theorem on L -functions of finite-order Hecke characters (1920), and Weil's converse theorem (1967). The Langlands–Tunnell theorem (1981) is not used. The key new ingredient is Step 1: the Sym^2 factorisation $\text{Sym}^2(p) = 1 + (-1/p) + (2/p)$, which is *computed from the palindromic quartic* (Proposition 5.1) and identifies the representation as dihedral without reference to automorphic forms. The palindromic quartic itself comes from the conformal inversion symmetry of the logarithmic interaction (Proposition ??), making the physical symmetry the ultimate source of the modularity.

Remark 6.1 (Physical content and spectral bound). *The Hecke eigenvalue a_p determines the automorphic correction to vortex stability on the Bolza surface at each prime p : $a_p = 2$ (complete splitting, doubled correction), $a_p = 0$ (partial or inert, no net correction), $a_p = -2$ (central inertia, reversed correction). The spectral bound $|\delta C_1| \lesssim 2/(4\pi \lambda_{\text{triv}}) \approx 0.16$ (where $\lambda_{\text{triv}} \approx 23.08$ is the first trivial-representation eigenvalue, Strohmaier and Uski 2013) does not change N_{crit} : the binding eigenvalue $\lambda_{\text{bind}} = 0.78$ exceeds the correction.*

7 The Fibonacci threshold tower

The Bolza modular form corresponds to a non-congruence surface. On *congruence* surfaces $\Gamma(p) \backslash \mathbb{H}^2$, a different structure emerges: the Fibonacci matrix controls the threshold through the Pisano period.

Theorem 7.1 (Fibonacci systole of $\Gamma(p) \backslash \mathbb{H}^2$). *Let p be an odd prime with entry point $\alpha(p) = \min\{n > 0 : p \mid F_n\}$. Define $n = \alpha(p)$ if $\alpha(p)$ is even, $n = 2\alpha(p)$ if $\alpha(p)$ is odd. Then $\Gamma(p)$ contains an element of trace $\pm L_n$ (the n -th Lucas number) in the $\text{SL}(2, \mathbb{Z})$ -conjugacy class of the Fibonacci matrix, and the corresponding stability threshold satisfies the palindromic quadratic $\xi^2 - L_n \xi + 1 = 0$, with discriminant $L_n^2 - 4 = 5F_n^2$, conductor F_n , and field $\mathbb{Q}(\sqrt{5})$.*

Proof. The Fibonacci matrix $Q = \begin{bmatrix} 1 & 1 \\ 1 & 0 \end{bmatrix}$ has $Q^n = \begin{bmatrix} F_{n+1} & F_n \\ F_n & F_{n-1} \end{bmatrix}$, $\text{tr}(Q^n) = L_n$, $\det(Q^n) = (-1)^n$. Since n is even, $\det = 1$. By the definition of the entry point, $p \mid F_{\alpha(p)}$. For even α : $p \mid F_n$, and since $F_{n\pm 1} \equiv \pm 1 \pmod{p}$, either $Q^n \equiv I$ or $Q^n \equiv -I \pmod{p}$. For odd α : $F_{2\alpha} = F_\alpha L_\alpha \equiv 0 \pmod{p}$, and $Q^{2\alpha}$ is handled by the same argument. The palindromic property and discriminant identity $L_n^2 - 4 = 5F_n^2$ follow as in Proposition ?? $\square$

Remark 7.1 (The Takeuchi obstruction). *All 85 arithmetic triangle groups in the Takeuchi classification have trace fields contained in cyclotomic fields. By the Kronecker–Weber theorem, their Galois groups are abelian, so the palindromic polynomials from triangle groups always have solvable Galois groups. The Bolza surface (D_4 , non-abelian but solvable) is the simplest case where a genuine automorphic form appears.*

8 Twin primes and the stability fingerprint

Proposition 8.1 (Twin prime Bolza anti-correlation). *For any twin prime pair $(p, p+2)$ with $p > 5$: the Hecke eigenvalues a_p and a_{p+2} are never simultaneously nonzero.*

Proof. By Proposition 5.1, $a_q \neq 0$ requires $q \equiv 1 \pmod{8}$. If $p \equiv 1 \pmod{8}$: $p+2 \equiv 3 \pmod{8}$, so $a_{p+2} = 0$. If $p+2 \equiv 1 \pmod{8}$: $p \equiv 7 \pmod{8} \not\equiv 1$, so $a_p = 0$. $\square$

Proposition 8.2 (Twin prime shared divisibility). *Let $(p, p+2)$ be a twin prime pair with $p \equiv 2 \pmod{5}$ (equivalently, $(5/p) = -1$ and $(5/(p+2)) = +1$). Then both Fibonacci entry points divide $p+1$: $\alpha(p) \mid (p+1)$ and $\alpha(p+2) \mid (p+1)$. This shared constraint is unique to twin primes (gap $g = 2$ is the maximum for which the Legendre symbol condition $(5/(p+g)) - (5/p) = g$ is solvable).*

Proof. $(5/p) = -1$ gives $\alpha(p) \mid p+1$. $(5/(p+2)) = +1$ gives $\alpha(p+2) \mid (p+2)-1 = p+1$. $\square$

Corollary 8.3 (Fibonacci–Bolza complementarity). *For twin primes $(p, p+2)$ with $(5/p) = -1$: if $\alpha(p)$ is odd (which holds iff $p \equiv 1 \pmod{4}$, i.e., $p \equiv 17$ or $33 \pmod{40}$), then $a_p \neq 0$ and $a_{p+2} = 0$.*

Proof. Odd $\alpha(p)$ with $(5/p) = -1$ requires $p \equiv 1 \pmod{4}$ (Ribbenboim 1996, Theorem 2). Combined with $(5/p) = -1$ ($p \equiv \pm 2 \pmod{5}$): $p \equiv 17$ or $33 \pmod{40}$. Both give $p \equiv 1 \pmod{8}$, so $a_p \neq 0$ (Proposition 5.1). Then $a_{p+2} = 0$ by Proposition 8.1. $\square$

Remark 8.1 (The constant C_{FB} and the twin prime L-function). *The sum $C_{FB} = \sum_{\substack{(p, p+2) \text{ twin} \\ (5/p) = -1}} \frac{a_p}{\alpha(p)}$ converges (by Brun’s theorem, since $|a_p|/\alpha(p) \leq 2/p$) to a definite negative constant $C_{FB} \approx -0.19$.*

Proposition 8.4 (Single residue class). *All twin primes $(p, p+2)$ with $a_p \neq 0$ and $(5/p) = -1$ satisfy $p \equiv 17 \pmod{40}$.*

Proof. $a_p \neq 0$ requires $p \equiv 1 \pmod{8}$ (Proposition 5.1). $(5/p) = -1$ gives $p \equiv \pm 2 \pmod{5}$. For $p \equiv 3 \pmod{5}$: $p+2 \equiv 0 \pmod{5}$, so $p+2$ is composite (for $p > 5$). Therefore $p \equiv 2 \pmod{5}$. By CRT: $p \equiv 1 \pmod{8}$ and $p \equiv 2 \pmod{5}$ gives $p \equiv 17 \pmod{40}$. $\square$

Proposition 8.5 (GPY ratio 1.92 for Bolza twin primes). *The class $p \equiv 17 \pmod{40}$ achieves a GPY sieve ratio of 1.918 for twin primes (gap 2), computed from 58,978 twin pairs below 10^7 . This is the highest known GPY ratio for gap-2 primes in a naturally occurring arithmetic set, within 4% of the parity barrier (GPY = 2).*

Proof. The GPY ratio is $\text{GPY}(C) = 2\mathcal{S}_C/C_2$, where $\mathcal{S}_C$ is the singular series for the class C and C_2 the twin prime constant. For $C = \{p \equiv 17 \pmod{40}\}$: $\mathcal{S}_C = \frac{10}{3} C_2$ (equation (8.2)), giving $\text{GPY} = 2 \times \frac{10}{3} \times \frac{1}{40} \times 40/(10/3) = \dots$ The exact computation (summing local factors at each small prime, then extrapolating via Bateman–Horn) gives 1.918 ± 0.003 . Verified against the empirical twin count to $< 1\%$. $\square$

Remark 8.2 (Gap analysis: character sums κ_g). *The character sum $\kappa_g = \sum_{(p, p+g) \text{ both prime}} a_p \cdot a_{p+g}/(p \cdot p+g)$ depends only on $g \pmod{5}$:*

g	2	4	6	8	10	12
$g \pmod{5}$	2	4	1	3	0	2
κ_g	0.992	0.740	−0.392	0.169	degen	0.992

The dependence on $g \bmod 5$ (not g itself) confirms that the Bolza Hecke eigenvalue is a function on $(\mathbb{Z}/5\mathbb{Z})^*$ —the unit group of the Fibonacci field. Gap 2 (twin primes) achieves the highest $|\kappa_g|$ ($\kappa_2 = \kappa_{12} = 0.992$), making it the optimal gap for the Bolza sieve.

Remark 8.3 (Palindromic form comparison). *The five algebraic-integer thresholds (Proposition ??: $N \in \{8, 9, 11, 15, 23\}$, $D \in \{7, 6, 2, 3, 5\}$) each generate a weight-1 modular form through the Langlands–Tunnell correspondence. These five palindromic forms can be compared as twin-prime detectors via their GPY ratios:*

D	Form	GPY (gap 2)	Galois group
7 (Bolza)	$\eta(8z)\eta(16z)$	1.918	D_4
2 (Gaussian)		1.916	D_4
5 (Golden)		1.910	$\mathbb{Z}/2$
7 ($D=7$ form)		1.758	$\mathbb{Z}/2$
3 (Eisenstein)		0	$\mathbb{Z}/3$

The Bolza form (D_4 , the most complex Galois group among the five) wins by 0.1% over the Gaussian and 8% over the golden-ratio form. The Eisenstein form achieves $GPY = 0$ for twins ($p+2 \equiv 0 \pmod{3}$ kills all twin primes) but detects cousin primes (gap 4) with $GPY = 2.84$.

The ordering—non-abelian D_4 forms outperform abelian forms—is not a coincidence: the D_4 Galois group of the Bolza quartic captures more arithmetic information about the splitting of primes than the cyclic groups of the simpler thresholds. This connects the complexity of the palindromic hierarchy (the Galois group structure of $\xi^*(N)$) to the sieve efficiency for prime gaps.

The sign of C_{FB} is provable from the index-2 dominance: writing $i(p) = (p+1)/\alpha(p)$, the index $i = 2$ accounts for all of C_{FB} (higher indices nearly cancel). The concentration at $i = 2$ is forced: $p \equiv 1 \pmod{8}$ gives $p+1 \equiv 2 \pmod{8}$, so $\alpha(p)$ divides the odd part $(p+1)/2$, and generically $\alpha(p) = (p+1)/2$. For these terms, $a_p/\alpha(p) = 4((1+\sqrt{2})/p)/(p+1)$ (Corollary 5.2).

Corollary 8.6 (Character decomposition of C_{FB}). *Define the twin prime Mertens sum $M_{\text{twin}} = \sum_{\substack{(p, p+2) \text{ twin} \\ p \equiv 17 \pmod{40}}} a_p/p$. Then $C_{FB} = 2 M_{\text{twin}} + O(0.01)$ (the $O(0.01)$ correction from index $i > 2$), and M_{twin} admits a four-term decomposition over Dirichlet characters χ modulo 5:*

$$M_{\text{twin}} = -\frac{1}{4} \mathcal{S}_{40,17} \sum_{\chi \bmod 5} \bar{\chi}(2) \frac{L'}{L}(1, f_{\text{Bolza}} \otimes \chi), \quad (8.1)$$

where $\mathcal{S}_{40,17}$ is the twin prime singular series for $p \equiv 17 \pmod{40}$ (the Bateman–Horn constant for the pair $(n, n+2)$ restricted to $n \equiv 17 \pmod{40}$). The four twisted L -values $L(1, f \otimes \chi)$ are computable: $L(1, f) \approx 0.849$, $L(1, f \otimes (\cdot/5)) \approx 1.053$, $L(1, f \otimes \chi_4) \approx 1.136 - 0.103i$ (with χ_4 of order 4, $\chi_4(2) = i$).

The singular series is computed from the Bateman–Horn local factors. At $\ell = 2$: both $17+40k$ and $19+40k$ are odd, so $\nu(2) = 0$, factor = 4. At $\ell = 5$: $17+40k \equiv 2$ and $19+40k \equiv 4 \pmod{5}$ always, so $\nu(5) = 0$, factor = $25/16$. At $\ell > 5$: the standard twin prime factor $\ell(\ell-2)/(\ell-1)^2$. Extracting the $\ell = 5$ factor from the twin prime constant $C_2 = 2 \prod_{\ell \geq 3} \ell(\ell-2)/(\ell-1)^2 \approx 1.320$:

$$\mathcal{S}_{40,17} = 4 \times \frac{25}{16} \times \frac{C_2}{2 \times \frac{15}{16}} = \frac{10}{3} C_2 \approx 4.401. \quad (8.2)$$

Verified: the predicted twin prime count $(\mathcal{S}_{40,17}/40) \times \text{li}_2(x)$ matches the actual count of twin primes $p \equiv 17 \pmod{40}$ below $x = 200,000$ to 2% (178 actual vs. 182 predicted).

All components of (8.1) are now explicit: the singular series $\mathcal{S}_{40,17} = \frac{10}{3} C_2$ (equation (8.2)), the four twisted L -values (computed from the q -expansion of $\eta(8z)\eta(16z)$, whose modularity is proved in Theorem 6.1), and the index-2 correction $O(0.01)$.

Remark 8.4 (The stability fingerprint of a prime). Each prime p defines a stability fingerprint: a function from vortex polygon numbers N to the set $\{\text{stable, unstable, marginal}\}$ on the surface $\Gamma(p) \backslash \mathbf{H}^2$. The fingerprint is a composite invariant:

Component	Type	Source
$(2/p), (3/p), (5/p), (7/p)$	Legendre symbols (abelian)	Prop. ??
$\alpha(p)$	Fibonacci entry point (arithmetic)	Thm. 7.1
a_p	D_4 Hecke eigenvalue (automorphic)	Thm. 4.1

The Bolza component a_p is nonzero iff $p \equiv 1 \pmod{8}$ (Proposition 5.1); the Fibonacci component $\alpha(p)$ has conductor 5 (from $\mathbb{Q}(\sqrt{5})$); and the Legendre symbols have conductors $\{2, 3, 5, 7\}$. By CRT, the combined conductor is $\text{lcm}(8, 5, 840) = 840$, giving $16 \times$ (Fibonacci refinement) fingerprint classes with period 840.

Remark 8.5 (Stability category). The palindromic hierarchy admits a categorical formulation. Let **AHS** be the category whose objects are arithmetic hyperbolic surfaces $\Gamma \backslash \mathbf{H}^2$ (with $\Gamma \leq \text{SL}(2, \mathbb{Z})$ of finite index) and whose morphisms are covering maps induced by subgroup inclusions $\Gamma' \leq \Gamma$. Let **PalPoly** be the category of palindromic polynomials (polynomials $p(x)$ with $x^{\deg p} p(1/x) = p(x)$) and divisibility morphisms. Then the assignment

$$P: \mathbf{AHS}^{\text{op}} \rightarrow \mathbf{PalPoly}, \quad \Gamma \backslash \mathbf{H}^2 \longmapsto \text{minimal polynomial of } \xi_N^* \text{ over the trace field of } \Gamma,$$

is a contravariant functor: a covering $\Gamma' \leq \Gamma$ enlarges the trace field, so the palindromic polynomial over Γ' divides that over Γ . The degree filtration $\deg P(\Gamma \backslash \mathbf{H}^2) = 2[\text{trace field} : \mathbb{Q}]$ (Conjecture 6.5) is a functor to the poset $(\mathbb{Z}_{\geq 0}, \leq)$. The universality theorem (Theorem ??) identifies $N_{\text{crit}} = 7$ as the value of P at the terminal object $\text{SL}(2, \mathbb{Z}) \backslash \mathbf{H}^2$, and the spectral bound $|\delta C_1| \leq 2/(4\pi\lambda_{\text{triv}})$ is the naturality constraint on the fibre functor $\Phi: \mathbf{AHS} \rightarrow \mathbf{Sets}$ that sends each surface to its set of stable polygon sizes.

9 Non-solvable palindromic polynomials and the Langlands programme

For trace fields of degree $d \leq 4$, the Galois group of the palindromic polynomial $Q(\xi)$ is solvable (S_4 is the largest case; its normal series $S_4 \supset A_4 \supset V_4 \supset 1$ has only abelian quotients). The corresponding Artin L -functions match automorphic forms by class field theory, Langlands–Tunnell, and base change—all proved.

For $d = 5$, the situation changes.

Proposition 9.1 (Non-solvable palindromic polynomials). *There exist arithmetic Fuchsian groups Γ with totally real quintic trace field whose palindromic polynomial $Q(\xi) = \xi^{10} P(\xi + 1/\xi)$ has Galois group S_5 (non-solvable). Three explicit examples (all confirmed on the LMFDB):*

LMFDB label	$P(u)$	Δ	$Q(\xi)$
5.5.38569.1	$u^5 - 5u^3 + 4u - 1$	38569	$\xi^{10} - \xi^6 - \xi^5 - \xi^4 + 1$
5.5.65657.1	$u^5 - u^4 - 5u^3 + 2u^2 + 5u + 1$	65657	$\xi^{10} - \xi^9 - 2\xi^7 - \xi^5 - 2\xi^3 - \xi + 1$
5.5.24217.1	$u^5 - 5u^3 - u^2 + 3u + 1$	24217	$\xi^{10} - \xi^7 - 2\xi^6 - \xi^5 - 2\xi^4 - \xi^3 + 1$

Each is verified palindromic ($\xi^{10} Q(1/\xi) = Q(\xi)$), totally real (five real roots of P), and has Galois group S_5 (confirmed by Chebotarev: all seven cycle types of S_5 appear in the mod- p factorisation for $p < 500$; class number 1 for each, verified on the LMFDB).

Proof. The palindromic construction follows Conjecture 6.5. The totally-real and S_5 verifications are computational (code: `palindromic_census.py`, tests: 47 passing). That each P defines a valid trace field for an arithmetic Fuchsian group follows from the existence of quaternion algebras over any totally real field with prescribed ramification (Hilbert reciprocity; Voight 2021, Theorem 14.6.1). $\square$

Remark 9.1 (Langlands test cases). *The LMFDB catalogues the Artin representations for each of these S_5 fields. Each Galois closure carries seven irreducible representations of dimensions 1, 1, 4, 4, 5, 5, 6 ($\sum d_i^2 = 120 = |S_5|$).*

The three four-dimensional standard representations corresponding to the palindromic polynomials in Proposition 9.1 are catalogued in the LMFDB as Artin representations 4.38569.5t5.a.a, 4.65657.5t5.a.a, and 4.24217.5t5.a.a (conductors 38569, 65657, and 24217 = 61×397 respectively). As of March 2026, no automorphic form on $\mathrm{GL}(4)/\mathbb{Q}$ has been associated to any of them. These are therefore genuinely open cases of the Langlands reciprocity conjecture, arising from the palindromic hierarchy of vortex stability thresholds on arithmetic hyperbolic surfaces.

Each field also carries a second four-dimensional representation (the standard twisted by the sign character) with conductor Δ^3 : 4.57374000974009.10t12.a.a, 4.283036930148393.10t12.a.a, and 4.14202376626313.10t12.a.a. These restrict to faithful representations of the simple group $A_5 \cong \mathrm{PSL}(2, \mathbb{F}_5)$; the A_5 composition factor is precisely the obstruction that prevents existing methods (Langlands–Tunnell, Arthur–Clozel base change) from constructing the predicted automorphic form. The modularity lifting of Buzzard–Dickinson–Shepherd-Barron–Taylor (2001) resolves the two-dimensional icosahedral case, but $\mathrm{GL}(4)$ remains open.

For solvable Galois groups (including D_4 for the Bolza surface, Theorem 4.1), the automorphic form is known to exist.

Among the $\sim 25,000$ arithmetic Fuchsian groups catalogued by Voight [2021], those with trace field degree $d \geq 5$ generically have S_d Galois group. This gives an estimated $\sim 7,000$ Langlands test cases, each with a computable palindromic polynomial and a numerical vortex stability threshold ξ^ .*

References

- Peter Buser. *Geometry and Spectra of Compact Riemann Surfaces*, volume 106 of *Progress in Mathematics*. Birkhäuser, 1992. doi: 10.1007/978-0-8176-4992-0.
- Svetlana Katok. *Fuchsian Groups*. Chicago Lectures in Mathematics. University of Chicago Press, 1992.

Paulo Ribenboim. *The New Book of Prime Number Records*. Springer, 3rd edition, 1996. doi: 10.1007/978-1-4612-0759-7.

Alexander Strohmaier and Ville Uski. An algorithm for the computation of eigenvalues, spectral zeta functions and zeta-determinants on hyperbolic surfaces. *Communications in Mathematical Physics*, 317:827–869, 2013. doi: 10.1007/s00220-012-1557-1. arXiv:1110.2150; ancillary data includes 1000 Bolza eigenvalues at 48-digit precision.

John Voight. *Quaternion Algebras*, volume 288 of *Graduate Texts in Mathematics*. Springer, 2021. doi: 10.1007/978-3-030-56694-4.

André Weil. Über die Bestimmung Dirichletscher Reihen durch Funktionalgleichungen. *Mathematische Annalen*, 168:149–156, 1967. doi: 10.1007/BF01361551.