A Pre-registered Test of the Transiting Planet Candidate TIC 402710280.01 at Its Predicted Transits of 2027

Rishit Vora
Independent Researcher
ORCID 0009-0003-8778-9883 rishitv@protonmail.com

Registered before any data of the tested epochs exist

1. Purpose

The preprint *A 36.34-day Sub-Neptune Candidate Transiting the Nearby K Dwarf TIC 402710280, Identified in TESS Full-Frame Images* reports a transit signal with period $P = 36.33986 \pm 0.00051$ d, detected in four transits recorded between 2021 and 2026. Because the period and transit shape were selected after those data were inspected, the archival data cannot test them independently. This document fixes, before the data exist, the predictions for the next observable transits, the data to be used, the processing, and the rules that map the result onto one of four outcomes. The analysis code (`prereg_test.py`) is frozen; its SHA-256 checksum is listed in `SHA256SUMS.txt`.

2. Predictions

The linear ephemeris of the preprint gives the mid-time of epoch E (counted from the Sector 39 transit) as

$$T_c(E) = T_0 + (E - 40)\, P, \qquad T_0 = 2460838.20080 \ \mathrm{BJD_{TDB}},$$

with uncertainty

$$\sigma_{\mathrm{pred}}(E) = \left[\sigma_{T_0}^2 + (E-40)^2 \sigma_P^2 + \sigma_{\mathrm{rms}}^2\right]^{1/2},$$

where $\sigma_{T_0} = 0.01192$ d, $\sigma_P = 0.00051$ d, and $\sigma_{\mathrm{rms}} = 21.8$ min is the rms of the four archival timing residuals, included because individual transits scatter about the linear ephemeris by that amount. The registered tests are:

Epoch	T_c ($\mathrm{BJD_{TDB}}$)	σ_{pred} (min)	Expected data
57	2461455.9785	30.4	TESS Sector 114 (2027 February 19)
59	2461528.6582	31.0	Ground-based photometry from Chile (2027 May 3, 03:41 UTC at La Silla

The expected depth at mid-transit is 2092 ppm and the expected duration is 5.27 h. Any later transit may be tested with the same code and rules; the result for each epoch is reported separately.

3. Data

For epoch 57, the data are, in order of preference: SPOC two-minute light curves (if the star is observed at that cadence), TESS-SPOC or QLP full-frame light curves, or a light curve extracted from the calibrated full-frame images with the code of the preprint's data record (`ffi_new.py`), without modification. For epoch 59, any ground-based light curve whose coverage satisfies the rule below. Cadences flagged by the data provider are removed before the test. The source used is reported with the outcome; no other source is substituted after a result has been obtained.

4. Processing

`prereg_test.py` reads the light curve, keeps cadences within ±0.8 d of the prediction, and normalises the flux by its median. The transit shape is fixed at the preprint's posterior medians ($R_p/R_\star = 0.04209$, $b = 0.224$, $a/R_\star = 53.63$, quadratic limb darkening $(0.393, 0.207)$), integrated over the exposure time. For each trial mid-time on a 0.5-minute grid spanning $\pm 3\sigma_{\rm pred}$, the code fits a quadratic baseline and a free transit amplitude by weighted least squares, with uniform weights set by the point-to-point scatter of the data outside the window. The best trial mid-time minimises χ^2; the detection statistic is $\Delta\chi^2 = \chi^2_{\rm baseline\ only} - \chi^2_{\rm best}$.

5. Outcomes

Recovered: $\Delta\chi^2 \geq 25$, a positive depth between 1046 and 3138 ppm (0.5–1.5 times the expected depth), and a best mid-time not at the edge of the $\pm 3\sigma_{\rm pred}$ grid.

Inconsistent: $\Delta\chi^2 \geq 25$ with a positive depth outside that range.

Excluded: no detection, the data cover at least 80% of the interval $T_c \pm (3\sigma_{\rm pred} + T_{14}/2)$, and at every trial mid-time the depth plus three times its uncertainty is below 1046 ppm.

Insufficient data: any other case.

An *excluded* result at epoch 57 or 59 would falsify the reported period. The measured mid-time and its offset from the prediction are reported in every case; an offset larger than 65 min ($3\sigma_{\rm rms}$) would independently confirm the timing scatter reported in the preprint.

6. Pixel-level check

If target pixel files are available for epoch 57, a difference image (out-of-transit minus in-transit) is formed and a point-spread function fitted to it, as in the preprint. The signal is *compatible with the target* if the fitted position lies within 3σ of the target's position measured on the same image, and is reported as such; this check does not change the photometric outcome.

7. Validation of the frozen code

Applied to the four archival transits of the data record, the code returns *recovered* for each (Sector 39: $O - C = +4.3$ min, depth 2203 ± 82 ppm; Sector 93: $O - C = -23.2$ min, depth 2028 ± 83 ppm;

Sector 100: $O - C = +6.9$ min, depth 2098 ± 98 ppm; Sector 102: $O - C = +35.5$ min, depth 2047 ± 87 ppm). Applied to transit-free data from Sector 101 shifted onto a predicted mid-time, it returns *excluded* (depths 86 ± 84 and -130 ± 90 ppm at two offsets), and *insufficient data* when the predicted transit falls in a data gap. Software: Python 3.13, numpy 2.5, batman 2.5.1.

8. Reporting

The outcome for each tested epoch, the data source, and the code output will be published as a new version of this record, regardless of the outcome.