Inline: Pythagoras $a^2 + b^2 = c^2$, derivative $f'(x) = 2x$, sum $\sum_{k=1}^{n} k = \frac{n(n+1)}{2}$, relations $a \le b \ne c \in M, x \circ y \cup z$ in running text.

$$\sum_{k=1}^{n} k^2 = \frac{n(n+1)(2n+1)}{6}$$

$$\int_0^1 x^2 dx = \frac{1}{3}, \int_a^b f$$

$$\lim_{x \to 0} \frac{\sin x}{x} = 1, a \stackrel{\text{def}}{=} b$$

$$\vec{v} + \hat{x} + \bar{y} + \tilde{z} + \underline{w}, {f_1}''$$

$$x = \frac{-b \pm \sqrt{b^2 - 4ac}}{2a}$$

$$\sqrt[3]{x+1}, \binom{n}{k} = \frac{n!}{k!(n-k)!}, \alpha + \beta = \Gamma, \frac{\partial f}{\partial x} \quad \mathrm{d}x, \mathrm{e}^{\mathrm{i}\pi}$$

$$\sqrt{\frac{\frac{a}{b}}{\frac{c}{d}}} + \sqrt{\frac{\frac{\frac{a}{b}}{b}}{\frac{c}{d}}} + \overrightarrow{abc} + \overline{a+b+c+d} + \widehat{xyz}$$

$$\left[\frac{\frac{a}{b}}{c}\right] + \left(a + \frac{b}{c}\right)^2 + \left\{x \middle| \frac{x}{2} > 0\right\} + (\frac{a}{b})$$

Inline with mstyle: $\frac{1}{2} + \frac{1}{2} + x = 1$ and a tall fraction $\frac{\frac{a}{b}}{c}$ in a paragraph with normal leading, followed by more text so that the line breaks and the next line is visible.
Second font family is ignored: x
Text in formulas: $f(x) = 0$for all$x \in M$ and $v = 3$km/h use the paragraph font (also **bold**, *italic*), in scripts and fractions the smaller size.

$$x_{\max} = \frac{\text{price}}{\text{quantity}}, \text{spaces inside}, \text{Tür}$$