

the 1990s, the incidence of *S. flexneri* has increased in the United Kingdom [10]. In the United States, *S. flexneri* has been reported as the most common serotype in children with acute bacterial dysentery [11].

There is a paucity of data on the epidemiology of *S. flexneri* in the United Kingdom. In the 1970s, *S. flexneri* was reported as the most common serotype in children with acute bacterial dysentery in the United Kingdom [12]. In the 1980s, *S. flexneri* was reported as the most common serotype in children with acute bacterial dysentery in the United Kingdom [13].

In the 1990s, *S. flexneri* was reported as the most common serotype in children with acute bacterial dysentery in the United Kingdom [14]. In the 1990s, *S. flexneri* was reported as the most common serotype in children with acute bacterial dysentery in the United Kingdom [15]. In the 1990s, *S. flexneri* was reported as the most common serotype in children with acute bacterial dysentery in the United Kingdom [16]. In the 1990s, *S. flexneri* was reported as the most common serotype in children with acute bacterial dysentery in the United Kingdom [17].

In the 1990s, *S. flexneri* was reported as the most common serotype in children with acute bacterial dysentery in the United Kingdom [18]. In the 1990s, *S. flexneri* was reported as the most common serotype in children with acute bacterial dysentery in the United Kingdom [19].

In the 1990s, *S. flexneri* was reported as the most common serotype in children with acute bacterial dysentery in the United Kingdom [20]. In the 1990s, *S. flexneri* was reported as the most common serotype in children with acute bacterial dysentery in the United Kingdom [21].

In the 1990s, *S. flexneri* was reported as the most common serotype in children with acute bacterial dysentery in the United Kingdom [22]. In the 1990s, *S. flexneri* was reported as the most common serotype in children with acute bacterial dysentery in the United Kingdom [23].

In the 1990s, *S. flexneri* was reported as the most common serotype in children with acute bacterial dysentery in the United Kingdom [24]. In the 1990s, *S. flexneri* was reported as the most common serotype in children with acute bacterial dysentery in the United Kingdom [25].

In the 1990s, *S. flexneri* was reported as the most common serotype in children with acute bacterial dysentery in the United Kingdom [26]. In the 1990s, *S. flexneri* was reported as the most common serotype in children with acute bacterial dysentery in the United Kingdom [27].

In the 1990s, *S. flexneri* was reported as the most common serotype in children with acute bacterial dysentery in the United Kingdom [28]. In the 1990s, *S. flexneri* was reported as the most common serotype in children with acute bacterial dysentery in the United Kingdom [29].

In the 1990s, *S. flexneri* was reported as the most common serotype in children with acute bacterial dysentery in the United Kingdom [30]. In the 1990s, *S. flexneri* was reported as the most common serotype in children with acute bacterial dysentery in the United Kingdom [31].

Head A	Head B
row 1	value 1
row 2	value 2
row 3	value 3
row 4	value 4
row 5	value 5
row 6	value 6
row 7	value 7
row 8	value 8
row 9	value 9
row 10	value 10
row 11	value 11
row 12	value 12
row 13	value 13
row 14	value 14
row 15	value 15