



Master in Physics at Aix-Marseille University,
Faculty of Science

Light pollution due to artificial light sources

Santiago Peña Martínez

santiago.pena-martinez@etu.univ-amu.fr

- 1 Introduction
- 2 Problem and Motivation
- 3 Numerical analysis
- 4 Initial calculations
- 5 Summary
- 6 Bibliography

KM3NeT as a neutrino telescope

New research infrastructure

- Discovery and subsequent observation high-energy neutrino sources in the universe



KM3NeT as a neutrino telescope

New research infrastructure

- Discovery and subsequent observation high-energy neutrino sources in the universe
- Measurement of neutrino mass hierarchy





KM3NeT as a neutrino telescope

New research infrastructure



- Discovery and subsequent observation high-energy neutrino sources in the universe
- Measurement of neutrino mass hierarchy
- Joint research with earth and sea sciences



Detection mechanism




Detection mechanism




Cherenkov radiation


Electromagnetic radiation emitted when a charged particle passes through a dielectric medium at a speed greater than the phase velocity of light in that medium.



Design



Several unknowns




Measuring the artificial light sources

A conversion between lumen and photons is needed.

Measuring the artificial light sources

A conversion between lumen and photons is needed.



Conversion factor

1 lumen = 4.11×10^{15} photons
at $\lambda = 555nm$

Water parameters

From Rayleigh scattering theory we have

$$L_{sca}(\lambda) = \left(\frac{\lambda}{550nm} \right)^{4.32} \quad (1)$$


$$\frac{dp_{sca}}{d\Omega}(\beta) = \frac{1}{4\pi} \frac{3}{3+b} (1 + b \cos^2 \beta) \quad (2)$$

Water parameters


From Rayleigh scattering theory we have

$$L_{sca}(\lambda) = \left(\frac{\lambda}{550nm} \right)^{4.32} \quad (1)$$

$$\frac{dp_{sca}}{d\Omega}(\beta) = \frac{1}{4\pi} \frac{3}{3+b} (1 + b \cos^2 \beta) \quad (2)$$




Water absorption length in
salt water.




Water molecular Rayleigh
scattering length.

PMT parameters




PMT quantum efficiency including glass and gel.




PMT angular acceptance.

PMT parameters




PMT quantum efficiency including glass and gel.




PMT angular acceptance.

Values taken from the official KM3NeT simulations.

Different setups



Different setups




The corresponding scattering differential probability and absorption probability are given by:

$$\frac{dP_{sca}}{ds}(\lambda, s) = \frac{e^{-s/L_{sca}(\lambda)}}{L_{sca}(\lambda)}, \quad (3)$$



$$P_{abs}(\lambda, s+x) = e^{-(s+x)/L_{abs}(\lambda)}, \quad (4)$$

Setup a)




$$R(\lambda, d, \delta) = \Phi_0(\lambda) QE_{PMT}(\lambda) \int_0^{\infty \approx 500} ds P_{abs}(\lambda, s + x) \frac{dP_{sca}}{ds}(\lambda, s) P_{PMT}(\alpha) \frac{dp_{sca}}{d\Omega}(\beta) \frac{A_{PMT}}{x^2} \quad (5)$$

Setup a)





$$R(\lambda, d, \delta) = \Phi_0(\lambda) Q E_{PMT}(\lambda) \int_0^{\infty \approx 500} ds P_{abs}(\lambda, s + x) \frac{dP_{sca}}{ds}(\lambda, s) P_{PMT}(\alpha) \frac{dp_{sca}}{d\Omega}(\beta) \frac{A_{PMT}}{x^2} \quad (5)$$

Setup b)



$$R(\lambda, d, \delta) = \Phi_0(\lambda) Q E_{PMT}(\lambda) \int_0^{\infty \approx 500} \int_0^{\pi/2} ds d\delta P_{abs}(\lambda, s+x) \frac{dP_{sca}}{ds}(\lambda, s) P_{PMT}(\alpha) \frac{dp_{sca}}{d\Omega}(\beta) \frac{A_{PMT}}{x^2} \quad (6)$$

Setup b)



$$R(\lambda, d, \delta) = \Phi_0(\lambda) Q E_{PMT}(\lambda) \int_0^{\infty \approx 500} \int_0^{\pi/2} ds d\delta P_{abs}(\lambda, s+x) \frac{dP_{sca}}{ds}(\lambda, s) P_{PMT}(\alpha) \frac{dp_{sca}}{d\Omega}(\beta) \frac{A_{PMT}}{x^2} \quad (6)$$

Setup c)


c)



$$R(\lambda, d) = \Phi_0(\lambda)QE_{PMT}(\lambda)P_{abs}(\lambda, d)P_{PMT}(0)\frac{A_{PMT}}{4\pi d^2}. \quad (7)$$

Setup c)

c)





$$R(\lambda, d) = \Phi_0(\lambda)QE_{PMT}(\lambda)P_{abs}(\lambda, d)P_{PMT}(0)\frac{A_{PMT}}{4\pi d^2}. \quad (7)$$

Lower limits wrap up

Background noise

Natural bioluminescence and natural radiation from ^{40}K is $\sim 6 - 7\text{kHz}$



Next steps

- Multiple scattering.
- Mie theory for particle scattering.
- Montecarlo simluations for single photons.

Bibliography I

- Silvia Adrian-Martinez et al. (2016). "Letter of intent for KM3NeT 2.0". In: *Journal of Physics G: Nuclear and Particle Physics* 43.8, p. 084001.
- JA Aguilar et al. (2005). "Transmission of light in deep sea water at the site of the ANTARES neutrino telescope". In: *Astroparticle Physics* 23.1, pp. 131–155.
- Hendrik Christoffel Hulst and Hendrik C van de Hulst (1981). *Light scattering by small particles*. Courier Corporation.
- Andre Morel and Hubert Loisel (1998). "Apparent optical properties of oceanic water: dependence on the molecular scattering contribution". In: *Applied Optics* 37.21, pp. 4765–4776.
- Theodore J Petzold (1972). *Volume scattering functions for selected ocean waters*. Tech. rep. Scripps Institution of Oceanography La Jolla Ca Visibility Lab.



**Thank you for
your attention**