

Homework 1: Machine numbers; Interpolation

Due Apr. 12th.

1. Floating point representation

For the `float` data type, write a program to **empirically** determine the following “Machine constants” for your computer:

- (a) The smallest ϵ such that $1.0 - \epsilon \neq 1.0$
- (b) The smallest ϵ such that $1.0 + \epsilon \neq 1.0$
- (c) The maximum representable number
- (d) The minimum representable positive number

Comment on why the numbers you get are expected based on the IEEE 754 representation.

2. Roundoff error

Numerically evaluate the expression $(1 - \cos(x))/x^2$ in double precision for values of x around 10^{-7} and smaller. Explain the difference between the numerical results and the analytic limit as $x \rightarrow 0$.

3. Interpolation

- (a) Write a program to read in a two column table from a file and perform linear interpolation at an arbitrary point. You may assume that the data is evenly spaced in the independent variable.
- (b) Use the program on the following input data: (available on the web site)

$x \quad y$

1	100
2	25
3	11.111111
4	6.25
5	4

and provide an estimate of y at $x = 4.75$.

- (c) Write a program using Neville’s algorithm to fit a 4th order polynomial to the above data and provide an estimate of y at $x = 4.75$.
- (d) The actual function tabulated above is $y = \frac{100}{x^2}$. Compare the actual value at $x = 4.75$ with the linear interpolation and the 4th order polynomial interpolation, and comment on why one is more accurate than the other.