

Homework 2: Ordinary differential equations

Due April 26.

1. Write a program (or programs) to integrate up to four sets of coupled differential equations using the Euler method, fourth order Runge-Kutta, and Leapfrog. (Note: Leapfrog only applies to special cases.) For Runge-Kutta, you may use a packaged routine such as available in `scipy` or Numerical Recipes. If you do use a packaged routine, **be sure to use one with a fixed timestep and order so that testing the convergence can be easily performed.**
2. Use your program to solve the differential equation for $x(t)$:

$$\frac{d^2x}{dt^2} + x = 0;$$

with the initial conditions $x(0) = 1, x'(0) = 0$. Note that this has the analytical solution: $x = \cos(t)$

- (a) Integrate the equation for $0 \leq t \leq 30$ using each of the methods, and step sizes of 1, .3, .1, .03, and .01. Comment on the behavior of the solutions.
 - (b) Plot $\log(|x_{\text{numerical}}(30) - x_{\text{exact}}(30)|)$ as a function of $\log(\text{stepsize})$ and check for the expected convergence of the error term.
3. Now try the two dimensional orbit described by the potential:

$$\Phi = -\frac{1}{\sqrt{1+x^2+y^2}}.$$

The orbits are given by the coupled differential equations:

$$\begin{aligned}\frac{d^2x}{dt^2} &= -\frac{x}{(1+x^2+y^2)^{3/2}}, \\ \frac{d^2y}{dt^2} &= -\frac{y}{(1+x^2+y^2)^{3/2}}.\end{aligned}$$

- (a) Integrate this for $0 \leq t \leq 100$ for the initial conditions $x = 1, y = 0, x' = 0, y' = .3$. Try this with either Leapfrog or Runge-Kutta and step sizes from .01 to 1. Plot x vs. y for these integrations.
- (b) Plot the energy $E = (x'^2 + y'^2)/2 + \Phi(x, y)$ as a function of time for your integrations.