

An Intro to CGA

Conformal Geometric Algebra

Spencer T. Parkin

Avalanche Software

September 10, 2012

Presentation Outline

In this presentation, we will...

Presentation Outline

In this presentation, we will...

- Introduce concepts from GA only as necessary,

Presentation Outline

In this presentation, we will...

- Introduce concepts from GA only as necessary,
- Introduce the generalized homogeneous model of geometry over GA,

Presentation Outline

In this presentation, we will...

- Introduce concepts from GA only as necessary,
- Introduce the generalized homogeneous model of geometry over GA,
- Define the specific conformal model of GA,

Presentation Outline

In this presentation, we will...

- Introduce concepts from GA only as necessary,
- Introduce the generalized homogeneous model of geometry over GA,
- Define the specific conformal model of GA,
- Find forms for all geometric primitives of the CGA model,

Presentation Outline

In this presentation, we will...

- Introduce concepts from GA only as necessary,
- Introduce the generalized homogeneous model of geometry over GA,
- Define the specific conformal model of GA,
- Find forms for all geometric primitives of the CGA model,
- Discuss the fundamental transformations of CGA.

Let \mathbb{V}^n denote an n -dimensional vector space.

Let \mathbb{V}^n denote an n -dimensional vector space.

Let $\{b_k\}_{k=1}^m$ be a set of m vectors taken from \mathbb{V}^n .

Let \mathbb{V}^n denote an n -dimensional vector space.

Let $\{b_k\}_{k=1}^m$ be a set of m vectors taken from \mathbb{V}^n .

Definition

We say the blade B , given by

$$B = \bigwedge_{k=1}^m b_k = b_1 \wedge \cdots \wedge b_m,$$

is a non-zero m -blade if and only if $\{b_k\}_{k=1}^m$ is a linearly independent set of vectors.

Let \mathbb{V}^n denote an n -dimensional vector space.

Let $\{b_k\}_{k=1}^m$ be a set of m vectors taken from \mathbb{V}^n .

Definition

We say the blade B , given by

$$B = \bigwedge_{k=1}^m b_k = b_1 \wedge \cdots \wedge b_m,$$

is a non-zero m -blade if and only if $\{b_k\}_{k=1}^m$ is a linearly independent set of vectors.

Clearly, if $B \neq 0$, then we must have $\text{grade}(B) = m \leq n$.

Visualizing Euclidean Blades

Imagine an **infinite** m -dimensional hyper-plane.

Visualizing Euclidean Blades

Imagine an **infinite** m -dimensional hyper-plane.

Think of B as a **finite** m -dimensional hyper-plane.

Visualizing Euclidean Blades

Imagine an **infinite** m -dimensional hyper-plane.

Think of B as a **finite** m -dimensional hyper-plane.

Attitude matters, **position** does not.

Visualizing Euclidean Blades

Imagine an **infinite** m -dimensional hyper-plane.

Think of B as a **finite** m -dimensional hyper-plane.

Attitude matters, **position** does not.

Non-Euclidean blades require more imagination!

Visualizing Euclidean Blades

Imagine an **infinite** m -dimensional hyper-plane.

Think of B as a **finite** m -dimensional hyper-plane.

Attitude matters, **position** does not.

Non-Euclidean blades require more imagination!

Our geometric arguments will not require us to visualize the homogeneous representation space.

Building Intuition About Euclidean Blades

Let $v_{||}$ denote the orthogonal **projection** of v down onto B .

Building Intuition About Euclidean Blades

Let v_{\parallel} denote the orthogonal **projection** of v down onto B .

Let $v_{\perp} = v - v_{\parallel}$ denote the orthogonal **rejection** of v from B .

Building Intuition About Euclidean Blades

Let v_{\parallel} denote the orthogonal **projection** of v down onto B .

Let $v_{\perp} = v - v_{\parallel}$ denote the orthogonal **rejection** of v from B .

For any vector $v \in \mathbb{V}^n$, we have

$$v \wedge B = (v_{\parallel} + v_{\perp}) \wedge B = v_{\perp} \wedge B,$$

$$v \cdot B = (v_{\parallel} + v_{\perp}) \cdot B = v_{\parallel} \cdot B.$$

Building Intuition About Euclidean Blades

Let v_{\parallel} denote the orthogonal **projection** of v down onto B .

Let $v_{\perp} = v - v_{\parallel}$ denote the orthogonal **rejection** of v from B .

For any vector $v \in \mathbb{V}^n$, we have

$$v \wedge B = (v_{\parallel} + v_{\perp}) \wedge B = v_{\perp} \wedge B,$$

$$v \cdot B = (v_{\parallel} + v_{\perp}) \cdot B = v_{\parallel} \cdot B.$$

$$\text{grade}(v \wedge B) = \text{grade}(B) + 1$$

$$\text{grade}(v \cdot B) = \text{grade}(B) - 1$$

Building Intuition About Euclidean Blades

Let v_{\parallel} denote the orthogonal **projection** of v down onto B .

Let $v_{\perp} = v - v_{\parallel}$ denote the orthogonal **rejection** of v from B .

For any vector $v \in \mathbb{V}^n$, we have

$$v \wedge B = (v_{\parallel} + v_{\perp}) \wedge B = v_{\perp} \wedge B,$$

$$v \cdot B = (v_{\parallel} + v_{\perp}) \cdot B = v_{\parallel} \cdot B.$$

$$\text{grade}(v \wedge B) = \text{grade}(B) + 1$$

$$\text{grade}(v \cdot B) = \text{grade}(B) - 1$$

We may also imagine $v \cdot B = (v \wedge B^*)^*$, where B^* is the **complement** of B with respect to \mathbb{V}^n .

Blades May Represent Vector Sub-Spaces

Recall that $B = b_1 \wedge \cdots \wedge b_m$.

Blades May Represent Vector Sub-Spaces

Recall that $B = b_1 \wedge \cdots \wedge b_m$.

Definition

For any $v \in \mathbb{V}^n$, we say that

$$v \in B \text{ if and only if } v \in \text{span}\{b_k\}_{k=1}^m.$$

Blades May Represent Vector Sub-Spaces

Recall that $B = b_1 \wedge \cdots \wedge b_m$.

Definition

For any $v \in \mathbb{V}^n$, we say that

$$v \in B \text{ if and only if } v \in \text{span}\{b_k\}_{k=1}^m.$$

Definition

If $v \notin B$, then $v \in B^*$, which represents the complement $(\mathbb{V}^n - \text{span}\{b_k\}_{k=1}^m) \cup \{0\}$.

Membership in Vector Spaces and Dual Vector Spaces

If $B \neq 0$, then $v \in B$ if and only if $v \wedge B = 0$.

Membership in Vector Spaces and Dual Vector Spaces

If $B \neq 0$, then $v \in B$ if and only if $v \wedge B = 0$.

Proof.

The set $\{b_k\}_{k=1}^m$ is linearly independent while the set $\{v\} \cup \{b_k\}_{k=1}^m$ is linearly dependent. □

Membership in Vector Spaces and Dual Vector Spaces

If $B \neq 0$, then $v \in B$ if and only if $v \wedge B = 0$.

Proof.

The set $\{b_k\}_{k=1}^m$ is linearly independent while the set $\{v\} \cup \{b_k\}_{k=1}^m$ is linearly dependent. □

If $B \neq 0$, then $v \in B^*$ if and only if $v \cdot B = 0$.

Membership in Vector Spaces and Dual Vector Spaces

If $B \neq 0$, then $v \in B$ if and only if $v \wedge B = 0$.

Proof.

The set $\{b_k\}_{k=1}^m$ is linearly independent while the set $\{v\} \cup \{b_k\}_{k=1}^m$ is linearly dependent. □

If $B \neq 0$, then $v \in B^*$ if and only if $v \cdot B = 0$.

Proof.

Notice that $0 = v \cdot B = (v \wedge B^*)^*$ if and only if $v \wedge B^* = 0$. □

Blades May Represent Geometries

Let \mathbb{R}^n denote n -dimensional Euclidean space.

Blades May Represent Geometries

Let \mathbb{R}^n denote n -dimensional Euclidean space.

Let $p : \mathbb{R}^n \rightarrow \mathbb{G}(\mathbb{V}^n)$ be a vector-valued function of a Euclidean point.

Blades May Represent Geometries

Let \mathbb{R}^n denote n -dimensional Euclidean space.

Let $p : \mathbb{R}^n \rightarrow \mathbb{G}(\mathbb{V}^n)$ be a vector-valued function of a Euclidean point.

Definition

We say that B **directly** represents a geometry as the set of all points

$$G(B) = \{x \in \mathbb{R}^n | p(x) \in B\}.$$

Blades May Represent Geometries

Let \mathbb{R}^n denote n -dimensional Euclidean space.

Let $p : \mathbb{R}^n \rightarrow \mathbb{G}(\mathbb{V}^n)$ be a vector-valued function of a Euclidean point.

Definition

We say that B **directly** represents a geometry as the set of all points

$$G(B) = \{x \in \mathbb{R}^n | p(x) \in B\}.$$

Definition

We say that B **dually** represents a geometry as the set of all points

$$G^*(B) = \{x \in \mathbb{R}^n | p(x) \in B^*\}.$$

Blades May Represent Geometries

Let \mathbb{R}^n denote n -dimensional Euclidean space.

Let $p : \mathbb{R}^n \rightarrow \mathbb{G}(\mathbb{V}^n)$ be a vector-valued function of a Euclidean point.

Definition

We say that B **directly** represents a geometry as the set of all points

$$G(B) = \{x \in \mathbb{R}^n | p(x) \in B\}.$$

Definition

We say that B **dually** represents a geometry as the set of all points

$$G^*(B) = \{x \in \mathbb{R}^n | p(x) \in B^*\}.$$

Note that $G(B) = G^*(B^*)$ and $G^*(B) = G(B^*)$.

We Can Combine Geometries

For any two blades $A, B \in \mathbb{G}(\mathbb{V}^n)$ such that $A \wedge B \neq 0$, we have

$$G(A) \cup G(B) \subseteq G(A \wedge B).$$

We Can Combine Geometries

For any two blades $A, B \in \mathbb{G}(\mathbb{V}^n)$ such that $A \wedge B \neq 0$, we have

$$G(A) \cup G(B) \subseteq G(A \wedge B).$$

Proof.

$$\begin{aligned} p(x) \in A \text{ or } p(x) \in B \\ \implies p(x) \in A \wedge B \end{aligned}$$



We Can Combine Geometries

For any two blades $A, B \in \mathbb{G}(\mathbb{V}^n)$ such that $A \wedge B \neq 0$, we have

$$G(A) \cup G(B) \subseteq G(A \wedge B).$$

Proof.

$$\begin{aligned} p(x) \in A \text{ or } p(x) \in B \\ \implies p(x) \in A \wedge B \end{aligned}$$



Let $C \subseteq A \wedge B$ represent the smallest vector sub-space such that $p(x) \in C$. Then we might have $C \not\subseteq A$ and $C \not\subseteq B$.

We Can Intersect Geometries

For any two blades $A, B \in \mathbb{G}(\mathbb{V}^n)$ such that $A \wedge B \neq 0$, we have

$$G^*(A) \cap G^*(B) = G^*(A \wedge B).$$

We Can Intersect Geometries

For any two blades $A, B \in \mathbb{G}(\mathbb{V}^n)$ such that $A \wedge B \neq 0$, we have

$$G^*(A) \cap G^*(B) = G^*(A \wedge B).$$

Proof.

$$\begin{aligned} & p(x) \in A^* \text{ and } p(x) \in B^* \\ \text{iff } & p(x) \notin A \text{ and } p(x) \notin B \\ \text{iff } & p(x) \notin A \wedge B \\ \text{iff } & p(x) \in (A \wedge B)^* \end{aligned}$$



The Homogeneous Nature Of The Model

For any non-zero scalar λ , we have $G(B) = G(\lambda B)$.

The Homogeneous Nature Of The Model

For any non-zero scalar λ , we have $G(B) = G(\lambda B)$.

For any blade B , there is a scalar λ such that λB is a homogenized form.

The Homogeneous Nature Of The Model

For any non-zero scalar λ , we have $G(B) = G(\lambda B)$.

For any blade B , there is a scalar λ such that λB is a homogenized form.

If B is the result of some geometric operation, then such a λ has geometric significance WRT to that operation.

The Geometric Product

Definition

For any vector $v \in \mathbb{V}^n$ and any blade $B \in \mathbb{G}(\mathbb{V}^n)$, we define

$$vB = v \cdot B + v \wedge B.$$

Let $\{v_k\}_{k=1}^m$ be any set of m vectors.

Let $\{v_k\}_{k=1}^m$ be any set of m vectors.

Definition

We say the element $V \in \mathbb{G}(\mathbb{V}^n)$, given by

$$V = \prod_{k=1}^m v_k,$$

is a versor if and only if for all k , the vector v_k^{-1} exists.

Let $\{v_k\}_{k=1}^m$ be any set of m vectors.

Definition

We say the element $V \in \mathbb{G}(\mathbb{V}^n)$, given by

$$V = \prod_{k=1}^m v_k,$$

is a versor if and only if for all k , the vector v_k^{-1} exists.

Notice that $v^{-1} = v/\sqrt{v^2} = v/|v|$.

The Inverse And The Reverse Of Versors

Definition

Given the versor $V = v_1 \dots v_m$, we define

$$\tilde{V} = \prod_{k=1}^m v_{m-k+1}.$$

The Inverse And The Reverse Of Versors

Definition

Given the versor $V = v_1 \dots v_m$, we define

$$\tilde{V} = \prod_{k=1}^m v_{m-k+1}.$$

The inverse V^{-1} of V is therefore given by

$$V^{-1} = \frac{\tilde{V}}{V\tilde{V}}.$$

Versors form a group under the geometric product.

The Versor Group

Versors form a group under the geometric product.

Proof.

Associativity follows from the associativity of the geometric product.

The scalar 1 is the **identity** versor.

For every versor V , there exists an **inverse** V^{-1} such that $VV^{-1} = V^{-1}V = 1$. □

Conjugation by versors is **outermorphic**!

Properties Of Versors

Conjugation by versors is **outermorphic**!

Recall that $B = b_1 \wedge \cdots \wedge b_m$. We then have

$$VBV^{-1} = \bigwedge_{k=1}^m Vb_k V^{-1}.$$

Properties Of Versors

Conjugation by versors is **outermorphic**!

Recall that $B = b_1 \wedge \cdots \wedge b_m$. We then have

$$VBV^{-1} = \bigwedge_{k=1}^m Vb_k V^{-1}.$$

Conjugation by versors is **grade preserving**!

Properties Of Versors

Conjugation by versors is **outermorphic**!

Recall that $B = b_1 \wedge \cdots \wedge b_m$. We then have

$$VBV^{-1} = \bigwedge_{k=1}^m Vb_k V^{-1}.$$

Conjugation by versors is **grade preserving**!

For any vector $v \in \mathbb{V}^n$, we have $VvV^{-1} \in \mathbb{V}^n$, therefore, we have $\text{grade}(B) = \text{grade}(VBV^{-1})$.

Versors May Represent Transformations

It follows that versors may be used to represent transformations of geometry as versors conjugated with blades representative of geometry.

Versors May Represent Transformations

It follows that versors may be used to represent transformations of geometry as versors conjugated with blades representative of geometry.

Given $G(B)$, it will be interesting to investigate $G(VBV^{-1})$.

The Specifics Of The Conformal Model

Replace \mathbb{R}^n with \mathbb{V}^n .

The Specifics Of The Conformal Model

Replace \mathbb{R}^n with \mathbb{V}^n .

Embed \mathbb{V}^n in \mathbb{V}^{n+2} as a **Euclidean** vector sub-space of a **non-Euclidean** vector space.

The Specifics Of The Conformal Model

Replace \mathbb{R}^n with \mathbb{V}^n .

Embed \mathbb{V}^n in \mathbb{V}^{n+2} as a **Euclidean** vector sub-space of a **non-Euclidean** vector space.

Let $o, \infty \in \mathbb{V}^{n+2}$ be vectors such that $o \cdot o = \infty \cdot \infty = 0$ and $o \cdot \infty = \infty \cdot o = -1$ and for all $v \in \mathbb{V}^n$, we have $v \cdot o = v \cdot \infty = 0$.

The Specifics Of The Conformal Model

Replace \mathbb{R}^n with \mathbb{V}^n .

Embed \mathbb{V}^n in \mathbb{V}^{n+2} as a **Euclidean** vector sub-space of a **non-Euclidean** vector space.

Let $o, \infty \in \mathbb{V}^{n+2}$ be vectors such that $o \cdot o = \infty \cdot \infty = 0$ and $o \cdot \infty = \infty \cdot o = -1$ and for all $v \in \mathbb{V}^n$, we have $v \cdot o = v \cdot \infty = 0$.

Definition

Define $p : \mathbb{V}^n \rightarrow \mathbb{G}(\mathbb{V}^{n+2})$ as

$$p(x) = o + x + \frac{1}{2}x^2\infty.$$

The Specifics Of The Conformal Model

Replace \mathbb{R}^n with \mathbb{V}^n .

Embed \mathbb{V}^n in \mathbb{V}^{n+2} as a **Euclidean** vector sub-space of a **non-Euclidean** vector space.

Let $o, \infty \in \mathbb{V}^{n+2}$ be vectors such that $o \cdot o = \infty \cdot \infty = 0$ and $o \cdot \infty = \infty \cdot o = -1$ and for all $v \in \mathbb{V}^n$, we have $v \cdot o = v \cdot \infty = 0$.

Definition

Define $p : \mathbb{V}^n \rightarrow \mathbb{G}(\mathbb{V}^{n+2})$ as

$$p(x) = o + x + \frac{1}{2}x^2\infty.$$

Having **invented** this specific model, what we are now able to **discover** about it is almost endless!

Points in n -dimensional Space

For any $c \in \mathbb{V}^n$, the vector $p(c)$ both **dually** and **directly** represents the point c in space.

Points in n -dimensional Space

For any $c \in \mathbb{V}^n$, the vector $p(c)$ both **dually** and **directly** represents the point c in space.

That is, $G(p(c)) = G^*(p(c)) = \{c\}$.

The function $p(x)$ factors out of the equation

$$(x - c)^2 - r^2 = 0$$

as the alternative equation

$$p(x) \cdot \left(p(c) - \frac{1}{2}r^2\infty \right) = 0.$$

The function $p(x)$ factors out of the equation

$$(x - c)^2 - r^2 = 0$$

as the alternative equation

$$p(x) \cdot \left(p(c) - \frac{1}{2}r^2\infty \right) = 0.$$

Points are degenerate spheres, or spheres with radius zero.

n -dimensional Dual Hyper-Spheres

The function $p(x)$ factors out of the equation

$$(x - c)^2 - r^2 = 0$$

as the alternative equation

$$p(x) \cdot \left(p(c) - \frac{1}{2}r^2\infty \right) = 0.$$

Points are degenerate spheres, or spheres with radius zero.
We may refer to $p(c)$ as a **round** point.

Generating All Dual Rounds Of CGA

Let $\{\sigma_k\}_{k=1}^m$ be m spheres of dimension n having a non-empty and non-degenerate intersection.

Generating All Dual Rounds Of CGA

Let $\{\sigma_k\}_{k=1}^m$ be m spheres of dimension n having a non-empty and non-degenerate intersection.

Then the blade B , given by

$$B = \bigwedge_{k=1}^m \sigma_k,$$

dually represents an $(n - m + 1)$ -dimensional hyper-sphere.

Generating All Dual Rounds Of CGA

Let $\{\sigma_k\}_{k=1}^m$ be m spheres of dimension n having a non-empty and non-degenerate intersection.

Then the blade B , given by

$$B = \bigwedge_{k=1}^m \sigma_k,$$

dually represents an $(n - m + 1)$ -dimensional hyper-sphere.

Rounds with zero radius give us tangent points!

All Rounds Of CGA For 3-dimensional Space




Figure : 3 Rounds, 3 Circles and 1 Point-Pair

$(n - 1)$ -dimensional Dual Hyper-Planes

The function $p(x)$ factors out of the equation

$$(x - c) \cdot v = 0$$

as the alternative equation

$$p(x) \cdot (v + (c \cdot v)\infty) = 0.$$

Generating All Dual Flats Of CGA

Let $\{\pi_k\}_{k=1}^m$ be m planes of dimension $n - 1$ having a **non-empty** and **non-degenerate** intersection.

Generating All Dual Flats Of CGA

Let $\{\pi_k\}_{k=1}^m$ be m planes of dimension $n - 1$ having a **non-empty** and **non-degenerate** intersection.

Then the blade B , given by

$$B = \bigwedge_{k=1}^m \pi_k,$$

dually represents an $(n - m)$ -dimensional hyper-plane.

Generating All Dual Flats Of CGA

Let $\{\pi_k\}_{k=1}^m$ be m planes of dimension $n - 1$ having a **non-empty** and **non-degenerate** intersection.

Then the blade B , given by

$$B = \bigwedge_{k=1}^m \pi_k,$$

dually represents an $(n - m)$ -dimensional hyper-plane.

Flats at infinity are **free blades**.

Generating All Dual Flats Of CGA

Let $\{\pi_k\}_{k=1}^m$ be m planes of dimension $n - 1$ having a **non-empty** and **non-degenerate** intersection.

Then the blade B , given by

$$B = \bigwedge_{k=1}^m \pi_k,$$

dually represents an $(n - m)$ -dimensional hyper-plane.

Flats at infinity are **free blades**.

0-dimensional **flats** are called **flat points**.

All Flats Of CGA For 3-dimensional Space




Figure : 3 Planes, 3 Lines, 1 Flat-Point

A Generalization Of Coplanarity

Definition

For $m \geq 0$, a set of $m + 2$ points $\{x_k\}_{k=1}^{m+2} \subset \mathbb{V}^n$ are **co- m -hyper-planar** if...

For $m = 0$, the points are identical,

For $m = 1$, the points are collinear,

For $m = 2$, the points are coplanar,

For $m = 3$, the points are co-hyper-planar,

etc...

A Condition For Linear Independents Of Points

Lemma

For $m \geq 1$, if $m + 1$ points $\{x_k\}_{k=1}^{m+1} \subset \mathbb{V}^n$ are non-co- $(m - 1)$ -hyper-planar, then $\{p(x_k)\}_{k=1}^{m+1}$ is a linearly independent set.

A Condition For Linear Independents Of Points

Lemma

For $m \geq 1$, if $m + 1$ points $\{x_k\}_{k=1}^{m+1} \subset \mathbb{V}^n$ are non-co- $(m - 1)$ -hyper-planar, then $\{p(x_k)\}_{k=1}^{m+1}$ is a linearly independent set.

The proof is not hard, but too big for this slide.

Generating All Direct Rounds Of CGA

Let $m \geq 1$. For $m + 1$ points $\{x_k\}_{k=1}^{m+1} \subset \mathbb{V}^n$ on an m -dimensional hyper-sphere that are non-co- $(m - 1)$ -hyper-planar, the blade B , given by

$$B = \bigwedge_{k=1}^{m+1} p(x_k)$$

directly represents the m -dimensional hyper-sphere.

Generating All Direct Rounds Of CGA

Let $m \geq 1$. For $m + 1$ points $\{x_k\}_{k=1}^{m+1} \subset \mathbb{V}^n$ on an m -dimensional hyper-sphere that are non-co- $(m - 1)$ -hyper-planar, the blade B , given by

$$B = \bigwedge_{k=1}^{m+1} p(x_k)$$

directly represents the m -dimensional hyper-sphere.

Proof.

Let the $(n - m + 1)$ -blade A **dually** represent the m -dimensional hyper-sphere determined by the points. If A **dually** represents this sphere, then A^* **directly** represents this sphere. Therefore, we need to show that there exists $\lambda \in \mathbb{R}$ such that $A^* = \lambda B$. For all k , we have $p(x_k) \in A^*$ and $p(x_k) \in B$. By our lemma, $\{p(x_k)\}_{k=1}^m$ is a linearly independent set. Lastly, $\text{grade}(B) = m + 1 = n + 2 - (n - m + 1) = n + 2 - \text{grade}(A) = \text{grade}(A^*)$. □

A Generalization Of Cospherical

Definition

For $m \geq 1$, a set of $m + 1$ points $\{x_k\}_{k=1}^{m+1} \subset \mathbb{V}^n$ are **co- m -hyper-planar** if...

For $m = 1$, the points are co-point-pair (distinct),

For $m = 2$, the points are co-circular,

For $m = 3$, the points are co-spherical,

For $m = 4$, the points are co-hyper-spherical,

etc...

Generating Almost All Direct Flats Of CGA

Let $m \geq 1$. For $m + 2$ points $\{x_k\}_{k=1}^{m+2} \subset \mathbb{V}^n$ on an m -dimensional hyper-plane that are (1) non-co- $(m - 1)$ -hyper-planar and (2) non-co- m -hyper-spherical, the blade B , given by

$$B = \bigwedge_{k=1}^{m+2} p(x_k),$$

directly represents the m -dimensional hyper-plane.

Generating Almost All Direct Flats Of CGA

Let $m \geq 1$. For $m + 2$ points $\{x_k\}_{k=1}^{m+2} \subset \mathbb{V}^n$ on an m -dimensional hyper-plane that are (1) non-co- $(m - 1)$ -hyper-planar and (2) non-co- m -hyper-spherical, the blade B , given by

$$B = \bigwedge_{k=1}^{m+2} p(x_k),$$

directly represents the m -dimensional hyper-plane.

Proof.

By (1), there exists the $(n - m)$ -blade A dually representative of the m -dimensional hyper-plane. By (2), $B \neq 0$. Lastly, $\text{grade}(B) = m + 2 = n + 2 - (n - m) = n + 2 - \text{grade}(A) = \text{grade}(A^*)$. \square

Generating All Direct Flats Of CGA

Let $m \geq 0$. For $m + 1$ points $\{x_k\}_{k=1}^{m+1} \subset \mathbb{V}^n$ on an m -dimensional hyepr-plane that are non-co- $(m - 1)$ -hyper-planar, the blade B , give by

$$B = \infty \wedge \bigwedge_{k=1}^{m+1} p(x_k),$$

directly represents the m -dimensional hyper-plane.

Generating All Direct Flats Of CGA

Let $m \geq 0$. For $m + 1$ points $\{x_k\}_{k=1}^{m+1} \subset \mathbb{V}^n$ on an m -dimensional hyper-plane that are non-co- $(m - 1)$ -hyper-planar, the blade B , give by

$$B = \infty \wedge \bigwedge_{k=1}^{m+1} p(x_k),$$

directly represents the m -dimensional hyper-plane.
The proof, again, is not hard, but can't fit here.

Quiz Time!

Question: Given a **dual** line L and a point P not on L , how do I find the **dual** plane N containing L and P ?

Quiz Time!

Question: Given a **dual** line L and a point P not on L , how do I find the **dual** plane N containing L and P ?

Answer: $N = (P \wedge L^*)^* = P \cdot L$.

Quiz Time!

Question: Given a **dual** line L and a point P not on L , how do I find the **dual** plane N containing L and P ?

Answer: $N = (P \wedge L^*)^* = P \cdot L$.

Question: Given a **dual** circle C and a point P not on C or on the plane determined by C , how do I find the **dual** sphere S containing C and P ?

Quiz Time!

Question: Given a **dual** line L and a point P not on L , how do I find the **dual** plane N containing L and P ?

Answer: $N = (P \wedge L^*)^* = P \cdot L$.

Question: Given a **dual** circle C and a point P not on C or on the plane determined by C , how do I find the **dual** sphere S containing C and P ?

Answer: $S = (P \wedge C^*)^* = P \cdot C$.

Quiz Time!

Question: Given a **dual** line L and a point P not on L , how do I find the **dual** plane N containing L and P ?

Answer: $N = (P \wedge L^*)^* = P \cdot L$.

Question: Given a **dual** circle C and a point P not on C or on the plane determined by C , how do I find the **dual** sphere S containing C and P ?

Answer: $S = (P \wedge C^*)^* = P \cdot C$.

Question: Let S be a **dual** sphere that intersects a **dual** plane N in more than one point, and let P be a point on S but not on N . Then, if P' is the reflection of P about N , what is the **dual** sphere reflection S' of S about N ?

Quiz Time!

Question: Given a **dual** line L and a point P not on L , how do I find the **dual** plane N containing L and P ?

Answer: $N = (P \wedge L^*)^* = P \cdot L$.

Question: Given a **dual** circle C and a point P not on C or on the plane determined by C , how do I find the **dual** sphere S containing C and P ?

Answer: $S = (P \wedge C^*)^* = P \cdot C$.

Question: Let S be a **dual** sphere that intersects a **dual** plane N in more than one point, and let P be a point on S but not on N . Then, if P' is the reflection of P about N , what is the **dual** sphere reflection S' of S about N ?

Answer:

$$S' = (P' \wedge (S \wedge N)^*)^* = P' \cdot (S \wedge N) = (P' \cdot S)N - (P' \cdot N)S.$$

The Fun Just Doesn't Stop!

Question: Let S **dually** represent the planet Saturn and let the R **directly** represent one of Saturn's rings. If this ring fell out of orbit, let the **direct** circle F on the surface of S approximate the debris field. What is F ?

The Fun Just Doesn't Stop!

Question: Let S **dually** represent the planet Saturn and let the R **directly** represent one of Saturn's rings. If this ring fell out of orbit, let the **direct** circle F on the surface of S approximate the debris field. What is F ?

Answer:

$$F = (S \wedge (R \wedge \infty)^*)^* = S \cdot (R \wedge \infty) = (S \cdot R)\infty - (S \cdot \infty)R.$$

The Fun Just Doesn't Stop!

Question: Let S **dually** represent the planet Saturn and let the R **directly** represent one of Saturn's rings. If this ring fell out of orbit, let the **direct** circle F on the surface of S approximate the debris field. What is F ?

Answer:

$$F = (S \wedge (R \wedge \infty)^*)^* = S \cdot (R \wedge \infty) = (S \cdot R)\infty - (S \cdot \infty)R.$$

Question: Let $\{N_k\}_{k=0}^3$ be 4 **dual** planes forming the sides of a non-degenerate tetrahedron, and let S be a **dual** sphere containing all 4 vertices of the tetrahedron. If all planes are known, how do we find S ? If S is known and all planes save one, how do we find the remaining plane?

The Fun Just Doesn't Stop!

Question: Let S **dually** represent the planet Saturn and let the R **directly** represent one of Saturn's rings. If this ring fell out of orbit, let the **direct** circle F on the surface of S approximate the debris field. What is F ?

Answer:

$$F = (S \wedge (R \wedge \infty)^*)^* = S \cdot (R \wedge \infty) = (S \cdot R)\infty - (S \cdot \infty)R.$$

Question: Let $\{N_k\}_{k=0}^3$ be 4 **dual** planes forming the sides of a non-degenerate tetrahedron, and let S be a **dual** sphere containing all 4 vertices of the tetrahedron. If all planes are known, how do we find S ? If S is known and all planes save one, how do we find the remaining plane?

Answer: Not very pretty!

Transformations Of Direct Geometry By Versors

Recall the **outermorphic** property of versors!

Transformations Of Direct Geometry By Versors

Recall the **outermorphic** property of versors!

Then, if the m -blade B **directly** represents any geometry, (**except a flat point**), then there exists m points $\{x_k\}_{k=1}^m \subset \mathbb{V}^n$ such that

$$VBV^{-1} = V \left(\bigwedge_{k=1}^m p(x_k) \right) V^{-1} = \bigwedge_{k=1}^m Vp(x_k)V^{-1}.$$

Transformations Of Direct Geometry By Versors

Recall the **outermorphic** property of versors!

Then, if the m -blade B **directly** represents any geometry, (**except a flat point**), then there exists m points $\{x_k\}_{k=1}^m \subset \mathbb{V}^n$ such that

$$VBV^{-1} = V \left(\bigwedge_{k=1}^m p(x_k) \right) V^{-1} = \bigwedge_{k=1}^m Vp(x_k)V^{-1}.$$

Therefore, if we understand how V transforms $p(x)$ for any $x \in \mathbb{V}^n$, then we can predict how V transforms any geometry, (**except flat points**)!

Transformations Of Direct Geometry By Versors

Recall the **outermorphic** property of versors!

Then, if the m -blade B **directly** represents any geometry, (**except a flat point**), then there exists m points $\{x_k\}_{k=1}^m \subset \mathbb{V}^n$ such that

$$VBV^{-1} = V \left(\bigwedge_{k=1}^m p(x_k) \right) V^{-1} = \bigwedge_{k=1}^m Vp(x_k)V^{-1}.$$

Therefore, if we understand how V transforms $p(x)$ for any $x \in \mathbb{V}^n$, then we can predict how V transforms any geometry, (**except flat points**)!

A versor may or may not leave ∞ invariant under conjugation.

Transformations Of Dual Geometry By Versors

If the m -blade B *dually* represents any geometry, then we can write

$$VBV^{-1} = V(B^*)^*V^{-1} = (VB^*V^{-1})^*,$$

relating this to what we know about the transformation of *directly* represented geometries.

Types Of Transformations By Versors

All conformal transformations can be represented by **versors**!
Some of these include...

- **Translations,**
- **Rotations,**
- **Dilations,**
- **Transversions.**

Surprise!

Blades representative of geometry are also **versors** representative of transformations!

Surprise!

Blades representative of geometry are also **versors** representative of transformations!

- **Planar reflections** are represented by a dual planes!

Surprise!

Blades representative of geometry are also **versors** representative of transformations!

- **Planar reflections** are represented by a dual planes!
- **Spherical reflections** are represented by dual spheres!

Surprise!

Blades representative of geometry are also **versors** representative of transformations!

- **Planar reflections** are represented by a dual planes!
- **Spherical reflections** are represented by dual spheres!
- **Translations** & **Rotations** are two successive planar reflections in well chosen planes.

Surprise!

Blades representative of geometry are also **versors** representative of transformations!

- **Planar reflections** are represented by a dual planes!
- **Spherical reflections** are represented by dual spheres!
- **Translations** & **Rotations** are two successive planar reflections in well chosen planes.
- **Dilations** & **Transversions** are two successive spherical reflections in well chosen spheres.

Surprise!

Blades representative of geometry are also **versors** representative of transformations!

- **Planar reflections** are represented by a dual planes!
- **Spherical reflections** are represented by dual spheres!
- **Translations** & **Rotations** are two successive planar reflections in well chosen planes.
- **Dilations** & **Transversions** are two successive spherical reflections in well chosen spheres.

Corollary: We can use versors to transform transformations!

Surprise!

Blades representative of geometry are also **versors** representative of transformations!

- **Planar reflections** are represented by a dual planes!
- **Spherical reflections** are represented by dual spheres!
- **Translations** & **Rotations** are two successive planar reflections in well chosen planes.
- **Dilations** & **Transversions** are two successive spherical reflections in well chosen spheres.

Corollary: We can use versors to transform transformations!

Note: Points are null, (non-invertible), and therefore, planar and spherical reflections **generate** the versor group of all transformations.

Planar Reflections

Translations

Rotations

Spherical Reflections

Dilations

Transversions

The End

Thank you for your time. Any questions?