

On The Multiplicative Inverse Of Multivectors With Respect To The Geometric Product

Spencer T. Parkin

February 4, 2013

We begin by extending the idea of linear independent among vectors to that of blades.

Definition 0.1. *For a given set $\{A_i\}_{i=1}^n$ of n non-zero blades of various grades, we say that it is a linearly independent set if there does not exist a linear combination L , given by*

$$L = \sum_{i=1}^k \alpha_i A_i, \quad (1)$$

of these blades, where $L = 0$ and not all scalars in $\{\alpha_i\}_{i=1}^n$ are zero.

It is clear that if no two blades in $\{A_i\}_{i=1}^n$ are of the same grade, then $\{A_i\}_{i=1}^n$ must be a linearly independent set. If this is not the case, however, the set may not be linearly independent. If a set of blades is not linearly independent, we will call it linearly dependent.

Definition 0.2. *For any given set of n non-zero blades $\{A_i\}_{i=1}^n$, we say that it is an irreducible set if there does not exist a set of m non-zero blades $\{B_i\}_{i=1}^m$, with $m < n$, such that*

$$\sum_{i=1}^n A_i = \sum_{i=1}^m B_i. \quad (2)$$

A non-irreducible set will be referred to as reducible. In the context of Definition 0.2, we will refer to $\{B_i\}_{i=1}^m$ as a reduction of the set $\{A_i\}_{i=1}^n$.

We now find that the irreducible sets are the linearly independent sets.

Lemma 0.1. *A given set of n non-zero blades $\{A_i\}_{i=1}^n$ is irreducible if and only if it is linearly independent.*

Proof. Clearly the lemma goes through in the case $n = 1$. Therefore, we will consider now only the cases $n > 1$. We prove the contrapositive in each direction.

Let $\{A_i\}_{i=1}^n$ be a linearly dependent set. Then, without loss of generality, we may write

$$A_n = \sum_{i=1}^{n-1} \alpha_i A_i. \quad (3)$$

Then, for each blade in $\{B_i\}_{i=1}^{n-1}$, letting $B_i = (\alpha_i + 1)A_i$, we have

$$\sum_{i=1}^n A_i = \sum_{i=1}^{n-1} B_i, \quad (4)$$

showing that $\{A_i\}_{i=1}^n$ is a reducible set.

Now assume that $\{A_i\}_{i=1}^n$ is a reducible set. Let $\{B_i\}_{i=1}^m$ be a linearly independent reduction of this set. (To find such a set, take any reduction of $\{A_i\}_{i=1}^n$ and combine any pair of linearly dependent blades until the set is linearly independent.) Considering now $\{B_i\}_{i=1}^m$ to be a basis for a linear space, we must show that $\{A_i\}_{i=1}^n$ also spans this space to establish its linear dependence. Hmm... \square

For the discussion to follow, we will find it convenient to think of a multivector M as a sum

$$M = \sum_{i=1}^n A_i, \quad (5)$$

where the set of blades $\{A_i\}_{i=1}^n$ is an irreducible set. (Clearly a multivector can always be written in terms of such a set.) Furthermore, we will overload the notation M as denoting both the multivector M and the linear space spanned by the set of blades $\{A_i\}_{i=1}^n$. It will therefore make sense to think of multivectors in M .

Consider now the multiplicative inverse of a multivector M with respect to the geometric product. The geometric product not being generally commutative, we will restrict our attention in this paper to the inverse M^{-1} of M , such that

$$1 = MM^{-1}. \quad (6)$$

Such an inverse may or may not exist. If it does exist, it is clearly unique by the zero-product property of the geometric product.

Lemma 0.2. *For any invertible multivector M , if it can be written in terms of the blades in the irreducible set $\{A_i\}_{i=1}^n$, then its inverse M^{-1} is a linear combination of the blades in this set.*

Proof. Given such a multivector M , it is natural to write M^{-1} as

$$M^{-1} = M_0 + M_1, \quad (7)$$

where $M_0 \in M$ and $\text{span}(M_1) \cap \text{span}(M) = \emptyset$. Put another way, for all multivectors $A \in M_0$, we have $A \in M$; and for all multivectors $A \in M_1$, we have $A \notin M$. Clearly M^{-1} takes on such a form, because any multivector may be written in such a form with respect to M . In the case of M^{-1} , however, we must show that $M_1 = 0$ and that $\text{span}(M_0) = \text{span}(M)$.

To that end, notice that for all $B \in M_1$, it is clear that there does not exist a blade $A \in M$ such that $\langle AB \rangle_0$ is a non-zero scalar. It follows that $\langle MB \rangle_0 = 0$, and then that $\langle MM_1 \rangle_0 = 0$. It then follows that

$$1 = MM^{-1} = \langle MM^{-1} \rangle_0 = \langle MM_0 \rangle_0 + \langle MM_1 \rangle_0 \quad (8)$$

if and only if $\langle MM_0 \rangle_0 = 1$ and $\langle MM_1 \rangle_0 = 0$. Now notice that there does not exist a multivector $A \in M_0$ and a multivector $B \in M_1$ such that

$$MA + MB = 0. \quad (9)$$

It follows that for all integers $i > 0$, we must have $\langle MM_0 \rangle_i = 0$ and $\langle MM_1 \rangle_i = 0$. We can now say that

$$1 = MM^{-1} = MM_0 + MM_1 \quad (10)$$

if and only if $MM_0 = 1$ and $MM_1 = 0$, and therefore $M_1 = 0$ by the zero product property.

What we want to show now is that $\text{span}(M_0) = \text{span}(M)$. It is clear by definition that $\text{span}(M_0) \subseteq \text{span}(M)$. \square

Knowing the form of which M^{-1} takes on in terms of M brings us a long way towards formulating M^{-1} in terms of M . Writing $M = \sum_{i=1}^n A_i$ and

then $M^{-1} = \sum_{i=1}^n \alpha_i A_i^{-1}$, we may now expand equation (6) to arrive at the following system of equations.

$$1 = \sum_{i=1}^n \alpha_i \tag{11}$$

$$0 = \sum_{i=1}^n \sum_{j=i+1}^n (\alpha_i A_j A_i^{-1} + \alpha_j A_i A_j^{-1}). \tag{12}$$

Recall that for any non-zero blade A_i that A_i^{-1} and A_i are scalar multiples of one another. We may therefore apply Lemma 0.2 in writing M^{-1} as $\sum_{i=1}^n \alpha_i A_i^{-1}$.