

# Section 3.8 Exercises

## Herstein's Topics In Algebra

Spencer T. Parkin

April 18, 2016

### Thoughts

I had some trouble with a part of the proof of Theorem 3.8.1. For integers  $a$  and  $n$ , the division algorithm can give us integers  $t$  and  $r$  such that  $a = tn + r$  with  $|r| < |n|$ , but Herstein asserts that such can be found where  $|r| \leq |n|/2$ . Well, what about  $a = 40$  and  $n = 7$ ? Here,  $t = 5$  and  $r = 5$ , and clearly  $5 \leq 7/2$  does not hold.

### Problem 1

Find all the units in  $J[i]$ .

In the field of complex numbers  $\mathbb{C}$ , multiplicative inverses are unique. That is, letting  $a + bi \in \mathbb{C}$ , we have

$$(a + bi)^{-1} = \frac{a - bi}{a^2 + b^2}.$$

Now since  $\mathbb{C}$  contains  $J[i]$ , this too must hold true. Thus, if  $x = a + bi \in J[i]$  is a unit, we must have  $d(x) | \Re(x)$  and  $d(x) | \Im(x)$ . But this is only possible for  $x = 1, -1, i, -i$ .

### Problem 2

If  $a + bi$  is not a unit of  $J[i]$  prove that  $a^2 + b^2 > 1$ .

Perhaps Herstein forgot to exclude  $0 + 0i$ .

If  $a$  and  $b$  are non-zero, then by Problem 1, they're not units, and  $d(a + bi) > 1$ . If  $a = 0$ , then by Problem 1,  $|b| > 1$ . Similarly, if  $b = 0$ , then  $|a| > 1$ .

### Problem 3

Find the greatest common divisor in  $J[i]$  of  $3 + 4i$  and  $4 - 3i$ , then  $11 + 7i$  and  $18 - i$ .

Let's consider for a moment the general problem of finding  $\gcd(a + bi, c + di)$ . Since  $J[i]$  is a Euclidean ring with unit element, we know by Theorem 3.7.1 that it is a principle ideal ring. Then by Lemma 3.7.1, we know that any greatest common divisor of  $a + bi$  and  $c + di$  is a generator of the ideal  $A$  given by

$$A = \{u(a + bi) + v(c + di) | u, v \in \mathbb{Z}\}.$$

That generator being  $x + yi$ , we must have

$$A = \{(x + yi)(u + vi) | u, v \in \mathbb{Z}\}.$$

But I'm not sure how to formulate  $x$  and  $y$  as a function of  $a, b, c, d$ , if that is even possible.