

On The Use Of Blades As Representatives Of Geometry

Spencer T. Parkin

Abstract. Abstract goes here...

Keywords. Key words go here...

1. Introduction And Motivation

In many models of geometry based upon geometric algebra (see []), blades are used to represent geometries. Seeing a great deal of commonality between these models, a formal treatment of this idea deserves to be given in an abstract setting.¹ To the author's knowledge, this is the first treatment of its kind.

2. Foundation

To lay the foundation of our work, we introduce \mathbb{V} as denoting an m -dimensional vector space generating a geometric algebra denoted by \mathbb{G} . We leave the signature of this geometric algebra unspecified, but in cases where a proof depends upon signature, one is given as either euclidean or non-euclidean.² The scalars of \mathbb{V} , (and therefore \mathbb{G}), are taken from the field of real numbers denoted by \mathbb{R} .³ We will let \mathbb{R}^n denote n -dimensional euclidean space,⁴ and let \mathbb{B} denote the set of all blades taken from \mathbb{G} . Lastly, we will let $p : \mathbb{R}^n \rightarrow \mathbb{V}$ be an unspecified function we'll use in the following definition which launches us forth into a general theory of models of geometry based upon blades in

¹For example, the use of ideals of polynomial rings as representatives of algebraic sets is studied in the setting of abstract algebra.

²The Gram-Schmidt orthogonalization process is applicable to all blades taken from and only from geometric algebras having euclidean signatures.

³To be more abstract, we could have used any field with characteristic 1, but there will be no foreseeable advantage to doing so in this paper.

⁴Some models of geometry find affine space to be the natural space within which to work, but this will not be the case in this paper.

a geometric algebra. In our abstract setting, the definition of this function does not matter. All that matters is that it is a function.

Definition 2.1 (Direct And Dual Representation). For any blade $B \in \mathbb{B}$, we say that B directly represents the set of all points $x \in \mathbb{R}^n$ such that $p(x) \wedge B = 0$, and say that B dually represents the set of all points $x \in \mathbb{R}^n$ such that $p(x) \cdot B = 0$. For convenience, we introduce the following functions using set-builder notation.

$$\begin{aligned}\hat{g}(B) &= \{x \in \mathbb{R}^n | p(x) \wedge B = 0\} \\ \dot{g}(B) &= \{x \in \mathbb{R}^n | p(x) \cdot B = 0\}\end{aligned}$$

From Definition 2.1, it's important to take away the realization that a given blade $B \in \mathbb{B}$ represents two geometries simultaneously; namely, $\hat{g}(B)$ and $\dot{g}(B)$. Which geometry we choose to think of B as being a representative of at any given time is completely arbitrary.⁵

It should also be clear from Definition 2.1 that the geometry represented by a blade B , (directly or dually), remains invariant under any non-zero scaling of the blade B . Something interesting happens, however, when we take the dual of B , as our first lemma shows.

Lemma 2.2 (Dual Something). *For any subset S of \mathbb{R}^n , if there exists $B \in \mathbb{B}$ such that $\hat{g}(B) = S$, then $\dot{g}(BI) = S$, where I is the unit pseudo-scalar of \mathbb{G} . Similarly, if there exists $B \in \mathbb{B}$ such that $\dot{g}(B) = S$, then $\hat{g}(BI) = S$.*

Proof. The first of these two statements is proven by

$$0 = p(x) \wedge B = -(p(x) \cdot BI)I \iff p(x) \cdot BI = 0,$$

while the second is proven by

$$p(x) \cdot B = 0 \iff 0 = (p(x) \cdot B)I = p(x) \wedge BI.$$

See identities (3.5) and (3.6) of Section 3. □

In words, Lemma 2.2 is telling us that for a single given geometry, the algebraic relationship between a blade directly (dually) representative of that geometry, and a blade dually (directly) representative of that geometry, is simply that, up to scale, they are duals of one another.

Of course, there will also be a geometric relationship between the geometry that is directly represented by a single given blade $B \in \mathbb{B}$, and the geometry that is dually represented by B , but this depends upon the definition of our function p , which we choose, in this paper, to leave open to speculation.

⁵In some literature on geometric algebra, a blade B intended to represent some piece of geometry directly or dually is referred to as a “geometry” or a “dual geometry,” respectively. This is confusing and not practiced in this paper. A blade is a blade; and when we refer to geometry, we will use proper language in identifying what represents it and how it does so. In this paper, a geometry is a subset of \mathbb{R}^n that can be represented dually or directly by some blade $B \in \mathbb{B}$ under Definition 2.1.

3. Useful Identities

In this section we give a number of useful algebraic identities that would otherwise distract us from the flow of the paper if given in the main body. This section is not intended as a complete review of geometric algebra. See [] for such a review.

Letting $v \in \mathbb{V}$ and $B \in \mathbb{B}$, recall that

$$vB = v \cdot B + v \wedge B. \quad (3.1)$$

Also recall that

$$v \wedge B = \frac{1}{2}(vB + (-1)^{\text{grade}(B)}Bv), \quad (3.2)$$

$$v \cdot B = \frac{1}{2}(vB - (-1)^{\text{grade}(B)}Bv). \quad (3.3)$$

Realizing that $\text{grade}(I) = m$, and that by (3.1), we have $vI = v \cdot I$, we can use equation (3.3) to establish the commutativity of vectors in \mathbb{V} with the unit psuedo-scalar I as

$$vI = -(-1)^m Iv. \quad (3.4)$$

Using equation (3.4) in conjunction with equation (3.3), we find that

$$(v \cdot B)I = v \wedge BI. \quad (3.5)$$

(In verifying this identity, it helps to realize that for any integer k , $(-1)^k = (-1)^{-k}$.) Replacing B in equation (3.5) with BI , we find that

$$v \wedge B = -(v \cdot BI)I. \quad (3.6)$$

References

Spencer T. Parkin
102 W. 500 S.,
Salt Lake City, UT 84101
e-mail: spencerparkin@outlook.com