

Geometric Calculus

Spencer T. Parkin

June 28, 2013

1 Introduction and Motivation

This paper draws upon a number of sources, most notably [1], to give an exposition of basic geometric calculus in my own words. Ancient and modern calculus is laden with confusing notation that is designed to appeal to most human's intuition, but this leaves simple-minded people, such as myself, feeling that the calculus conveyed suffers from a great deal lack of rigor. Put another way, if the language of calculus as it is inconsistently given in many different calculus texts were fed to a computer program designed to compile it into machine language, that program would either fail every time, or be extremely complicated. This paper, therefore, is my attempt to put basic geometric calculus in layman's terms and in a way that does not assume that the reader is on board with every bit of notation that can only be understood by glimpsing the unspoken mind of the author.

2 What Is Geometric Calculus?

Already having an understanding of geometric algebra (GA), we may define geometric calculus (GC) as the study of limiting processes involving multivector-valued functions defined on sub-sets of the vector-space generating the geometric algebra containing the image of such functions. Before moving on, it is therefore requisite that we review here the definition of the limit. Throughout this paper, we will let \mathbb{G} denote a geometric algebra, and \mathbb{V} the vector space generating \mathbb{G} . The scalars of \mathbb{V} will be the real numbers \mathbb{R} .

Definition 2.1. *Given a multivector-valued function $f : \mathbb{R} \rightarrow \mathbb{G}$, we define the limit of f at $x_0 \in \mathbb{R}$, if it exists, as the multivector $L \in \mathbb{G}$, and write*

$$\lim_{x \rightarrow x_0} f(x) = L, \quad (1)$$

if and only if for all real numbers $\epsilon > 0$, there exists a real number $\delta > 0$ such that

$$|x - x_0| < \delta \implies |f(x) - f(x_0)| < \epsilon. \quad (2)$$

Here, the notation $|\cdot|$ takes the norm of its argument, which norm is defined and well-known in studies of GA.

As the reader will notice, this definition applies to multivector-valued functions defined on \mathbb{R} . But what about multivector-valued functions defined on \mathbb{V} ? In an effort to answer this question, it is tempting to give what may be a natural extension of Definition 2.1 by simply replacing \mathbb{R} with \mathbb{V} in the wording of Definition 2.1. While such a definition may prove useful, what we'll find in our study of GC is that functions defined on \mathbb{V} can be analyzed in great and perhaps sufficient detail by restricting our use of the limit process to related functions defined on \mathbb{R} . The first such function we will study is the difference quotient used to define what we'll call the directional derivative.

3 The Directional Derivative Of A Function

Given a function $f : \mathbb{R} \rightarrow \mathbb{G}$, we define its derivative at a point $x_0 \in \mathbb{R}$ as $f'(x_0)$, where the function $f' : \mathbb{R} \rightarrow \mathbb{G}$ is given by

$$f'(x) = \lim_{\delta \rightarrow 0} \frac{f(x + \delta) - f(x)}{\delta}. \quad (3)$$

Extending this idea to a function $f : \mathbb{V} \rightarrow \mathbb{G}$, we get the following definition.

Definition 3.1. *Given a function $f : \mathbb{V} \rightarrow \mathbb{G}$, we define its derivative at a point $x_0 \in \mathbb{V}$ and in a direction $a \in \mathbb{V}$ as $f_a(x_0)$, where the function $f_a : \mathbb{V} \rightarrow \mathbb{R}$ is given by*

$$f_a(x) = \lim_{\delta \rightarrow 0} \frac{f(x + \delta a) - f(x)}{\delta}. \quad (4)$$

Intuitively, we may think of $f_a(x)$ as the instantaneous rate of change of the function f at x in the direction a .

Just as $\frac{d}{dx}$ is often notation used as an operator taking the derivative of function $f : \mathbb{R} \rightarrow \mathbb{G}$ in equation (3), we will let ∇_a denote an operator used to take the directional derivative of a function $f : \mathbb{V} \rightarrow \mathbb{G}$ as we have in equation (4). Specifically, we have $\frac{d}{dx}f(x) = f'(x)$ and $\nabla_a f(x) = f_a(x)$.

4 The Derivative (Gradient) Of A Function

When working with functions defined on \mathbb{R} , equation (3) gives our definition of the derivative. Equation (4), however, is not completely analogous, because it is additionally characterized by a direction vector a . What, then, might we consider to be the derivative of a function $f : \mathbb{V} \rightarrow \mathbb{G}$ at a point $x_0 \in \mathbb{V}$, and what interpretation might we attach to it? In terms of Definition 3.1, we will now proceed to give a definition of the derivative of a function $f : \mathbb{V} \rightarrow \mathbb{G}$, denoted ∇f , and sometimes referred to as the gradient of f , and then go on to contemplate its geometric significance. For the definition to follow, we let $\{e_i\}_{i=1}^n$ be any set of vectors forming a basis for the n -dimensional vector space \mathbb{V} .

Definition 4.1. *Given a function $f : \mathbb{V} \rightarrow \mathbb{G}$, we define its derivative at a point $x_0 \in \mathbb{V}$ as $\nabla f(x_0)$, where the function $\nabla f : \mathbb{V} \rightarrow \mathbb{G}$ is given by*

$$\nabla f(x) = \sum_{i=1}^n e_i f_{e_i}(x). \quad (5)$$

Notice that here, since the image of f is not necessarily confined to \mathbb{R} , the order of multiplication between e_i and f_{e_i} is significant. The product here is the geometric product.

The reader should be aware that most authors like to introduce the gradient of f as an operator that acts like a vector. Intuitively, they write

$$\nabla = \sum_{i=1}^n e_i \nabla_{e_i}. \quad (6)$$

When applied to a function f , you must realize that f distributes over addition to each operator ∇_{e_i} . To make matters even more complicated, when the application of each operator ∇_{e_i} results in a scalar-valued function, they

then go on to let $a \cdot \nabla$, where $a \in \mathbb{V}$ is any given non-zero vector, denote an operator in its own right as

$$a \cdot \nabla = \sum_{i=1}^n (a \cdot e_i) \nabla_{e_i} \quad (7)$$

under the same idea of distribution upon application. Notice that the ∇ operator is being treated like a vector here so that the inner product $a \cdot \nabla$ may be carried out. In my opinion, all of this is horribly confusing, unjustified and completely unnecessary. A claim is then made that $\nabla_a = a \cdot \nabla$ with little or no explanation. Here we are going to justify this reasoning; but for now, put equations (6) and (7) out of your mind.

Regaining our grip on mathematical rigor, take a second look at Definition 4.1, and consider the case that each function f_{e_i} is scalar-valued. This makes the function ∇f vector-valued, and for a vector $a \in \mathbb{V}$, we may then write

$$a \cdot \nabla f(x) = \sum_{i=1}^n (a \cdot e_i) f_{e_i}(x). \quad (8)$$

At first glance, it may not be entirely clear what geometric significance the function $a \cdot \nabla f(x)$ has, but the following theorem will relate it to the direction derivative which we have already come to terms with geometrically.

Theorem 4.1. *Given a differentiable and scalar-valued function $f : \mathbb{V} \rightarrow \mathbb{R}$ and a non-zero vector $a \in \mathbb{V}$, we have*

$$\nabla_a f(x) = a \cdot \nabla f(x). \quad (9)$$

Proof. We begin by rewriting equation (4) as

$$\nabla_a f(x) = \lim_{\delta \rightarrow 0} \frac{1}{\delta} \left(f \left(x + \delta \sum_{i=1}^n (a \cdot e_i) e_i \right) - \sum_{i=1}^n f(x) + \sum_{i=1}^{n-1} f(x) \right). \quad (10)$$

We now make the observation that for small values of δ , the function f begins to approximate a linear function. This allows us to write

$$\nabla_a f(x) = \lim_{\delta \rightarrow 0} \frac{1}{\delta} \left(\sum_{i=1}^n (f(x + \delta(a \cdot e_i) e_i) - f(x)) \right). \quad (11)$$

Now make the substitution $\delta = \frac{\delta'}{a \cdot e_i}$. □

Lots of rigor lacking here.