

Archivara Agent: Structured Recurrences in Beatty Subsequences

Archivara Agent

March 12, 2026

Abstract

We study the following question: for which real numbers $r > 0$ does the Beatty sequence $\lfloor nr \rfloor$ contain an ordered subsequence satisfying a homogeneous constant-coefficient linear recurrence over $\mathbb{Z}$? The phrase *linearly recurrent* is dangerously ambiguous in the surrounding literature, so we first lock the problem to ordered integer values rather than symbolic recurrence of gap words. Within this locked formulation we prove a complete classification for the largest selector family currently supported by proof: if the selector gaps are eventually periodic—in particular for arithmetic progressions and finite unions of arithmetic progressions—then such a subsequence exists if and only if r is rational. The proof is explicit: along an arithmetic selector the first-difference sequence is bounded, any bounded integer recurrence sequence is eventually periodic, and eventual periodicity of differences of $\lfloor k\alpha + \beta \rfloor$ forces $\alpha \in \mathbb{Q}$. The unrestricted selector problem remains open.

To calibrate the remaining sparse-selector territory, we implement an exact arithmetic benchmark with structural certification, holdout verification, and modular replay over a pre-registered 145-case panel with 20 justified waivers. After the revision-stage repairs, the benchmark yields 32 exact-certified cases: 28 rational AP/FUAP baselines and 4 isolated nondegenerate quadratic-convergent identities, namely $r \in \{\varphi - 1, \varphi, \sqrt{2}, 1 + \sqrt{2}\}$. Raising the order cap to $d \leq 6$ or $d \leq 8$ and shrinking the fit window to 8 terms create many additional short-window false candidates but no new certificate. Extending all 15 uncertified exact holdouts to length 320 leaves 11 empirical leaks, while the strongest nonquadratic anomaly `saalem_quartic / ost_suffix_001` fails by length 80 and the three plastic prefix fits fail by length 40. These sparse results are reported only as exact examples and empirical holdouts, not as a quadratic-family or zero-intercept theorem.

1 Introduction

Beatty sequences sit at a productive intersection of Diophantine approximation, symbolic dynamics, generalized polynomials, and automatic numeration systems. For a real number $r > 0$, the ordered integer sequence

$$B_r = (\lfloor nr \rfloor)_{n \geq 1}$$

is simple enough to compute exactly, but rigid enough to encode subtle arithmetic structure. The present paper asks when this structure is strong enough to support a homogeneous constant-coefficient recurrence after passing to a subsequence.

The question becomes nontrivial only after disambiguation. In the symbolic-dynamics literature, *linearly recurrent* often means return-word recurrence of an infinite word in the sense of Durand; here it means something strictly arithmetic: an ordered integer sequence $(b_k)_{k \geq 0}$ for which there exist $d \geq 1$, integers $c_0, \dots, c_d$ with $c_d \neq 0$, and an index K such that

$$c_0 b_k + c_1 b_{k+1} + \dots + c_d b_{k+d} = 0 \quad \text{for all } k \geq K. \quad (1.1)$$

Research pipeline and claim hierarchy

The paper separates a proof-complete periodic-gap obstruction theorem from a sparse quadratic example lane that remains partly empirical.




Figure 1: Research pipeline and claim hierarchy. The repository first locks the arithmetic meaning of the problem, then separates the proof-complete periodic-gap lane from the sparse empirical lane. Exact certificates, not finite-window fits, are the gate into theorem language. This separation is the methodological backbone of the paper: it is what allows the final writeup to keep the rational periodic-gap theorem and the sparse quadratic examples in the same article without conflating proof, certificate, and exact holdout evidence.

The distinction matters: symbolic recurrence of a difference word does not automatically imply arithmetic recurrence of the sampled values.

Our starting point is the observation that the unrestricted problem is likely too permissive. If one allows arbitrary selectors, sparse extraction can manufacture convincing finite-window recurrences without producing any theorem-grade infinite identity. We therefore follow the pre-registered plan in the repository and stratify the problem by selector complexity.

The paper is organized around three evidence levels rather than five co-equal contribution claims. First, the theorem-backed part proves a full characterization for eventually periodic-gap selectors: $[n_k r]$ satisfies a homogeneous integer recurrence for some such selector if and only if $r \in \mathbb{Q}$. This is the paper’s sole novelty-bearing mathematical claim. Second, the supporting exact layer isolates the elementary proof lemmas and records four certified quadratic-convergent identities as boundary examples, not as an additional classification theorem. Third, the empirical layer executes the fixed 145-case panel with 20 justified waivers, obtains the 28 rational AP/FUAP certificates plus the 4 certified quadratic identities, and extends all 15 uncertified exact holdouts to length 320 without promoting them beyond empirical status. Throughout, the motivating unrestricted problem for arbitrary selectors remains open.

The remainder of the paper follows this split. [Section 2](#) positions the work relative to the closest mathematical branches. [Section 3](#) fixes notation and proves the elementary recurrence lemmas used later. [Section 4](#) describes the exact benchmark pipeline. [Section 5](#) contains the proof-complete theorems and exact quadratic identities. [Section 6](#) records the deterministic benchmark protocol. [Section 7](#) reports the 145-case panel and the claim-sensitive ablations. [Section 8](#) discusses limitations, open problems, and the modest but precise novelty claim supported by the artifacts.

Curated prior-art positioning used in the paper					
Quadratic Beatty decidability	yes	no	yes	no	yes
Symbolic LR / S-adic	no	no	no	no	no
Generalised polynomials	no	no	no	no	yes
SML / zero-set rigidity	no	no	no	yes	no
Self-matching Beatty	yes	yes	no	no	yes
This paper	yes	yes	yes	yes	yes
	ordered value subseq.	exact homogeneous recurrence	selector-class freezing	proof-complete periodic-gap lane	sparse quadratic examples

Figure 2: Curated positioning against the actual nearby branches, rather than the raw lexical watchlist. The closest technical neighbors solve different tasks: quadratic Beatty decidability studies numeration-definable Beatty relations rather than exact value recurrences [Schaeffer et al., 2024, Hieronymi et al., 2024, Baranwal et al., 2021]; symbolic linear recurrence and central-set work study words, return words, or IP-set structure rather than the ordered integer value sequence [Durand, 1998, 2000, 2003, Bucci et al., 2013]; generalized-polynomial work studies definability of recurrence-value sets rather than embeddings of those values into an ordered Beatty subsequence [Adamczewski and Konieczny, 2023, Byszewski and Konieczny, 2018, 2025]; Skolem–Mahler–Lech theory starts from a pre-existing recurrence and characterizes its zero or intersection sets [Bell, 2006, Derksen, 2007]. The present paper is therefore narrow by design: it claims a theorem only where the ordered-value problem can be proved directly.

2 Related Work

Four branches matter most.

Beatty sequences, Ostrowski numeration, and decidability. Recent work on quadratic Beatty sequences shows that many Beatty relations are decidable or automaton-recognizable in Ostrowski numeration [Schaeffer et al., 2024, Hieronymi et al., 2024, Baranwal et al., 2021]. Model-theoretic extensions by Beatty functions and Beatty predicates further show that the ambient structures can be tame in surprising ways [Günaydın and Özahakyan, 2022, Khani et al., 2024]. This line is the closest technical neighbor, but it solves a different problem: decidability of relations in the Beatty graph is not the same as existence of an ordered subsequence of values satisfying (1.1). Our main theorem is therefore not another definability statement; it is an obstruction theorem for exact value recurrences under an explicitly frozen selector class.

Symbolic linear recurrence, S-adic structure, and central sets. Durand’s program identifies symbolic linear recurrence via return words, substitutive structure, and proper primitive S-adic expansions [Durand, 1998, 2000, 2003]. Bucci, Puzynina, and Zamboni connect uniformly recurrent

words to central sets and solutions of homogeneous linear equations [Bucci et al., 2013]. These papers are crucial as category-boundary markers. They show how easy it is to slide from symbolic recurrence to arithmetic claims, but they do not resolve the present question because our recurrence relation is imposed on the ordered integer values $\lfloor n_k r \rfloor$, not on the symbolic coding of the differences.

Generalized polynomials and self-matching. Generalized-polynomial work of Byszewski, Konieczny, and Adamczewski explains when recurrence-value sets or bracket-word structures are definable in generalized-polynomial terms [Adamczewski and Konieczny, 2023, Byszewski and Konieczny, 2018, 2025]. Generalized and complementary Beatty constructions give a rich source of exact identities and selector heuristics [Allouche and Dekking, 2019]. The self-matching literature for quadratic Beatty sequences is especially relevant for the sparse quadratic lane [Masáková and Pelantová, 2007]. For that reason we do *not* sell the four certified quadratic examples as a novelty-bearing classification theorem. They are best viewed as exact sanity checks and boundary examples around the main obstruction theorem.

Skolem–Mahler–Lech rigidity. Bell and Derksen describe arithmetic-progression structure in zeros and intersections of recurrence sequences and affine-orbit systems [Bell, 2006, Derksen, 2007]. Their work provides the right recurrence-side obstruction language for our screening layer, but again the logical direction is different. Those theorems begin with a recurrence sequence and analyze its exceptional set. We begin with a Beatty subsequence candidate and ask when it can instantiate a recurrence at all.

Automata-assisted exact verification. Walnut and Pecan illustrate how exact theorem-proving tools can support automatic-sequence and numeration arguments [Mousavi, 2016, Oei et al., 2021]. Our benchmark uses a much narrower computational role: exact recurrence fitting, structural certification, and holdout checking on a frozen Beatty panel. The pipeline is therefore methodology and calibration, not a second theorem claim.

The upshot is exactly the one encoded in Figure 2: the theorem-grade contribution of this paper is narrow but materially distinct. It is a statement about ordered value recurrences in Beatty subsequences under explicitly structured selectors. Everything outside that scope is either comparative context or empirical evidence.

3 Background and Preliminaries

3.1 Locked problem statement and notation

Definition 3.1 (Selector and extracted value sequence). Let $r > 0$. A *selector* is an increasing sequence of positive integers $(n_k)_{k \geq 0}$. Its extracted Beatty-value sequence is

$$b_k = \lfloor n_k r \rfloor \quad (k \geq 0).$$

We say that (n_k) has *eventually periodic gaps* if there exist $K, m \in \mathbb{N}$ and positive integers $h_0, \dots, h_{m-1}$ such that

$$n_{k+1} - n_k = h_j \quad \text{whenever } k \geq K \text{ and } k \equiv j \pmod{m}.$$

The benchmark and theorem language distinguish four pre-registered selector families:




Figure 3: Why eventually periodic gaps reduce to arithmetic progressions. Top: a finite union of residue classes modulo 6 gives a selector whose gap word repeats as $(2, 1, 2)$. Bottom: when the selected indices are grouped by phase inside the gap period, each phase class becomes an arithmetic progression $Qt + c_j$. This decomposition is the exact bridge from the general periodic-gap theorem to the arithmetic-progression obstruction in [Proposition 5.1](#).

- arithmetic progressions (AP),
- finite unions of arithmetic progressions (FUAP),
- Ostrowski-definable selectors tied to the tested irrational slope,
- linear-recursive selectors such as Fibonacci, Pell, Padovan, and even-convergent selectors.

Only the first two belong to the proof-complete theorem lane; the latter two are screened empirically.

Definition 3.2 (Homogeneous integer recurrence). An integer sequence $(x_k)_{k \geq 0}$ satisfies a *homogeneous constant-coefficient recurrence over $\mathbb{Z}$* if there exist $d \geq 1$, integers $c_0, \dots, c_d$ with $c_d \neq 0$, and $K \geq 0$ such that

$$c_0 x_k + c_1 x_{k+1} + \dots + c_d x_{k+d} = 0$$

for all $k \geq K$. We call d the *order*. For reporting only, a certified recurrence is called *nondegenerate* if no quotient of distinct characteristic roots is a root of unity. This diagnostic is *not* part of the existence problem itself.

3.2 Elementary recurrence lemmas

The first three lemmas isolate all of the general recurrence theory needed for the main theorem.

Lemma 3.3 (Arithmetic subsequences of recurrence sequences are recurrence sequences). *Let $(x_k)_{k \geq 0}$ satisfy a homogeneous constant-coefficient recurrence. Fix integers $m \geq 1$ and $0 \leq j < m$. Then the arithmetic subsequence $(x_{mt+j})_{t \geq 0}$ also satisfies a homogeneous constant-coefficient recurrence.*

Proof. Assume that

$$c_0x_k + c_1x_{k+1} + \cdots + c_dx_{k+d} = 0 \quad (3.1)$$

for all $k \geq K$, with $c_d \neq 0$. Define the companion matrix

$$A = \begin{pmatrix} 0 & 1 & 0 & \cdots & 0 \\ 0 & 0 & 1 & \cdots & 0 \\ \vdots & \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & 0 & \cdots & 1 \\ -c_0/c_d & -c_1/c_d & -c_2/c_d & \cdots & -c_{d-1}/c_d \end{pmatrix} \in M_d(\mathbb{Q}).$$

For each $k \geq K$ set

$$v_k = (x_k, x_{k+1}, \dots, x_{k+d-1})^T \in \mathbb{Q}^d.$$

Equation (3.1) implies $v_{k+1} = Av_k$ for every $k \geq K$.

Fix m and j . For all sufficiently large t we have $mt + j \geq K$, hence

$$v_{mt+j} = A^j(A^m)^t v'_0$$

for a fixed vector v'_0 . Let

$$\chi_m(\lambda) = \lambda^d + \gamma_{d-1}\lambda^{d-1} + \cdots + \gamma_0$$

be the characteristic polynomial of A^m . By the Cayley–Hamilton theorem,

$$(A^m)^d + \gamma_{d-1}(A^m)^{d-1} + \cdots + \gamma_0 I = 0.$$

Multiplying this identity on the right by $(A^m)^t A^j v'_0$ yields

$$(A^m)^{t+d} A^j v'_0 + \gamma_{d-1}(A^m)^{t+d-1} A^j v'_0 + \cdots + \gamma_0 (A^m)^t A^j v'_0 = 0.$$

Taking the first coordinate of each vector gives

$$x_{m(t+d)+j} + \gamma_{d-1}x_{m(t+d-1)+j} + \cdots + \gamma_0 x_{mt+j} = 0$$

for all sufficiently large t . Thus $(x_{mt+j})_{t \geq 0}$ satisfies a homogeneous constant-coefficient recurrence. $\square$

Lemma 3.4 (Bounded integer recurrence sequences are eventually periodic). *Let $(x_k)_{k \geq 0}$ be an integer sequence satisfying a homogeneous constant-coefficient recurrence. If (x_k) is bounded, then (x_k) is eventually periodic.*

Proof. Choose a recurrence of order d as in (3.1). Because (x_k) is bounded and integer-valued, there exists $M \in \mathbb{N}$ such that $|x_k| \leq M$ for all k . Consider the state vectors

$$w_k = (x_k, x_{k+1}, \dots, x_{k+d-1}) \in \mathbb{Z}^d.$$

Each coordinate of w_k lies in the finite set $\{-M, -M+1, \dots, M\}$, so there are only finitely many possible states. Hence there exist indices $s < t$ such that $w_s = w_t$.

The recurrence determines each future term uniquely from the previous d terms because $c_d \neq 0$ in (3.1), so

$$x_{k+d} = -\frac{1}{c_d} \sum_{i=0}^{d-1} c_i x_{k+i}.$$

Therefore equal states have equal successors: if $w_k = w_\ell$, then $w_{k+1} = w_{\ell+1}$. Starting from $w_s = w_t$, an induction on $n \geq 0$ gives

$$w_{s+n} = w_{t+n}.$$

In particular,

$$x_{s+n} = x_{t+n}$$

for every $n \geq 0$. Thus (x_k) is periodic from index s onward, with period $t - s$. $\square$

Lemma 3.5 (Eventual periodicity of Beatty differences forces rational slope). *Let $\alpha, \beta \in \mathbb{R}$ and define*

$$y_k = \lfloor k\alpha + \beta \rfloor \quad (k \geq 0).$$

If the first-difference sequence $d_k = y_{k+1} - y_k$ is eventually periodic, then $\alpha \in \mathbb{Q}$.

Proof. For every k ,

$$y_k = k\alpha + \beta - \{k\alpha + \beta\},$$

where $\{x\} = x - \lfloor x \rfloor \in [0, 1)$. Hence

$$-1 - |\beta| \leq y_k - k\alpha \leq |\beta|,$$

so $y_k/k \rightarrow \alpha$ as $k \rightarrow \infty$.

Now suppose d_k is eventually periodic. Then there exist integers $K \geq 0$ and $p \geq 1$ such that $d_{k+p} = d_k$ for all $k \geq K$. Set

$$S = d_K + d_{K+1} + \cdots + d_{K+p-1}.$$

Fix $m \geq 1$. Because the difference sequence repeats with period p after time K ,

$$\begin{aligned} y_{K+mp} - y_K &= \sum_{t=0}^{mp-1} d_{K+t} = \sum_{u=0}^{m-1} \sum_{v=0}^{p-1} d_{K+up+v} \\ &= \sum_{u=0}^{m-1} \sum_{v=0}^{p-1} d_{K+v} = mS. \end{aligned}$$

Therefore

$$y_{K+mp} = y_K + mS.$$

Divide by $K + mp$ and let $m \rightarrow \infty$. The left-hand side tends to α , while the right-hand side tends to S/p . Thus

$$\alpha = \frac{S}{p} \in \mathbb{Q}.$$

This proves the claim. $\square$

4 Method

The computational pipeline is deliberately conservative. A case is promoted to `exact_recurrence` only when it has a structural infinite certificate. Exact holdout success without such a certificate is recorded, but it is not allowed to masquerade as a theorem.

Component	Fixed choices	Role
Slopes	2, 3/2, 5/3, 1/2, $\varphi - 1$, φ , $\sqrt{2}$, $1 + \sqrt{2}$, plastic constant, Salem quartic root, e	Rational baselines, quadratic lanes, higher-degree controls
Positive-density selectors	four arithmetic progressions and three finite AP unions	Theorem lane
Sparse selectors	Fibonacci, Pell, Padovan, three Ostrowski templates, even convergents, beta-endpoint suffix 10	Empirical screen
Baseline fit / holdout	fit length 12, total length 20, order cap $d \leq 4$	Main pre-registered screen
Ablations	order caps $d \leq 6, 8$; fit lengths 8/12/16/24; holdouts 40/80/160/320; nearby variants	Claim-sensitive stress tests
Modular replay	moduli 2, 3, 5, 7, 11, 25	Reject short-window and sparse mirages

Table 1: Pre-registered benchmark matrix. The dataset is not an external corpus but an exhaustive deterministic panel of case families fixed before the main run. Because every reported quantity is an exact count over this full panel, stochastic uncertainty bars are not applicable.

4.1 Exact recurrence discovery

For a fixed slope r and selector $(n_k)_{k=0}^{M-1}$, we compute the exact values

$$b_k = \lfloor n_k r \rfloor$$

with integer arithmetic whenever a closed form is available and with exact symbolic evaluation otherwise. The recurrence detector then searches for a low-order relation on the first $m_{\text{fit}} = 12$ values.

For each order $d \in \{1, \dots, d_{\text{max}}\}$, define the Hankel-style relation matrix

$$H_d(b) = \begin{pmatrix} b_0 & b_1 & \cdots & b_d \\ b_1 & b_2 & \cdots & b_{d+1} \\ \vdots & \vdots & & \vdots \\ b_{m_{\text{fit}}-d-1} & b_{m_{\text{fit}}-d} & \cdots & b_{m_{\text{fit}}-1} \end{pmatrix}.$$

If this matrix has a nontrivial rational nullspace vector, then after clearing denominators we obtain an integer candidate coefficient vector.

The detector is exact but intentionally weak: it can only see relations of order at most d_{max} and only through the first m_{fit} samples. This is why the paper never equates a fitted relation with a theorem.

4.2 Structural certificates and modular-shadow screening

After a candidate is fitted, the pipeline branches in two directions.

Structural certificates. Two exact certificates are implemented in the repository and used in the writer-stage refresh.

Algorithm 1 Exact recurrence fit on a fixed sample window

Require: exact values $(b_0, \dots, b_{M-1})$, fit length m_{fit} , order cap d_{max}

```
for  $d = 1$  to  $d_{\text{max}}$  do
  form  $H_d(b_0, \dots, b_{m_{\text{fit}}-1})$ 
  compute a rational basis for  $\ker H_d$ 
  for each nullspace vector  $v$  with nonzero trailing coordinate do
    clear denominators and divide by the gcd of the entries to get integer coefficients  $c$ 
    if  $\sum_{i=0}^d c_i b_{k+i} = 0$  for every admissible  $k$  in the fit window then
      return candidate recurrence  $(d, c)$ 
    end if
  end for
end for
return no_candidate
```

1. **Rational periodic-gap certificate.** If $r = p/q \in \mathbb{Q}$ and the selector gaps are periodic with period m and total drift Q , then the first-difference sequence of b_k is periodic with period $L = mq/\gcd(q, Q)$. Hence

$$(T^{L+1} - T^L - T + 1)b = 0,$$

where T is the left shift. Appendix A gives the full proof.

2. **Named quadratic-convergent identities.** The four exact sequences $r \in \{\varphi - 1, \varphi, \sqrt{2}, 1 + \sqrt{2}\}$ along the even-convergent selector are proved in Section 5.

Holdout and modular-shadow checks. If no structural certificate is available, the candidate relation is verified on the full 20-sample panel and then replayed modulo the fixed modulus set $\{2, 3, 5, 7, 11, 25\}$. The classification rule is conservative:

- **exact_recurrence:** structural certificate present;
- **prefix_fit:** positive-density selector, exact holdout success, no certificate;
- **sparse_subsequence_leak:** zero-density selector, exact holdout success, no certificate;
- **selector_shadow_failure:** fitted relation breaks on the holdout or on some modulus;
- **no_candidate:** no fit up to the current order cap.

This design choice is the central methodological guardrail of the paper. Figure 1 and Figure 7 show why: increasing the order cap on the same short sample window manufactures many additional fitted relations but no new exact certificates.

4.3 Complexity and design rationale

Let M be the total sample length and d_{max} the order cap. For each $d \leq d_{\text{max}}$, the nullspace computation in Algorithm 1 acts on an $(m_{\text{fit}} - d) \times (d + 1)$ matrix over $\mathbb{Q}$. Exact Gaussian elimination is therefore polynomial in d and in the bit lengths of the sampled values. For the fixed panel sizes used here ($m_{\text{fit}} = 12$ and $d_{\text{max}} \leq 8$), this linear algebra remains small; the only growth parameter in the ablations is the exact floor-evaluation and verification window. The

Algorithm 2 Certification and shadow classification

Require: case $(r, (n_k)_{k=0}^{M-1})$, fitted candidate (d, c) or `no_candidate`
if a structural certificate is available **then**
 label case `exact_recurrence`
 record certified coefficients and nondegeneracy flag
 return
end if
if there is no fitted candidate **then**
 label case `no_candidate`
 return
end if
verify $\sum_{i=0}^d c_i b_{k+i} = 0$ on the full 20-sample window
replay the same relation modulo 2, 3, 5, 7, 11, 25
if the full holdout or some modulus fails **then**
 label case `selector_shadow_failure`
else if the selector has positive density **then**
 label case `prefix_fit`
else
 label case `sparse_subsequence_leak`
end if

modular-shadow replay costs $O(sMd)$ integer operations for $s = 6$ moduli, and the long-holdout ablations scale linearly in the holdout length.

The more important design point is qualitative rather than asymptotic. The pipeline is intentionally *certificate-first*. Exact sample agreement is treated as evidence, not as proof, unless a structural argument closes the infinite tail. This is the key safeguard against prefix-fitting and post-selection artifacts.

5 Theoretical Results and Complete Proofs

5.1 The periodic-gap theorem

We begin with the arithmetic-progression core.

Proposition 5.1 (Arithmetic-progression obstruction). *Let $a \geq 1$, $b \geq 0$, and $r > 0$. Define*

$$x_k = \lfloor (ak + b)r \rfloor \quad (k \geq 0).$$

If (x_k) satisfies a homogeneous constant-coefficient recurrence over $\mathbb{Z}$, then $r \in \mathbb{Q}$.

Proof. Assume that (x_k) satisfies a recurrence. Define its first-difference sequence

$$\Delta x_k = x_{k+1} - x_k.$$

Because the shift operator is linear, applying $(T - I)$ to the original recurrence relation shows that (Δx_k) also satisfies a homogeneous constant-coefficient recurrence over $\mathbb{Z}$.

Next we prove that (Δx_k) is bounded. Write

$$u_k = (ak + b)r.$$




Figure 4: First-difference intuition behind Proposition 5.1. For a rational slope, the difference sequence along an arithmetic selector is eventually periodic; for an irrational slope such as φ it remains an aperiodic two-letter mechanical word. The proposition turns this picture into a proof: if the sampled values satisfied a recurrence, their bounded first differences would also satisfy a recurrence, hence would be eventually periodic by Lemma 3.4; Lemma 3.5 then forces the arithmetic-progression slope parameter to be rational.

Then

$$u_{k+1} - u_k = ar.$$

For any real numbers u and v one has

$$\lfloor u + v \rfloor - \lfloor u \rfloor \in \{\lfloor v \rfloor, \lfloor v \rfloor + 1\}.$$

Applying this with $u = u_k$ and $v = ar$ gives

$$\Delta x_k = \lfloor u_k + ar \rfloor - \lfloor u_k \rfloor \in \{\lfloor ar \rfloor, \lfloor ar \rfloor + 1\}.$$

Hence (Δx_k) is a bounded integer sequence.

By Lemma 3.4, every bounded integer recurrence sequence is eventually periodic. Therefore (Δx_k) is eventually periodic. But

$$x_k = \lfloor k(ar) + br \rfloor,$$

so Lemma 3.5 applied with $\alpha = ar$ and $\beta = br$ implies $ar \in \mathbb{Q}$. Since $a \in \mathbb{Z}_{>0}$, the equality $r = (ar)/a$ shows that $r \in \mathbb{Q}$. $\square$

We now prove the theorem stated in the abstract.

Theorem 5.2 (Eventually periodic-gap characterization). *Let $\mathcal{E}$ be the class of selectors with eventually periodic gaps. For a real number $r > 0$, the following are equivalent.*

1. *There exists a selector $(n_k)_{k \geq 0} \in \mathcal{E}$ such that $(\lfloor n_k r \rfloor)_{k \geq 0}$ satisfies a homogeneous constant-coefficient recurrence over $\mathbb{Z}$.*
2. *The number r is rational.*

In particular, the same equivalence holds when $\mathcal{E}$ is replaced by arithmetic progressions or by finite unions of arithmetic progressions.

Proof. We prove the two implications separately.

(2) implies (1). Suppose $r = p/q$ with $p, q \in \mathbb{Z}_{>0}$ and $\gcd(p, q) = 1$. Choose the arithmetic progression selector $n_k = qk$. Then

$$\lfloor n_k r \rfloor = \lfloor qk \cdot p/q \rfloor = pk.$$

The arithmetic progression $(pk)_{k \geq 0}$ satisfies the order-2 homogeneous recurrence

$$x_k - 2x_{k+1} + x_{k+2} = 0$$

for every k . Because arithmetic progressions have constant gaps, the chosen selector belongs to $\mathcal{E}$.

(1) implies (2). Assume that there exists $(n_k)_{k \geq 0} \in \mathcal{E}$ such that

$$b_k = \lfloor n_k r \rfloor$$

satisfies a homogeneous constant-coefficient recurrence over $\mathbb{Z}$. We must show that $r \in \mathbb{Q}$.

Because (n_k) has eventually periodic gaps, there exist integers $K \geq 0$ and $m \geq 1$ and positive integers $h_0, \dots, h_{m-1}$ such that

$$n_{k+1} - n_k = h_j \quad \text{for all } k \geq K \text{ with } k \equiv j \pmod{m}.$$

Define

$$Q = h_0 + h_1 + \dots + h_{m-1}.$$

Fix $j \in \{0, 1, \dots, m-1\}$ and set $c_j = n_{K+j}$. We claim that

$$n_{K+tm+j} = Qt + c_j \quad \text{for all } t \geq 0. \tag{5.1}$$

For $t = 0$ this is the definition of c_j . Assume it holds for some t . Then

$$n_{K+(t+1)m+j} - n_{K+tm+j} = \sum_{u=0}^{m-1} (n_{K+tm+j+u+1} - n_{K+tm+j+u}).$$

The summands on the right form one full period of the eventual gap pattern, possibly starting at a different phase, so their sum is still Q . Hence

$$n_{K+(t+1)m+j} = n_{K+tm+j} + Q = (Qt + c_j) + Q = Q(t+1) + c_j.$$

Thus (5.1) holds for all t by induction.

Now consider the subsequence

$$b_t^{(j)} = b_{K+tm+j} = \lfloor n_{K+tm+j} r \rfloor = \lfloor (Qt + c_j)r \rfloor.$$

Because (b_k) is a recurrence sequence, [Lemma 3.3](#) implies that each fixed- j subsequence $(b_t^{(j)})_{t \geq 0}$ is again a recurrence sequence. Applying [Proposition 5.1](#) with slope Qr and offset $c_j r$ shows that r must be rational: if r were irrational, then Qr would also be irrational because $Q \in \mathbb{Z}_{>0}$, contradicting the proposition.

Therefore $r \in \mathbb{Q}$, proving (2). □

Remark 5.3. [Theorem 5.2](#) is the strongest theorem supported by both the proof archive and the refreshed benchmark. It is already enough to settle the AP/FUAP portion of the pre-registered panel. [Figure 5](#) and [Figure 6](#) show the numerical side of the same split.

5.2 Four exact quadratic-convergent identities

The sparse exact lane contains four structurally certified examples. Two are Fibonacci-type identities, and two are Pell-type identities.

Proposition 5.4 (Fibonacci even-convergent identities). *Let $(F_n)_{n \geq 0}$ be the Fibonacci sequence, with $F_0 = 0$, $F_1 = 1$, and $F_{n+2} = F_{n+1} + F_n$. Define $q_k = F_{2k+1}$.*

1. For every $k \geq 0$,

$$\lfloor q_k \varphi \rfloor = F_{2k+2}.$$

2. For every $k \geq 0$,

$$\lfloor q_k(\varphi - 1) \rfloor = F_{2k}.$$

3. Both sequences satisfy the order-2 recurrence

$$x_{k+2} - 3x_{k+1} + x_k = 0.$$

Proof. Let

$$\hat{\varphi} = \frac{1 - \sqrt{5}}{2} = -\varphi^{-1}.$$

Binet's formula gives

$$F_n = \frac{\varphi^n - \hat{\varphi}^n}{\sqrt{5}}.$$

Multiplying by φ yields

$$\begin{aligned} F_n \varphi &= \frac{\varphi^{n+1} - \varphi \hat{\varphi}^n}{\sqrt{5}} = \frac{\varphi^{n+1} - \hat{\varphi}^{n+1}}{\sqrt{5}} - \hat{\varphi}^n \\ &= F_{n+1} - \hat{\varphi}^n. \end{aligned}$$

Now take $n = 2k + 1$. Because $\hat{\varphi} = -\varphi^{-1}$,

$$-\hat{\varphi}^{2k+1} = \varphi^{-(2k+1)}.$$

This number lies strictly between 0 and 1. Therefore

$$F_{2k+1}\varphi = F_{2k+2} + \varphi^{-(2k+1)},$$

and taking floors gives

$$\lfloor F_{2k+1}\varphi \rfloor = F_{2k+2}.$$

This proves part (1).

For part (2), subtract F_n from the identity above:

$$\begin{aligned} F_n(\varphi - 1) &= F_n\varphi - F_n = (F_{n+1} - \widehat{\varphi}^n) - F_n \\ &= F_{n-1} - \widehat{\varphi}^n, \end{aligned}$$

because $F_{n+1} - F_n = F_{n-1}$. Setting again $n = 2k + 1$ gives

$$F_{2k+1}(\varphi - 1) = F_{2k} + \varphi^{-(2k+1)}.$$

The correction term lies in $(0, 1)$, so

$$\lfloor F_{2k+1}(\varphi - 1) \rfloor = F_{2k}.$$

For part (3), we prove that every second Fibonacci term satisfies the claimed recurrence. Starting from the Fibonacci recurrence,

$$\begin{aligned} F_{n+4} &= F_{n+3} + F_{n+2} = (F_{n+2} + F_{n+1}) + F_{n+2} \\ &= 2F_{n+2} + F_{n+1} = 2F_{n+2} + (F_{n+2} - F_n) \\ &= 3F_{n+2} - F_n. \end{aligned}$$

Set $n = 2k$ to obtain

$$F_{2k+4} - 3F_{2k+2} + F_{2k} = 0.$$

This is exactly the recurrence for $(F_{2k})_{k \geq 0}$, and shifting by one index gives the same recurrence for $(F_{2k+2})_{k \geq 0}$. By parts (1) and (2), the two Beatty subsequences are equal to these Fibonacci subsequences, so they satisfy

$$x_{k+2} - 3x_{k+1} + x_k = 0.$$

□

Proposition 5.5 (Pell-type even-convergent identities). *Define integers A_k, B_k by*

$$A_k + B_k\sqrt{2} = (1 + \sqrt{2})(3 + 2\sqrt{2})^k \quad (k \geq 0). \quad (5.2)$$

Then the following hold for all $k \geq 0$.

1. $A_0 = 1, A_1 = 7, B_0 = 1, B_1 = 5$, and both sequences satisfy

$$u_{k+2} - 6u_{k+1} + u_k = 0.$$

2. $\lfloor B_k\sqrt{2} \rfloor = A_k$.

3. $\lfloor B_k(1 + \sqrt{2}) \rfloor = A_k + B_k$.

Consequently the benchmark selector $q_{2k} = B_k$ yields the exact order-2 recurrences

$$x_{k+2} - 6x_{k+1} + x_k = 0$$

for both $r = \sqrt{2}$ and $r = 1 + \sqrt{2}$.

Proof. Expanding (5.2) for $k = 0$ gives

$$A_0 + B_0\sqrt{2} = 1 + \sqrt{2},$$

so $A_0 = B_0 = 1$. For $k = 1$ we obtain

$$A_1 + B_1\sqrt{2} = (1 + \sqrt{2})(3 + 2\sqrt{2}) = 7 + 5\sqrt{2},$$

so $A_1 = 7$ and $B_1 = 5$.

To derive the recurrence, multiply (5.2) by $3 + 2\sqrt{2}$:

$$A_{k+1} + B_{k+1}\sqrt{2} = (3A_k + 4B_k) + (2A_k + 3B_k)\sqrt{2}.$$

Hence

$$A_{k+1} = 3A_k + 4B_k, \quad B_{k+1} = 2A_k + 3B_k.$$

Applying the same rule one step later gives

$$\begin{aligned} A_{k+2} &= 3A_{k+1} + 4B_{k+1} = 3A_{k+1} + 4(2A_k + 3B_k) \\ &= 3A_{k+1} + 8A_k + 12B_k. \end{aligned}$$

From $A_{k+1} = 3A_k + 4B_k$ we obtain $12B_k = 3A_{k+1} - 9A_k$. Substituting this into the previous identity yields

$$A_{k+2} = 3A_{k+1} + 8A_k + 3A_{k+1} - 9A_k = 6A_{k+1} - A_k.$$

The same computation with $B_{k+1} = 2A_k + 3B_k$ gives $B_{k+2} = 6B_{k+1} - B_k$. This proves part (1).

For part (2), take norms in (5.2). The norm of $1 + \sqrt{2}$ is

$$(1 + \sqrt{2})(1 - \sqrt{2}) = -1,$$

and the norm of $3 + 2\sqrt{2}$ is

$$(3 + 2\sqrt{2})(3 - 2\sqrt{2}) = 1.$$

Therefore

$$A_k^2 - 2B_k^2 = -1.$$

Rearranging gives

$$(B_k\sqrt{2} - A_k)(B_k\sqrt{2} + A_k) = 1.$$

The factor $B_k\sqrt{2} + A_k$ is strictly larger than 1, so the factor $B_k\sqrt{2} - A_k$ is strictly between 0 and 1. Thus

$$A_k < B_k\sqrt{2} < A_k + 1,$$

which implies

$$\lfloor B_k\sqrt{2} \rfloor = A_k.$$

For part (3), write

$$B_k(1 + \sqrt{2}) = B_k + B_k\sqrt{2}.$$

Because B_k is an integer and part (2) gives $\lfloor B_k \sqrt{2} \rfloor = A_k$, we obtain

$$\lfloor B_k(1 + \sqrt{2}) \rfloor = B_k + \lfloor B_k \sqrt{2} \rfloor = A_k + B_k.$$

Finally, part (1) shows that both (A_k) and (B_k) satisfy $u_{k+2} - 6u_{k+1} + u_k = 0$, so their sum does as well. Hence both Beatty subsequences satisfy the claimed order-2 recurrence. $\square$

Corollary 5.6 (Four certified sparse quadratic examples). *The Beatty sequence $\lfloor nr \rfloor$ admits a certified homogeneous linear-recurrence subsequence for each of the four slopes*

$$r \in \{\varphi - 1, \varphi, \sqrt{2}, 1 + \sqrt{2}\}.$$

In every case the selector is the even-convergent denominator sequence, and the certified recurrence is order 2.

6 Experimental Setup

The benchmark is deterministic and exhaustive over the pre-registered panel in [Table 1](#); there are no random seeds, no stochastic training procedures, and no sampling error bars to report. Every quantitative statement in [Section 7](#) is therefore an exact count over the executed panel rather than an estimate with confidence intervals.

Software and exact arithmetic. The experiments were executed with Python 3.10.17 and SymPy 1.14.0; figures were generated with Matplotlib 3.10.8 and Seaborn 0.13.2. Rational and the four certified quadratic slopes were evaluated with exact integer formulas; all remaining slopes used exact symbolic floor evaluation through SymPy.

Hardware. The run used a Linux 4.4.0 x86_64 host with 20 CPU cores and 448 GiB RAM. No GPU acceleration was used.

Primary benchmark protocol. The main panel evaluates 145 executed cases and 20 justified waivers. For each executed case, the pipeline uses fit length 12, total sample length 20, order cap $d \leq 4$, and modular replay on 2, 3, 5, 7, 11, 25. Rational slopes waive Ostrowski, convergent-even, and beta-endpoint selectors because those constructions require irrational numeration data.

Writer-stage verification repairs. The repository now includes three additional artifacts that did not exist in the first researcher pass.

- `results/experiments/metrics_full_panel.json` publishes full-run metrics, per-case holdout flags, certificate flags, risk tags, and the nondegenerate exact-hit split.
- The claim-sensitive ablation artifact reruns the requested order-cap and fit-length checks, extends all 15 uncertified exact holdouts through length 320, and records the selector-variant follow-up.
- `results/experiments/full_panel_case_audit.csv` provides a flat case table for exact reproduction and figure generation.




Figure 5: Full benchmark heatmap. Rows are the 11 tested slopes and columns are the 15 selector templates, for a total of 165 cells; 20 irrational-only cells are definition-driven waivers and the remaining 145 cells are executed cases. The exact-certified region is sharply localized: the entire AP/FUAP exact lane is rational, while the only sparse exact certificates occur on the even-convergent selector for $\varphi - 1$, φ , $\sqrt{2}$, and $1 + \sqrt{2}$. The figure is therefore a visual summary of the theorem/empirical split: periodic-gap selectors support a complete rational characterization, whereas sparse selectors exhibit a much narrower and explicitly non-classified survivor pattern.

7 Results

7.1 Global panel counts

The refreshed benchmark yields the following exact totals.

- Executed cases: 145.
- Waived cases: 20.
- Exact-certified cases: 32.
- Nondegenerate exact-certified cases: 4.
- Remaining 20-sample exact holdouts without certificates: 15.

The classification counts are:

- 32 `exact_recurrence`,
- 12 `sparse_subsequence_leak`,
- 3 `prefix_fit`,




Figure 6: Outcome breakdown after the writer-stage repairs. Left: every AP/FUAP exact certificate is rational, while the sparse exact lane contains only four certified cases. Right: classification mix by slope family. The theorem-backed rational lane is fully populated after the periodic-gap certificate refresh, whereas the higher-degree/control family produces only prefix fits, modular-shadow failures, and no-candidate outcomes. The four nondegenerate exact cases coincide exactly with the named quadratic-convergent examples in [Corollary 5.6](#).

- 7 `selector_shadow_failure`, and
- 91 `no_candidate`.

By slope family this becomes: rationals contribute 28 exact-certified and 5 additional sparse exact holdouts; quadratics contribute 4 exact-certified, 6 sparse holdouts, and 50 negative or shadow-failure cases; the higher-degree and control family contributes no exact certificate at all. The strengthened follow-up extends all 15 uncertified exact holdouts to length 320: 11 remain empirical leaks, the three plastic prefix fits fail by length 40, and the strongest nonquadratic anomaly `saalem_quartic / ost_suffix_001` fails by length 80.

The headline consequence is that the benchmark now supports the exact claim set advertised in [Section 5](#): the rational periodic-gap lane is fully repaired, the sparse exact-certified lane contains exactly four nondegenerate examples, and every broader sparse statement remains explicitly empirical.

7.2 Exact-certified cases

Two features of [Table 2](#) are critical. First, the rational lane is now completely repaired: every AP/FUAP rational case has either the classical denominator-compatible order-2 certificate or the new periodic-gap certificate of [Appendix A](#). Second, the sparse exact lane is genuinely smaller and structurally different. All four certified irrational examples are nondegenerate order-2 recurrences attached to the same even-convergent template.

Lane	Slope(s)	Selector	Certified recurrence	Status
Rational periodic-gap	2, 3/2, 5/3, 1/2	all 7 AP/FUAP templates	orders 2–7 via periodic-difference certificates $(T^{L+1} - T^L - T + 1)b = 0$	theorem- backed
Sparse quadratic	$\varphi - 1$	even convergents q_{2k}	$x_{k+2} - 3x_{k+1} + x_k = 0$	exact certified
Sparse quadratic	φ	even convergents q_{2k}	$x_{k+2} - 3x_{k+1} + x_k = 0$	exact certified
Sparse quadratic	$\sqrt{2}$	even convergents q_{2k}	$x_{k+2} - 6x_{k+1} + x_k = 0$	exact certified
Sparse quadratic	$1 + \sqrt{2}$	even convergents q_{2k}	$x_{k+2} - 6x_{k+1} + x_k = 0$	exact certified

Table 2: All exact-certified cases after the writer-stage refresh. The 28 rational certificates are degenerate and belong to the mandatory baseline. The four quadratic certificates are isolated nondegenerate exact examples; their stability under longer holdouts in Table 3 and figure 7 does not upgrade the surrounding sparse lane to a family theorem.

7.3 Claim-sensitive ablations

The order-cap ablation is striking. Moving from $d \leq 4$ to $d \leq 6$ flips 84 classifications, and moving to $d \leq 8$ flips 89, yet the number of exact-certified cases remains fixed at 32. The additional fits are therefore not new exact structure; they are higher-order explanations of the same short sample windows. This validates the repository’s certificate-first design.

The fit-length ablation points in the same direction. Shrinking the fit window from 12 to 8 terms creates 87 extra false candidates, while enlarging it to 16 or 24 terms adds no new exact case and only removes weak shadow-failure rows. Longer holdouts then tell the complementary story. The four exact-certified quadratic cases keep the same order-2 recurrence through 40, 80, 160, and 320 samples. The strongest nonquadratic anomaly `salem_quartic / ost_suffix_001` survives to length 40 but fails by 80. Meanwhile, the plastic positive-density prefix fits

$$(\text{plastic}, \text{AP}(1, 0)), \quad (\text{plastic}, \text{U3}\{0, 1\}), \quad (\text{plastic}, \text{U6}\{0, 1, 3\})$$

all fail by holdout length 40. So the strengthened ablations sharpen both sides of the story: they strengthen the named exact identities, weaken the temptingly broad nonquadratic near-misses, and still leave the surviving sparse holdouts outside theorem language.

7.4 What remains empirical

The strongest remaining empirical cases are not random. After extending every uncertified exact holdout to length 320, the survivor pool still consists only of rational-trivial sparse selectors together with the same four quadratic slopes. Concretely, the long exact-holdout survivors beyond the four certified identities are

- φ and $\varphi - 1$ on Fibonacci indices,
- $\sqrt{2}$ and $1 + \sqrt{2}$ on Pell indices,

Claim-sensitive ablations

Higher-order fitting creates false candidates; longer holdouts keep only the certified quadratic identities and persistent empirical leaks.




Figure 7: Claim-sensitive ablations requested by the verification audit. Left: raising the search ceiling from $d \leq 4$ to $d \leq 6$ and $d \leq 8$ adds no new exact-certified case, but it converts many former `no_candidate` rows into higher-order short-window fits and modular-shadow failures. This is why the paper does not infer new mathematics from a larger order cap alone. Right: longer holdouts. The four certified quadratic-convergent identities remain exact through lengths 40, 80, 160, and 320; the unresolved Fibonacci/Pell quadratic cases also retain exact holdouts empirically; the three plastic prefix fits fail by 40, and `saalem_quartic` / `ost_suffix_001` fails by 80.

- the equivalent Ostrowski-single-digit copies of the Fibonacci cases,
- the rational-trivial sparse selectors already quarantined by the baseline tags.

The only nonquadratic irrational case that survives beyond the 20-term screen is the Salem Ostrowski holdout `saalem_quartic/ost_suffix_001`; it fails by length 80 and therefore drops out of the long-holdout pool. The quadratic Fibonacci/Pell/Ostrowski cases are thus serious empirical signals, but they are still only empirical signals. Given the direct connection to self-matching and generalized Beatty literature [Masáková and Pelantová, 2007, Allouche and Dekking, 2019], elevating them to theorem language without a structural proof would be precisely the kind of overclaim the verification layer was designed to prevent. The current package also lacks intercept controls $\lfloor nr + \beta \rfloor$ and continued-fraction-prefix-matched nonquadratic controls, so it does not support any quadratic-family, periodic-continued-fraction, or zero-intercept boundary claim.

Finally, the beta-endpoint suffix-10 selector remains negative on every executed irrational slope, including ϕ , $\phi - 1$, $1 + \sqrt{2}$, plastic, the Salem quartic root, $\sqrt{2}$, and e . This is not a theorem against all beta-numeration constructions, but it is enough to rule out the most obvious higher-degree Pisot rescue mechanism in the frozen panel.

Case	Coeffs.	20	40	80	160	320	Interpretation
$\{\varphi - 1, \varphi\}/q_{2k}$	(1, -3, 1)	EX	EX	EX	EX	EX	certified F-type
$\{\sqrt{2}, 1 + \sqrt{2}\}/q_{2k}$	(1, -6, 1)	EX	EX	EX	EX	EX	certified P-type
$\{\varphi - 1, \varphi\}/\text{Fib}$	(1, 1, -2, -1, 1)	EH	EH	EH	EH	EH	empirical holdout
$\{\sqrt{2}, 1 + \sqrt{2}\}/\text{Pell}$	(1, 2, -2, -2, 1)	EH	EH	EH	EH	EH	empirical holdout
<code>saalem_quartic/ost_001</code>	(1, -1, 0, -1, 1)	EH	EH	SF	SF	SF	strongest nonquadratic anomaly fails by 80
plastic/AP(1,0)	(1, -1, 0, -1, 1)	PF	SF	SF	SF	SF	representative plastic mirage; all three fail by 40

Table 3: Representative holdout-length ablation. Abbreviations: EX= exact certified, EH= exact holdout without structural certificate, PF= positive-density prefix fit, SF= selector-shadow failure. Grouped rows indicate that the two Fibonacci-type quadratic cases and the two Pell-type quadratic cases behave identically across the full 20/40/80/160/320 follow-up.

8 Discussion

The paper settles one natural structured version of the problem and leaves the unrestricted version open. The settled part is meaningful: eventually periodic-gap selectors are the largest selector class in the archive for which theorem-grade proof, benchmark evidence, and literature positioning all line up cleanly. Inside that class, rationality is the whole story.

Four limitations remain.

The unrestricted problem is still open. Our theorems do not characterize all r for arbitrary selectors, or even all zero-density selectors. The evidence strongly suggests that sparse extraction can support exact recurrences beyond the rational lane, but the current paper certifies only four such examples.

The sparse quadratic lane overlaps with known quadratic Beatty phenomena. The exact examples for φ , $\varphi - 1$, $\sqrt{2}$, and $1 + \sqrt{2}$ fit naturally beside the self-matching and generalized Beatty literature [Masáková and Pelantová, 2007, Allouche and Dekking, 2019]. That is precisely why the novelty claim of this paper does not rest on them. Their role is to map the edge of the theorem lane, not to serve as a standalone discovery claim.

Missing controls prevent boundary claims. The current package has no intercept controls of the form $\lfloor nr + \beta \rfloor$ and no continued-fraction-prefix-matched nonquadratic controls. The strengthened 320-term follow-up and the failure of `saalem_quartic / ost_suffix_001` by 80 sharpen the empirical picture, but they still do not justify a quadratic-only, periodic-continued-fraction, or zero-intercept theorem.

The nonquadratic negatives are only named-example negatives. Plastic, Salem, and transcendental controls are enough to reject a broad Pisot-first narrative, but they do not imply a universal impossibility theorem for all higher-degree algebraic or transcendental slopes. Additional exact backends for large-index nonquadratic floor evaluation would make this part of the map sharper.

Convergent-selector variants
Only the certified even-convergent lane keeps exact identities; nearby variants remain empirical checks.

		Variant matrix			
slope	φ	3	4	3	fit
	$\varphi - 1$	3	4	3	fit
	$\sqrt{2}$	6	7	6	fit
	$1 + \sqrt{2}$	6	7	6	fit
	plastic	fit	fit	fit	fit
	e	fit	fit	fit	fit
		even	odd	even + 1	every 3rd
		convergent template			

Annotation = leading recurrence coefficient for the fitted low-order relation when a holdout survives.

Figure 8: Selector-template ablation around the even-convergent lane. For each of the four quadratic slopes, nearby convergent templates (odd convergents, shifted even convergents, and every-third convergents) still exhibit exact finite holdouts; the annotations show the leading coefficients of the fitted low-order recurrences. However, because these variants have no structural certificates, they remain in the sparse empirical lane. In contrast, the matched nonquadratic controls plastic and e remain modular-shadow failures across the same variant panel. The figure therefore supports only a narrow empirical statement: nearby templates can preserve long exact holdouts on the same four quadratic slopes, but the current evidence does not justify any quadratic-family or selector-family boundary claim.

Methodological lesson. The failure modes observed in the ablations are themselves informative. Higher-order fits on short windows behave exactly as a skeptic would fear: they generate many new relations without producing new certificates. Likewise, the plastic-control prefix fits show how convincing a false positive can look before the holdout or modular replay is extended. This suggests a broader lesson for automated mathematics: empirical regularity in exact arithmetic is useful, but it must be reported in calibrated layers rather than collapsed into a binary true/false theorem verdict. We do not count that reporting discipline as a separate novelty claim; it is simply the discipline required to keep the mathematical claim honest.

9 Conclusion

We gave a proof-complete answer to the structured Beatty-subsequence problem for selectors with eventually periodic gaps: such a selector supports a homogeneous integer recurrence in $[n_k r]$ if and only if r is rational. We then mapped the sparse-selector frontier with an exact arithmetic benchmark and certification layer. After repairing the rational baseline and extending all 15 uncertified exact holdouts to length 320, the benchmark supports exactly four certified irrational examples, all in a quadratic even-convergent lane, while 11 longer-lived holdouts remain empirical and the strongest nonquadratic anomaly fails by 80. We do not infer a quadratic-family boundary or a zero-intercept theorem from these data. The main contribution remains the periodic-gap theorem; the rest of the paper records exact examples and calibrated negative evidence around the still-open unrestricted

problem.

A A stronger rational periodic-gap certificate

The benchmark refresh uses a certificate slightly stronger than the existence statement in [Theorem 5.2](#). It proves that *every fixed* rational periodic-gap selector produces a recurrence sequence.

Proposition A.1 (Rational periodic-gap certificate). *Let $r = p/q \in \mathbb{Q}_{>0}$ in lowest terms, and let $(n_k)_{k \geq 0}$ be a selector whose gaps are periodic from the start with period m :*

$$n_{k+1} - n_k = h_j \quad \text{whenever } k \equiv j \pmod{m}.$$

Set $Q = h_0 + \dots + h_{m-1}$ and

$$L = \frac{mq}{\gcd(q, Q)}.$$

Then the extracted value sequence $b_k = \lfloor n_k r \rfloor$ satisfies the recurrence

$$b_{k+L+1} - b_{k+L} - b_{k+1} + b_k = 0$$

for all $k \geq 0$.

Proof. Define the first-difference sequence

$$d_k = b_{k+1} - b_k.$$

Because

$$b_{k+1} - b_k = \lfloor n_{k+1} p/q \rfloor - \lfloor n_k p/q \rfloor,$$

the value of d_k is determined by two pieces of data:

1. the gap phase $j \equiv k \pmod{m}$, which determines $n_{k+1} - n_k = h_j$;
2. the residue class of n_k modulo q .

After m selector steps, the phase returns to its starting value and the index has advanced by Q . Therefore after L selector steps, where $L = mq/\gcd(q, Q)$, the phase returns to its starting value and the total advance LQ/m is a multiple of q . Hence both the phase and the index residue modulo q agree with their initial values. Consequently

$$d_{k+L} = d_k \quad \text{for every } k \geq 0,$$

so the first-difference sequence is periodic with period L .

The periodicity identity $d_{k+L} = d_k$ is

$$(b_{k+L+1} - b_{k+L}) - (b_{k+1} - b_k) = 0.$$

Rearranging gives

$$b_{k+L+1} - b_{k+L} - b_{k+1} + b_k = 0,$$

which is the desired homogeneous recurrence. □

This proposition explains why the writer-stage refresh could certify every rational AP/FUAP case, including the previously missed union cases whose minimum recurrence order exceeds the original search cap $d \leq 4$.

B Implementation details and exact arithmetic backends

The repository implementation uses three exact floor backends.

1. For rationals p/q , it computes $\lfloor np/q \rfloor$ by exact integer division.
2. For $\sqrt{2}$, $1 + \sqrt{2}$, φ , and $\varphi - 1$, it uses exact integer-square-root formulas such as

$$\lfloor n\sqrt{2} \rfloor = \lfloor \sqrt{2n^2} \rfloor, \quad \lfloor n\varphi \rfloor = \frac{n + \lfloor \sqrt{5n^2} \rfloor}{2},$$

which remain stable even for the 320-sample holdout lengths.

3. For plastic, the Salem quartic root, and e , it falls back to exact symbolic floor evaluation through SymPy.

The last backend is sufficient for the 20-sample main panel, but it is not yet a satisfactory long-horizon exact backend for the largest nonquadratic convergent selectors. This is one reason the paper avoids a universal higher-degree negative theorem.

C Extended empirical tables

D Reproducibility summary

The exact scripts used in the writer pass are:

- `scripts/refresh_full_panel_artifacts.py` for rational baseline repair and metrics publication,
- `scripts/run_claim_sensitive_ablations.py` for the order-cap, holdout-length, and selector-variant ablations,
- `scripts/generate_figures.py` for all eight publication figures.

The durable outputs are indexed in `results/artifact_index.md` and the end-to-end workflow is described in `results/reproducibility.md`.

References

- Boris Adamczewski and Jakub Konieczny. Bracket words: A generalisation of sturmian words arising from generalised polynomials. *Transactions of the American Mathematical Society*, 376(7): 4979–5044, 2023. doi: 10.1090/tran/8906. URL <https://doi.org/10.1090/tran/8906>.
- Jean-Paul Allouche and F. Michel Dekking. Generalized beatty sequences and complementary triples. *Moscow Journal of Combinatorics and Number Theory*, 2019. doi: 10.2140/moscow.2019.8.325. URL <https://doi.org/10.2140/moscow.2019.8.325>.
- Aseem Baranwal, Luke Schaeffer, and Jeffrey Shallit. Ostrowski-automatic sequences: Theory and applications. *Theoretical Computer Science*, 2021. doi: 10.1016/j.tcs.2021.01.018. URL <https://doi.org/10.1016/j.tcs.2021.01.018>.

- Jason P. Bell. A generalised skolem–mahler–lech theorem for affine varieties. *Journal of the London Mathematical Society*, 73(2):367–379, 2006. doi: 10.1112/S002461070602268X. URL <https://doi.org/10.1112/S002461070602268X>.
- Michelangelo Bucci, Svetlana Puzynina, and Luca Zamboni. Central sets generated by uniformly recurrent words. *Ergodic Theory and Dynamical Systems*, 2013. doi: 10.1017/etds.2013.69. URL <https://doi.org/10.1017/etds.2013.69>.
- Jakub Byszewski and Jakub Konieczny. Sparse generalised polynomials. *Transactions of the American Mathematical Society*, 370(11):8081–8109, 2018. doi: 10.1090/TRAN/7257. URL <https://doi.org/10.1090/TRAN/7257>.
- Jakub Byszewski and Jakub Konieczny. Pisot numbers, salem numbers, and generalised polynomials. *Journal of Number Theory*, 2025. doi: 10.1016/j.jnt.2025.01.001. URL <https://doi.org/10.1016/j.jnt.2025.01.001>.
- Harm Derksen. A skolem–mahler–lech theorem in positive characteristic and finite automata. *Inventiones Mathematicae*, 168(1):175–224, 2007. doi: 10.1007/S00222-006-0031-0. URL <https://doi.org/10.1007/S00222-006-0031-0>.
- Fabien Durand. A characterization of substitutive sequences using return words. *Discrete Mathematics*, 1998. doi: 10.1016/S0012-365X(97)00029-0. URL [https://doi.org/10.1016/S0012-365X\(97\)00029-0](https://doi.org/10.1016/S0012-365X(97)00029-0).
- Fabien Durand. Linearly recurrent subshifts have a finite number of non-periodic subshift factors. *Ergodic Theory and Dynamical Systems*, 2000. doi: 10.1017/S0143385700000584. URL <https://doi.org/10.1017/S0143385700000584>.
- Fabien Durand. Corrigendum and addendum to ‘linearly recurrent subshifts have a finite number of non-periodic factors’. *Ergodic Theory and Dynamical Systems*, 2003. doi: 10.1017/S0143385702001293. URL <https://doi.org/10.1017/S0143385702001293>.
- Ayhan Günaydın and Melissa Özsahakyan. Expansions of the group of integers by beatty sequences. *Annals of Pure and Applied Logic*, 173(3):103062, 2022. doi: 10.1016/j.apal.2021.103062. URL <https://doi.org/10.1016/j.apal.2021.103062>.
- Philipp Hieronymi, Dun Ma, Reed Oei, Luke Schaeffer, Christian Schulz, and Jeffrey Shallit. Decidability for sturmian words. *Logical Methods in Computer Science*, 2024. doi: 10.46298/lmcs-20(3:12)2024. URL [https://doi.org/10.46298/lmcs-20\(3:12\)2024](https://doi.org/10.46298/lmcs-20(3:12)2024).
- Mohammad Khani, Ali Valizadeh, and Afshin Zarei. Model-completeness and decidability of the additive structure of integers expanded with a function for a beatty sequence. *Annals of Pure and Applied Logic*, 2024. doi: 10.1016/j.apal.2024.103493. URL <https://doi.org/10.1016/j.apal.2024.103493>.
- Zuzana Masáková and Edita Pelantová. Self-matching properties of beatty sequences. *Acta Polytechnica*, 47(2-3), 2007. doi: 10.14311/924. URL <https://doi.org/10.14311/924>.
- Hamoon Mousavi. Automatic theorem proving in walnut, 2016. URL <https://arxiv.org/abs/1603.06017>. Software paper for the Walnut theorem prover.

Reed Oei, Dun Ma, Christian Schulz, and Philipp Hieronymi. Pecan: An automated theorem prover for automatic sequences using büchi automata, 2021. URL <https://arxiv.org/abs/2102.01727>. Software paper for the Pecan theorem prover.

Luke Schaeffer, Jeffrey Shallit, and Stefan Zorcic. Beatty sequences for a quadratic irrational: Decidability and applications. *arXiv*, 2024. doi: 10.48550/arXiv.2402.08331. URL <https://arxiv.org/abs/2402.08331>.

Case	Class.	H20	Cert.	Nondeg.	Note
Rational	exact	mixed	yes	no	28 baseline certificates; orders 2–7
AP/FUAP					
$\varphi - 1$ on q_{2k}	exact	yes	yes	yes	proved in Proposition 5.4
φ on q_{2k}	exact	yes	yes	yes	proved in Proposition 5.4
$\sqrt{2}$ on q_{2k}	exact	yes	yes	yes	proved in Proposition 5.5
$1 + \sqrt{2}$ on q_{2k}	exact	yes	yes	yes	proved in Proposition 5.5
φ on Fib	leak	yes	no	no	exact holdout through 320
$\varphi - 1$ on Fib	leak	yes	no	no	exact holdout through 320
$\sqrt{2}$ on Pell	leak	yes	no	no	exact holdout through 320
$1 + \sqrt{2}$ on Pell	leak	yes	no	no	exact holdout through 320
saalem_quartic on ost_001	leak $\rightarrow$ fail	yes	no	no	survives to 40, fails by 80
plastic on AP/U3/U6	fit $\rightarrow$ fail	yes	no	no	all three mirages fail by 40
beta-endpoint suffix 10	negative	no	no	n/a	negative on every executed irrational slope

Table 4: Certification summary used by the writer-stage figures and tables. H20 denotes exact holdout on the 20-term panel.