

After your denture surgery

ALVEOPLASTY · IMMEDIATE DENTURE DELIVERY

Keep this guide handy
Call our office with any questions

Your remaining teeth were removed and the bone was smoothed so your new denture can fit comfortably. A soft liner inside the denture cushions the healing tissues. You're getting used to a new denture while your gums and bone are still healing — this takes time and some patience. Most people are surprised by how fast they adapt, even when the first days feel strange.

FIRST 24 HRS	DAYS 1-7	WEEKS 2-8	MONTHS 3-6
Keep denture in. Rest. Ice. Soft, cool foods.	Swelling peaks day 2-3. Remove denture for cleaning only.	Gums shrink as they heal. Adjustments and relines as needed.	Bone fully heals. Time for a final reline or new denture.

Leave the denture in for the first 24 hours

The denture acts as a pressure bandage over the sockets and bone. It helps stop bleeding, reduces swelling, and keeps the tissue in its healing shape. **Do not remove it for the first 24 hours, even if it feels tight or uncomfortable.** If you take it out too early, the tissue can swell so much that the denture won't go back in.

1 The first 24 hours

BLEEDING

Some oozing is normal for the first day — the denture itself applies pressure and helps with this. If bleeding is heavy, bite gently on gauze placed *over* the denture for 30-45 minutes. Keep your head elevated, including when sleeping.

ICE FOR SWELLING

- 20 minutes on, 20 minutes off, for the first 24-48 hours
- Swelling peaks at 48-72 hours — not the day of surgery
- Bruising on the face or neck can appear days later and is normal
- From day 3, switch to warm moist compresses

REST & ACTIVITY

- Take it easy for 2-3 days. No exercise or heavy lifting for 5 days.
- Sleep with your head elevated on an extra pillow for the first 2-3 nights.
- Keep the denture in overnight for the first night.

2 After the first 24 hours — denture care

REMOVING THE DENTURE

After the first 24 hours, you can start removing the denture — **only for cleaning**, then put it right back in. Clean it gently over a folded towel or a sink filled with water (so it doesn't break if dropped).

CLEANING THE DENTURE

- Use a soft brush and mild soap or denture cleaner — **not regular toothpaste** (too abrasive for the soft liner)
- Rinse thoroughly before putting it back in
- At night for the first few weeks, keep it in to maintain tissue shape — unless we've told you otherwise

⚠ **Care for the soft liner inside your denture.** The pink or clear cushion material (Lynal or similar) is delicate. **Never use hot water** — it warps the liner. **Do not use denture adhesive** on the soft liner — it doesn't bond properly and can degrade the material. Rinse with cool water only.

Living with your new denture

IMMEDIATE DENTURE RECOVERY · CONTINUED

3 Medications, rinsing & hygiene

Most patients need pain medication for 3–5 days. A combination of **ibuprofen + acetaminophen** is usually more than enough — see your **Medication Guide** for dosing.

SALT WATER RINSES (START TOMORROW)

Mix ½ tsp salt in 8 oz warm water. Rinse gently **after each meal** and at bedtime. Take the denture out to rinse, and rinse the denture separately. Let water fall out of your mouth — don't spit forcefully.

IF PRESCRIBED A CHLORHEXIDINE RINSE

Start it 2 days before surgery and use it twice daily, morning and night, for the first 2 weeks after (skip the evening of surgery — no rinsing for 24 hours). Take the denture out first, let the rinse pool, tilt gently, let it fall out. Don't eat or drink for 30 minutes after.

4 Eating & talking

EATING

- First few days: cool, soft foods only — yogurt, pudding, applesauce, soft eggs
- Days 3–7: add soft pasta, fish, cooked vegetables
- Cut food into small pieces and chew slowly, using both sides evenly
- Avoid sticky foods (caramel, bread crusts) — they pull at the denture
- No hot or spicy foods for 72 hours

TALKING

- Speech will feel strange for the first week or two
- Reading aloud at home for 10–15 min a day helps your tongue adapt faster
- Extra saliva at first is normal — your body is adjusting to the denture
- "S" and "F" sounds are the hardest — they come back within days

5 What's normal vs. what's not

NORMAL (EXPECTED)

- ✓ Swelling peaking at 48–72h, then improving
- ✓ Bruising of the face, jaw, or neck appearing days later
- ✓ The denture feeling bulky or too big in your mouth
- ✓ Speaking, eating, and smiling feeling unfamiliar
- ✓ Extra saliva for the first few days
- ✓ A sore spot or two developing over the first week — we'll adjust at your follow-up
- ✓ The denture getting looser as your gums shrink during healing
- ✓ Slight bad taste for a few days

CALL OUR OFFICE

- ✗ Severe pain not controlled by medication
- ✗ Pain, swelling, or bad taste that returns *after* you'd started to feel better
- ✗ Heavy bleeding that won't stop after 30–45 min of pressure
- ✗ The denture won't go back in after you removed it
- ✗ A specific sharp pain that won't resolve — likely a sore spot that needs adjustment
- ✗ Fever over 101°F
- ✗ Foul smell or pus from the gums

i About looseness. Your denture is snug today because the tissue is swollen and the ridge is still full. As the swelling goes down and the bone settles over the next 3–6 months, the denture will gradually feel looser. This is expected. We'll do adjustments along the way, and once healing is complete (usually around 3–6 months), you'll get a final reline or a brand new denture that fits the fully healed ridge.

6 Follow-up visits

Your first follow-up is typically within **24–48 hours** to check healing and adjust any sore spots. Expect several short visits over the next few months for small adjustments and relines as your tissues change shape. These are a normal part of getting an immediate denture — not a sign something went wrong.

When to call us

- Severe pain that your medication doesn't help at all
- Pain, swelling, or bad taste returning after you'd started to feel better
- The denture won't go back in after removal
- A sharp, specific sore spot (we can adjust it)
- Heavy bleeding that won't stop after 30–45 min of pressure
- Fever over 101°F

URGENT — DON'T WAIT

- Difficulty breathing or swallowing
- Rapidly spreading swelling to your neck or around your eye

Call our office right away for the on-call provider. If you truly cannot breathe or swallow, call 911.

The first two weeks are the hardest. Most people who were nervous about this procedure tell us afterward that it went better than they expected — and that the adjustment got easier every day.

Acknowledgment & signature

I acknowledge that I have received and understand these instructions, and all my questions have been answered. I agree to follow them and to call the office with any questions or concerns.

PATIENT SIGNATURE

DATE

PATIENT PRINTED NAME

DATE OF BIRTH

PARENT / LEGAL GUARDIAN / CAREGIVER SIGNATURE (IF SIGNING FOR THE PATIENT)

DATE

PRINTED NAME

RELATIONSHIP TO PATIENT