

# Honours Diary 2020

Jarvis Carroll

March 23, 2020

## Notation

In this diary unless explicitly stated within a section, I have been using the notation specified by Nielsen and Chuang, with the following additions:

- Implicit quantifiers for index variables such as  $i, j, k$ . (Nielsen and Chuang seem to do this actually, perhaps dropping more than I do)
  - $\{x_i\} = \{x_i \mid i \in I\}$ ,  $\{|x_i\rangle\} = \{|x_i\rangle \mid i \in I\}$  etc.
  - $(x_i) = (x_1, x_2, \dots, x_n)$
  - $\sum_i$  in place of  $\sum_{i \in I}$
  - $\forall i$  in place of  $\forall i \in I$

## March 13

Set up TeXstudio and basic document structure.

### Exercise 2.1

Linear Dependence, show that  $(1, -1)$ ,  $(1, 2)$  and  $(2, 1)$  are linearly dependent.

$$\begin{aligned} & (1, -1) + (1, 2) - (2, 1) \\ &= (1 + 1 - 2, -1 + 2 - 1) \\ &= (0, 0) \end{aligned}$$

### Exercise 2.2

Matrix representations: Suppose  $V$  is a vector space with basis vectors  $|0\rangle$  and  $|1\rangle$ , and  $A$  is a linear operator from  $V$  to  $V$  such that  $A|0\rangle = |1\rangle$  and  $A|1\rangle = |0\rangle$ . Give a matrix representation for  $A$ , with respect to the input basis  $|0\rangle, |1\rangle$ , and the output basis  $|0\rangle, |1\rangle$ . Find input and output bases which give rise to a different matrix representation of  $A$ .

Equation 2.12 gives us the defining property of matrix representations:

$$A|v_j\rangle = \sum_i A_{ij}|w_i\rangle$$

This gives us a pair of vector equations:

$$|1\rangle = A|0\rangle = A_{00}|0\rangle + A_{10}|1\rangle$$

$$|0\rangle = A|1\rangle = A_{01}|0\rangle + A_{11}|1\rangle$$

By linear independence of  $|0\rangle, |1\rangle$ , it follows that

$$\begin{array}{ll} A_{00} = 0 & A_{01} = 1 \\ A_{10} = 1 & A_{11} = 0 \end{array}$$

i.e.  $A$  has the matrix representation:

$$A = \begin{bmatrix} 0 & 1 \\ 1 & 0 \end{bmatrix}$$

## March 16

### Exercise 2.3

$$A : V \rightarrow W$$

$$B : W \rightarrow X$$

$$V = \text{span}\{|v_i\rangle\} \text{ etc.}$$

At this point we are already identifying  $A, B$  with their matrix representations. We would like to show equality between the function composition  $B \circ A$  and the matrix product of the corresponding matrix representations,  $B \times A$ . Once we have done this we will be able to identify both of these concepts as simply the expression  $BA$ , but for now we will use the explicit operators  $\circ$  and  $\times$ .

Our goal then is to show that the matrix representation of  $B \circ A$  is  $B \times A$ .

$$\begin{aligned}
& \sum_i (B \circ A)_{ij} |x_i\rangle && \text{(arbitrary column of matrix } B \circ A) \\
&= (B \circ A)|v_j\rangle && \text{(Definition of matrix representation)} \\
&= B(A|v_j\rangle) && \text{(Definition of composition)} \\
&= B\left(\sum_k A_{kj} |w_k\rangle\right) && \text{(Matrix representation)} \\
&= \sum_k A_{kj} (B|w_k\rangle) && \text{(Linearity of B)} \\
&= \sum_k A_{kj} \left(\sum_i B_{ik} |x_i\rangle\right) && \text{(Matrix representation)} \\
&= \sum_i \left(\sum_k B_{ik} A_{kj}\right) |x_i\rangle && \text{(distribution)} \\
&= \sum_i (B \times A)_{ij} |x_i\rangle && \text{(matrix product)}
\end{aligned}$$

So by linear independence of  $|x_i\rangle$  we know that  $(B \circ A)_{ij} = (B \times A)_{ij}$  for arbitrary  $i, j$ , i.e. the matrix representation of  $B \circ A$  is the matrix product  $B \times A$ .

#### Exercise 2.4

$$I : V \rightarrow V$$

$$I|x_i\rangle = |x_i\rangle$$

We would like to show that  $I_{ij} = \delta_{ij}$ .

$$\begin{aligned}
& \sum_i I_{ij} |x_i\rangle \\
&= I|x_j\rangle \\
&= |x_j\rangle \\
&= \sum_i \delta_{ij} |x_i\rangle
\end{aligned}$$

Again by linear independence of  $\{|x_i\rangle\}$  we have  $I_{ij} = \delta_{ij}$ .

#### Exercise 2.5

$$((y_i), (z_i)) = \sum_i y_i^* z_i$$

We need to prove 3 properties. First linearity, taking  $|v\rangle = (v_j) = (v_1, v_2, \dots, v_n)$  and  $|w_i\rangle = (w_{ij}) = (w_{i1}, w_{i2}, \dots, w_{in})$ .

For this we define

$$|z\rangle = (z_j) = \sum_i \lambda_i |w_i\rangle$$

Then by the definitions of sum and scalar product in  $\mathbb{C}^n$  we observe

$$\begin{aligned} & z_j \\ &= \left( \sum_i \lambda_i |w_i\rangle \right)_j \\ &= \sum_i (\lambda_i |w_i\rangle)_j \\ &= \sum_i \lambda_i w_{ij} \end{aligned}$$

With this linearity falls out.

$$\begin{aligned} & (|v\rangle, |z\rangle) \\ &= ((v_j), (z_j)) \\ &= \sum_j v_j^* z_j \\ &= \sum_j v_j^* \left( \sum_i \lambda_i w_{ij} \right) \\ &= \sum_i \lambda_i \left( \sum_j v_j^* w_{ij} \right) \\ &= \sum_i \lambda_i ((v_j), (w_{ij})) \\ &= \sum_i \lambda_i (|v\rangle, |w_i\rangle) \end{aligned}$$

Next we prove conjugate symmetry.

$$\begin{aligned} & (|w\rangle, |v\rangle)^* \\ &= ((w_i), (v_i))^* \\ &= \left( \sum_i w_i^* v_i \right)^* \\ &= \sum_i w_i^{**} v_i^* \\ &= \sum_i v_i^* w_i \\ &= ((v_i), (w_i)) \\ &= (|v\rangle, |w\rangle) \end{aligned}$$

Finally positivity:

$$\begin{aligned}
 & (|v\rangle, |v\rangle) \\
 &= ((v_i), (v_i)) \\
 &= \sum_i v_i^* v_i \\
 &= \sum_i |v_i|^2
 \end{aligned}$$

Clearly this expression is at least 0, with equality when  $v_i = 0 \ \forall i$ , i.e. when  $|v\rangle = (0, 0, \dots, 0)$ .

Therefore the operator  $(\cdot, \cdot)$  is an inner product on the vector space  $\mathbb{C}^n$ .