

Safe Mobilization Prescribing in Acute Care – Physiotherapy

SAFE PRESCRIPTION OF MOBILIZING PATIENTS IN ACUTE CARE SETTINGS WHAT TO ASSESS, WHAT TO MONITOR, WHEN NOT TO MOBILIZE AND HOW TO MOBILIZE AND PROGRESS

PURPOSE, SCOPE & DISCLAIMER: The purpose of this document is to provide physical therapists with guidance on safe mobilization of the patient in acute care settings. This decision-making guide is evidence informed and where there is insufficient evidence, expert informed. It is not intended to replace the clinician's clinical reasoning skills and interprofessional collaboration. Mobilization, for the purposes of this document, has been defined as "To work towards the functional task of locomotion"

WHAT TO ASSESS	
The chart <ul style="list-style-type: none"> Medical history. Premorbid level of function (e.g., mobility aids), activity and exercise response. Primary diagnosis. Medications. Investigations, lab work (e.g., Hgb, RBC, Blood sugar, ECG, fluid/electrolytes). Risk factors and lifestyle conditions. Physician orders re specific restrictions on mobilization. 	The patient, family and team member <ul style="list-style-type: none"> Multisystem review (e.g., cognition, respiratory, cardiac, musculoskeletal & neuro systems) Level of cooperation. Ask patient what he/she currently feels about mobilization concerns and readiness. Consider the impact of the illness or medical procedures & medications on the patient's mobility (e.g., weakness from disuse, incision, trauma, pain, equipment needs, e.g., walker). Coordinate with team members the timing of treatment with medication, availability of equipment and of personnel to optimize effectiveness.
WHEN TO CONSIDER NOT MOBILIZING ¹⁻⁶	
<p><i>*Please note: the cited values are not absolute criteria for withholding mobilization but are within the range of concern that could benefit from team discussion</i></p>	
Cardiovascular status <ul style="list-style-type: none"> Mean arterial pressure: <65^{1,3} or >110¹. BP: A drop in systolic pressure (>20 mm Hg) or below pre-exercise level OR a disproportionate rise i.e., >200 mm Hg for systolic or >110 mm Hg for diastolic⁴. HR: <40³ or >130^{3,6}; requiring temporary pacer. Hemodynamic: Administration of a new pressor e.g., inotropes agent¹; two or more pressor or frequent increase⁶; uncontrolled systemic hypertension; active bleeding.^{3,5} Acute or unstable cardiac status: New MI¹; dysrhythmia requiring new medications¹; active cardiac ischemia³; unstable rhythm³; intra aortic balloon.³ Pulmonary embolus: Discussion with physician required to determine suitability. Deep venous thrombosis: May mobilize as tolerated immediately after low molecular weight heparin (e.g., enoxaparin (lovenox®), dalteparin (fragmin®), tinzaparin (innohep®), nadroparin (fraxiparin®)) is given. If patient is on any other form of anticoagulation (e.g., IV heparin) please check mobility orders with the physician. Monitor patient for changes in pain, swelling, colour and sudden shortness of breath.⁵ 	<ul style="list-style-type: none"> F_{O₂}: >60%⁵. Ventilator issues: Decreased ventilatory support that could precipitate fatigue or increased ventilatory support. Ventilator asynchrony³: unsecure airway³; pressure control ventilation⁶; uncontrolled airway irritability. Uncontrolled asthma.
Respiratory Status <ul style="list-style-type: none"> SpO₂: <88%^{1,3} or undetermined cyanosis. RR: <5 or >40³. 	Neurological status <ul style="list-style-type: none"> Patient status: Severe agitation, distress, or combative^{2,3}; not able to understand instructions thus risking patient or therapist safety. ICP: Increased² i.e., >20 mm Hg, however, ICP needs to be considered in conjunction with cerebral compliance. Uncleared, unstable/non fixated spinal cord injury³ or head injury.
WHAT TO MONITOR DURING MOBILIZATION	
<p>Subjective: Dizziness, vertigo, shortness of breath, fatigue, nausea, pain *consider use of scales e.g., Borg scale of perceived exertion.</p> <p>Objective: Cognition, balance, perspiration, cyanosis, heart rate, oxygen saturation, respiratory rate and blood pressure and all other factors relevant to patient and mobility task, for example, cardiac rhythm in those patients when ECG is essential during mobilization.</p>	
HOW TO MOBILIZE AND PROGRESS ⁷	
Step 1 Prepare <ul style="list-style-type: none"> Note obstacles or challenges related to the patient and environment and plan appropriately (e.g., set up equipment – chairs, transfer belt, mobility aids, length of leads/lines). Determine whether the benefits outweigh the risk. Ensure pre-medication as indicated (analgesia, bronchodilators, oxygen). Obtain baseline vital signs (heart rate, blood pressure, oxygen saturation). Have objective end-points such as limits of blood pressure, heart rate, oxygen saturation and level of exertion pre-determined before mobilization. 	<ul style="list-style-type: none"> If postural hypotension is suspected, monitor BP and ask patient about lightheadedness at each phase of the mobilization i.e., sitting on edge of bed, standing, walking a few paces.
Step 2 Safety first <ul style="list-style-type: none"> Use proper body mechanics during transfer and allow gradual change from lying to upright position. Encourage circulation exercises i.e., foot and ankle, knee flexion/extension before commencing more demanding mobilization procedures. 	Step 3 When to quit while you are still ahead <ul style="list-style-type: none"> Monitor closely. Watch for signs of fatigue, pain, diaphoresis and intolerance during activity. Frequently ask patient how he/she feels. Evaluate patient's status at each progression to determine whether to continue or stop.
	Step 4 Monitor and progress <ul style="list-style-type: none"> Determine the limiting factor of the mobilization and any undesirable response(s). Use objective outcome measures to monitor progress e.g., ease of transfer, sitting duration, walking distance, HR, RR, oxygen saturation, Borg scales, and pain scales. After mobilization, monitor patient until vital signs have returned to pre-activity level.

This material has been prepared solely for use at Providence Health Care (PHC), Provincial Health Services Authority (PHSA) and Vancouver Coastal Health (VCH). PHC, PHSA and VCH accept no responsibility for use of this material by any person or organization not associated with PHC, PHSA and VCH. A printed copy of this document may not reflect the current electronic version.

HOW TO PROGRESS ^{1,3,8-13} *Continue to monitor vitals to guide progression*				
LEVEL (Morris')	LEVEL I	LEVEL II	LEVEL III	LEVEL IV
TARGET LEVEL OF CONSCIOUS (RASS) ¹⁴	RASS -5 to -2	RASS -2 to -1	RASS -1 to +1	RASS -1 to +1
STRENGTH CRITERIA FOR ENTERING THIS LEVEL			Able to move arm vs. gravity.	Able to move arm and leg vs. gravity.
TURNING AND BED MOBILITY	Q2H Patient to assist as able.	Q2H Same as Level I, plus: • Scooting/bridging • Supine ↔ sitting.	Q2H Gradual withdrawal of assistance. Initiation of training to promote patient's independence.	Q2H Focus on training to promote patient's independence.
POSITIONING AND DEVICES	Keep HOB >30°. Apply splints, other positioning devices as per OT/PT instructions. Focusing on preventing pressure ulcers, especially on heels and sacrum.	Same as Level I.	Same as Level I. Assess for seating needs.	Same as Level III.
EXERCISE PROGRAM	PROM ex to incorporate into patient care e.g., during washing, turns.	Encourage pt assist with ROM during patient care e.g., during washing, turns.	Same as Level II with more active involvement.	Same as Level III.
		Consider inclusion of: • Breathing exercises. • Stretching exercises. • Balance/coordination exercises for head, neck, and trunk.	Same as Level II with more active involvement. Consider inclusion of: • Arm ergometry.	Same as Level III with more active involvement. Consider inclusion of: • Weight bearing/ weight shifting exercises.
	Additional exercise/mobilization as per physio assessment.	Same as Level I.	Same as Level I.	Same as Level I.
MOBILIZATION	HOB ≥45° x 30-60 minutes BID, support to achieve midline head and trunk position.	High fowlers or cardiac chair position x 30-60 minutes TID.	Assist physio with dangle on side of bed. May need ceiling lift if patient heavy. Sitting balance exercises with physio as appropriate, 5 to 10 minutes to start. Initially OD, progress to BID as patient tolerates.	If dangle and stand at bedside successful, physio assesses ability to weight shift, ability to transfer to chair. Initial time in chair 30 minutes, progress per OT/PT assessment. Initially OD, progress to BID as patient tolerates.
		Mobilization may include* tilt table, dangle or to chair with mechanical lift pm. *Use caution for patients at risk of hypotension.	As per physio assessment of patient strength, assist physio with sit to stand, walking in place; +/- walker. Patients with neuro/ortho status precluding WB require individualized mobilization prescription.	If patient able to transfer to chair, tolerates well, physio assesses ambulation, begins walking practice with appropriate aids, increasing distance and frequency as patient tolerates.

Richmond Agitation Sedation Scale (RASS)¹⁴:

- +4 Combative; violent, immediate danger to staff
- +3 Very agitated; pulls or removes tubes/lines; aggressive
- +2 Agitated; frequent non-purposeful movement, fights ventilator
- +1 Restless; anxious but movement not aggressive or vigorous
- 0 Alert and Calm
- 1 Drowsy; not fully alert, sustained awakening (eye-opening/contact) to voice >10 sec
- 2 Light sedation; briefly awakens with eye contact to voice < 10 sec
- 3 Moderate sedation; Movement or eye opening to voice but no eye contact
- 4 Deep sedation; No response to voice but movement or eye opening to physical stimulation
- 5 Unarousable; No response to voice or physical stimulation

Developed by the SAFEMOB Task Force: Dr. Elizabeth Dean, Dr. Darlene Reid, Frank Chung, Simone Gruenig, Rosalyn Jones, Jocelyn Ross, Maylinda Urbina, Alison Hoens.

References

- Morris PE, Goad A, Thompson C, Taylor K, Harry B, Passmore L et al. Early intensive care unit mobility therapy in the treatment of acute respiratory failure. Critical Care Medicine. 2008; 36:2238-43.
- Stiller, K. Safety issues that should be considered when mobilizing critically ill patients. Crit Care Clin 2007; 23, 35-53.
- Schweickert WD, Pohlman MQ, Pohlman NS Nigos G, Pawlik AJ, Esbrook CL et al. Early physical and occupational therapy in mechanically ventilated, critically ill patients: an RCT. Lancet. 2009; 373:1874-82.
- ACSM Guidelines for Exercise Testing and Prescription. 8th edition. Lippincott Williams & Wilkins. Philadelphia 2010 pp.209-10.
- Timmerman, RA. A mobility protocol for critically ill adults. [DIMENS CRIT CARE NURS. 2007; 26(5): 175-9.
- Singh G, Fletcher R, Cunningham K, and Szivka M. Mobilization with a Deep Vein Thrombosis. Clinical Practice Guideline (DRAFT in process). Fraser Health Authority; 2009.
- Reid WD, Chung F. Clinical management notes and case histories in cardiopulmonary physical therapy. New Jersey: Slack; 2004.
- Chung F, Fletcher R, Lavioie K, Parent L, Perel D, Roy L, Urbina, M. Members of the physiotherapy professional practice council critical care practice stream for the Fraser Health Authority. Canada. Forthcoming. 2009.
- Bailey P, Thomsen GE, Spuhler VJ, Blair R, Jewkes J, Bezjian L, et al. Early activity is feasible and safe in respiratory failure patients. Crit Care Med. 2007; 35:139-45.
- Collard HR, Saint S, Matthay MA. Prevention of ventilator-associated pneumonia: an evidence-based systematic review. Ann Intern Med. 2003;138:494-501.
- Dodek P, Keenan S, Cook D, MD, Heyland D, Jacka M, Hand L et al. Evidence-based clinical practice guideline for the prevention of ventilator associated pneumonia. Ann Intern Med. 2004; 141:305-13.
- Needham D. Mobilizing patients in the intensive care unit: improving neuromuscular weakness and physical function. JAMA. 2008; 300:1685-90.
- Perme C, Chandrasekhar R. Early mobility and walking program for patients in intensive care units: creating a standard of care. Am J Crit Care. 2009; 18:212-21.
- Sessler CN, Gosnell M, Grap MJ, Brophy GT, O'Neal PV, Keane KA et al. The Richmond Agitation-Sedation Scale: validity and reliability. Am J Respir Crit Care Med 2002; 166: 1338-44.

ot no
l.

Effective Date:	March 1 2014
Posted Date:	
Last Revised:	
Last Reviewed:	
Approved By:	Professional Practice Leader Physiotherapy
	PHC
Owners:	Physiotherapy
	Professional Practice Leader Physiotherapy