

PEDIATRIC BIOCHEMICAL GENETICS HYPOGLYCEMIA MANAGEMENT ALGORITHM

NOT FOR USE FOR HYPOGLYCEMIA TREATMENT IN INFANTS <1 MONTH OF AGE

BLOOD GLUCOSE (BG) < 4.0 mmol/L (unless cut-off otherwise specified in MD orders)

1. Assess vital signs and level of consciousness (LOC)

2. If airway, breathing, vitals and/or LOC compromised, call CODE BLUE or 911 (as per site applicability) and initiate resuscitation

**Conscious & able to swallow/has enteral access
(includes GSD diet/tube feeds)**

1A Give 5 - 10 grams fast-acting carbohydrate orally

Suggestions for fast-acting carbohydrates:

- 6 grams =
 - 1-12 months: 30 mL of 20% sugar water*
 - 1 - 10 years: 1 packet saltine crackers or 1½ Dex4® tabs or 50 mL apple juice
- 9 grams =
 - >10 years: 2 packets saltine crackers or 2 Dex4® tabs or 80 mL apple juice

***20% sugar water: mix 3 × 4-gram packets table sugar in 59 mL sterile water; shake 1 minute.**

NOTE: All patients at risk of hypoglycemia should have hypoglycemia treatment supplies in fridge upon admission and during stay.

Tube feeds: If gastric, use juice, and flush pre/post with water. If post-pyloric (NJ), page MD.

2A Recheck blood glucose after 15 mins

- If BG < ordered cut-off, repeat **Step 1A** OR give next meal/snack/cornstarch dose if due. After repeating this twice, call MD
- If BG ≥ ordered cut-off, no further action, resume usual feeding schedule

UNCONSCIOUS (CODE BLUE and Resuscitation initiated) OR NPO/TPN

- Attempt to establish IV access within two minutes if not already available and page MD
- Glucagon **CONTRAINDICATED** in GSD

IV ACCESS

1B Obtain order from MD to give 2 mL/kg D10W IV bolus (maximum: 150mL)

2B Recheck blood glucose after 10 mins

If BG < ordered cut-off

- 3B Patient remains unconscious and/or NPO:**
- Continue with D10 IV fluids; discuss with MD and ask for new order if solution change is necessary
 - IV rate should be adjusted to ensure the BG remains between 4-6mmol/l

- 3C Patient is conscious and allowed/able to swallow:**
- Initiate **Step 1A** or give meal/snack/cornstarch dose if due

If BG ≥ ordered cut-off

- 3D Patient remains unconscious and/or NPO:**
- Initiate **Step 3B**

- 4D Patient is conscious and able to swallow:**
- Resume usual feeding schedule and discuss if any changes appropriate with MD

- 4B Recheck BG 1 hour post hypoglycemia treatment to ensure BG remains > MD order cut-off**
- If BG < ordered cut-off, repeat **Step 1B**.
 - If BG ≥ ordered cut-off, continue IV fluids until new MD orders

NO IV ACCESS

1C Call IV team & MD

Contact Biochemical Genetics On-Call if BG <3.4 mmol/L (unless cut-off otherwise specified in MD orders)

