## PROVIDENCE HEALTH CARE – HAZARDOUS DRUGS CONTROL MATRIX

		Group 1							
BC Hazardous Drug Control Matrix Nursing Section			Parenteral Oral Solid Oral Liquid Topical, Rectal & Implants and Inhalation Therapy						
			(IV, IVe, IM, SUBCUT, IT, IP)	(Tablet, Capsule)	Orai Liquiu	Vaginal	Ophthalmic		Illinatation merapy
LABELLING MEDICATION			Precautions Hazardous Drug GROUP 1 HD1						
TRANSPORT IN FACILITY			Do not use tube system						
			<ul> <li>If transporting drugs in a reusable outer container - two pairs of chemo-approved gloves are required</li> <li>If transporting drugs in a disposable outer container (e.g. plastic bag), PPE is not required</li> <li>Ensure a Hazardous Drugs spill kit accompanies drug transport or is readily available</li> </ul>						
PREPARATION OF DRUGS BY NURSING	DRUGS BY NURSING PROTECTIVE EQUIPMENT (PPE)		See Safe Work Procedures						
PRIMING IV LINES		Do not prime IV lines with hazardous drugs							
MEDICATION ADMINISTRATION	PPE	Gloves	two pairs of chemo- approved gloves  one pair of two pairs of chemo-approved gloves  two pairs of chemo-approved gloves						
Drugs in final dosage		Gown	chemo-approved gown None chemo-approved gown if risk of splash						chemo-approved gown
form (including when using CSTDs)		Eye/Face	eye/face protection	None	eye/face protection if risk of splash eye protec				
		N95	None N95 Required						
WORK SURFACE CLEANING			<ul> <li>Wear two pairs of chemo-approved gloves, chemo-gown, eye/face protection if risk of splash</li> <li>Use Accelerated Hydrogen Peroxide 0.5% (e.g. Accel Intervention wipes™)</li> </ul>						
PRECAUTIONARY PERIOD  Place precautionary signage at the bedside or entrance to client room  LABORATORY  LAUNDRY  SPILL MANAGEMENT  Hazardous drug spill kits in all areas where hazardous drugs are handled			The precautionary period is 48 hours following each administration of a Group 1 HD  During precautionary period  Required PPE: two pairs of chemo-approved gloves, chemo-approved gown, eye/face protection if risk of splash for any activities for which there is a risk of contact with BBF  Outside precautionary period  Follow routine precautions when handling blood and body fluids  During an individual's precautionary period all blood, urine and stool samples; other body fluids visibly contaminated with blood (except sputum) and tissues or organs not in fixatives must be labelled with a laboratory HD1 label  All laboratory specimens from a patient in the precautionary period are to be placed in an outer sealed plastic bag. Blood specimens may be tubed, and must be labelled and placed in two sealed bags, with the outer bag labelled  During precautionary period: All linen to be placed in Group 1 HD labelled laundry bag  Required PPE: two pairs of chemo-approved gloves, chemo-approved gown, eye/face protection if risk of splash  Outside precautionary period: All linen to be placed in regular laundry bag  Required PPE: Follow routine practices  Drug Waste  Manage spills according to Hazardous Drug Spill Cleanup Procedures for size of spill. See Appendix L in ECP  Manage spills according to Hazardous Drug Spill Cleanup Procedures for size of spill. See Appendix L in ECP						
WASTE MANAGEMENT Acute/Long Term Care (i.e. in facility)			Procedures for size of spill. S Controlled Substances: Pharmaceutical Waste Bin Non-Controlled Substance Cytotoxic Waste Bin Required PPE: two pairs of chemo-app chemo-approved gowr eye/face protection if ri	s: proved gloves if risk of splash	Cytotoxic Sharps Container	Where possible, d through the sewer disposed of throug leak-proof cytotoxi	. Where not gh sewer, use a	Cytotoxic con	ntainer

Acronyms: CSTD = Closed System Transfer Device IV = Intravenous IV = Intra