

Date: _____ Time: _____

Select only ONE option based on PAWSS score and age.

If a PAWSS score previously completed on this admission and patient is believed to be actively experiencing alcohol withdrawal, select ONE option based on age.

CIWA-Ar Monitoring and Benzodiazepine Administration

CIWA-Ar Score	<input type="checkbox"/> PAWSS 3 or Below * OR* Unable to Complete PAWSS Due to Patient Condition All Ages	<input type="checkbox"/> PAWSS 4 and Above *OR* Exhibiting Severe Alcohol Withdrawal *AND* 69 Years and Younger	<input type="checkbox"/> PAWSS 4 and Above *OR* Exhibiting Severe Alcohol Withdrawal *AND* 70 Years and Older	CIWA-Ar Assessment
0 to 9	No LORazepam	No LORazepam	No LORazepam	Q1H x 3, then Q6H x 24 hours, then Q24H x 72 hours
10 to 19	No LORazepam Notify Physician for Patient Re-Assessment	LORazepam 1 to 2 mg PO or sublingual or IM or IV Q1H PRN	LORazepam 0.5 to 1 mg PO or sublingual or IM or IV Q1H PRN	Q1H until score below 10
20 or greater	No LORazepam Notify physician for patient reassessment	LORazepam 2 mg PO or sublingual or IM or IV Q1H PRN Notify physician for patient reassessment	LORazepam 1 mg PO or sublingual or IM or IV Q1H PRN Notify physician for patient reassessment	Q1H until score below 20