

**Vancouver
CoastalHealth**
VA: VGH / UBC / GFS
VC: BP / Purdy / GPC

ADDRESSOGRAPH

COMPLETE OR REVIEW ALLERGY STATUS PRIOR TO WRITING ORDERS

**ACUTE LYMPHOBLASTIC LEUKEMIA (ALL 13-01)
INTRATHECAL CHEMOTHERAPY ORDERS - OUTPATIENT
(FOR CONSOLIDATION II & CONTINUATION)
Adult Ph-Negative ALL Patients (16-39 years)**

(Items with check boxes must be selected to be ordered)

(Page 1 of 1)

Date: _____ Time: _____

Time
Processed
RN/LPN Initials
Comments

LABORATORY:

Check INR, PTT, platelet count, AST, ALT, and bilirubin (total and direct) prior to each LP

MEDICATIONS:

BCCA Code for PCIS order entry: LKNOS

*Intrathecal chemotherapy orders require one prescriber signature only.***Intrathecal Chemotherapy Injections:** (Use preservative-free solutions only)

methotrexate 12 mg plus cytarabine 40 mg plus hydrocortisone 50 mg IT every 18 weeks until 2 years of continued complete remission as per completed INTRATHECAL CHEMOTHERAPY (#819) PRE-PRINTED order for each date below

Hold systemic methotrexate on the week that IT methotrexate is given

Week 0 (date): _____ (date = date of first LP given during CNS phase)

Week 18 (date): _____

Week 36 (date): _____

Week 54 (date): _____

Week 72 (date): _____

Week 90 (date): _____

Week 108 (date): _____

NOTES TO PRESCRIBER: (Unit Clerk/Pharmacy do not process – reminders to prescriber only)

First IT chemotherapy begins 18 weeks after the start of ALL 13-01 CNS CHEMOTHERAPY (#853) Orders and continues every 18 weeks throughout Consolidation II and Continuation phases

Give as close to 18 weeks as possible

Always give at the start of a chemo cycle

Modification: IF AST is greater than 8 x of upper limit of normal and/or direct bilirubin is greater than 23.9 micromol/L
OMIT methotrexate IT and administer IT cytarabine and hydrocortisone. (Physician to write orders)

Prescriber to verify with patient that prophylactic LMWH has been held for at least 12 hours prior to each LP