



Guideline BD-00-07-40112

Negative Pressure Wound Therapy (NPWT): Guideline for Adults and Children

Site Applicability

VCH and PHC sites where NPWT is approved for use

Practice Level

Profession	Advanced Skill		
FIOIESSIOII	(requiring additional education)		
Nurse Specialized in Wound, Ostomy and Continence (NSWOC), Nurse Practitioner (NP)	 Upon successful completion of additional NPWT education provided by their specialty, NSWOC may: Provide a client-specific order and establish the wound care treatment plan for other nurses to follow. Care for a wound with NPWT Follow this DST 		
Registered Nurse (RN)	 Upon successful completion of additional education listed under required education, RNs: Care for wound with NPWT when there is a client specific order in place. Develop a wound care treatment plan based on a client specific order. Follow this DST 		
Registered Psychiatric Nurse (RPN)	 Upon successful completion of additional education listed under required education, RPNs: Care for wound with NPWT when there is a client specific order and a wound treatment plan established by an RN. Follow this DST 		
Licensed Practical Nurse (LPN)	 Upon successful completion of additional education listed under required education, LPNs: Care for a wound with NPWT: when it is stable with a predictable healing trajectory, when there is a client specific order, AND when there is an established wound care treatment plan in place. Follow this DST 		

Required Education:

- RNs/RPNs/LPNs must successfully complete the following education:
 - Level One Monitor and Manage to care for any NPWT device:
 - o Additional Negative Pressure Wound Therapy: Monitor and Manage
 - Level Two: Dressing Application to apply a NPWT device:
 - o PHC: Negative Pressure Wound Therapy (VAC) Certification

This material has been prepared solely for use at Providence Health Care (PHC) and Vancouver Coastal Health (VCH). PHC and VCH accept no responsibility for use of this material by any person or organization not associated with PHC and VCH. A printed copy of this document may not reflect the current electronic version.

Effective Date: 18 September 2020 Page 1 of 3





Guideline BD-00-07-40112

- VCH Acute Sites: <u>VCH VAC Basic / Advanced Education (Acute Care Nursing)</u>
- VCH Community Sites: <u>VCH Introduction to Negative Pressure Wound Therapy</u> (Community Nursing)
- LPNs must also complete <u>Licensed Practical Nurses Care of Wounds with a Non-Visible Wound</u> Bed in addition to above learning modules.

Requirements

A Physician, NP, NSWOC or Wound Clinician order is required for the initiation of Negative Pressure Wound Therapy using either a reusable machine or a disposable device.

Need to Know

The Provincial Nursing Skin and Wound Committee (PNSWC) developed this DST to provide a provincial standard for all patients/clients/residents who are having NPWT.

This DST covers all NPWT except open abdominal therapy.

Guideline

The Provincial Nursing Skin and Wound Committee (PNSWC) developed this DST to provide a provincial standard for wound care for all patients, clients and residents who are undergoing Negative Pressure Wound Therapy (NPWT).

The document outlines the assessment, care and management to be followed for all NPWT devices.

The PNSWC has also developed specific procedural documents for each of the different devices (see Guidelines) and educational resources (see Patient and Family Education section).

Guideline: Negative Pressure Wound Therapy (NPWT) for Adults and Children

Documentation

See **Guideline**

Patient and Family Education

Education pamphlets available through Patient Health Information Materials

Evaluation

See Guideline

Related Documents

See **Guidelines**

References

See Guidelines

Appendices

See Guideline

This material has been prepared solely for use at Providence Health Care (PHC) and Vancouver Coastal Health (VCH). PHC and VCH accept no responsibility for use of this material by any person or organization not associated with PHC and VCH. A printed copy of this document may not reflect the current electronic version.

Effective Date: 18 September 2020 Page 2 of 3





Guideline BD-00-07-40112

Effective Date:	18-SEPTEMBER-2020		
Posted Date:	18-SEPTEMBER-2020		
Last Revised:	18-SEPTEMBER-2020		
Last Reviewed:	18-SEPTEMBER-2020		
Approved By:	PHC	VCH	
(committee or position)	Endorsed by: Professional Practice Standards Committee	Endorsed by: Vice President, Professional Practice and Chief Clinical Information Officer	
Owners:	PHC/VCH		
(optional)	The Provincial Nursing Skin and Wound Committee in collaboration with Regional Skin and Wound Committees from NHA, IHA, FHA, VIHA, PHSA and VCH/PHC NSWOC, Clinical Nurse Specialist, Skin and Wound, PHC NSWOC, Clinical Nurse Specialist, Wound Ostomy Continence, VCH Nurse Clinician, Burn Plastic Trauma Unit, VGH NSWOC, VGH NSWOC, LGH NSWOC, LGH NSWOC, Nurse Practitioner- Long Term Care, Residential Program, VCH NSWOC, Pender Community Health Centre, VCH NSWOC, Nursing Informatics, Clinical Lead (Pixalere) IMITS - Clinical Solutions - Community Applications, VCH Provincial Nursing Skin and Wound Committee, VCH		
	*NSWOC = Nurse Specializing in Wound Ostomy Continence		

This material has been prepared solely for use at Providence Health Care (PHC) and Vancouver Coastal Health (VCH). PHC and VCH accept no responsibility for use of this material by any person or organization not associated with PHC and VCH. A printed copy of this document may not reflect the current electronic version.

Effective Date: 18 September 2020 Page 3 of 3