

β -blocker, Intravenous (Metoprolol, Labetalol, Esmolol) Administration of First Dose in SPH CSICU

Site Applicability

PHC – SPH CSICU

Practice Level:

Specialized: limited to Registered Nurses in **CSICU** who have completed the certification package.

Requirements

Providers are responsible for ordering medication type, amount, frequency and maximum dosage.

Need to Know:

- In the early recovery period after cardiac surgery, IV beta- blockers may be used to manage tachyarrhythmia and hypertension. β -blockers prevent catecholamines from binding to β -adrenergic receptors throughout the body; thereby reducing heart rate and blood pressure.

Receptor	Location	Normal response to stimulation	Effects of beta-blockers
$\beta - 1$	Heart	<ul style="list-style-type: none"> Positive chronotrope Positive inotrope Automaticity Rate of conduction 	↓HR, contractility, MvO ₂ ↓BP ↓Irritability ↓AV conduction
$\beta - 2$	Bronchioles Blood vessels GI tract	<ul style="list-style-type: none"> Bronchodilation Vasodilation of muscle vasculature 	<ul style="list-style-type: none"> Bronchoconstriction Vasoconstriction of muscle vasculature Relaxation of GI smooth muscle
	Other	<ul style="list-style-type: none"> Drives K⁺ intracellular Renin release 	Serum K ⁺ ↓Blood pressure

This material has been prepared solely for use at Providence Health Care (PHC). PHC accepts no responsibility for use of this material by any person or organization not associated with PHC. A printed copy of this document may not reflect the current electronic version.

- β -blockers are a negative inotrope and are not usually administered to patients with cardiac index (CI) 2.2 or less, or to patients who require IV inotropes to keep their Cardiac Index over 2.2.
- Patients with asthma are at higher risk for developing bronchospasm and should be given non-cardioselective β -blockers (i.e. labetalol) with caution.
- Before initiating IV β -blockers, assess for and treat possible contributing factors for tachycardia and/or hypertension:
 - Hypothermia
 - Hypovolemia
 - Pain/anxiety/substance withdrawal
 - Hypoxia/hypercarbia
 - Use of β -1 agonists - (epinephrine, NORepinephrine, DOBUTamine, DOPamine)

Approved Medications:

Medication	Blocks	Indications (main uses)	Typical Doses
Labetalol	α -1, β -1, β -2	Hypertension	<i>Power plan order</i> 5 to 10 mg IV repeat Q 5 minutes until target HR/BP met *Max 50 mg/24h. Hold if HR less than 60
Metoprolol	β -1 selective	Hypertension, SVT, tachycardia, control ventricular response in A fib/flutter	1 to 2 mg IV may repeat every 2 to 5 mins until target HR/BP met (max 15 mg)
Esmolol	β -1	Short-acting test of Beta-blocker tolerance	10 to 20 mg IV push Q5MIN up to a total of 1 mg/kg in DIVIDED doses until target HR/BP met

Protocol

Assessment:

1. Initial Assessment:

- HR, BP, hemodynamic status (including Cardiac output/Cardiac Index, Ejection Fraction, history of COPD/asthma, any signs of ischemia, arrhythmias, increased bleeding).
- Review patient status with charge nurse.
- Ensure that there is a spare temporary pulse generator available at bedside. Check battery status.

2. Ongoing Assessment post intervention:

- Effectiveness of intervention:

The common targets are:

- HR 60 to 70 bpm or as indicated by MD
- SBP: 90 to 120 mmHg or as indicated by MD
- Cardiac Index over 2.2 (without inotropic support) or as indicated by MD

- Assess HR & BP Q15 minutes x 3 then Q1H
- Assess hemodynamic status 30 minutes after administration of beta-blocker & PRN.
- Ensure that there is a spare temporary pulse generator and cables available either at the bedside or in the room. Check battery status.

Interventions:

1. Ensure there is a Physician order

2. Review patient hemodynamics/History to determine whether IV Beta blocker use is appropriate:

- HR
- BP
- Hemodynamic status; including Cardiac Output /Cardiac Index and preload
- Ejection Fraction
- History COPD/asthma

3. Administer medication at appropriate rate.

4. Notify Physician if doses seem ineffective after 2 to 3 doses or if max ordered dose reached
5. If HR less than 60 **after** administration of beta-blocker and patient hemodynamically unstable initiate temporary pacing at the following settings:

Epicardial wires	Mode/Rate	mA	Sensitivity	
AV	DDD – 70 bpm	Atrial: 10 Vent: 10	Atrial: 0.5 Vent: 2.0	Atrial tracking: ON <i>If patient is in AF – turn A tracking OFF</i>
Ventricular	VVI – 70 bpm	Vent: 10	Vent: 2.0	N/A
Atrial	AAI – 70 bpm	Atrial: 10	Atrial: 0.5	N/A

6. If CI falls below 2.2 or systolic BP less than 90 mmHg:

- Hold subsequent doses of IV beta- blocker
- Notify MD and request alternative medication to manage HR or BP
- Consider initiating positive inotrope to increase CI over 2.2 & systolic BP over 90 mmHg

Documentation:

Cerner:

1. Document vital signs in Adult Critical care Quick View
2. Document medication in MAR and “add comment “as needed to capture patient assessment

References:

1. Lexicomp Online Lexi-Drugs Product Monograph: Labetalol, Metoprolol and Esmolol: Accessed October, 2023. Wolters Kluwer Health
2. Parenteral Drug Therapy Manual-Providence Health Care. (Oct, 2023) Metoprolol, Labetolol and Esmolol Monograph

Persons/Groups Consulted:

Cardiac Surgery SPH

Cardiac Anesthesia SPH

CSICU Pharmacist

CSICU staff

Author:

Nurse Educator CSICU

Effective Date:	November 2004
Posted Date:	21-NOV-2023
Last Revised:	21-NOV-2023
Last Reviewed:	21-NOV-2023
Approved By:	PHC
<i>(committee or position)</i>	Professional Practice Standards Committee
Owners:	PHC
<i>(optional)</i>	Cardiac Surgery