

# Enhanced Recovery After Surgery (ERAS) for Minimally Invasive Mitral Valve Repair or Replacement Surgery Clinical Pathway

## Site Applicability

Vancouver General Hospital

## Overall ERAS Goals:

- ↓ stress response to surgery
- Improve patient experience
- ↓ complications and length of stay

## Specific ERAS Goals:

1. Gum chewing x 15–60 minutes while awake, several times/day
2. Advance DAT from POD 0
3. Discontinue CVC POD 2
4. Discontinue indwelling urinary catheter POD 1 by noon
5. Saline lock IV POD 1 or IV at TKVO when drinking greater than or equal to 600mL/12hr
6. Capillary Blood Glucose TID, HS and sliding scale insulin as ordered. If patient non-diabetic and all glucometer readings are less than 8.1mmol/L x 24 hrs, may discontinue glucometer
7. Ondansetron 4mg IV Q8H X 3 doses. First dose 8 hours after intra-op dose.
8. Mobility goals:  
POD 0: Dangle on edge of bed (if extubated) and hemodynamically stable with RN &/or PT  
POD 1: Walk to bathroom and sit up for meals as tolerated; Walk x 3 (minimum 5 meters/walk) a day  
POD 2: Walk to bathroom and sit up for meals; Walk in hallway minimum 3 times per day

## Pathway Patient Goals:

Patient will recover from surgery with an expected 4 – 6 day length of stay (LOS) and experience a safe discharge home.

1. Post-operative complications will be prevented by:
  - Extubation within 4 hours post-op
  - Transfer out of CSICU POD 1
  - Discharge home by POD 4
2. Patient will report pain below 3/10 or adequate for mobilizing and DB+C exercises
3. Effective discharge planning and teaching provided to patient and caregivers for a safe discharge

## Inclusion Criteria

- All patients having **minimally invasive** mitral valve repair or replacement surgery

## Exclusion Criteria

- Open mitral valve procedures, aortic valve, tricuspid, coronary artery bypass surgery, ascending aorta repair, descending aorta repair, TEVAR, robotic assisted procedures

#### CLINICAL PATHWAY

---

##### **Removal Criteria**

- When significant deviations from expected outcomes are noted

##### **Instructions**

- Review pathway once per shift for patient care goals and expected outcomes
- A variance must be documented when expected outcomes have not been met or interventions not given. The variance is documented in the Electronic Health Record (Cerner) each shift until resolved.

# CLINICAL PATHWAY

| Post- Operative Day 0 (CSICU)   |  |
|---|--|
| CARE CATEGORIES<br>DAILY TASKS AND ACTIVITIES   | EXPECTED OUTCOMES  |
| <b>Safety</b>   | <ul style="list-style-type: none"> <li>Beside safety check completed</li> </ul>  |
| <b>Key Diagnostics &amp; Other Assessments</b> <ul style="list-style-type: none"> <li>Blood work, ABG's as per orders</li> <li>Glucose monitoring as per orders</li> <li>Portable CXR, ECG (unless A-V, or V paced) on admission to unit</li> </ul>   | <ul style="list-style-type: none"> <li>The results of the following are within acceptable range: CBC, electrolytes, urea, creatinine, glucose, coagulation status</li> <li>CXR completed and reviewed by MD</li> <li>ECG completed and reviewed by MD</li> </ul>   |
| <b>Central Nervous System</b> <ul style="list-style-type: none"> <li>Sedation and analgesic administered as per orders</li> <li>Monitor the patient as per Pain Assessment and Documentation Standards</li> <li>Delirium screening as per nursing standard</li> </ul>   | <ul style="list-style-type: none"> <li>Patient reports pain control as adequate or 3/10</li> <li>Complete pain assessment as per Pain Assessment and Documentation Standards</li> <li>Nerve block assessment and safety check completed as per protocol</li> <li>No evidence of delirium</li> </ul>                      |
| <b>Cardiovascular System</b> <ul style="list-style-type: none"> <li>Nursing assessment frequency as per nursing standard</li> <li>Maintain CI above 2.2 L/min/m<sup>2</sup></li> <li>Maintain SBP 90 to 120 mmHg (unless otherwise specified)</li> <li>Maintain HR as per order</li> <li>Temporary pacing as per nursing standards</li> <li>Monitor CT drainage with vital signs</li> </ul> | <ul style="list-style-type: none"> <li>Patient in stable cardiac rhythm</li> <li>Normothermic within 2 hours post-op</li> <li>Hemodynamically stable as per Critical Care goals ordered in Patient Care section in Cerner</li> <li>CT drainage less than 150 mL/h for the first 4 hrs; then less than 50 mL/h</li> </ul> |
| <b>Respiratory System</b> <ul style="list-style-type: none"> <li>Maintain PaO<sub>2</sub> above 80 mm Hg</li> <li>Maintain SpO<sub>2</sub> above 92% as per respiratory standard</li> <li>Assess weaning criteria respiratory standard</li> <li>Extubate within 4 hours post-op</li> </ul>  | <ul style="list-style-type: none"> <li>Lung sounds within normal parameters for patient</li> <li>Chlorhexidine mouthwash pre/post extubation</li> <li>Extubate within 4 hours post-op or as assessed</li> </ul>  |
| <b>Gastrointestinal System</b> <ul style="list-style-type: none"> <li>NPO</li> <li>Complete "Adult Swallowing Screen" post extubation as per dysphagia assessment protocol</li> <li>Start diet as ordered if safe post extubation and dysphagia screen</li> </ul>   | <ul style="list-style-type: none"> <li>Nausea and vomiting absent or controlled with antiemetic</li> <li>Nursing Bedside Swallow Screen completed</li> <li>Tolerating clear fluids post extubation and dysphagia screening</li> <li>Gum chewing (15 mins TID) when awake post extubation</li> </ul>                      |
| <b>Genitourinary System</b> <ul style="list-style-type: none"> <li>Maintain urine output between 0.5 to 1 mL/kg/h</li> <li>CAUTI precaution (no dependent loop, secured catheter, change collecting container daily and label)</li> </ul>   | <ul style="list-style-type: none"> <li>Urine output is between 0.5 to 1 mL/kg/h</li> <li>Secure catheter and provide pericare/catheter care Q Shift</li> </ul>   |
| <b>Skin</b> <ul style="list-style-type: none"> <li>Complete skin assessment as per Braden Risk and Skin Assessment with first repositioning post-op</li> <li>Dressings assessed as per nursing standard</li> </ul>  | <ul style="list-style-type: none"> <li>Skin integrity assessed as per Braden Risk Assessment</li> <li>Keep dressings dry and intact. Do not change dressing until POD#3/as per order, unless</li> </ul>  |

# CLINICAL PATHWAY

|  |  |
|--|--|
|  | saturated, otherwise outline drainage with a pen and reinforce as needed.  |
| <b>Mobility</b> <ul style="list-style-type: none"> <li>Falls Risk Assessment prior to first mobilization</li> <li>Dangle and stand if extubated and hemodynamically stable</li> </ul>  | <ul style="list-style-type: none"> <li>Fall prevention assessment completed and care plan in place if indicated</li> <li>Dangle on edge of bed if able with RN or PT</li> </ul>  |
| <b>Medications</b> <ul style="list-style-type: none"> <li>Inotropes titrated to maintain hemodynamic parameters as per orders</li> <li>Insulin sliding scale as ordered</li> <li>Analgesics as ordered</li> </ul>                | <ul style="list-style-type: none"> <li>Inotropes weaned off</li> <li>Blood glucose as per protocol</li> <li>Achieve adequate (or 3/10) pain control with minimal opioids</li> </ul>  |
| <b>Consult</b> <ul style="list-style-type: none"> <li>As needed: POPS</li> </ul>   | <ul style="list-style-type: none"> <li>Consults performed as ordered</li> </ul>  |
| <b>Patient/Family Teaching</b> <ul style="list-style-type: none"> <li>Oriented to plan of care for the next 24 hours</li> <li>Pain scale and use of analgesics</li> <li>Deep breathing and coughing (ICOUGH protocol)</li> </ul> | <ul style="list-style-type: none"> <li>Patient and family understand plan of care</li> <li>Patient &amp; family understand pain control management</li> <li>Patient &amp; family participate in deep breathing and coughing (ICOUGH protocol)</li> </ul> |

## CLINICAL PATHWAY

| Post-Operative Day 1 (CSICU / WARD)  |  |
|--|--|
| CARE CATEGORIES<br>DAILY TASKS AND ACTIVITIES  | EXPECTED OUTCOMES  |
| <b>Safety</b>  | <ul style="list-style-type: none"> <li>Beside safety check completed</li> </ul>  |
| <b>Key Diagnostics &amp; Other Assessments</b> <ul style="list-style-type: none"> <li>Blood work as per orders</li> <li>Nursing assessment frequency as per unit standard: Q4H and PRN</li> <li>In CSICU: vital signs Q1H</li> <li>On ward: vital signs Q4H x 24 hours</li> </ul>  | <ul style="list-style-type: none"> <li>Blood work results are within acceptable range</li> <li>Temp 36° to 37.5° C</li> </ul>  |
| <b>Central Nervous System</b> <ul style="list-style-type: none"> <li>Analgesic administered as ordered</li> <li>Monitor the patient as per Pain Assessment and Documentation Standards</li> <li>Delirium screening as per nursing standard</li> </ul>  | <ul style="list-style-type: none"> <li>Patient reports pain control as adequate or 3/10</li> <li>Complete pain assessment as per Pain Assessment and Documentation Standards</li> <li>Nerve block assessment and safety check completed as per protocol</li> <li>No evidence of delirium</li> </ul>  |
| <b>Cardiovascular System</b> <ul style="list-style-type: none"> <li>Remove PA lines, chest tube(s) and arterial line if hemodynamically stable, as per nursing standard and MD orders</li> <li>Epicardial pacing and care of wires as per nursing standards and as MD orders</li> <li><b>Ward:</b> ECG strips Q12H or with a change in rhythm</li> <li>CXR 2 hours following chest tube removal</li> </ul> | <ul style="list-style-type: none"> <li>Patient in stable intrinsic cardiac rhythm</li> <li>Hemodynamically stable as per Critical Care Goals ordered in Patient Care section in Cerner</li> <li>Invasive monitoring lines removed</li> <li>Chest tubes removed if less than 100ml for over 4 hours</li> <li>Epicardial wires capped</li> <li>Chest X-Ray 2 hours post CT removal and reviewed by MD</li> </ul> |
| <b>Respiratory</b> <ul style="list-style-type: none"> <li>Wean from O<sub>2</sub> and maintain SpO<sub>2</sub> above 92%</li> <li>Deep breathing &amp; coughing Q1H (spirometer)</li> <li>Mouth care: AM and HS + PRN (pneumonia prevention)</li> </ul>  | <ul style="list-style-type: none"> <li>No signs of respiratory complications</li> <li>Patient reminded of mouth care: AM and HS + PRN</li> </ul>   |
| <b>Gastrointestinal System</b> <ul style="list-style-type: none"> <li>Complete "Adult Swallowing Screen" post extubation as per dysphagia assessment protocol if not completed on POD 0</li> <li>Regular Diet (+/- Diabetes no sugar added if diabetic)</li> </ul>   | <ul style="list-style-type: none"> <li>Nursing Bedside Swallow Screen completed</li> <li>Tolerating prescribed regular diet as tolerated</li> <li>Gum chewing (15 mins TID) when awake post extubation</li> <li>No nausea &amp; vomiting</li> <li>Bowel protocol initiated</li> <li>Saline lock IV once patient is drinking well (i.e. 600mL/ 12 hour)</li> </ul>  |
| <b>Genitourinary System</b> <ul style="list-style-type: none"> <li>Daily weight</li> <li>Remove Foley catheter before 12:00pm unless contraindicated</li> <li>In + out catheterization as per order</li> </ul>   | <ul style="list-style-type: none"> <li>Foley removed</li> <li>Voiding without difficulty</li> <li>Patient has an adequate fluid balance. Refer to intake and output documentation</li> <li>Measure urine output Q4H until Foley catheter is removed</li> </ul>   |

# CLINICAL PATHWAY

|   |   |
|---|---|
| <b>Skin</b> <ul style="list-style-type: none"> <li>Dressing assessment and care daily</li> </ul>  | <ul style="list-style-type: none"> <li>Skin integrity assessed as per Braden Risk Assessment</li> <li>No evidence of skin breakdown</li> <li>Dressings dry and intact. Do not change dressing until POD#3/as per order, unless saturated</li> </ul>   |
| <b>Mobility</b> <ul style="list-style-type: none"> <li>Falls Risk Assessment as per protocol</li> <li>Mobilize as per ERAS pathway</li> <li>Up in chair for all meals as tolerated</li> </ul>   | <ul style="list-style-type: none"> <li>Fall prevention assessment completed and careplan in place if indicated</li> <li>Walk x 3 (minimum 5 meters/walk) a day</li> <li>Up in chair for meals as tolerated</li> </ul>   |
| <b>Medications</b> <ul style="list-style-type: none"> <li>Wean inotropes off</li> <li>IV insulin infusion or sliding scale insulin as ordered</li> <li>Assess and initiate anticoagulation as ordered</li> <li>Analgesics as ordered</li> <li>Resume pre-op medications as appropriate</li> </ul>             | <ul style="list-style-type: none"> <li>Inotropes weaned off (document time)</li> <li>Blood glucose as per protocol</li> <li>Anticoagulation as per orders</li> <li>Achieve adequate (or 3/10) pain control with minimal opioids</li> </ul>  |
| <b>Consults</b> <ul style="list-style-type: none"> <li>As needed: Psychiatry, Endocrine, Social Work</li> </ul>   | <ul style="list-style-type: none"> <li>Seen by consultants as ordered</li> </ul>  |
| <b>Patient/Family Teaching</b> <ul style="list-style-type: none"> <li>Oriented to plan of care for the next 24 hours</li> <li>Patient and family reviewing ERAS booklet</li> </ul> <b>Review:</b> <ul style="list-style-type: none"> <li>Pain scale and use of analgesics</li> <li>ICOUGH protocol</li> </ul> | <ul style="list-style-type: none"> <li>Patient and family understand plan of care</li> <li>Patient &amp; family participate in the ICOUGH protocol</li> <li><b>ERAS Minimally Invasive Mitral Valve Surgery Booklet:</b> <ul style="list-style-type: none"> <li>Patient and family have this booklet at bedside or electronic version</li> <li>Reviewed and reinforced pain management</li> </ul> </li> </ul> |
| <b>Discharge Planning</b> <ul style="list-style-type: none"> <li>Discuss length of stay</li> <li>Discuss goals for the day (i.e.: exercises, pain management, rest)</li> <li>Initiate teaching as applicable: anticoagulation, smoking cessation, endocarditis, HF</li> </ul>                                 | <ul style="list-style-type: none"> <li>Assess patient need for additional services</li> <li><b>ERAS Minimally Invasive Mitral Valve Surgery Booklet:</b> <ul style="list-style-type: none"> <li>Patient and family have ERAS booklet at bedside or electronic version</li> <li>Patient reviewed daily goals</li> <li>Patient is aware of discharge criteria</li> </ul> </li> </ul>                            |

# CLINICAL PATHWAY

| Post-Operative Day 2 (Ward)   |  |
|---|--|
| CARE CATEGORIES<br>DAILY TASKS AND ACTIVITIES   | EXPECTED OUTCOMES  |
| <b>Safety</b>   | <ul style="list-style-type: none"> <li>Beside safety check completed</li> </ul>  |
| <b>Key Diagnostics &amp; Other Assessments</b> <ul style="list-style-type: none"> <li>Blood work as ordered</li> <li>MD/NP or Pharmacist to determine target INR and required anticoagulation</li> <li>Nursing assessment Q8H and PRN</li> </ul>                                    | <ul style="list-style-type: none"> <li>MD/NP aware of abnormal results</li> <li>Anticoagulation, INR discussed; target set</li> <li>Temp 36° to 37.5° C</li> </ul>   |
| <b>Central Nervous System</b> <ul style="list-style-type: none"> <li>Delirium screening as per nursing standard</li> <li>Monitor the patient as per Pain Assessment and Documentation Standards</li> <li>Analgesic administered as ordered</li> </ul>                               | <ul style="list-style-type: none"> <li>No evidence of delirium</li> <li>Complete pain assessment as per Pain Assessment and Documentation Standards</li> <li>Patient reports pain control as adequate or 3/10</li> </ul>                               |
| <b>Cardiovascular System</b> <ul style="list-style-type: none"> <li>Vital signs Q6H (0200h assessment at RN discretion)</li> <li>ECG strips Q12H or with a change in rhythm</li> <li>Epicardial pacing and care of wires as per nursing standard and as per MD/NP orders</li> </ul> | <ul style="list-style-type: none"> <li>Vital signs within normal limits for patient</li> <li>Patient in stable intrinsic cardiac rhythm</li> <li>Epicardial wires capped</li> </ul>  |
| <b>Respiratory System</b> <ul style="list-style-type: none"> <li>Wean from O<sub>2</sub> and maintain SpO<sub>2</sub> above 92%</li> </ul>  | <ul style="list-style-type: none"> <li>Mouth care of each meals</li> <li>No signs of respiratory complications</li> </ul>  |
| <b>Gastrointestinal System</b> <ul style="list-style-type: none"> <li>Regular Diet (+/- Diabetes no sugar added if diabetic)</li> <li>If no BM x 24 hrs, follow protocol</li> </ul>   | <ul style="list-style-type: none"> <li>Tolerating prescribed diet</li> <li>Gum chewing (15 mins TID) when awake</li> <li>No nausea &amp; vomiting</li> <li>Bowel movement daily</li> <li>Discontinue CVC if in situ. Insert a peripheral IV</li> </ul> |
| <b>Genitourinary System</b> <ul style="list-style-type: none"> <li>Daily weight</li> </ul>  | <ul style="list-style-type: none"> <li>Voiding without difficulty</li> <li>Patient has an adequate fluid balance. Refer to intake and output documentation</li> </ul>  |
| <b>Skin</b> <ul style="list-style-type: none"> <li>Incision assessment and care daily</li> </ul>  | <ul style="list-style-type: none"> <li>No evidence of skin breakdown</li> <li>Skin integrity assessed as per Braden Risk Assessment</li> <li>Dressings dry and intact. Do not change dressing until POD#3/as per order, unless saturated</li> </ul>    |
| <b>Mobility</b> <ul style="list-style-type: none"> <li>Falls Risk Assessment as per protocol</li> <li>Mobilize as per ERAS pathway</li> <li>May shower (insulate epicardial wires)</li> </ul>   | <ul style="list-style-type: none"> <li>Fall prevention assessment completed and care plan in place if indicated</li> <li>Walk in hallway minimum 3 times a day</li> <li>Up in chair for meals or TID and to washroom PRN</li> </ul>                    |
| <b>Medications</b> <ul style="list-style-type: none"> <li>Anticoagulation initiated as ordered</li> <li>Glycemic control as per orders</li> <li>Analgesics as ordered</li> <li>Diuresis to target weight as per orders</li> </ul>   | <ul style="list-style-type: none"> <li>Anticoagulation initiated as per MRP orders</li> <li>Blood glucose as protocol</li> <li>Achieve adequate (or 3/10) pain control with minimal opioids</li> </ul>   |

## CLINICAL PATHWAY

|  |   |
|--|---|
| <b>Consults</b> <ul style="list-style-type: none"> <li>Reassess need for additional consults</li> </ul>  | <ul style="list-style-type: none"> <li>No additional consults required</li> <li>New consults initiated as ordered</li> </ul>  |
| <b>Patient/Family Teaching Review:</b> <ul style="list-style-type: none"> <li>Incision care</li> <li>Mood changes</li> <li>Deep breathing and coughing</li> <li>Pain management</li> <li>Delirium</li> </ul>   | <b>ERAS Minimally Invasive Valve Surgery Booklet:</b> <ul style="list-style-type: none"> <li>Patient and family have reviewed this booklet with nurse and understand post-op care and management</li> <li>Reviewed and reinforced pain management</li> </ul>  |
| <b>Discharge Planning</b> <ul style="list-style-type: none"> <li>Discuss length of stay</li> <li>Goals of the day</li> <li>Who is your support person when you are going to be home?</li> <li>Continue teaching as applicable: anticoagulation, smoking cessation, endocarditis, HF</li> </ul> | <ul style="list-style-type: none"> <li>Discussion on these topics took place</li> <li><b>ERAS Minimally Invasive Mitral Valve Surgery Booklet:</b> <ul style="list-style-type: none"> <li>Patient and family have reviewed daily goals</li> <li>Patient is aware of discharge criteria</li> </ul> </li> </ul> |



# CLINICAL PATHWAY

| Post-Operative Day 3   |   |
|--|---|
| CARE CATEGORIES<br>DAILY TASKS AND ACTIVITIES  | EXPECTED OUTCOMES   |
| <b>Safety</b>  | <ul style="list-style-type: none"> <li>Beside safety check completed</li> </ul>   |
| <b>Key Diagnostics &amp; Other Assessments</b> <ul style="list-style-type: none"> <li>Blood work as per orders</li> <li>MD/NP or pharmacist to determine target INR and required anticoagulation</li> <li>Nursing assessment Q12H and PRN</li> </ul>     | <ul style="list-style-type: none"> <li>MD/NP aware of abnormal results</li> <li>Temp 36° to 37.5° C</li> <li>INR at target</li> </ul>   |
| <b>Central Nervous System</b> <ul style="list-style-type: none"> <li>Delirium screening as per nursing standard</li> <li>Monitor the patient as per Pain Assessment and Documentation Standards</li> <li>Analgesic administered as per orders</li> </ul> | <ul style="list-style-type: none"> <li>No evidence of delirium</li> <li>Complete Pain assessment as per Pain Assessment and Documentation Standards</li> <li>Patient reports pain control as adequate or 3/10</li> </ul>                            |
| <b>Cardiovascular System</b> <ul style="list-style-type: none"> <li>Vital signs Q8H</li> <li>ECG strips Q12H and with a change in rhythm</li> <li>Epicardial pacing and care of wires as per nursing standards and as MD/NP orders</li> </ul>            | <ul style="list-style-type: none"> <li>Vital signs within normal limits for patient</li> <li>Patient in stable intrinsic cardiac rhythm</li> <li>Epicardial pacing wires removed by MD/NP</li> </ul>  |
| <b>Respiratory System</b> <ul style="list-style-type: none"> <li>Maintain SpO<sub>2</sub> above 92% on room air</li> </ul>   | <ul style="list-style-type: none"> <li>Mouth care after each meals</li> <li>No signs of respiratory complications</li> </ul>  |
| <b>Gastrointestinal System</b> <ul style="list-style-type: none"> <li>Regular Diet (+/- Diabetes no sugar added if diabetic)</li> <li>If no BM x 48 hrs, follow protocol and notify MD/NP</li> </ul>   | <ul style="list-style-type: none"> <li>Tolerating diet</li> <li>Gum chewing (15 mins TID) when awake</li> <li>No nausea &amp; vomiting</li> <li>Bowel movement daily</li> </ul>   |
| <b>Genitourinary System</b> <ul style="list-style-type: none"> <li>Daily weight</li> </ul>   | <ul style="list-style-type: none"> <li>Voiding without difficulty</li> <li>Patient has an adequate fluid balance. Refer to intake and output documentation</li> </ul>   |
| <b>Skin</b> <ul style="list-style-type: none"> <li>Incision assessment and care daily</li> </ul>   | <ul style="list-style-type: none"> <li>Skin integrity assessed as per Braden Risk Assessment</li> <li>No evidence of skin breakdown</li> <li>Mepilex removed incision cleaned, well approximated, dry + intact. Incision left exposed</li> </ul>    |
| <b>Mobility</b> <ul style="list-style-type: none"> <li>Falls Risk Assessment as per protocol</li> <li>May shower (insulate epicardial wires)</li> <li>Encourage independent mobilization</li> </ul>  | <ul style="list-style-type: none"> <li>Fall prevention assessment completed and care plan in place if indicated</li> <li>Walk independently in hallway minimum 3 times per day</li> <li>Up in chair for meals or TID and to washroom PRN</li> </ul> |
| <b>Medications</b> <ul style="list-style-type: none"> <li>Anticoagulation as per orders</li> <li>Glycemic control as per orders</li> <li>Analgesics as ordered</li> <li>Diuresis to target weight as per orders</li> </ul>                               | <ul style="list-style-type: none"> <li>Achieve adequate (or 3/10) pain control with minimal opioids</li> </ul>  |
| <b>Consults</b> <ul style="list-style-type: none"> <li>As needed: Psychiatry, SW, Pastoral Care, Nephrology</li> </ul>   | <ul style="list-style-type: none"> <li>No additional consults required</li> <li>New consults initiated as ordered</li> </ul>  |

This material has been prepared solely for use at Vancouver Coastal Health (VCH). VCH accept no responsibility for use of this material by any person or organization not associated with VCH. A printed copy of this document may not reflect the current electronic version.

CLINICAL PATHWAY

|  |   |
|--|---|
| <p><b>Patient/Family Teaching</b></p> <ul style="list-style-type: none"> <li>Review previous topics and educate as needed</li> </ul> <p><b>New topics:</b></p> <ul style="list-style-type: none"> <li>Heart Healthy diet</li> <li>Resuming sex</li> <li>Risk factor counseling</li> <li>Returning to work</li> <li>Driving</li> <li>Sleep Hygiene</li> <li>Activity after discharge</li> </ul> | <ul style="list-style-type: none"> <li><b>ERAS Minimally Invasive Mitral Valve Surgery Booklet:</b> <ul style="list-style-type: none"> <li>Patient and family have reviewed this booklet with nurse and understand post-op care and management</li> </ul> </li> </ul>   |
| <p><b>Discharge Planning</b></p> <ul style="list-style-type: none"> <li>Discuss transportation plans</li> <li>Arrange PT/OT equipment PRN</li> <li>Coordinate TST needs with CML</li> </ul>  | <ul style="list-style-type: none"> <li>Discussion on these topics took place</li> <li><b>ERAS Minimally Invasive Mitral Valve Surgery Booklet:</b> <ul style="list-style-type: none"> <li>Patient and family have reviewed daily goals</li> <li>Patient is aware of discharge criteria</li> </ul> </li> </ul> |

# CLINICAL PATHWAY

| Post-Operative Day 4   |  |
|--|--|
| CARE CATEGORIES<br>DAILY TASKS AND ACTIVITIES  | EXPECTED OUTCOMES  |
| <b>Safety</b>  | <ul style="list-style-type: none"> <li>Beside safety check completed</li> </ul>  |
| <b>Key Diagnostics &amp; Other Assessments</b> <ul style="list-style-type: none"> <li>MD/NP or pharmacist to determine target INR and required anticoagulation</li> <li>Nursing assessment Q12H and PRN</li> <li>Vital signs Q12H <b>unless</b> otherwise indicated</li> </ul>   | <ul style="list-style-type: none"> <li>INR at target</li> <li>Temp 36° to 37.5° C</li> </ul>   |
| <b>Central Nervous System</b> <ul style="list-style-type: none"> <li>Delirium screening as per nursing standard</li> <li>Monitor the patient as per Pain Assessment and Documentation Standards</li> <li>Analgesic administered as per orders</li> </ul>   | <ul style="list-style-type: none"> <li>No evidence of delirium</li> <li>Complete pain assessment as per Pain Assessment and Documentation Standards</li> <li>Patient reports pain control as adequate or 3/10</li> </ul>   |
| <b>Cardiovascular System</b> <ul style="list-style-type: none"> <li>ECG strips Q12H or with a change in rhythm</li> <li>Epicardial pacing wires removed by MD/NP with nursing care as per standard (IV saline lock remains until discharge)</li> <li>Discontinue telemetry if NSR x 24 hours</li> </ul>                                    | <ul style="list-style-type: none"> <li>Vital signs within normal limits for patient</li> <li>Patient in stable intrinsic cardiac rhythm</li> <li>Epicardial pacing wires removed</li> </ul>  |
| <b>Respiratory</b> <ul style="list-style-type: none"> <li>Maintain SpO<sub>2</sub> above 92% on room air</li> </ul>  | <ul style="list-style-type: none"> <li>Mouth care after each meals</li> <li>No signs of respiratory complications</li> </ul>   |
| <b>Gastrointestinal System</b> <ul style="list-style-type: none"> <li>Diet as ordered</li> <li>If no BM x 72 hrs, follow protocol and notify MD/NP</li> </ul>  | <ul style="list-style-type: none"> <li>Tolerating diet</li> <li>No nausea &amp; vomiting</li> <li>Gum chewing (15 mins TID) when awake</li> <li>Bowel movement daily</li> </ul>  |
| <b>Genitourinary System</b> <ul style="list-style-type: none"> <li>Daily weight</li> </ul>   | <ul style="list-style-type: none"> <li>Voiding without difficulty</li> </ul>   |
| <b>Skin</b> <ul style="list-style-type: none"> <li>Incision assessment and care daily</li> <li>Discuss removal of surgical clips/staples with MD/NP</li> <li>Remove chest tube sutures 3 days following chest tube removal if site is dry and well approximated or as otherwise directed by MD/NP. Apply steri-strips as needed</li> </ul> | <ul style="list-style-type: none"> <li>Skin integrity assessed as per Braden Risk Assessment</li> <li>No evidence of skin breakdown</li> <li>Surgical incision well approximated, dry and intact</li> <li>Chest tube sutures removed, incision well approximated, and sutures applied</li> </ul> |
| <b>Mobility</b> <ul style="list-style-type: none"> <li>Falls Risk Assessment as per protocol</li> <li>Encourage mobilization</li> </ul>  | <ul style="list-style-type: none"> <li>Fall prevention assessment completed and care plan in place if indicated</li> <li>Walk independently in hallway minimum 3 times a day</li> <li>Up in chair for meals or TID and to washroom PRN</li> </ul>  |
| <b>Medications</b> <ul style="list-style-type: none"> <li>Anticoagulation as per orders</li> <li>Glycemic control as per pre-printed orders</li> <li>Analgesics as ordered</li> <li>Diuresis to target weight as per orders</li> </ul>   | <ul style="list-style-type: none"> <li>Achieve adequate (or 3/10) pain control with minimal opioids</li> </ul>   |
| <b>Consults</b> <ul style="list-style-type: none"> <li>Reassess need for additional consults</li> </ul>  | <ul style="list-style-type: none"> <li>No additional consults required</li> <li>New consults initiated as ordered</li> </ul>   |

# CLINICAL PATHWAY

|  |   |
|--|---|
| <p><b>Patient/Family Teaching</b></p> <ul style="list-style-type: none"> <li>• Review any topics patient and family have questions about and education as needed</li> <li>• <b>Review:</b> Medications</li> <li>• Cardiac Rehab program</li> <li>• When to call doctor or 911</li> </ul> | <ul style="list-style-type: none"> <li>• <b>ERAS Minimally Invasive Mitral Valve Surgery Booklet:</b> <ul style="list-style-type: none"> <li>○ Patient and family have reviewed this booklet with nurse and understand post-op care and management</li> </ul> </li> </ul>   |
| <p><b>Discharge Planning</b></p> <ul style="list-style-type: none"> <li>• Discharge teaching</li> <li>• Provide: “My care Plan”, and discharge prescription if applicable</li> <li>• Patient aware of follow up information</li> </ul>   | <ul style="list-style-type: none"> <li>• Discharge teaching done</li> <li>• Documentation given</li> <li>• <b>ERAS Minimally Invasive Mitral Valve Surgery Booklet:</b> <ul style="list-style-type: none"> <li>○ Patient and family reviewed daily goals</li> <li>○ Patient is aware of discharge criteria</li> </ul> </li> </ul> |

## CLINICAL PATHWAY

| Post-Operative Additional Day  |   |
|--|---|
| CARE CATEGORIES<br>DAILY TASKS AND ACTIVITIES  | EXPECTED OUTCOMES   |
| <b>Safety</b>  | <ul style="list-style-type: none"> <li>Beside safety check completed</li> </ul>   |
| <b>Key Diagnostics &amp; Other Assessments</b> <ul style="list-style-type: none"> <li>MD/NP or pharmacist to determine target INR and required anticoagulation</li> </ul>  | <ul style="list-style-type: none"> <li>INR at target</li> </ul>   |
| <b>Central Nervous System</b> <ul style="list-style-type: none"> <li>Delirium screening as per nursing standard</li> <li>Monitor the patient as per Pain Assessment and Documentation Standards</li> <li>Analgesic administered as per orders</li> </ul>         | <ul style="list-style-type: none"> <li>No evidence of delirium</li> <li>Complete pain assessment as per Pain Assessment and Documentation Standards</li> <li>Patient reports pain control as adequate or 3/10</li> </ul>  |
| <b>Cardiovascular System</b> <ul style="list-style-type: none"> <li>Nursing assessment Q12H and PRN</li> <li>Vital signs Q12H</li> <li>Epicardial pacing and care of wires as per nursing standards and as MD/NP orders</li> </ul>                               | <ul style="list-style-type: none"> <li>Vital signs within normal limits for patient</li> <li>Epicardial pacing wires removed by MD/NP</li> <li>Patient in stable intrinsic cardiac rhythm</li> </ul>  |
| <b>Respiratory</b>   | <ul style="list-style-type: none"> <li>Mouth care after each meals</li> <li>No signs of respiratory complications</li> </ul>  |
| <b>Gastrointestinal System</b>   | <ul style="list-style-type: none"> <li>Tolerating diet</li> <li>No nausea &amp; vomiting</li> <li>Gum chewing (15 mins TID) when awake</li> <li>Bowel movement daily</li> </ul>   |
| <b>Genitourinary System</b> <ul style="list-style-type: none"> <li>Daily weight</li> </ul>   | <ul style="list-style-type: none"> <li>Voiding without difficulty</li> </ul>  |
| <b>Skin</b> <ul style="list-style-type: none"> <li>Surgical incision exposed to air</li> </ul>   | <ul style="list-style-type: none"> <li>Skin integrity assessed as per Braden Risk Assessment</li> <li>No evidence of skin breakdown</li> <li>Incisions dry and intact</li> </ul>  |
| <b>Mobility</b> <ul style="list-style-type: none"> <li>Falls Risk Assessment as per protocol</li> <li>Activity as tolerated</li> <li>Independent personal care</li> </ul>  | <ul style="list-style-type: none"> <li>Fall prevention assessment completed and care plan in place if indicated</li> <li>Patient independent with personal care and walking as tolerated</li> </ul>   |
| <ul style="list-style-type: none"> <li><b>Medications</b></li> <li>Anticoagulation as per orders</li> <li>Glycemic control as per orders</li> <li>Analgesics as ordered</li> <li>Diuresis to target weight as per orders</li> </ul>                              | <ul style="list-style-type: none"> <li>Achieve adequate (or 3/10) pain control with minimal opioids</li> </ul>  |
| <b>Consults</b> <ul style="list-style-type: none"> <li>Reassess need for additional consults</li> </ul>  | <ul style="list-style-type: none"> <li>No additional consults required</li> <li>New consults initiated as ordered</li> </ul>  |
| <b>Patient/Family Teaching</b> <ul style="list-style-type: none"> <li>Review any topics patient and family have questions about and educate as needed</li> <li>Review: Medications</li> <li>Cardiac Rehab program</li> <li>When to call doctor or 911</li> </ul> | <ul style="list-style-type: none"> <li><b>ERAS Minimally Invasive Mitral Valve Surgery Booklet:</b> <ul style="list-style-type: none"> <li>Patient and family have reviewed this booklet with nurse and understand post-op care and management</li> </ul> </li> </ul> |

# CLINICAL PATHWAY

|   |   |
|---|---|
| <b>Discharge Planning</b> <ul style="list-style-type: none"> <li>Discharge teaching</li> <li>Provide: "My care Plan", and discharge prescription if applicable</li> <li>Patient aware of follow up information</li> </ul> | <ul style="list-style-type: none"> <li>Discharge teaching done</li> <li>Documentation given</li> <li><b>ERAS Minimally Invasive Mitral Valve Surgery Booklet:</b> <ul style="list-style-type: none"> <li>Patient and family reviewed daily goals</li> <li>Patient is aware of discharge criteria</li> </ul> </li> </ul> |
|---|---|

Developed By

|                        |   |
|------------------------|---|
| <b>Effective Date:</b> |   |
| <b>Posted Date:</b>    |   |
| <b>Last Revised:</b>   |   |
| <b>Last Reviewed:</b>  |   |
| <b>Approved By:</b>    | VCH   |
|                        | <b>Endorsed By:</b> <ul style="list-style-type: none"> <li>Gurdip Bhatti, Clinical Nurse Educator, Cardiac Surgery Intensive Care Unit, VGH</li> <li>Tina Oye, Clinical Nurse Educator, Cardiac Ward, VGH</li> </ul>                                    |
|                        | <b>Final Sign Off:</b>  |
| <b>Owners:</b>         | VCH   |
|                        | Cardiac Surgery<br><b>Developer Lead(s):</b> <ul style="list-style-type: none"> <li>Gurdip Bhatti, Clinical Nurse Educator, Cardiac Surgery Intensive Care Unit, VGH</li> <li>Tina Oye, Clinical Nurse Educator, Cardiac Ward, VGH</li> <li></li> </ul> |