

Anaphylaxis: Initial Emergency Management (Adult and Pediatric)

Site Applicability

All VCH & PHC sites – Acute, Community and Long-term Care

Practice Level

Profession	Advanced Skill (requiring additional education)
RN and RPN	Nurse Independent Activity: Upon successful completion of additional education , may: <ul style="list-style-type: none"> Administer epinephrine IM to treat anaphylaxis Administer oxygen to manage hypoxia Approved sites in Acute Care only: Emergency performance of venipuncture to establish intravenous (IV) access
LPN	Nurse Independent Activity: Upon successful completion of additional education , may: <ul style="list-style-type: none"> Administer epinephrine IM to treat anaphylaxis Administer oxygen to maintain SpO₂ above 92 percent With an order: <ul style="list-style-type: none"> Approved sites in Acute Care only: Emergency performance of venipuncture to establish intravenous (IV) access

Education

- [Understanding Autonomous Practice and Nurse Independent Activities \(NIA\) or Nurse-Initiated Protocols \(NIP\)](#) on LearningHub (Recommended)
- Provincial Education [Anaphylaxis – Initial Emergency Treatment by Nurses \(Adult and Pediatric\)](#) on LearningHub (**Required**)
- Approved sites in Acute Care performing venipuncture to establish IV access: with education per [IV Therapy, Peripheral: Insertion, Care and Maintenance \(BD-00-12-40080\)](#)

Requirements

- Nurse Independent Activities (NIAs) for RNs, RPNs, and LPNs is supported as per policy:
 - [Nurse Independent Activities \(NIA\) and Nurse Initiated Protocols \(NIP\) \(BCD-11-11-40001\)](#)
- Physician or Nurse Practitioner (NP) orders override the use of NIA.

This material has been prepared solely for use at Providence Health Care (PHC) and Vancouver Coastal Health (VCH). PHC and VCH accept no responsibility for use of this material by any person or organization not associated with PHC and VCH. A printed copy of this document may not reflect the current electronic version.

- Patients who have been treated for anaphylaxis must have immediate definitive follow up by a physician or NP. In the community and ambulatory settings, patients who have been treated for anaphylaxis must be transferred to hospital by Emergency Health Services (EHS).

Procedure

Follow Provincial Clinical Decision Support Tool [Anaphylaxis: Initial Emergency Treatment by Nurses \(Adult and Pediatric\)](#) - see [Appendix A](#) in this document for the provincial DST if web link broken.

Documentation

- Document care per site specific standards; in addition, document the NIAs as per Policy: [Nurse Independent Activities \(NIA\) and Nurse-Initiated Protocols \(NIP\)](#).
- Cerner: Document in the NIA PowerForms:
 - *NIA Anaphylaxis: Initial Treatment (Adult)*
 - *PED NIA Anaphylaxis: Initial Treatment (Pediatric)*
- Document any patient teaching provided.
- Ensure client allergy status is updated in the client health record.
- If anaphylaxis is believed to be related to drug administration, including immunization, report the adverse events:
 - Vaccines: BC Centre for Disease Control
 - [Adverse Event Following Immunization \(AEFI\) Case Report Form](#)
 - Medications other than vaccines:
 - [VCH: BCPSLS](#)
 - [PHC: BCPSLS](#)

Related Documents

- [Policy: Nurse Independent Activities \(NIA\) and Nurse Initiated Protocols \(NIP\) \(BCD-11-11-40001\)](#)
- [Post-Anaphylaxis teaching and discharge instructions: \(FF.400.AL54\)](#)
- Additional information about Anaphylaxis:
 - [Lexicomp Monograph](#)
 - [Pathophysiology \(Up-to-Date\): Anaphylaxis Pathophysiology](#)
 - [World Allergy Organization Anaphylaxis Guidelines](#)

References

See references in Provincial Clinical Decision Support Tool [Anaphylaxis: Initial Emergency Treatment by Nurses \(Adult and Pediatric\)](#)

Appendices

[Appendix A: Anaphylaxis: Initial Emergency Treatment by Nurses \(Adult and Pediatric\) Clinical Decision Support Tool](#)

First Released Date:	16-OCT-2020	
Posted Date:	16-OCT-2020	
Last Revised:	26-MAR-2021	
Last Reviewed:	16-OCT-2020	
Approved By: <i>(committee or position)</i>	PHC	VCH
	Endorsed By: Professional Practice Standards Committee	Endorsed By: (Regional SharePoint 2nd Reading) Health Authority Profession Specific Advisory Council Chairs (HAPSAC) Health Authority and Area Specific Interprofessional Advisory Council Chairs (HAIAC) Operations Directors Professional Practice Directors Final Sign Off: Vice President, Professional Practice and Chief Clinical Information Officer, VCH
Owners: <i>(optional)</i>	PHC / VCH	
	Developer Lead(s): <ul style="list-style-type: none"> Practice Initiatives Lead, Professional Practice, VC Practice Initiatives Lead, Professional Practice, VC Development Team members: <ul style="list-style-type: none"> Regional Immunization CRN, Communicable Disease Control, VCH Primary Care Clinical Educator, Primary Care, VC 	

Date of Approval/Review/Revision

Original Posted Date: August 16, 2018

First Posted Date: August 16, 2018

Revised Date: October 16, 2020

This material has been prepared solely for use at Providence Health Care (PHC) and Vancouver Coastal Health (VCH). PHC and VCH accept no responsibility for use of this material by any person or organization not associated with PHC and VCH. A printed copy of this document may not reflect the current electronic version.

Appendix A: Anaphylaxis: Initial Emergency Treatment by Nurses (Adult and Pediatric)

Provincial Clinical Decision Support Tool

October 16, 2020

Original source of document: [Anaphylaxis Provincial DST](#)

Contents

1.0	Site Applicability	2
2.0	Scope of Practice Limits and Conditions	2
2.1	Conditions on Practice	2
A.	Administration of epinephrine	2
B.	Administration of oxygen	2
3.0	Policy Statement	2
4.0	Need to Know and Key Points	3
5.0	Assessment and Nursing Diagnosis of Anaphylaxis	3
6.0	Anaphylaxis Response Kit Contents	6
	Client Owned Auto-injector of Epinephrine	6
7.0	Injection Location Considerations	6
8.0	Intervention: Initial Emergency Treatment of Anaphylaxis	7
9.0	Client / Family Education	8
10.0	Documentation	8
11.0	Reporting	8
	References	9
	Appendix A: Common Triggers for Anaphylaxis	11
	Appendix B: EPINEPHrine 1mg/ml Dose by Age Table	12

Maintained by:	HPA Leads (Nursing) – Anaphylaxis Working Group					
Issuing Authority:	Island Health – Professional Practice and Learning and Performance Support					
Last Revised:	2021-Mar-26	Last Reviewed:	2019-Feb-01	First Issued:	2019-Feb-01	Page 1 of 16

1.0 Site Applicability

If anaphylaxis is suspected, based upon clinical presentation and possible exposure to a trigger, treatment should be provided as outlined in this document.

All Acute, Community, Long Term Care sites and non-hospital settings

Exception: when an alternate practice standard or procedure, clinical decision support tool or medical order is in effect for initial emergency treatment of anaphylaxis.

Note: All nurses (RN, RPN and LPN) who immunize without an order must follow the Decision Support Tools (DST) – BC Communicable Disease Control Manual, Chapter 2: Immunization; Part 3 Management of Anaphylaxis in a Non-Hospital Setting established by the BC Centre for Disease Control (BCCDC). The clinical direction in both of these documents is identical.

In this document *client* refers to patient, client or resident.

2.0 Scope of Practice Limits and Conditions

This decision support tool is intended for use by nurses.

Following an assessment and nursing diagnosis of anaphylaxis, Registered Nurses (RN), Registered Psychiatric Nurses (RPN) and Licensed Practical Nurses (LPN) may carry out the initial emergency management of anaphylaxis:

- regardless of the causative agent;
- without an order;
- across all healthcare settings (i.e. hospital and non-hospital)

2.1 Conditions on Practice

A. Administration of epinephrine

Prior to administering epinephrine for the emergency management of anaphylaxis:

- 1) RN, RPN and LPN must follow this decision support tool.
- 2) RPN and LPN must successfully complete additional education [**Anaphylaxis Initial Emergency Treatment by Nurses (Adult and Pediatric)**]. Available on [Learning Hub](#)

This education is recommended for RNs.

B. Administration of oxygen

Health Authority or employer practice limits may apply.

If available in the practice setting, prior to initiating oxygen therapy:

- 1) LPNs must follow a decision support tool and complete additional education.

3.0 Policy Statement

Clients who have been treated for anaphylaxis must have immediate follow up by a physician or Nurse Practitioner.

In the community, long term care and ambulatory settings, clients who have been treated for anaphylaxis must be transferred to hospital via ambulance.

Maintained by:	HPA Leads (Nursing) – Anaphylaxis Working Group					
Issuing Authority:	Island Health – Professional Practice and Learning and Performance Support					
Last Revised:	2021-Mar-26	Last Reviewed:	2019-Feb-01	First Issued:	2019-Feb-01	Page 2 of 16

4.0 Need to Know and Key Points

Anaphylaxis occurs with exposure to a trigger (see [Appendix A](#)) in a susceptible individual. Onset of symptoms usually occurs in minutes but can occur hours after exposure to a trigger.

Death from anaphylaxis *may* occur as a result of severe respiratory complications, cardiovascular collapse, or both.

Early administration of intramuscular (IM) Epinephrine is first line treatment for anaphylaxis to prevent death and there is no known equivalent substitute.

There is no contra-indication to epinephrine administration in anaphylaxis.

Epinephrine:

- Acts on smooth muscle of the bronchial tree reducing bronchospasm
- Counteracts histamine-induced vasodilation
- Increases cardiac output
- Reduces histamine release

Note: Diphenhydramine (Benadryl) is **NOT INDICATED** in anaphylaxis.

Antihistamines are not indicated as initial first line treatment in the emergency management of anaphylaxis as there is no effect on respiratory or cardiovascular symptoms and they are of little clinical importance in life-threatening anaphylaxis based on current evidence.

H1 antihistamines (e.g. Benadryl) relieve localized and less severe systemic allergic reactions and the only useful clinical effect is the improvement of itch and hives.^{4, 13, 14}

5.0 Assessment and Nursing Diagnosis of Anaphylaxis

Early recognition of anaphylaxis is essential to ensure timely intervention.

Assess the client for signs and symptoms of anaphylaxis. These generally involve two or more body systems. See **Table 1** for clinical scenarios and body systems involved.

IMPORTANT: Anaphylaxis can occur **without** presence of hives.

Table 1: Clinical Scenarios and Body Systems Involved with Anaphylaxis.

Clinical Scenario	(1) No Clear Trigger	(2) Suspected Trigger (new food, drug or immunization)	(3) Accidental Exposure to Known Allergen (Same as column 2 or ↓ BP)
SIGNS and SYMPTOMS <i>Onset Minutes to Hours</i>	Skin or mucosal or both + Plus at least one of the following: <ul style="list-style-type: none"> • Respiratory Compromise • Reduced Blood Pressure or Associated Symptoms 	TWO OR MORE of the following: <ul style="list-style-type: none"> • Skin or mucosal • Respiratory Compromise • Reduced Blood Pressure or Associated S/S • Persistent GI 	REDUCED BLOOD PRESSURE ONLY <ul style="list-style-type: none"> • Child – Low Systolic or decrease greater than 30 percent • Adult – Systolic 90 or decrease greater than 30 percent from baseline

Refer to Image 1: **World Allergy Organization Anaphylaxis Guideline Poster** on page 4 of this document for a detailed description of signs and symptoms to inform the assessment and nursing diagnosis of anaphylaxis.

Maintained by:	HPA Leads (Nursing) – Anaphylaxis Working Group				
Issuing Authority:	Island Health – Professional Practice and Learning and Performance Support				
Last Revised:	2021-Mar-26	Last Reviewed:	2019-Feb-01	First Issued:	2019-Feb-01

Image 1: World Allergy Organization Anaphylaxis Guideline Poster

Anaphylaxis is highly likely when any one of the following three criteria is fulfilled:

1 Sudden onset of an illness (minutes to several hours), with involvement of the skin, mucosal tissue, or both (e.g. generalized hives, itching or flushing, swollen lips-tongue-uvula)

AND AT LEAST ONE OF THE FOLLOWING:

Sudden respiratory symptoms and signs
(e.g. shortness of breath, wheeze, cough, stridor, hypoxemia)

Sudden reduced BP or symptoms of end-organ dysfunction (e.g. hypotonia [collapse], incontinence)

OR 2 Two or more of the following that occur suddenly after exposure to a *likely allergen or other trigger** for that patient (minutes to several hours):

Sudden skin or mucosal symptoms and signs
(e.g. generalized hives, itch-flush, swollen lips-tongue-uvula)

Sudden respiratory symptoms and signs
(e.g. shortness of breath, wheeze, cough, stridor, hypoxemia)

Sudden reduced BP or symptoms of end-organ dysfunction (e.g. hypotonia [collapse], incontinence)

Sudden gastrointestinal symptoms (e.g. crampy abdominal pain, vomiting)

OR 3 Reduced blood pressure (BP) after exposure to a *known allergen*** for that patient (minutes to several hours):

Infants and children: low systolic BP (age-specific) or greater than 30% decrease in systolic BP***

Adults: systolic BP of less than 90 mm Hg or greater than 30% decrease from that person's baseline

* For example, immunologic but IgE-independent, or non-immunologic (direct mast cell activation)
 ** For example, after an insect sting, reduced blood pressure might be the only manifestation of anaphylaxis; or, after allergen immunotherapy, generalized hives might be the only initial manifestation of anaphylaxis.
 *** Low systolic blood pressure for children is defined as less than 70 mm Hg from 1 month to 1 year, less than (70 mm Hg + [2 x age]) from 1 to 10 years, and less than 90 mm Hg from 11 to 17 years. Normal heart rate ranges from 80-140 beats/minute at age 1-2 years; from 80-120 beats/minute at age 3 years; and from 70-115 beats/minute after age 3 years. In infants and children, respiratory compromise is more likely than hypotension or shock, and shock is more likely to be manifest initially by tachycardia than by hypotension.

(Adapted from Canadian Pediatric Society, 2018)

Anaphylaxis must be distinguished from fainting (vasovagal syncope) and anxiety (panic attack). See Table 2: Signs and Symptoms of Anaphylaxis versus Fainting and Anxiety.

Maintained by:	HPA Leads (Nursing) – Anaphylaxis Working Group				
Issuing Authority:	Island Health – Professional Practice and Learning and Performance Support				
Last Revised:	2021-Mar-26	Last Reviewed:	2019-Feb-01	First Issued:	2019-Feb-01

Table 2: Signs and Symptoms of Anaphylaxis versus Fainting and Anxiety

	Anaphylaxis	Fainting	Anxiety
Definitions	A potentially life threatening allergic reaction that is rapid in onset and progression of symptoms.	Temporary unconsciousness caused by diminished blood supply to the brain due to painful stimuli or emotional reaction.	Protective physiological state recognized as fear, apprehension, or worry
Onset	<ul style="list-style-type: none"> rapid onset and progression of symptoms occurs minutes to hours after exposure to trigger recovery dependent on response to treatment 	<ul style="list-style-type: none"> sudden onset occurs before, during or shortly after trigger (e.g. sight of the needle) recovery occurs within 1 to 2 minutes 	<ul style="list-style-type: none"> sudden onset occurs before, during, or shortly after trigger (e.g. sight of the needle) recovery generally occurs within 1 to 2 minutes
Skin/Mucosal	<ul style="list-style-type: none"> localized subcutaneous (or sub mucosal) swelling and tingling to face and mouth hives – may be delayed warm, itchy, red and blotchy 	<ul style="list-style-type: none"> pale excessive perspiration cold, clammy 	<ul style="list-style-type: none"> pale excessive perspiration cold, clammy
Respiratory	<ul style="list-style-type: none"> labored breathing - hoarse voice, throat tightness, rapid breathing, wheezing, coughing, nasal flaring, nasal and chest congestion rhinitis (stuffy or runny nose, itchy watery eyes and sneezing) shortness of breath, stridor, retractions, chest pain and cyanosis 	<ul style="list-style-type: none"> breathing normal or shallow, irregular and labored 	<ul style="list-style-type: none"> breathing rapid and shallow (hyperventilation) breath-holding in children
Cardiovascular	<ul style="list-style-type: none"> weak and rapid pulse hypotension alone after an exposure can represent anaphylaxis hypotension is less common in children shock 	<ul style="list-style-type: none"> slow, steady pulse decreased systolic and diastolic 	<ul style="list-style-type: none"> rapid pulse normal or elevated systolic
Gastrointestinal	<ul style="list-style-type: none"> nausea, vomiting, diarrhea abdominal pain or cramping dysphagia (difficulty swallowing) drooling in children 	<ul style="list-style-type: none"> nausea 	<ul style="list-style-type: none"> nausea
Other	<ul style="list-style-type: none"> anxious or feeling of “impending doom” sudden lack of energy (lethargy) in children quietness or sleepiness in children headache, light-headedness or dizziness decreased level of consciousness uterine cramps 	<ul style="list-style-type: none"> fearfulness light-headedness dizziness numbness, weakness sometimes accompanied by brief clonic seizure activity 	<ul style="list-style-type: none"> fearfulness light-headedness dizziness numbness, weakness tingling around lips and spasm in the hands and feet associated with hyperventilation

NOTE: Bolded text indicates symptoms specific to **pediatric** clients.

Maintained by:	HPA Leads (Nursing) – Anaphylaxis Working Group				
Issuing Authority:	Island Health – Professional Practice and Learning and Performance Support				
Last Revised:	2021-Mar-26	Last Reviewed:	2019-Feb-01	First Issued:	2019-Feb-01

6.0 Anaphylaxis Response Kit Contents

- Copy of this Decision Support Tool
- 4 ampoules of EPINEPHrine 1mg/mL
- 4 - 1 mL syringes
- Needles (25 to 27 gauge)
 - 4 - 1 inch
 - 4 - 1½ inch
- Alcohol swabs

Check medication vials and equipment. Replace if outdated.

Client Owned Auto-injector of Epinephrine

A client's own auto-injector of epinephrine may be used to administer epinephrine in situations where a delay in administration of epinephrine may occur (e.g. kit not readily available).


Clinical judgement is required.

Note: EpiPen® delivers 0.3 mg of epinephrine and EpiPen Junior® delivers 0.15 mg of epinephrine.

7.0 Injection Location Considerations

The correct site of intramuscular (IM) administration of epinephrine is ALWAYS the vastus lateralis located at the middle third of the *lateral thigh* [See image 1].

Image 1: Vastus Lateralis Injection Site



A, Landmarks for vastus lateralis site. **B**, Administering intramuscular injection in vastus lateralis site. (From Perry, A.G., Potter, P.A., Ostendorf, W.R. [Eds.]. [2018]. *Clinical nursing skills & techniques* [9th ed.]. St. Louis: Elsevier.)


Notes:

- If immunization or medication has been administered to both legs, give epinephrine IM at least 2.5 cm (1 inch) from original injection site.
- Alternate legs with multiple epinephrine IM doses.
- Administration of epinephrine through the clothing is acceptable in emergency situations.

Maintained by:	HPA Leads (Nursing) – Anaphylaxis Working Group				
Issuing Authority:	Island Health – Professional Practice and Learning and Performance Support				
Last Revised:	2021-Mar-26	Last Reviewed:	2019-Feb-01	First Issued:	2019-Feb-01

8.0 Intervention: Initial Emergency Treatment of Anaphylaxis

Perform Simultaneously



Maintained by:	HPA Leads (Nursing) – Anaphylaxis Working Group				
Issuing Authority:	Island Health – Professional Practice and Learning and Performance Support				
Last Revised:	2021-Mar-26	Last Reviewed:	2019-Feb-01	First Issued:	2019-Feb-01

9.0 Client or Family Education

Clients who have experienced anaphylaxis and their families should receive education including:

- Information on anaphylaxis being a potentially life threatening allergic reaction
- Avoiding triggers
- The emergency steps to take in case of anaphylaxis:
 - Carry your prescribed epinephrine auto-injector
 - Administer prescribed epinephrine auto-injector immediately
 - If in a hospital, long term care home or other institution: summon help
 - If in the community, or at home: call an ambulance
- Follow up with your physician, nurse practitioner, allergist or other appropriate resource to:
 - Develop an anaphylaxis action plan including self-administration of epinephrine
 - Obtain medical identification (e.g. MedicAlert®)
 - Undergo testing to confirm triggers
 - Explore ways that sensitivities to triggers may be reduced

Suggested client teaching resources:

- Health Link BC. (2018) [Severe Allergic Reaction \(Anaphylaxis\)](#)
- Food Allergy Canada. (2017) [Allergic Reactions Could You Save a Life](#)

10.0 Documentation

- Document the following in the health record:
 - Assessment
 - Nursing diagnosis of the condition
 - Interventions carried out
 - Client's response
 - Follow-up actions, transfer of care and client teaching
- Enter client allergy status as per employer policies to ensure communication to other providers.

11.0 Reporting

Suspected anaphylaxis adverse reactions related to drug or immunization administration are reported as required by employer policy:

- Patient Safety Learning System (PSLS) or equivalent (as required)
- Long Term Care settings must report adverse event to licensing body (as required)
- If the trigger for anaphylaxis is the result of an immunization, complete the following:
 - BCCDC [Worksheet for Events Managed as Anaphylaxis Following Immunization](#)
 - BCCDC [Adverse Event Following Immunization \(AEFI\) Case Report Form](#)

Maintained by:	HPA Leads (Nursing) – Anaphylaxis Working Group				
Issuing Authority:	Island Health – Professional Practice and Learning and Performance Support				
Last Revised:	2021-Mar-26	Last Reviewed:	2019-Feb-01	First Issued:	2019-Feb-01

References

1. Simons, F.E. et al. (2011). World Allergy Organization guidelines for the assessment and management of anaphylaxis. *World Allergy Organization Journal*. 4:13-37. Retrieved from <http://www.waojournal.org/content/4/2/13>
2. Simons, F.E. et al. (2015). 2015 Update of the evidence base: World Allergy Organization anaphylaxis guidelines. *World Allergy Organization Journal*. 8:32. Retrieved from <https://waojournal.biomedcentral.com/articles/10.1186/s40413-015-0080-1>
3. Campbell, R.L., Kelso, J.M. Anaphylaxis: acute diagnosis. *UpToDate*. Retrieved April 19th, 2017 from https://www.uptodate.com/contents/anaphylaxis-acute-diagnosis?source=see_link
4. Campbell, R.L., Kelso, J.M. Anaphylaxis: emergency treatment. *UpToDate*. Retrieved June 8, 2018 from https://www.uptodate.com/contents/anaphylaxis-emergency-treatment?source=search_result&search=anaphylaxis&selectedTitle=2~150
5. BC Centre for Disease Control. (2013). Section V – Management of Anaphylaxis in a Non-Hospital Setting. *Communicable Disease Control Immunization Program*. Vancouver: BC Centre for Disease Control. Retrieved from http://www.bccdc.ca/resource-gallery/Documents/Guidelines%20and%20Forms/Guidelines%20and%20Manuals/Epid/CD%20Manual/Chapter%20-%20Imms/SectionV_Anaphylaxis.pdf
6. Canadian Society of Allergy and Clinical Immunology. (2016). *Anaphylaxis in Schools and Other Settings 3rd Edition Revised*. Canadian Society of Allergy and Clinical Immunology. Retrieved from <http://resources.allergyaware.ca/download/anaphylaxis-3rd-edition-revised.pdf>
7. College of Registered Nurses of British Columbia. (2017). *Scope of Practice for Registered Nurses*. Vancouver: College of Registered Nurses of British Columbia. Retrieved from <https://www.crnbc.ca/Standards/Lists/StandardResources/433ScopeforRegisteredNurses.pdf>
8. College of Registered Psychiatric Nurses of British Columbia. (2017). *Scope of Practice for Registered Psychiatric Nurses: Standards, Limits and Conditions*. Vancouver: College of Registered Psychiatric Nurses of British Columbia. Retrieved from <http://www.crpnb.ca/wp-content/uploads/2017/02/2016-03-15-Revised-Scope-with-disclaimer-updated-Feb-13-2017.pdf?v=410>
9. College of Licensed Practical Nurses of British Columbia. (2017). *Scope of Practice for Licensed Practical Nurses*. Vancouver: College of Licensed Practical Nurses of British Columbia. Retrieved from <https://www.clpnbc.org/Documents/Practice-Support-Documents/Scope-of-Practice-ONLINE.aspx>
10. Health Canada. (2016). *Food Allergies: Information for Consumers*. Retrieved from <http://www.hc-sc.gc.ca/fn-an/securit/allerg/fa-aa/index-eng.php>
11. Epinephrine. (n.d.). In *Lexicomp*. Retrieved from <https://online.lexi.com/lco/action/home>
12. Muraro, A. et al. (2014). Anaphylaxis: guidelines from the European Academy of Allergy and Clinical Immunology. *Allergy*. 29: 1026-1045. Retrieved from <http://onlinelibrary.wiley.com/doi/10.1111/all.12437/full>

Maintained by:	HPA Leads (Nursing) – Anaphylaxis Working Group				
Issuing Authority:	Island Health – Professional Practice and Learning and Performance Support				
Last Revised:	2021-Mar-26	Last Reviewed:	2019-Feb-01	First Issued:	2019-Feb-01

13. Sheikh, A., Broek, V.T., Brown, S.G.A. and Simons, F.E.R. (2007). *H₁-antihistamines for the treatment of anaphylaxis: Cochrane systematic review*. retrieved on June 8, 2018 from <https://onlinelibrary.wiley.com/doi/epdf/10.1111/j.1398-9995.2007.01435.x>
14. Australasian Society of Clinical Immunology and Allergy (2017). *Acute management of anaphylaxis*. Retrieved on June 8, 2018 from <https://www.allergy.org.au/images/stories/pospapers/ASCIAGuidelinesAcuteManagementAnaphylaxis2017Updated.pdf>
15. Medication Administration: Subcutaneous Injection. (2017, November). In *Elsevier clinical skills*. Retrieved from <https://epm601.elsevierperformancemanager.com/Personalization/Home?VirtualName=interiorhealth-canada>
16. Medication Administration: Intramuscular Injection. (2018, May). In *Elsevier clinical skills*. Retrieved from <https://epm601.elsevierperformancemanager.com/Personalization/Home?VirtualName=interiorhealth-canada>
17. Canadian Pediatric Society. (2018) *Position statement Emergency treatment of anaphylaxis in infants and children*. Retrieved August 15, 2018 from <https://www.cps.ca/en/documents/position/emergency-treatment-anaphylaxis>

Maintained by:	HPA Leads (Nursing) – Anaphylaxis Working Group					
Issuing Authority:	Island Health – Professional Practice and Learning and Performance Support					
Last Revised:	2021-Mar-26	Last Reviewed:	2019-Feb-01	First Issued:	2019-Feb-01	Page 10 of 16

Appendix A: Common Triggers for Anaphylaxis

The true global rate of occurrence of anaphylaxis from all triggers in the general population is unknown because of under-recognition by patients and caregivers and under-diagnosis by healthcare professionals.¹

Numerous nursing and health care activities expose patients to common triggers for anaphylaxis. Below are some selected common triggers for anaphylaxis. Triggers for individuals can vary widely and the below list is not exhaustive.

Food	Environment	Medications or Other
<ul style="list-style-type: none"> • Peanuts • Tree nuts • Soy • Seafood • Shellfish • Milk • Eggs • Mustard • Wheat • Sesame 	<ul style="list-style-type: none"> • Venom from stinging insects • Horse • Latex 	<ul style="list-style-type: none"> • Antibiotics especially β lactams (including penicillins, cephalosporins, carbapenems) • Non-steroidal anti-inflammatories (including ASA) • Biologic medications including immunizations • Chemotherapy • Radiocontrast media (x-ray dye) • Dextrans • Substances found in illicit drugs • Blood and Blood Products • Latex Gloves

Maintained by:	HPA Leads (Nursing) – Anaphylaxis Working Group					
Issuing Authority:	Island Health – Professional Practice and Learning and Performance Support					
Last Revised:	2021-Mar-26	Last Reviewed:	2019-Feb-01	First Issued:	2019-Feb-01	Page 11 of 16

Appendix B: EPINEPHrine 1mg/ml Dose by Age Table

NOTE: Use dose by weight whenever possible.

Dose by Age (Dosing is 0.01 mg/kg to maximum of 0.5 mg)		
Age	Dose in mg	Volume (1 mg/mL)
2 to 6 months	0.07 mg	0.07 mL
7 to 12 months	0.1 mg	0.1 mL
13 months to 4 years	0.15 mg	0.15 mL
5 years	0.2 mg	0.2 mL
6 to 9 years	0.3 mg	0.3 mL
10 to 13 years	0.4 mg	0.4 mL
Greater than or equal to 14 years	0.5 mg	0.5 mL

Maintained by:	HPA Leads (Nursing) – Anaphylaxis Working Group					
Issuing Authority:	Island Health – Professional Practice and Learning and Performance Support					
Last Revised:	2021-Mar-26	Last Reviewed:	2019-Feb-01	First Issued:	2019-Feb-01	Page 12 of 16



Adapted from:

Vancouver Coastal Health and Providence Health Care Professional Practice. (September 2017)
Anaphylaxis: Initial Emergency Treatment Decision Support Tool. (Draft)

Developed by:

Paula Araujo, Regional Practice Leader, Clinical Process, Interior Health
Cathy Czechmeister, Lead, Professional Practice Strategy - Nursing, Northern Health
Sheila Gordon-Payne, Lead, Professional Practice Strategy – Nursing, Northern Health
Melina Kason, Practice Consultant, Professional Practice, Island Health

Reviewed by:

Dr. Monika Naus MD FRCPC, Medical Director, Communicable Diseases and Immunization Service, BC Centre for Disease Control
Dr. Scott B. Cameron MD, PhD, FRCPC, Clinical Instructor, Division of Allergy and Immunology, Department of Pediatrics, University of British Columbia
Christine Halpert, Senior Practice Leader, Communicable Diseases and Immunization Service, BC Centre for Disease Control
Stephanie Meier, Public Health Resource Nurse, Communicable Diseases and Immunization Service, BC Centre for Disease Control
BC Provincial Health Authorities Representatives from Fraser Health, Interior Health, Vancouver Coastal Health, Provincial Health Services Authority, Providence Health

Endorsed by:

British Columbia Chief Nursing Officer Council

Maintained by:	HPA Leads (Nursing) – Anaphylaxis Working Group					
Issuing Authority:	Island Health – Professional Practice and Learning and Performance Support					
Last Revised:	2021-Mar-26	Last Reviewed:	2019-Feb-01	First Issued:	2019-Feb-01	Page 13 of 16