Two Stage Cluster Sampling Simulation Results

Simulation Codes by Nengak Emmanuel Goltong

2024-03-17

Simulation Results

This simulation exercise consist of the following steps:

- 1. Six independent variables (normally distributed) are simulated (a total of N times M)
- 2. The simulated data is then split into M distinct clusters each of size N. The variables Y, X and Z are constructed following the relationship defined in the work of Maji et al. only that here, the error component is added.
- 3. A random sample of m (or m' then m) clusters is selected out of the M total clusters. This is called the first sampled units (fsu).
- 4. A random sample of n (or n' then n) units are sampled from each of m selected clusters. This is called the second sampled units (ssu).
- 5. All the different estimators of the Mean Square Error are calculated based on the observed data and compared.
- 6. Steps 3 to 5 are repeated a hundred times for each specific case and the estimates of Mean Square Error are all saved in arrays after which the means are calculated and compared.
- 7. Coding can be fun and challenging. What do you think?

Table 1. Artificial Population study (when r is not varying)

Case A, Procedure I (N = 10, M = 10, n' = 7, n = 5, m = 7)

Ibrahim LOSS_Maji LOSS_Ibrahim
7.7131 -140.7166 -147.7131
.8227 -134.2475 -154.8227
.1662 -127.8759 -166.1662
.8098 -121.5513 -180.8098

Table 2. Artificial Population study (when r is not varying)

Case B, Procedure I (N = 10, M = 10, n' = 7, n = 5, m = 7)

р	q	PRE_Maji	PRE_Ibrahim	LOSS_Maji	LOSS_Ibrahim
0.05	0.95	287.1852	309.8038	-187.1852	-209.8038
0.1	0.9	276.1314	325.5487	-176.1314	-225.5487
0.15	0.85	265.463	344.4662	-165.463	-244.4662
0.2	0.8	255.1155	367.6047	-155.1155	-267.6047

Table 3. Artificial Population study (when r is not varying)

Procedure II (N = 10, M = 10, n = 5, m' = 8, m = 7)

р	q	PRE_Maji	PRE_Ibrahim	LOSS_Maji	LOSS_Ibrahim
0.05	0.95	228.7942	120.1068	-128.7942	-20.10682
0.1	0.9	234.1931	148.4815	-134.1931	-48.48153
0.15	0.85	253.0895	211.4755	-153.0895	-111.4755
0.2	0.8	1767.786	264.3053	-1667.786	-164.3053

Table 4. Artificial Population study (when r is varying)

Case A, Procedure I (N = 10, M = 10, n' = 7, n = 4, m = 7)

p1	q1	p2	q2	p3	q3	p4	q4	PRE_MajPRE_IbrahimLOSS_MajiLOSS_	<u>Ibrahim</u>
0.05	0.95	0.05	0.95	0.05	0.95	0.1	0.9	278.6145 254.9173 -178.6145 -154.9	9173
0.05	0.95	0.05	0.95	0.1	0.9	0.05	0.95	278.7064 255.0313 -178.7064 -155.0)313
0.05	0.95	0.05	0.95	0.1	0.9	0.1	0.9	276.8989 254.4278 -176.8989 -154.4	4278
0.1	0.9	0.1	0.9	0.1	0.9	0.05	0.95	275.0897 254.0803 -175.0897 -154.0	0803
0.1	0.9	0.15	0.85	0.05	0.95	0.05	0.95	274.8001 253.6213 -174.8001 -153.6	3213
0.1	0.9	0.1	0.9	0.1	0.9	0.15	0.85	271.5271 253.2227 -171.5271 -153.2	2227

Table 5. Artificial Population study (when r is varying)

Case B, Procedure I (N = 10, M = 10, n' = 7, n = 4, m = 7)

p1	q1	p2	q2	p3	q3	p4	q4	PRE_Maj	PRE_Ibrahi	imLOSS_Maji	LOSS_Ibrahim
0.05	0.95	0.05	0.95	0.05	0.95	0.1	0.9	293.0767	265.191	-193.0767	-165.191
0.05	0.95	0.05	0.95	0.1	0.9	0.05	0.95	292.7693	264.9849	-192.7693	-164.9849
0.05	0.95	0.05	0.95	0.1	0.9	0.1	0.9	290.5565	264.5681	-190.5565	-164.5681
0.1	0.9	0.1	0.9	0.1	0.9	0.05	0.95	287.7565	263.9043	-187.7565	-163.9043
0.1	0.9	0.15	0.85	0.05	0.95	0.05	0.95	287.5897	263.2503	-187.5897	-163.2503
0.1	0.9	0.1	0.9	0.1	0.9	0.15	0.85	283.4256	263.4702	-183.4256	-163.4702

Table 6. Artificial Population study (when r is varying)

Procedure II (N = 10, M = 10, n = 4, m' = 8, m = 7)

p1	q1	p2	q2	р3	q3	p4	q4	PRE_Ma	j₽RE_Ibrahi	mLOSS_Maji	LOSS_Ibrahim
0.05	0.95	0.05	0.95	0.05	0.95	0.1	0.9	266.1098	122.7567	-166.1098	-22.75673
0.05	0.95	0.05	0.95	0.1	0.9	0.05	0.95	266.8984	122.7514	-166.8984	-22.75141
0.05	0.95	0.05	0.95	0.1	0.9	0.1	0.9	269.011	128.5508	-169.011	-28.55079
0.1	0.9	0.1	0.9	0.1	0.9	0.05	0.95	272.7992	136.7851	-172.7992	-36.7851
0.1	0.9	0.15	0.85	0.05	0.95	0.05	0.95	271.8519	134.3659	-171.8519	-34.3659
0.1	0.9	0.1	0.9	0.1	0.9	0.15	0.85	280.343	159.5629	-180.343	-59.5629