

Human Genome Architecture, Reference Cartography, & The Biophysics of Disease

A Comprehensive Study Monograph on Diploid Phasing, Non-Coding Dark Matter, the T2T-CHM13 Revolution, 30th-Power Epistasis, and Translational CRISPR Medicine.

Instructor: Prof. Sergei Pond, Ph.D. **Lecture:** Lecture 2

Primary Reference: Nurk et al. *Science* (2022)

Format: Course Reading Monograph & Exam Study Guide

| 1. Diploid Architecture & The Phasing Dilemma

The human somatic genome consists of approximately **6.1 billion base pairs** (6.1×10^9 bp) distributed across 46 chromosomes (23 homologous pairs). While introductory genetics often conceptualizes an individual's genome as an abstract single string of letters, human clinical biology is fundamentally **diploid**. Every individual inherits one complete set of 23 maternal chromosomes and one set of 23 paternal chromosomes.

Modern short-read Next-Generation Sequencing (NGS)—the workhorse of clinical diagnostics, generating 150 bp paired-end reads—routinely collapses this diploid reality into an unphased list of variant calls. An NGS pipeline reports that a patient is heterozygous for Variant 1 and heterozygous for Variant 2 within the same gene. Crucially, short reads **cannot physically connect mutations** that are separated by more than 300–500 base pairs.

THE CLINICAL PHASING PARADOX (SPOT CHECK 1)

Consider a patient evaluated for an autosomal recessive metabolic disorder. Short-read exome sequencing identifies two severe loss-of-function variants in a metabolic enzyme:

Exon 2 c.120G>A (p.Arg40Ter) and Exon 9 c.840C>T (Frameshift) .

Without phase resolution, clinical prediction is impossible:

- **In Trans (Compound Heterozygous):** Variant 1 is on the maternal chromosome; Variant 2 is on the paternal chromosome. Both alleles are non-functional. The patient produces **0% functional enzyme** and develops catastrophic metabolic failure.
- **In Cis (Same Chromosome):** Both mutations reside on the maternal chromosome. The paternal homolog remains **100% wild-type and functional**. The patient produces normal enzyme levels, exhibits normal serum biochemistry, and is a completely healthy, asymptomatic carrier!

To resolve phase, clinical geneticists must either perform **parental trio sequencing** (confirming maternal vs. paternal origin) or utilize **long-read single-molecule sequencing** (PacBio HiFi or Oxford Nanopore >20 kb) that physically spans both exons on continuous DNA molecules.

At the population level, the non-random association of alleles at different loci is termed **Linkage Disequilibrium (LD)**. For two biallelic loci with alleles A/a and B/b , the coefficient of linkage disequilibrium D is defined as:

$$D = p_{AB} - p_A p_B, \quad D' = \frac{D}{D_{\max}}, \quad r^2 = \frac{D^2}{p_A(1 - p_A)p_B(1 - p_B)}$$

Recombination breaks down LD across generational time at a rate proportional to the recombination fraction c between loci:

$$D_t = D_0(1 - c)^t$$

Where $c \approx 0.01$ corresponds to 1 centimorgan (cM), roughly 1 Megabase (Mb) in human nuclear DNA. Across generations, ancestral founder haplotypes fragment into localized LD blocks,

dictating the resolution of Genome-Wide Association Studies (GWAS).

| 2. The C-Value Enigma & Non-Coding Anatomy

In the early era of molecular biology, geneticists hypothesized that an organism's total genomic DNA mass (the *C-value*) would scale proportionally with its biological complexity. This hypothesis was thoroughly refuted by the **C-Value Paradox** (now known as the *C-Value Enigma*).

THE C-VALUE FALLACY

The single-celled amoeba *Polychaos dubium* possesses a genome of **~670 billion base pairs (670 Gb)**—over 200 times larger than the human genome (3.055 Gb). The marbled lungfish (*Protopterus aethiopicus*) maintains a 130 Gb genome. If DNA mass dictated morphological sophistication, amoebae and lungfish would represent the apex of evolutionary complexity.

The human haploid genome consists of **3,054,815,472 base pairs** (Nurk et al. 2022). Remarkably, only **~19,900 protein-coding genes** are encoded within this mass. The entire human protein-coding exome occupies approximately **34 Mb (~1.1% to 1.5%)** of the sequence.

Genomic Fraction	Percentage (%)	Megabases (Mb)	Biological Function & Clinical Significance
Protein-Coding Exons	~1.5%	~34 Mb	~19,900 genes; target of clinical Whole Exome Sequencing (WES).
Introns	~38.0%	~1,150 Mb	Alternative splicing, transcription factor binding, microRNAs, snoRNAs.
Retrotransposons (LINEs)	~21.0%	~640 Mb	Autonomous retrotransposons (LINE-1); drive insertional mutagenesis.
Retrotransposons (SINEs/Alu)	~13.0%	~400 Mb	Non-autonomous elements; high sequence homology causes unequal crossing over.
LTR Retrotransposons / HERVs	~8.5%	~260 Mb	Human Endogenous Retroviruses; regulate placental syncytin and innate immunity.
DNA Transposons	~3.0%	~90 Mb	Inactive cut-and-paste remnants; historical genome architecture drivers.
Satellite DNA & Centromeres	~8.0–10.0%	~250 Mb	Alpha-satellites, Higher-Order Repeats (HORs), kinetochore assembly, rDNA.
Enhancers & Promoters	~5.0–8.0%	~180 Mb	Tissue-specific cis-regulatory modules, GATA1, CTCF boundary loops.

Human morphological and cognitive complexity arises not from raw gene counts, but from **combinatorial regulatory networks**: tissue-specific enhancers, alternative promoter usage, expansive alternative splicing (generating >100,000 distinct proteomic isoforms), non-coding regulatory RNAs, and high-order 3D chromatin conformation (Topologically Associating Domains, or TADs).

| 3. Reference Cartography: GRCh38 to T2T-CHM13

Every clinical diagnostic report, variant call, and RNA-seq alignment relies upon a **reference genome**. The reference acts as the coordinate cartography against which patient sequencing reads are compared. For two decades, the scientific community operated under incomplete drafts.

The original Human Genome Project utilized bacterial artificial chromosome (BAC) clone-by-clone sequencing, while Celera Genomics employed whole-genome shotgun sequencing. In 2013, the Genome Reference Consortium released **GRCh38** (hg38). Although widely heralded as the gold standard, GRCh38 was a **mosaic composite derived from multiple anonymous donors** (predominantly RPCI-11, a donor of mixed African-European ancestry).

⚠️ **THE LEGACY 8% VOID IN GRCH38**

Despite extensive refinement, GRCh38 contained **hundreds of unsequenced gaps encompassing ~8% of the genome (~200–240 Mb)**. These unsequenced regions consisted of:

- **Centromeric Satellite Arrays:** Millions of bases of repeating 171 bp alpha-satellite Higher-Order Repeats (HORs).
- **Acrocentric p-Arms:** The short arms of chromosomes 13, 14, 15, 21, and 22, containing hundreds of tandem copies of the 45S ribosomal DNA (rDNA) arrays.
- **Segmental Duplications:** Low-copy repeats (>90% sequence identity spanning >1 kb) that collapsed during short-read assembly algorithms.

In April 2022, the **Telomere-to-Telomere (T2T) Consortium** published the complete, gapless sequence of a human genome: **T2T-CHM13** (Nurk et al. 2022, *Science*).

The breakthrough required two fundamental innovations:

Biological Simplification (CHM13): The consortium sequenced a **complete hydatidiform mole (CHM13)**. When an anucleate ovum is fertilized by a single sperm that subsequently duplicates its chromosomes, the resulting blastocyst is genetically **monoploid (46,XX with identical paternal copies)**. Sequencing CHM13 eliminated the mathematical nightmare of resolving heterozygous single-nucleotide differences between maternal and paternal homologs.

Long-Read Sequencing Synergy: The assembly combined:

Pacific Biosciences (PacBio) HiFi: Circular consensus sequencing yielding reads of 15–25 kb with **>99.9% base-call accuracy** (Q30+), capable of distinguishing subtle differences between paralogous repeats.

Oxford Nanopore Ultra-Long Reads: Continuous single-molecule reads spanning **100 kb to >1 Mb**, providing the topological scaffolding required to traverse entire centromeric satellite arrays.

🏆 **T2T-CHM13 MILESTONE METRICS**

T2T-CHM13 added **238 Mb of previously unmapped sequence**, discovered **115 novel protein-coding genes**, characterized **3,604 newly resolved genes** (predominantly segmental duplications), and closed every single telomere, centromere, and rDNA array across all 24 human chromosomes.

| 4. Paralog Misalignment & Ghost Pathogenic Calls

The transition from GRCh38 to T2T-CHM13 was not an abstract bioinformatic upgrade; it revolutionized clinical diagnostic fidelity.

When short sequencing reads (150 bp) are mapped to an incomplete reference, software aligners (BWA-MEM, Bowtie2) operate under the assumption that the reference represents the complete universe of genomic loci. If a patient's read originates from a gene copy that is **missing from the reference**, the aligner forces that read to bind to the single closest homologous sequence present in the assembly.

💡 THE "GHOST" VARIANT PARADOX (SPOT CHECK 2)

In a landmark clinical audit published in *Science* (Nurk et al. 2022), investigators examined variants cataloged in public clinical repositories (ClinVar, gnomAD) under GRCh38:

- Under **GRCh38**, a recurrent frameshift deletion in a medically important cardiovascular gene showed an apparent **12% carrier frequency** in healthy population cohorts. Cardiologists struggled to reconcile how an alleged "lethal penetrant frameshift" could occur in 1 out of every 8 healthy individuals.
- When the exact same raw FASTQ reads were re-aligned against **T2T-CHM13**, the carrier frequency collapsed to exactly **0.0%**!

The Mechanism: The true genome contains a non-functional, ancestral pseudogene duplication located inside an unsequenced gap in GRCh38. In healthy humans, this duplicate copy carries an endogenous 1 bp deletion. Under GRCh38, reads from this missing duplicate mapped directly to the functional primary gene, manufacturing an artificial "pathogenic mutation." T2T-CHM13 provided the true duplicate locus, allowing reads to map to their correct coordinate and instantly eliminating thousands of false-positive disease diagnoses.

| 5. The Globin Clusters & Sickle Cell Biophysics

No genetic system better illustrates the interplay between physical genomic architecture, stoichiometric gene expression, and biophysical epistasis than the human globin gene families.

Human hemoglobin is a heterotetramer composed of two α -like chains and two β -like chains ($\alpha_2\beta_2$). These chains are encoded on two distinct chromosomes:

The α -Globin Cluster (Chromosome 16p13.3): Arranged in developmental order: 5' – HBZ (Zeta) – HBM – HBA2 (Alpha-2) – HBA1 (Alpha-1) – HBQ1 – 3'. Normal individuals possess **four functional α -globin genes** (two per diploid homolog: $\alpha\alpha/\alpha\alpha$).

The β -Globin Cluster (Chromosome 11p15.4): Spanning ~50 kb, regulated by an upstream **Locus Control Region (LCR)**: 5' – HBE1 (Epsilon) – HBG2 (G-gamma) – HBG1 (A-gamma) – HBD (Delta) – HBB (Beta) – 3'.

Developmental Stage	Tetramer Formula	Name	Relative Abundance in Adults
Embryonic	$\zeta_2\epsilon_2, \alpha_2\epsilon_2, \zeta_2\gamma_2$	Gower 1, Gower 2, Portland	0% (Silenced in 1st trimester)
Fetal	$\alpha_2\gamma_2$	HbF (Fetal Hemoglobin)	<1.0% (Awakened therapeutically)
Adult (Major)	$\alpha_2\beta_2$	HbA (Adult Hemoglobin)	~96.0–97.0%
Adult (Minor)	$\alpha_2\delta_2$	HbA2	~2.0–3.0%

In 1949, Linus Pauling, Harvey Itano, and colleagues published their landmark paper establishing Sickle Cell Anemia as the world's first proven "**molecular disease.**" In 1957, Vernon Ingram demonstrated that this devastating pathology is caused by a single substitution:

Codon 6 of *HBB* : GAG (Glu, Negative) → GTG (Val, Hydrophobic)

In oxygenated hemoglobin (the *R*-state), the mutant Val6 residue is harmlessly exposed to solvent. However, when red blood cells traverse deoxygenating peripheral tissues, the tetramer transitions to the unliganded *T*-state. This conformational shift exposes a hydrophobic receptor pocket on the partner β -chain formed by residues **Phenylalanine 85 (Phe85) and Leucine 88 (Leu88)**.

The hydrophobic Val6 of one deoxygenated tetramer inserts directly into the Phe85/Leu88 hydrophobic pocket of an adjacent tetramer. This stereospecific lock drives self-assembly into **14-**

strand helical crystalline polymer fibers that distort flexible biconcave erythrocytes into rigid, sickle-shaped sickles.

| 6. 30th-Power Epistasis & Capillary Transit

Why do patients with sickle cell disease not suffer instantaneous, continuous fatal thrombosis throughout their entire circulatory system? The answer lies in the extraordinary non-linear physical chemistry of **nucleation kinetics** discovered by William Eaton and James Hofrichter at the NIH.

Hemoglobin polymerization does not occur immediately upon deoxygenation. It requires a **homogeneous nucleation event**: approximately 30 independent HbS tetramers must spontaneously collide, align with atomic precision, and overcome unfavorable entropic barriers to form a critical thermodynamic nucleus.

THE EATON-HOFRICHTER DELAY TIME LAW

Once the critical nucleus forms, fiber elongation proceeds almost instantaneously. The time preceding fiber growth is termed the **delay time (τ_d)**. Eaton and Hofrichter proved that delay time is inversely proportional to the **30th to 35th power** of intracellular hemoglobin concentration (C_{Hb}):

$$\tau_d \propto \left(\frac{1}{C_{\text{Hb}}} \right)^\gamma, \quad \gamma \approx 30 - 35$$

This extreme 30th-power dependence creates a biological **transit race**:

The average microvascular capillary transit time in human tissues is $t_{\text{transit}} \approx 0.5$ to 1.0 second.

If the delay time τ_d exceeds 1.0 s, red blood cells traverse the narrow capillary bed, re-enter the venous circulation, and return to the pulmonary capillary bed to become re-oxygenated *before polymerization can begin*.

If $\tau_d < 1.0$ s, polymers form inside the capillary lumen, rigidifying the erythrocyte, occluding microvessels, starving downstream tissues of oxygen, and initiating catastrophic vaso-occlusive crises (VOC), acute chest syndrome, and ischemic stroke.

THE EPISTATIC PARADOX: ALPHA-THALASSEMIA RESCUE (SPOT CHECK 3)

In clinical hematology, Patient A is homozygous for the classic sickle mutation ($HbSS$). They suffer debilitating vaso-occlusive crises monthly and show high stroke velocity on transcranial Doppler.

Patient B is also homozygous for the exact same HBB mutation ($HbSS$), yet suffers **almost zero pain crises and zero stroke risk**. Genomic profiling reveals that Patient B co-inherited a two-gene deletion in the α -globin cluster on Chromosome 16 ($- \alpha / - \alpha$, α -thalassemia trait).

The Biophysical Mechanism: Normal red cells have a mean corpuscular hemoglobin concentration (MCHC) of ~ 34 g/dL, yielding $\tau_d \approx 0.8$ s (losing the capillary transit race). Co-inheriting α -thalassemia impairs α -chain synthesis, creating an intracellular globin chain imbalance that reduces cellular hemoglobin concentration by just **10%** (diluting C_{Hb} to $0.90 \times C_0$).

Applying the 30th-power kinetic law:

$$\tau_{\text{thal}} = \tau_0 \times \left(\frac{1}{0.90} \right)^{30} = 0.8 \text{ s} \times (1.111)^{30} = 0.8 \text{ s} \times 23.75 \approx \mathbf{19.0 \text{ seconds!}}$$

A modest 10% dilution in protein concentration expands the polymerization delay time by **nearly 24-fold** (from 0.8 s to 19 s). The red cell easily traverses the 1-second capillary passage long before sickling begins, completely aborting vascular occlusion!

| 7. Balanced Polymorphism & Malaria Resistance

Given the lethal nature of untreated homozygous sickle cell disease ($HbSS$), classic Mendelian selection would rapidly purge the allele from human populations. Yet, across Sub-Saharan Africa, the Mediterranean basin, and India, the sickle allele (HbS) is maintained at carrier frequencies of **10% to 25%**.

In 1949, J.B.S. Haldane formulated the **Malaria Hypothesis**: severe endemic infectious diseases exert powerful selective pressures that remodel the human genome. Sickle cell trait (*HbAS*) is the definitive paradigm of **heterozygote advantage (overdominance)**.

Genotype	Phenotype	Malaria Fitness	Sickle Fitness
AA	Normal HbA	Lethal Cerebral Malaria	Normal
AS	Sickle Trait	90% Malaria Protection	Asymptomatic
SS	Sickle Disease	N/A	Vaso-occlusion / Stroke

When relative fitness satisfies $W_{AS} > W_{AA}$ and $W_{AS} > W_{SS}$, natural selection establishes a stable polymorphic equilibrium frequency $\hat{q}$ for the deleterious allele:

$$\hat{q} = \frac{s}{s + t}$$

The Cellular Mechanism of Malaria Protection: The intra-erythrocytic parasite *Plasmodium falciparum* metabolizes host hemoglobin, generating lactic acid and consuming oxygen. In an *HbAS* heterozygote, this localized intra-cellular hypoxia accelerates selective sickling of *parasitized erythrocytes only*. Macrophages in the splenic reticuloendothelial system promptly identify and phagocytose these deformed cells, destroying the parasite before cerebral malaria can develop.

| 8. The CRISPR Revolution: Silencing BCL11A

For over seven decades, hematologists noted that newborn infants homozygous for the sickle mutation (*HbSS*) are entirely asymptomatic. Disease manifestations only begin between 6 and 12 months of life.

The reason is developmental globin switching: newborns produce **Fetal Hemoglobin (HbF, $\alpha_2\gamma_2$)**. The γ -globin chain contains a neutral **Glutamine at position 87 (Gln87)** in place of the β -chain's Thr87. Gln87 disrupts the lateral contact geometry required for sickle polymer assembly. A single

HbF tetramer incorporated into a growing HbS polymer terminates the chain, acting as a potent polymer poison.

Adult red cells express less than 1% HbF because a transcriptional repressor protein—**BCL11A (B-Cell Lymphoma/Leukemia 11A)**—binds directly to the promoters of the γ -globin genes (*HBG1/2*), shutting them down permanently.

TRANSLATIONAL BREAKTHROUGH: CASGEVY (EXAGAMGLOGENE AUTOTEMCEL)

Knocking out the *BCL11A* protein-coding sequence using CRISPR-Cas9 would reactivate HbF, but would also cause catastrophic immunological failure and neurodevelopmental defects because BCL11A is an essential transcription factor in B-cell maturation and brain development.

In 2013, Bauer, Orkin, and colleagues made a transformative discovery using ENCODE functional genomic tracks: BCL11A expression in the erythroid lineage is governed by a **tissue-specific enhancer located in Intron 2 (+58 kb from the TSS) on Chromosome 2p16.1** containing a core **GATA1 binding motif**.

In December 2023, the FDA approved **Casgevy**, the world's first CRISPR-Cas9 therapeutic:

1. Patient CD34+ hematopoietic stem cells are harvested via apheresis.
2. Electroporation delivers Cas9-sgRNA ribonucleoprotein complexes targeting the **erythroid-specific +58 kb GATA1 enhancer**.
3. A single double-strand break disrupts the enhancer, selectively extinguishing BCL11A expression *in erythrocytes only*, while preserving normal expression in B-cells and neurons.
4. Re-infused stem cells engraft, producing **>40% circulating HbF**—permanently curing patients of vaso-occlusive crises!

Therapeutic Agent	Modality	Molecular Target	Biophysical / Clinical Mechanism
Casgevy (<i>exa-cel</i>)	CRISPR-Cas9 Cell Therapy	<i>BCL11A</i> Intron 2 +58 kb GATA1 Enhancer	Permanently silences repressor; elevates protective HbF > 40% ; eliminates pain crises.
Lyfgenia (<i>lovo-cel</i>)	Lentiviral Gene Addition	Synthetic <i>HBB</i> ^{T87Q} globin cDNA	Transduces anti-sickling β -globin variant carrying fetal-like Gln87 residue.
Oxbryta (<i>voxelotor</i>)	Small Molecule Allosteric Modulator	N-terminal α -globin Val1 residue	Stabilizes high-affinity oxy-R-state; prevents deoxygenation-induced polymer formation.
Adakveo (<i>crizanlizumab</i>)	Monoclonal Antibody	Endothelial P-Selectin receptor	Inhibits erythrocyte-platelet-endothelial cell adhesion, reducing vaso-occlusive frequency.
Hydroxyurea	Oral Chemotherapeutic	Ribonucleotide Reductase / NO signaling	Modestly reactivates HbF (10–20%) through epigenetic stress erythropoiesis.