

Genomics in Medicine & The AI Frontier

Lecture 01 Study Monograph: Deconstructing Clinical Dogmas, Historical Demography, the Economics of Evidence-Based Medicine, and the Precision Revolution.

Instructor: Prof. Sergei Pond **Department:** Biology / iGEM

Format: Accessible Offline Reading Monograph

1. The Epistemic Crisis of "Normal"

Every medical chart begins with reference intervals: numbers highlighted in green for "normal" and red for "abnormal." However, in clinical pathology, a "normal range" is rarely a biological boundary; it is a statistical convention defined as the central **95% interval** ($\mu \pm 1.96\sigma$) of a reference population.

⚠ THE GAUSSIAN MYTH IN CLINICAL DIAGNOSTICS

Defining normality as 95% guarantees that exactly **5% of completely healthy people** ($\alpha = 0.05$) will be classified as "abnormal" on any single laboratory test by mathematical definition.

In modern multi-omic medicine, patients do not receive one test; they receive multiplex panels measuring 20 to 500 biomarkers simultaneously. When k independent tests are performed, the probability of obtaining *at least one false abnormal result* in a perfectly healthy individual scales exponentially:

$$P(\geq 1 \text{ False Positive}) = 1 - (1 - \alpha)^k = 1 - (0.95)^k$$

Number of Tests (k)	Clinical Context	Chance of ≥ 1 False Alarm
$k = 1$	Single Fasting Glucose	5.0%
$k = 14$	Comprehensive Metabolic Panel (CMP)	51.2% (Coin flip)
$k = 20$	Routine Executive Wellness Screen	64.2%
$k = 100$	Targeted Metabolomics Panel	99.4%
$k = 500$	Liquid Biopsy / Proteomic Screen	> 99.999% (Certainty)

 THE HIGH-DIMENSIONAL SCREENING PARADOX (SPOT CHECK 01)

If a liquid biopsy screens 500 proteins where 25 (5%) reflect early cancer, operating at 80% sensitivity (20 True Positives) and 95% specificity (23.75 False Positives on 475 healthy proteins), the **False Discovery Rate (FDR)** is:

$$\text{FDR} = \frac{\text{FP}}{\text{TP} + \text{FP}} = \frac{23.75}{20.0 + 23.75} \approx 54.3\%$$

More than half of all positive cancer alerts are false alarms. High-dimensional diagnostics without Bayesian priors lead directly to unnecessary invasive biopsies and patient panic.

I 2. Deep Time & Historical Demography

A pervasive myth in popular biology states: *"In pre-modern times, life expectancy was 35 years, meaning ancient humans aged rapidly and became biological elders in their mid-thirties."*

Historical demographic records (such as the 1750 London Parish Registers) reveal this to be a fundamental misinterpretation of mathematical averages driven by extreme right-skewed childhood mortality.

 DECONSTRUCTING THE 35-YEAR-OLD ELDER MYTH (SPOT CHECK 02)

A life expectancy at birth ($e_0 = 35$) was not caused by accelerated aging. It was driven by **~40% infant and child mortality before age 5** (due to cholera, dysentery, and smallpox). For an individual who survived past childhood (age 20), conditional life expectancy (e_{20}) was **62 to 70 years**.



Historical archival photograph of Louis Pasteur working with glassware and broth in his Paris laboratory (1885).

Louis Pasteur (1822–1895): Established germ theory, pasteurization, and attenuation of rabies and anthrax vaccines.



Historical portrait of Dr. Robert Koch seated in formal attire with spectacles (circa 1900).

Robert Koch (1843–1910): Isolated *B. anthracis*, *M. tuberculosis*, and *V. cholerae*; formalized Koch's Postulates.

3. The 7 Axes of Human Disease

Human pathology does not fit into a single reductionist framework. Historically and conceptually, human illnesses span seven distinct mechanobiological axes:

Axis	Primary Mechanism	Classic Paradigm
1. Pathogenic	Exogenous microbial invasion & virulence factors	Plague (<i>Y. pestis</i>), Cholera, HIV, SARS-CoV-2
2. Toxic / Environmental	Chemical poisoning & metabolic disruption	Lead neurotoxicity, Asbestos mesothelioma
3. Mechanical / Structural	Physical wear, trauma, shear stress, & ischemia	Atherosclerotic plaque rupture, osteoarthritis
4. Nutritional / Metabolic	Macronutrient excess or micronutrient deficiency	Scurvy, Type 2 Diabetes, Pellagra
5. Genetic / Congenital	Germline or somatic DNA variation	Sickle cell anemia, Cystic Fibrosis, CML
6. Autoimmune / Degenerative	Loss of self-tolerance & cellular senescence	Type 1 Diabetes, Rheumatoid Arthritis, Alzheimer's
7. Psychosocial / Stress	Chronic HPA axis hyperactivation & allostatic load	Major depression, stress cardiomyopathy

4. Ancient DNA & The ERAP2 Trade-off

In 1348, the Black Death (*Yersinia pestis*) wiped out 30–50% of the European population in fewer than five years. This created one of the most intense natural selection events in human history.

By extracting and sequencing ancient DNA from victims buried in East Smithfield plague pits in London before, during, and after the pandemic, geneticists discovered massive allele frequency shifts in immune loci—most notably the **ERAP2 (Endoplasmic Reticulum Aminopeptidase 2)** gene.

 THE EVOLUTIONARY SCAR OF SURVIVAL

Individuals carrying the protective *ERAP2* variant produced fully functional aminopeptidases that effectively cleaved bacterial antigens for MHC Class I presentation, conferring a **40% survival advantage** against the Black Death. However, in modern sanitized environments, this hyper-reactive variant is a major risk factor for **Crohn's disease and autoimmune colitis**.

| 5. Beyond Humans: Pathogen Phylodynamics

Genomic medicine does not stop at the human nuclear membrane. Modern infectious disease surveillance tracks evolutionary changes across viral and bacterial phylogenies in real time.

Using the molecular clock equation ($d = \mu \times t$), we infer transmission trees, detect cryptic zoonotic spillover events, and identify convergent escape mutations (e.g., SARS-CoV-2 Spike substitutions, H5N1 receptor-binding adaptations) before widespread clinical failure occurs.

| 6. Evidence-Based Medicine vs. The Precision Trap

For four decades, Evidence-Based Medicine (EBM) has relied on the double-blind Randomized Controlled Trial (RCT) as the gold standard. However, RCTs evaluate **population average effects**. In genetically heterogeneous populations, average benefits frequently mask individual non-response.

Pharmaceutical marketing heavily emphasizes **Relative Risk Reduction (RRR)**, which inflates perceived efficacy, while clinical reality is governed by **Absolute Risk Reduction (ARR)** and the **Number Needed to Treat (NNT)**.

$$ARR = Risk_{Control} - Risk_{Treated}, \quad NNT = \frac{1}{ARR}, \quad RRR = \frac{ARR}{Risk_{Control}}$$

💡 THE ASYMMETRIC NNT RATIO (SPOT CHECK 03)

Consider two trials reporting an identical **25% Relative Risk Reduction (RRR = 0.25)**:

- **Trial A (Unselected Population):** Event risk drops from 2.0% to 1.5% (ARR = 0.5%).
 $NNT_A = 1/0.005 = \mathbf{200}$. (You must treat 200 people for 5 years to prevent 1 event; 199 receive zero benefit).
- **Trial B (Genomic High-Risk Cohort):** Event risk drops from 20.0% to 15.0% (ARR = 5.0%).
 $NNT_B = 1/0.05 = \mathbf{20}$.

$$\text{Efficiency Ratio} = \frac{NNT_A}{NNT_B} = \frac{200}{20} = \mathbf{10.0\times}$$

Genomic enrichment increases clinical efficiency by an order of magnitude.

| 7. CML & The Gleevec Precision Revolution

Before 2001, Chronic Myeloid Leukemia (CML) was a fatal diagnosis with a 3–5 year median survival. Treatment consisted of non-specific cytotoxic chemotherapy (busulfan, hydroxyurea) and interferon, which produced debilitating toxicity without eradicating the malignant clone.

In 1960, Nowell and Hungerford identified the tiny **Philadelphia chromosome** in Philadelphia, PA. Decades later, molecular cloning proved this was a reciprocal translocation between chromosomes 9 and 22:



The BCR-ABL fusion protein is a constitutively active tyrosine kinase that continually phosphorylates downstream substrates without requiring external growth signals, driving uncontrolled granulocyte proliferation.

💊 RATIONAL DRUG DESIGN: THE IMATINIB (GLEEVEC) PARADIGM

Rather than poisoning dividing cells, biochemists modeled the inactive ATP-binding pocket of the ABL kinase domain and synthesized **Imatinib (STI-571 / Gleevec)**. Gleevec forms precise hydrogen bonds within the pocket, locking the enzyme in an inactive conformation. 5-year survival jumped from <30% to **>90%** with minimal side effects.

| 8. Pharmacogenomic Case Studies in Practice

Today, genomic stratification determines efficacy and toxicity across billions in prescription drug spending:

Drug	Indication	Biomarker	Population Frequency	Clinical Impact
Plavix (<i>clopidogrel</i>)	Post-stent / Stroke Antiplatelet	CYP2C19 (*2, *3) Loss-of- Function	~15% Europeans ~30% East Asians	Prodrug cannot be activated; causes 3.5× fatal stent thrombosis . Requires switch to Prasugrel.
Keytruda (<i>pembrolizumab</i>)	Solid Tumors / Melanoma	MSI-H / dMMR or TMB ≥ 10	~2–4% Pan- cancer solid tumors	Hypermutated tumors generate dense neoantigens; achieves >50% objective response across tissue types.
Herceptin (<i>trastuzumab</i>)	Invasive Breast Cancer	ERBB2 / HER2 Amplification	~15–20% Breast tumors	Mandatory companion diagnostic (IHC 3+/FISH+). In unamplified patients, causes cardiotoxicity with zero benefit.
Tagrisso (<i>osimertinib</i>)	Non-Small Cell Lung Cancer	EGFR T790M Gatekeeper Mutation	~50–60% of relapses on 1st-gen TKIs	Overcomes steric hindrance from Darwinian resistance mutations at \$17,000/month.