

# **Chocolate Feast**



| Problem Submissions Leaderboard Discussions Editorial Topics |
|--|
|--|

#### **Problem Statement**

Little Bob loves chocolate, and he goes to a store with \$N in his pocket. The price of each chocolate is \$C. The store offers a discount: for every M wrappers he gives to the store, he gets one chocolate for free. How many chocolates does Bob get to eat?

## **Input Format:**

The first line contains the number of test cases, T.

T lines follow, each of which contains three integers, N, C, and M.

## **Output Format:**

Print the total number of chocolates Bob eats.

## **Constraints:**

1 < T < 1000

 $2 < N < 10^5$ 

1 < C < N

 $2 \leq M \leq N$ 

#### Sample input

```
3
10 2 5
12 4 4
6 2 2
```

## **Sample Output**



#### **Explanation**

In the first case, he can buy 5 chocolates with \$10 and exchange the 5 wrappers to get one more chocolate. Thus, the total number of chocolates is 6.

In the second case, he can buy 3 chocolates for \$12. However, it takes 4 wrappers to get one more chocolate. He can't avail the offer and hence the total number of chocolates remains 3.

In the third case, he can buy 3 chocolates for \$6. Now he can exchange 2 of the 3 wrappers and get 1 additional piece of chocolate. Now he can use his 1 unused wrapper and the 1 wrapper of the new piece of chocolate to get one more piece of chocolate. So the total is 5.

Copyright © 2015 HackerRank. All Rights Reserved

**Related Topics** 

Div Mod

Submissions: 38418

Max Score: 25

Difficulty: Easy

More



Join us on IRC at #hackerrank on freenode for hugs or bugs.

Contest Calendar | Blog | Scoring | Environment | FAQ | About Us | Support | Careers | Privacy Policy | Request a Feature