

Abstract

Interspecific competition plays an important role in structuring communities but few studies have been done on herbivores. We present a competition model that predicts the outcome of competition between herbivore species competing for plants. Our model imbeds well-known concepts of the resource-ratio theory, such as the minimum level of resources, consumption vectors, and the quotas of resources required in the biomass. However, unlike traditional approaches that focused on plants as resources, we suggest that chemical elements and energy bounded in plant biomass represent the ultimate resources that herbivores compete for. Our model shows that the outcomes of competition between herbivores result from two main processes: the minimal requirement of resource for herbivores (R^*), and spatial segregation of resources embedded into different plants. The first process (minimal requirement) follows the classical R^* rule and determines the competitive exclusion principle. On the other hand, foraging strategy of herbivores can allow coexistence because resources are spatially segregated. Put together, these two processes determine how herbivore species can coexist, since resources are bounded. Hence, a plant community rich in one resource should support an herbivore community rich in the same resource, leading to a bottom-up effect.

Introduction

Interspecific competition for resources is thought to play an important role in structuring communities (Gause, 1934; Tilman, 1987). Modelling approaches have proven helpful to predict competitive outcomes, when resources are well identified and the number of different resources is limited. Hence, the theory is particularly well suited for autotrophic organisms (e.g., algae or plants) competing for essential nutrients (Tilman, 1982). Indeed, nutrients (i.e., chemical elements constituting biomass) are well-identified non-substitutable resources, and usually, competition is acute for only a few limiting nutrients, such as, for instance, phosphorous (P), nitrogen (N), or potassium (K). In short, models predict that niche segregation along resource ratios should promote species coexistence. For example, plants with low N:P requirement are more likely to coexist with plants with high N:P requirement. Tilman (1980) provided a graphical representation of these competition-coexistence processes based on Zero-Net-Growth Isoclines and consumption vectors, usually referred to as the resource-ratio theory. In the last few decades, the resource-ratio theory has helped popularize the use of competition models to predict competitive outcomes in experimental set ups and in semi-natural conditions.

However, the transfer of such a theory to higher trophic levels is not obvious. The main reason is that for heterotrophic organisms, resources are not easy to characterize. Consider for instance the case of herbivores. If one considers that plant species are the resources herbivores compete for, questions arises: to what extent does a given herbivore require more of a given plant species than another? Should plant species be considered essential non-substitutable resources, or substitutable resources? For example, the diet of the red deer (*Cervus elaphus*) in Europe includes

145 different plant species (Gebert and Verheyden-Tixier, 2001). How many of these
species are really essential for red deer, and how many of the essential resources
are substitutable? This question is a research topic in itself. As a consequence,
predicting competitive outcomes in heterotrophic communities with simple models
remains extremely challenging and restricted to very specific cases (Murray and
Baird, 2008). This limitation may prevent significant progress in our understanding
of community structure at higher trophic levels.

In accordance with the theory of biological stoichiometry (Sterner and Elser,
2002), we argue here that nutrients and energy contained in plant biomass rather
than the plants themselves are the resources which limit the growth rate of herbivores
and for which herbivores compete for. In this case, most of the upper-cited challenges
disappear. Indeed, nutrients and energy clearly are essential non-substitutable re-
sources for heterotrophs. For example, nitrogen is required in proteins and nucleic
acids, and phosphorous is required in phospholipids, nucleic acids, or bones. The
number of essential nutrients required for heterotrophic organism biomass of het-
erotrophs does not exceed 26 (Sterner and Elser, 2002). The fact that herbivore's
growth rate may be limited by food quality (e.g., the level of nitrogen or phosphorus
in plant biomass) rather than food quantity has been widely documented (Sterner
et al., 1992; Hessen, 1992; Urabe and Watanabe, 1992). Moreover, it is possible to de-
termine the requirements of a given herbivore for a given element, using metabolism
studies (Mould and Robbins, 1981). In addition, it is clear that the ratios of nutri-
ents required in the biomass vary across herbivore species. For example, the C:N:P
ratio markedly varies among zooplanktonic herbivores (Andersen and Hessen, 1991;
Sterner et al., 1992). In table 1, we show that such a variation is observed as well
in terrestrial herbivores. Based on these observations, a competition model based
on niche segregation across herbivore species along energy and nutrient axes appears
perfectly relevant. All these arguments support the idea that the resource ratio the-
ory could be applied to herbivores if one considers that just like plants, herbivores
are ultimately limited by energy and nutrients (figure 1).

Yet, a major challenge remains: while plants take up energy and different nu-
trients separately, herbivores consume these resources already packaged within the
plant biomass they ingest. This constraint precludes a straightforward use of the re-
source ratio theory. In this paper, we introduce a theoretical framework that adapts
the classical resource-ratio theory to the specific case of herbivores consuming nutri-
ents and energy that are bounded in fixed ratios in plant biomass. We show that it
is possible to predict the competitive outcomes based on plant stoichiometry, and on
herbivore feeding strategies.

The model

Simulations and graphics were performed with the R software (R Development Core
Team, 2013). Equations for isoclines and consumption vector slopes were found
analytically, as well as stability of equilibrium points. Results were also tested with
simulations using Runge-Kutta 4 approximation, from package deSolve (Soetaert
et al., 2010).

86 The model includes p herbivore species competing for k resources embedded into
 87 n plant species. For each resource u , each herbivore has its own specific require-
 88 ment, and each plant contains its own resource availability. These requirements and
 availabilities are represented by quotas (i.e., quantity of resource per biomass unit).
 90 Hence, for an herbivore i , the biomass dynamic writes:

$$\frac{dH_i}{dt} = \left(\text{Min} \left\{ \frac{\sum_{j=1}^n g_{ij} V_j Q_{vjR1}}{Q_{hiR1}}, \dots, \frac{\sum_{j=1}^n g_{ij} V_j Q_{vjRk}}{Q_{hiRk}} \right\} - m_i \right) H_i \quad (1)$$

Where H_i is herbivore i biomass, V_j is plant j biomass, g_{ij} is consumption rate of
 92 plant j by herbivore i , m_i is mortality rate of herbivore i , Q_{vjRk} is the quota of
 resource k into plant j biomass (for example: g R_k /kg of plant j), and Q_{hiRk} is
 94 quota of resource k into herbivore i biomass (for example: g R_k /kg of herbivore i).
 The ratio between herbivore and plant quotas sets influence of this chemical element
 96 on growth limitation for this herbivore. According to Liebig's law of minimum, the
 least available resource relative to herbivore requirements is assumed to be growth
 98 limiting.

Plant biomass is:

$$\frac{dV_j}{dt} = S_j - a_j V_j - \sum_{i=1}^p g_{ij} V_j H_i \quad (2)$$

100 Where S_j is a function representing the increase of plant biomass (gross supply), a_j
 is intrinsic loss per capita, such as senescence, the last term on the right represents
 102 consumption by herbivores.

Resource availability is described as the plant biomass multiplied by the plant quota
 104 of the considered element.

$$R_u = \sum_{j=1}^n V_j Q_{vjRu} \quad (3)$$

Table 2 represents a review of the state variables and the parameters used in
 106 the model. Indeed, resources are provided by several plants, and each plant provides
 several resources. The first case is a spatial segregation of resources, while the second
 108 case is a stoichiometric balance issue between plant and herbivore.

Spatial segregation of resources

110 A given resources may be provided by only one plant or several plants, which creates a
 spatial segregation of resources. First of all, consider a simple model with p herbivore
 112 species feeding on one plant species embedding a single resource. In case of one
 resource involved, this resource (R) is assumed to be growth limiting.

114 For a matter of simplicity, in case of one plant and one resource, resource quota
 in plant biomass (Q_{vjRk}) is written Q_{vR} , and resource quota in herbivore i biomass
 116 (Q_{hiRk}) is written Q_{hiR} . At steady state, for any herbivore i , the model leads to the
 following equations (see appendix 1 for details):

$$\overline{H}_i = \frac{g_i Q_{vR} S - a Q_{hiR} m}{g_i Q_{hiR} m} \quad (4)$$

Therefore, at equilibrium, herbivore biomass increases with plant availability (supply) and resource quota in plant biomass (Q_{vR}), while it decreases with herbivore mortality and resource quota in herbivore biomass (Q_{hiR}). Resource availability at steady state is:

$$\overline{R}_{hi} = \overline{V} Q_{vR} = \frac{S}{a + g_i \overline{H}_i} Q_{vR} \quad (5)$$

This \overline{R}_{hi} represents level of resource remaining available for another herbivore, it also represents the minimal threshold of resource availability for herbivore species i (i.e., if resource availability is below this threshold this herbivore species cannot maintain a population at steady state). Indeed, it can be assimilated to Tilman's R^* . Therefore, considering two herbivores (species 1 and species 2), if $g_1 \overline{H}_1 > g_2 \overline{H}_2$, species 1 depletes resource below the threshold of species 2, and species 2 is excluded. In other words, the competitive exclusion principles holds, that is, no more than one herbivore species can persist on one limiting resource if this resource is provided by a single plant.

However, consider now that this resource (R) is provided by n plants. Herbivores have access to the resource through the consumption of several species of plants, depending on their feeding strategy. At steady state, it is possible to calculate \overline{R}_{hi} for each herbivore i (see appendix 2 for details):

$$\overline{R}_{hi} = \sum_{j=1}^n \frac{S_j}{a_j + g_{ij} \overline{H}_i} Q_{v1R} \quad (6)$$

For simplicity, consider the case where two herbivore species (H_1 and H_2) compete for one resource (R) embedded into two plants species (V_1 and V_2). Extending equation 6, resource availabilities for each herbivore species at steady state are:

$$\overline{R}_{h1} = \frac{S_1}{a_1 + g_{11} \overline{H}_1} Q_{v1R} + \frac{S_2}{a_2 + g_{12} \overline{H}_1} Q_{v2R} \quad (7)$$

and

$$\overline{R}_{h2} = \frac{S_1}{a_1 + g_{21} \overline{H}_2} Q_{v1R} + \frac{S_2}{a_2 + g_{22} \overline{H}_2} Q_{v2R} \quad (8)$$

Clearly, the competition outcome depends on the foraging strategies of the competitors, more specifically, on the efficiencies of the consumption functions (g_{ij}). If $g_{11} > g_{21}$ and $g_{12} > g_{22}$, herbivore 1 consumes both plants more efficiently than its competitor. Thus, the competitive exclusion principle holds, and herbivore 2 is excluded. The reverse situation ($g_{11} < g_{21}$ and $g_{12} < g_{22}$) leads to the exclusion of herbivore 1. However, if the two herbivores are specialized on different plant species $g_{11} > g_{21}$ and $g_{12} < g_{22}$, or if $g_{11} < g_{21}$ and $g_{12} > g_{22}$, coexistence is possible. In other words, despite the fact that herbivores compete for a single resource, the competitive exclusion principle does not necessarily hold. Indeed, plants species create a spatial segregation of the resource, and the specialization of herbivores on different plant species, similar to niche segregation in space, makes coexistence possible.

Bottom-up stoichiometry

Resource ratio in a given plant interacts with resource ratio of the herbivore, thus constraining persistence of herbivores. It is a bottom-up effect mediated by stoichiometry. Consider p herbivores feeding on one plant species embedding k resources. According to the Liebig's law of the minimum expressed by the minimum function in equation 1, for each herbivore i the interplay between resource requirement and resource availability will determine which resource is the most limiting for growth (Grover, 1997). Depending on their requirements, and on the ratios of resources available in plant biomass, two herbivores feeding on a single plant species may not be limited by the same resource. Thus, requirement differences among herbivores can be interpreted as niche segregation. For a single herbivore i , feeding on the plant, the level of limiting resource R_u (with $u \in [1, k]$) in plant biomass at equilibrium writes (see appendix 3 for details):

$$\bar{R}_{uhi} = \bar{V}|_{Ru} Q_{vRu} = \frac{m_i}{g_i} Q_{hiRu} \quad (9)$$

Where $\bar{V}|_{Ru}$ is plant biomass at steady state when R_u is limiting. Similarly to cases 1 and 2, this level represents the level of resource available to another herbivore and is therefore a key driver of competitive outcomes. It can be graphically represented by a zero net growth isocline (ZNGI) in the phase space of the k resources in plant biomass. One can also represent how resources are consumed by the herbivore with a consumption vector in this phase space (Tilman, 1980). As for the classical resource-ratio theory (Tilman, 1982), we will show here that this graphical representation is a valuable tool to address the competitive outcomes between herbivore species.

For simplicity, consider two herbivore species (H_1 and H_2) competing for two resources (R_1 and R_2) embedded into a single plant species (V). According to equation 9, the slopes of the ZNGIs only depend on herbivore parameters, which are constant. Therefore, the ZNGIs are parallel to the axis on the phase plan $\{R_1, R_2\}$. According to the classic resource-ratio theory (Tilman, 1982), a first necessary condition for coexistence at equilibrium is that the ZNGIs of the two herbivores cross one another. A second necessary condition relies on the difference between the slopes of the consumption vectors of the competitors (Tilman, 1982). The consumption vectors graphically illustrate how resource consumption drives the levels of resources from resource supply to equilibrium levels. In the phase plan, resource supply is represented by the supply point S , that is, the total amount of resources at equilibrium in absence of consumption:

$$S_{R_1} = Q_{vR_1}(S - a\bar{V}) \quad (10)$$

$$S_{R_2} = Q_{vR_2}(S - a\bar{V}) \quad (11)$$

For a given herbivore i , the consumption vector \vec{C}_i writes:

$$\vec{C}_i = \begin{pmatrix} g_i \bar{V} \bar{H}_i Q_{vR_1} \\ g_i \bar{V} \bar{H}_i Q_{vR_2} \end{pmatrix} = g_i \bar{V} \bar{H}_i \begin{pmatrix} Q_{vR_1} \\ Q_{vR_2} \end{pmatrix} \quad (12)$$

The vectors slope is determined by the ratios of resources embedded in plant biomass. An important consequence is that all herbivores have the same slope of consumption

vector when feeding on a single given plant. This constraint precludes coexistence,
 188 and the resource-ratio embedded in plant biomass determines which of the com-
 petitor will displace the other (figure 2). Hence, despite niche segregation on the
 190 requirement axe, two herbivores cannot coexist on two resources. This result is in
 contrast with the classic resource-ratio theory, which stipulates that two consumer
 192 may coexist on two resources. The reason is that classically, consumers are assumed
 to control the slope of their consumption vector (Tilman, 1982). This assumption
 194 is valid for primary producers, which take up essential nutrients independently in
 their environment, in the form of dissolved chemical molecules, but not for herbi-
 196 vores, which take up essential nutrients already bounded in plant biomass. Therefore,
 niche segregation alone cannot promote coexistence. Only herbivore that can match
 198 plant stoichiometry the best can persist and outcompete other herbivores.

Case 4: several plant species and several resources

In this section, we present the general case of the model, where herbivores compete
 200 for multiple resources embedded into multiple plants. Assuming that nutrient quo-
 202 tas differ among plant species, the consumption of a given plant is not equivalent
 to consumption of another plant. As in case 2, herbivores may control their diet by
 204 allocating more time/energy to feed on specific plants rather than others. There-
 fore, each herbivore may find a specific pathway to collect resources (Simpson and
 206 Raubenheimer, 1995; Raubenheimer and Simpson, 1999). The foraging strategy of
 a given herbivore is graphically represented by the consumption vector, which is the
 208 combination of the consumption vectors for all the plant species consumed. As a
 consequence, the slope of the consumption vector depends on the foraging strategy
 210 of the herbivore, which can adjust the slope by consuming more efficiently a given
 species versus the others. Thus, niche segregation among herbivore species is a com-
 212 bination of segregation in space (as in case 2) and segregation on the requirements (as
 in case 3). As we shall see, this niche segregation allows for coexistence of multiple
 214 herbivores on multiple resources.

To simplify, we consider here the case of two herbivores competing for two plants
 216 embedding two resources (see appendix 4 for details). The resource supply is repre-
 sented in the phase plan by the supply point S, which is the total amount of resources
 218 at equilibrium in plants 1 and 2, in absence of consumption:

$$S_{R1} = Q_{v1R1}(S_{v1} - a\bar{V}_1) + Q_{v2R1}(S_{v2} - a\bar{V}_2) \quad (13)$$

$$S_{R2} = Q_{v1R2}(S_{v1} - a\bar{V}_1) + Q_{v2R2}(S_{v2} - a\bar{V}_2) \quad (14)$$

220 For a given herbivore i , the ZNGI slope ($\alpha|_{Ru}$) for each resource (R_u) is:

$$\alpha|_{Ru} = \frac{\bar{V}_1|_{Ru}Q_{v1R2} + \bar{V}_2|_{Ru}Q_{v2R2}}{\bar{V}_1|_{Ru}Q_{v1R1} + \bar{V}_2|_{Ru}Q_{v2R1}} \quad (15)$$

Note that the slope depends on $\{\bar{V}_1, \bar{V}_2\}$, which themselves depend both on the
 222 herbivore and the plant parameters (including the supply parameters). Hence, the
 ZNGI slope is not parallel to the axis.

224 However, it is possible to define boundary ZNGI. Considering that herbivore i can
 226 consume both plants, its foraging strategy will lie between exclusive consumption of
 plant 1, on one side, and exclusive consumption of plant 2, on the other side. Thus,
 boundary ZNGI slope writes (see appendix 4 for details):

$$\bar{R}_u = \frac{Q_{hiRu}m_i}{g_{ij}} \quad (16)$$

228 This isocline \bar{R}_u represents the ZNGI in the case where herbivore i consumes only
 plant j and is limited by nutrient u . In case on two plants consumed, the real \bar{R}_u
 230 will lie between two boundary ZNGI (one for each plant).

By looking at boundary ZNGI equation (16), it appears that ZNGI slopes de-
 232 pends only on herbivore constant parameters. Therefore, it is parallel to the axis.
 Moreover, two herbivores having similar quotas or different feeding strategies would
 234 have different ZNGI. Thus, if boundary ZNGI of two herbivores cross each other,
 coexistence might be possible.

236 Next, competition outcome will depend on the relative orientation of the con-
 sumption vector of herbivores (Tilman, 1980). However, vector slope calculation is
 238 uneasy in the case of packaged resources because vector slope depends on plant sup-
 ply points. Hence, for a given herbivore, vectors do not have a constant slope valid
 240 for all supply conditions. This is why instead of consumption vectors, the model
 allows the calculation of boundary vectors that determine an area in the phase plan
 242 where each herbivore species could coexist with an other one. It is possible to deter-
 mine a boundary relation between plant supply points (S_1 and S_2) that allows each
 244 herbivore to consume both resources when one resource is limiting, and leads to \bar{V}_1
 and \bar{V}_2 at steady state. These boundary supply points are S_{1hi} and S_{2hi} for plant 1
 246 and 2 respectively (see appendix 4). Hence, boundary vector slopes are:

$$\vec{C}_{h1} = \begin{pmatrix} \bar{S}_{1h1} Q_{v1R1} + \bar{S}_{2h1} Q_{v2R1} \\ \bar{S}_{1h1} Q_{v1R2} + \bar{S}_{2h1} Q_{v2R2} \end{pmatrix} \quad (17)$$

$$\vec{C}_{h2} = \begin{pmatrix} \bar{S}_{1h2} Q_{v1R1} + \bar{S}_{2h2} Q_{v2R1} \\ \bar{S}_{1h2} Q_{v1R2} + \bar{S}_{2h2} Q_{v2R2} \end{pmatrix} \quad (18)$$

248 These two boundary vectors play a similar role as consumption vectors in Tilman's
 model. However, as resources are packaged into plants, vector slopes depend on the
 250 way herbivore species feed on plants containing resources. Hence, assuming plant 1
 is richer in resource 1 ($Q_{v1R1} > Q_{v2R1}$), and plant 2 is richer in resource 2 ($Q_{v1R2} <$
 252 Q_{v2R2}), and assuming herbivore 1 is more limited by resource 1 ($Q_{h1R1} > Q_{h1R2}$),
 and herbivore 2 is more limited by resource 2 ($Q_{h2R1} < Q_{h2R2}$), three different strate-
 254 gies can be considered. The first one occurs when each herbivore species consumes
 preferentially the plant species which gives the greater quantity of the most limiting
 256 resource for this herbivore species, and which is less limiting for its competitor (i.e.,
 $g_{11} > g_{12}$ and $g_{21} < g_{22}$). In that case, if we consider the boundary vectors for
 258 herbivore 1 and for herbivore 2, we can define several zones on the phase plan (see
 figure 3). The zone between the two boundary vectors represents the supply condi-
 260 tions allowing stable coexistence of the two herbivore species. Thus, each herbivore

consumes the most profitable plant (i.e., quotas and consumption functions follow a similar trend), but each herbivore is specialized (more or less) on a different plant than its competitor, which promotes coexistence.

The second case represents the reverse situation, where each herbivore species consumes preferentially the plant species which provides greater quantity of the most limiting resource for its competitor (i.e., $g_{11} < g_{12}$ and $g_{21} > g_{22}$). Hence, the zone between the boundary vectors does not allow coexistence (see figure 4), and in many cases equation 32A does not have any realistic solution (with both herbivore species persisting). Thus, competitive exclusion is the general outcome.

The third case occurs when herbivore species are not selective at all: they consume plant species according to their respective biomass. In that case, both herbivores consume resources in the same way (i.e., $g_{11} = g_{21}$ and $g_{12} = g_{22}$). Hence, there is no segregation in herbivore consumption: all herbivore species have the same consumption vector, which is similar to case 3, where only one plant species is present (see figure 2). Therefore, the trajectory will either cross herbivore 1 ZNGI first, and this herbivore will be excluded, or the trajectory will cross herbivore 2 ZNGI first, and this herbivore will be excluded. Only a particular case, where the trajectory leads exactly on the crossing point, allows coexistence.

If we release the previous assumptions on herbivore and plant quotas (i.e., inequality for herbivore requirements, and inequality for plant profitability), two other situations can potentially occur. First, both herbivores can have the same quotas (i.e., $Q_{h1R1} = Q_{h2R1}$ and $Q_{h1R2} = Q_{h2R2}$) but different consumption functions. In that case, one plant will be more interesting for both herbivores (i.e., the plant that is richer for the most limiting resource). Hence, the herbivore that can consume this plant the most efficiently will exclude its competitor. Second, both plants can have the same quotas (i.e., $Q_{v1R1} = Q_{v2R1}$ and $Q_{v1R2} = Q_{v2R2}$). In that case, the system is similar to the case where only one plant is present (case 3), which leads to competitive exclusion.

Last, due to resource packaging into plants, the supply point is constrained. Resource ratio within each plant will define a slope:

$$\alpha_{vj} = \frac{Q_{vjR2}}{Q_{vjR1}} \quad (19)$$

In case of two plants, supply point will lie between the two extreme slopes (α_{v1} and α_{v2}) defining a feasibility cone. Therefore, the whole phase plan is not available. Resource supply occurs only within this feasibility cone, which adds another constraint on herbivore persistence and competition.

Discussion

Unlike classical studies on herbivore competition we consider here that the resource limiting herbivores growth is not plant biomass but rather, the essential nutrients contained in plant biomass.

According to our results, the fundamental difference between herbivore competition and plant competition is that resources are not independent, which leads to

specific constraints for herbivores. Thus, several differences exist between classical
 302 resource-ratio theory of competition usually applied to plants (Tilman, 1980) and
 our model for herbivores. First, if only one plant species is available, coexistence is
 304 almost impossible even if herbivore compete for several resources. Second, ZNGI can
 be represented with boundary ZNGI delimiting resource levels at steady state.

306 Third, consumption vectors are not informative because their slope varies with
 supply points. However, boundary vectors can be calculated, and they play a similar
 308 role as usual consumption vectors. Last, due to this packaging effect of resources
 and to plant quotas, resource availability itself is constrained. Hence, part of the
 310 phase plan might be unavailable. This is a major difference with models assuming
 a total independence between resources (such as N and P for plants) within which
 312 availabilities of resources can vary independently from one another. When resources
 are packaged, they are not independent. Thus, some supply couples (e.g., large
 314 quantity of R_1 and almost no R_2 available) is unlikely because plants will provide
 both resources (according to their own ratio). Hence, parameter space allowing
 316 herbivore persistence can be narrower than predicted by its ZNGI and vector.

More generally, existing theories about coexistence between herbivores are based
 318 on niche segregation. This segregation may depend on space utilization or on rela-
 tionship between body size and metabolism (Owen-Smith and Novellie, 1982). Our
 320 approach is different and allows us to disentangle the different mechanisms of niche
 segregation. We argue that coexistence between herbivores can occur by two ways,
 322 which are diversity between foraging strategies as well as stoichiometric diversity
 between niches. Diversity between foraging strategies is a spatially niche segregation
 324 within which each herbivore species consumes one plant species more than others,
 assuming that this plant species has the best profitability for this herbivore species.
 326 If this spatial segregation is total, with each herbivore species specialized on one
 plant species which is different from other competitors, coexistence occurs without
 328 other constraint than herbivore species-specific persistence. The second way is niche
 segregation based on resource ratios (i.e., a bottom-up stoichiometric component).
 330 Although the packaging effect creates a supplementary level of complexity, divergence
 with classic resource-ratio models should not impede the rising of a clear conclusion:
 332 coexistence is favoured by requirement segregation as well as consumption segre-
 gation; only one of them (i.e., different requirements and similar consumptions, or
 334 similar requirements and different consumptions) is not enough. We retrieve here
 the two classical components of the niche theory (Chase and Leibold, 2003): species
 336 requirement (Hutchinson, 1957) and species impact (Elton, 1927; Macarthur and
 Levins, 1967). However, while these two components are clearly separated for com-
 338 petition between plants (i.e., ZNGI represent species requirements, and vectors rep-
 resent species impact), these two components are merged for herbivore competition
 340 (boundary ZNGIs and boundary vectors involve both aspects).

Nonetheless, our model allows us to disentangle the relative effects of the bottom-
 342 up stoichiometric component (i.e., plant and herbivore resource quotas) and the non-
 stoichiometric component (i.e., spatial segregation) of the competitive interaction.
 344 Indeed, the foraging strategy of the competing herbivores will mainly determine
 the competitive outcome. Non-selective herbivores (representing case 2 and case
 346 4 with non-selective behavior) will show a strong bottom-up stoichiometric effect.

Therefore, coexistence is unlikely. For example, non-selective zooplanktonic filters, such as cladocerans (limited by P) and copepods (limited by N), usually do not coexist: either one group or the other dominates according the N:P ratio of the consumed algae (Andersen and Hessen, 1991; Hessen, 1992; Sterner et al., 1992; Elser et al., 1996; Koski, 1999). A similar assumption can be made for non-selective terrestrial grazer herbivores (Albon and Langvatn, 1992) for which coexistence would be unlikely. On the other hand, selective herbivores, specialized on a few plants, would show a strong spatial component and a weaker stoichiometric component of the interaction (representing case 2 and case 4 with selective behavior). Therefore, they should be more prone to coexist.

Concerning the stoichiometric component, our assumption is that a stoichiometric diversity between niches exists among herbivore species, and field studies tend to show it. For aquatic herbivores, such as zooplankton, it seems that grazers (especially *Daphnia sp.*) are P-limited, while copepods seem to be N-limited (see above). This stoichiometric diversity exists for terrestrial herbivores either (see table 1). Chemical requirements for wild herbivores are mostly unknown, but for mammal herbivores, numerous studies have used data from cattle as proxy for diet requirement for wild species (Voeten and Prins, 1999). Nevertheless, data begins to be available. For example, N requirements have been estimated for fawns (Smith et al., 1975), yearlings (Holter et al., 1979) and adults (Asleson et al., 1996) of white-tailed deer (*Odocoileus virginianus*), as well as P requirement for white-tailed deer (Grasman and Hellgren, 1993) and moose (Schwartz et al., 1987). Moreover, males from big species should be more concerned by P-limitation (Grasman and Hellgren, 1993). More generally, it seems that nutrient requirement and absorption constraint depend on body size and digestive system (Janis, 1976).

Concerning the spatial component and the foraging strategy, several studies have been done, mainly on terrestrial herbivore species. It seems that most of migrated species adopt an extraction maximizing strategy: through seasons, they move from places to places that have a large amount of nutritive quality elements (Albon and Langvatn, 1992). There, they can find plant communities with a high proportion of nutritional plant species. They adopt this strategy rather than selecting nutritious species within communities (Ben-Shahar and Coe, 1992). On the other hand, resident species mostly adopt a demand minimizing strategy: sometimes, they have a low metabolic rate, and flexible breeding period, which allow them to decrease demand in energy during dry period (Murray, 1991). Moreover, for continental herbivore species, the diet quality decreases when body size increases, especially during dry season (Codron et al., 2007).

However, there are some limits to this model. We pay only attention to the requirement for a given herbivore species. But we do not consider what happens in case of an overconsumption of a non-limiting nutrient. In fact, we can have a more physiological approach. Considering the case where an herbivore species consuming a plant species that gives a small amount of a limiting nutrient and a large amount of a non-limiting nutrient, one can argue that an excretion cost might exist for this nutrient. This can limit the amount of plant consumed by this herbivore species.

Finally, this model gives an interesting conclusion. The stoichiometric constraints should go up through the trophic chain. It means that, on a soil that is poor in a

given nutrient (e.g., nitrogen), we should find plant species that can survive with a
 394 poor availability for this nutrient. Then, they contain a small amount of this nutrient
 and they will sustain herbivore species that are poor in this nutrient, leading to a
 396 strong bottom-up effect.

Appendix 3

398 When p herbivores compete for k resources embedded into 1 plant, equations 1, 2
 and 3 respectively become:

$$\frac{dH_i}{dt} = \left(\text{Min} \left\{ \frac{Q_{vR1}}{Q_{hiR1}}, \dots, \frac{Q_{vRk}}{Q_{hiRk}} \right\} g_i V - m_i \right) H_i \quad (20A)$$

400

$$\frac{dV}{dt} = S - aV - \sum_{i=1}^p g_i V H_i \quad (21A)$$

$$R_k = V Q_{vRk} \quad (22A)$$

402 At steady state, it is possible to determine resource availability for each herbivore i .
 Assuming that R_u is a limiting resource, with $u \in [1, k]$.

$$\bar{V}|_{R_u} = \frac{m_i}{g_i} \frac{Q_{hiRu}}{Q_{vRu}} \quad (23A)$$

404 Where $\bar{V}|_{R_u}$ is the steady state plant biomass when R_u is limiting. \bar{R}_u is:

$$\bar{R}_u = \bar{V}|_{R_u} Q_{vRu} = \frac{m_i}{g_i} Q_{hiRu} \quad (24A)$$

Appendix 4

406 In a case of two herbivore species (H_1 and H_2) competing for two resources (R_1 and
 R_2) embedded into two plant species (V_1 and V_2), the system considered becomes:

$$\frac{dH_i}{dt} = \left(\text{Min} \left\{ \frac{g_{i1} V_1 Q_{v1R1} + g_{i2} V_2 Q_{v2R1}}{Q_{hiR1}}, \frac{g_{i1} V_1 Q_{v1R2} + g_{i2} V_2 Q_{v2R2}}{Q_{hiR2}} \right\} - m_i \right) H_i \quad (25A)$$

408

$$\frac{dV_j}{dt} = S_j - a_j V_j - g_{1j} V_j H_1 - g_{2j} V_j H_2 \quad (26A)$$

$$R_1 = q_{v1R1} V_1 + q_{v2R1} V_2 \quad (27A)$$

410

$$R_2 = q_{v1R2} V_1 + q_{v2R2} V_2 \quad (28A)$$

Each herbivore has two zero net growth isoclines (ZNGI), but their calculation is
 412 more complicate because of multiple sources for R_1 and R_2 . However, it is possible
 to define boundary ZNGI. Considering that herbivore i can consume both plants,

its foraging strategy will lie between exclusive consumption of plant 1, on one side, and exclusive consumption of plant 2, on the other side. Thus, for an herbivore i consuming each plant j (with $j \in [1, 2]$) and limited by each resource u (with $u \in [1, 2]$), equation 25A becomes

$$\frac{dH_i}{dt} = \left(\frac{g_{ij}V_jQ_{vjRu}}{Q_{hiRu}} - m_i \right) H_i \quad (29A)$$

At steady state, $\bar{R}_u = \bar{V}_j * Q_{vjRu}$. Thus,

$$\frac{g_{ij}\bar{V}_jQ_{vjRu}}{Q_{hiRu}} - m_i = \frac{g_{ij}\bar{R}_u}{Q_{hiRu}} - m_i = 0 \quad (30A)$$

which leads to

$$\bar{R}_u = \frac{Q_{hiRu}m_i}{g_{ij}} \quad (31A)$$

\bar{R}_u represents the slope of the boundary ZNGI for herbivore i consuming plant j and limited by resource u . In case of two plants and two resources, each herbivore will have two boundary ZNGI for each resources. The real value of \bar{R}_u will lie between the two boundary ZNGI.

It is possible to calculate \bar{V}_1 and \bar{V}_2 that are the equilibrium biomass of plant 1 and 2 respectively and are solutions of the following system:

$$\begin{cases} g_{11}\bar{V}_1Q_{v1r1} + g_{12}\bar{V}_2Q_{v2r1} = m_1Q_{h1r1} \\ g_{21}\bar{V}_1Q_{v1r2} + g_{22}\bar{V}_2Q_{v2r2} = m_2Q_{h2r2} \end{cases} \quad (32A)$$

If system (32A) has two realistic solutions (i.e., $\bar{V}_1 > 0$ and $\bar{V}_2 > 0$), and if $\bar{H}_1 > 0$ and $\bar{H}_2 > 0$, which means that coexistence occurs at that point, then this point is an equilibrium point. Knowing \bar{V}_1 and \bar{V}_2 at this equilibrium point, it is possible to determine a boundary relation between plant supply points (S_1 and S_2) that allows herbivore species 1 to consume both resources when R_1 is limiting, and leads to \bar{V}_1 and \bar{V}_2 at steady state. Similarly, we can determine a relation that allows herbivore 2 to consume both resources when R_2 is limiting, and leads to \bar{V}_1 and \bar{V}_2 . The relations are:

$$\bar{S}_{2h1} = (\bar{S}_{1h1} - a_1) \frac{g_{12}\bar{V}_2}{g_{11}\bar{V}_1} + a_2\bar{V}_2 \quad (33A)$$

$$\bar{S}_{2h2} = (\bar{S}_{1h2} - a_1) \frac{g_{22}\bar{V}_2}{g_{21}\bar{V}_1} + a_2\bar{V}_2 \quad (34A)$$

Where S_{1h1} and S_{2h1} are the boundary supply points for plant 1 and 2 respectively for herbivore 1. Similarly, S_{1h2} and S_{2h2} are boundary supply points for herbivore 2.

References

Albon, S. D., and R. Langvatn. 1992. Plant phenology and the benefits of migration in a temperate ungulate. *Oikos* 65:502–513.

- 440 Andersen, T., and D. O. Hessen. 1991. Carbon, nitrogen, and phosphorus content of
freshwater zooplankton. *Limnology and Oceanography* 36:807–814.
- 442 ARC. 1980. The nutrient requirements of ruminant livestock. Agricultural Research
Council. CBA, Slough, UK .
- 444 Asleson, M. A., E. C. Hellgren, and L. W. Varner. 1996. Nitrogen requirements for
antler growth and maintenance in white-tailed deer. *Journal of Wildlife Manage-*
446 *ment* 60:744–752.
- Ben-Shahar, R., and M. J. Coe. 1992. The relationships between soil factors, grass
448 nutrients and the foraging behavior of wildbeest and zebra. *Oecologia* 90:422–428.
- Chase, J., and M. Leibold. 2003. Ecological niches: linking classical and contempo-
450 rary approaches. University of Chicago Press.
- Codron, D., J. A. Lee-Thorp, M. Sponheimer, J. Codron, D. De Ruiter, and J. S.
452 Brink. 2007. Significance of diet type and diet quality for ecological diversity of
african ungulates. *Journal of Animal Ecology* 76:526–537.
- 454 Elser, J. J., D. R. Dobberfuhl, N. A. MacKay, and J. H. Schampel. 1996. Organism
size, life history, and n:p stoichiometry. *Bioscience* 46:674–684.
- 456 Elton, C. 1927. Animal ecology. University of Chicago Press.
- Gause, G. 1934. The Struggle for Existence. Williams and Wilkins ed., Baltimore,
458 Maryland. 163 pp.
- Gebert, C., and H. Verheyden-Tixier. 2001. Variations of diet composition of red
460 deer (*cervus elaphus* l.) in europe. *Mammal Review* 31:189–201.
- Grasman, B. T., and E. C. Hellgren. 1993. Phosphorus nutrition in white-tailed deer:
462 Nutrient balance, physiological responses, and antler growth. *Ecology* 74:2279–
2296.
- 464 Grover, J. 1997. Resource Competition. Chapman & Hall ed., London. 332 pp.
- Hessen, D. O. 1992. Nutrient element limitation of zooplankton production. *Ameri-*
466 *can Naturalist* 140:799–814.
- Holter, J. B., H. H. Hayes, and S. H. Smith. 1979. Protein requirement of yearling
468 white-tailed deer. *Journal of Wildlife Management* 43:872–879.
- Hutchinson, G. E. 1957. Population studies - animal ecology and demography - con-
470 cluding remarks. *Cold Spring Harbor Symposia on Quantitative Biology* 22:415–
427.
- 472 Janis, C. 1976. The evolutionary strategy of equidae and origins of rumen and cecal
digestion. *Evolution* 30:757–774.

- 474 Koski, M. 1999. Carbon : Nitrogen ratios of baltic sea copepods - indication of
mineral limitation? *Journal of Plankton Research* 21:1565–1573.
- 476 Ludwig, F., T. E. Dawson, H. de Kroon, F. Berendse, and H. H. T. Prins. 2003.
Hydraulic lift in acacia tortilis trees on an east african savanna. *Oecologia* 134:293–
478 300.
- Macarthur, R., and R. Levins. 1967. Limiting similarity convergence and divergence
480 of coexisting species. *American Naturalist* 101:377–385.
- Menard, C., P. Duncan, G. Fleurance, J. Y. Georges, and M. Lila. 2002. Compara-
482 tive foraging and nutrition of horses and cattle in european wetlands. *Journal of
Applied Ecology* 39:120–133.
- 484 Mould, E. D., and C. T. Robbins. 1981. Nitrogen metabolism in elk. *Journal of
Wildlife Management* 45:323–334.
- 486 Murray, M. G. 1991. Maximizing energy retention in grazing ruminants. *Journal of
Animal Ecology* 60:1029–1045.
- 488 Murray, M. G., and D. R. Baird. 2008. Resource-ratio theory applied to large herbi-
vores. *Ecology* 89:1445–1456.
- 490 Owen-Smith, N., and P. Novellie. 1982. What should a clever ungulate eat. *American
Naturalist* 119:151–178.
- 492 Raubenheimer, D., and S. J. Simpson. 1999. Integrating nutrition: a geometrical
approach. *Entomologia Experimentalis Et Applicata* 91:67–82.
- 494 Schwartz, C. C., W. L. Regelin, and A. W. Franzmann. 1987. Protein digestion in
moose. *Journal of Wildlife Management* 51:352–357.
- 496 Simpson, S. J., and D. Raubenheimer. 1995. The geometric analasis of feeding and
nutrition: a user’s guide. *Journal of Insect Physiology* 41:545–553.
- 498 Smith, S. H., J. B. Holter, H. H. Hayes, and H. Silver. 1975. Protein requirement of
white-tailed deer fawns. *Journal of Wildlife Management* 39:582–589.
- 500 Soetaert, K., T. Petzoldt, and R. W. Setzer. 2010. Solving differential equations in
r: Package desolve. *Journal of Statistical Software* 33:1–25.
- 502 Sterner, R., and J. Elser. 2002. *Ecological stoichiometry: the biology of elements
from molecules to the biosphere*. Princeton Univ Press, New Yersey, USA. 584 pp.
- 504 Sterner, R. W., J. J. Elser, and D. O. Hessen. 1992. Stoichiometric relationships
among producers, consumers and nutrient cycling in pelagic ecosystems. *Biogeo-
506 chemistry* 17:49–67.
- Tilman, D. 1980. Resources - a graphical-mechanistic approach to competition and
508 predation. *American Naturalist* 116:362–393.

- . 1982. *Resource Competition and Community Structure*, vol. 17. Princeton
510 Univ Press, New Jersey, USA. 296 pp.
- . 1987. The importance of the mechanisms of interspecific competition. *Amer-*
512 *ican Naturalist* 129:769–774.
- Treydte, A. C., I. M. A. Heitkonig, and F. Ludwig. 2009. Modelling ungulate depen-
514 dence on higher quality forage under large trees in african savannahs. *Basic and*
Applied Ecology 10:161–169.
- 516 Urabe, J., and Y. Watanabe. 1992. Possibility of n or p limitation for planktonic
cladocerans: an experimental test. *Limnology and Oceanography* 37:244–251.
- 518 Voeten, M., and H. Prins. 1999. Resource partitioning between sympatric wild and
domestic herbivores in the tarangire region of tanzania. *Oecologia* 120:287–294.

Table 1: Minimum daily intake requirements of digestible protein (DP) and phosphorus (P) for four grazer species of different body weight (BW in kg). Modified from Treydte et al. (2009).

Intake (mg/kg BW/day)	Model scenario source	Warthog 83 kg BW	Wildebeest 143 kg BW	Zebra 271 kg BW	Buffalo 481 kg BW
DProtein	ARC (1980)	730	550	430	380
	Menard et al. (2002)	970	850	720	630
	Ludwig et al. (2003)	1040	910	770	680
P	ARC (1980)	9	14	18	20
	Menard et al. (2002)	13	27	37	42
DProtein:P ratio		83	38	23	18

Table 2: State variables and parameters used in the model. Dimensions for each variable and parameter are based on mass (M) and time (T).

Symbols	Definitions	Dimensions
State Variables		
H_i	Herbivore i biomass	M
V_j	Plant j biomass	M
R_k	Resource k mass	M
Parameters		
g_{ij}	plant j consumption rate by herbivore i	$T^{-1}.M^{-1}$
m_i	herbivore i biomass mortality rate <i>per capita</i>	T^{-1}
S_j	plant j biomass supply	$M.T^{-1}$
a_j	plant j biomass natural loss rate <i>per capita</i>	T^{-1}
Q_{hiRu}	resource u quota for herbivore i	dimensionless
Q_{vjRu}	resource u quota for plant j	dimensionless




Figure 1: Each herbivore species has its own resource ratio between resource 1 (R_1) and resource 2 (R_2). This species feeds on different plant species with different resource ratios. Herbivore requirements as well as their resource consumptions are key factors for their persistence.




Figure 2: Phase plan with two herbivore species feeding on one plant species. Black lines are herbivore 1 ZNGI, dotted lines are herbivore 2 ZNGI, and grey arrow is the common consumption vector. Interpretations are quite similar with those from Tilman's model (Tilman, 1982). Roman numbers represent different zones for supply point position. Zone I and zone V do not provide sufficient amounts in R_1 and R_2 respectively. None herbivore species can live in these conditions. Zone II contains enough quantities of R_1 for herbivore 1, but not for herbivore 2. Zone III potentially has enough quantities of both resources for both herbivore species. In zone IV, only herbivore 2 can survive because there is not enough R_2 for herbivore 1. Supply point represents total amount of resources due to plant biomass rebuilding. Both herbivore species sample resources through the same consumption vector. In this example, the trajectory cross herbivore 2 ZNGY first and then herbivore 1 ZNGI. The crossing point with herbivore 1 ZNGI becomes an equilibrium point (E). Herbivore 1 is limited by R_2 . Herbivore 2 is excluded.




Figure 3: Phase plan for two coexisting herbivores. Lines represent boundary ZNGI, arrows are boundary vectors, and grey lines bound feasible supply conditions according to plants considered (i.e., feasibility cone). Continuous ZNGI and vector belong to herbivore 1, and dotted ZNGI and vector belong to herbivore 2. Zone I and zone VII contain not enough R_1 and R_2 respectively. Neither herbivore 1 nor herbivore 2 can survive. Zone II contains not enough R_1 for herbivore 2. Zone III does not allow herbivore 2 to survive if herbivore 1 is present. Zone V does not allow herbivore 1 to survive if herbivore 2 is present. Zone VI contains not enough R_2 for herbivore 1. If the supply point is within zone IV both herbivore can survive together: it is coexistence. Then, the cross zone of the two ZNGIs is a stable equilibrium point (SE).




Figure 4: Phase plan for competitive exclusion. Representation of ZNGI and vectors is similar to figure 3, except for zone IV. This zone generally leads to exclusion of herbivore 1 or 2. The crossing zone of ZNGIs either can be a non stable equilibrium (NSE) or does not allow coexistence of the two herbivore species.