Data from: Influence of intra- and interspecific variation in predator-prey body size ratios on trophic interaction strengths

Cuthbert, Ross N., Queen's University Belfast, https://orcid.org/0000-0003-2770-254X

Wasserman, Ryan J., Botswana International University of Science and Technology

Dalu, Tatenda, University of Venda

Kaiser, Horst, Rhodes University

Weyl, Olaf L. F., SAIAB

Dick, Jaimie T. A., Queen's University Belfast

Sentis, Arnaud, Aix-Marseille University

McCoy, Mike W., East Carolina University

Alexander, Mhairi E., University of the West of Scotland

rossnoelcuthbert@gmail.com, ryanwas21@gmail.com, dalutatenda@yahoo.co.uk, h.kaiser@ru.ac.za, o.weyl@saiab.ac.za, j.dick@qub.ac.uk, arnaud.sentis@gmail.com, mccoym@ecu.edu, mhairi.alexander@uws.ac.uk

Publication date: June 1, 2020

Publisher: Dryad

https://doi.org/10.5061/dryad.7m0cfxppt

Citation

Cuthbert, Ross N. et al. (2020), Data from: Influence of intra- and interspecific variation in predator-prey body size ratios on trophic interaction strengths, v6, Dataset,

https://doi.org/10.5061/dryad.7m0cfxppt

Abstract

1. Predation is a pervasive force that structures food webs and directly influences ecosystem functioning. The relative body sizes of predators and prey may be an important determinant of

Dryad Page 1 of 3

interaction strengths. However, studies quantifying the combined influence of intra- and interspecific variation in predator-prey body size ratios are lacking.

- 2. We use a comparative functional response approach to examine interaction strengths between three size classes of invasive bluegill and largemouth bass towards three scaled size classes of their tilapia prey. We then quantify the influence of intra- and interspecific predator-prey body mass ratios on the scaling of attack rates and handling times.
- 3. Type II functional responses were displayed by both predators across all predator and prey size classes. Largemouth bass consumed more than bluegill at small and intermediate predator size classes, whilst large predators of both species were more similar. Small prey were most vulnerable overall, however differential attack rates among prey were emergent across predator sizes. For both bluegill and largemouth bass, small predators exhibited higher attack rates towards small and intermediate prey sizes, whilst larger predators exhibited greater attack rates towards large prey. Conversely, handling times increased with prey size, with small bluegill exhibiting particularly low feeding rates towards medium-large prey types. Attack rates for both predators peaked unimodally at intermediate predator-prey body mass ratios, whilst handling times generally shortened across increasing body mass ratios.
- 4. We thus demonstrate effects of body size ratios on predator-prey interaction strengths between key fish species, with attack rates and handling times dependent on the relative sizes of predator-prey participants.
- 5. Considerations for intra- and interspecific body size ratio effects are critical for predicting the strengths of interactions within ecosystems and may drive differential ecological impacts among invasive species as size ratios shift.

Funding

National Research Foundation, South Africa, Award: 110507

National Research Foundation, South Africa, Award: 117700

Alexander von Humboldt-Stiftung, Award: N/A

U.S. National Science Foundation, Award: 1556743

References

This dataset is supplement to https://doi.org/10.1002/ece3.6332

Dryad Page 2 of 3

Keywords

predator-prey interaction, size ratios, bluegill, largemouth bass, tilapia

Files

1 files for this dataset

data.csv 6.45 kB application/vnd.ms-excel

License

This work is licensed under a <u>CCO 1.0 Universal (CCO 1.0) Public Domain</u> <u>Dedication</u> license.



This releases your work to the public domain for any use.

Dryad Page 3 of 3