

AI AgriYield Predictor: Final Advisory Report

1. Farm Overview & Predicted Yield

State: California

Crop: Almonds

N-P-K: 180-50-150 kg/ha

Soil pH: 6.5

Rainfall: 600 mm

Temperature: 25 C

Predicted Yield: 2350.50 kg/acre

2. Agronomic Advisory Plan

Yield Benchmarking

The predicted yield of 2,350.50 kg/acre (approximately 5,181 lbs/acre) represents an exceptionally high performance benchmark for California almond production.

* Regional Context: The standard average yield for mature almond orchards in California typically ranges between 900 kg/acre and 1,150 kg/acre (2,000 to 2,500 lbs/acre) of nut meats.

* Evaluation: A target of 2,350.50 kg/acre places this operation in the top tier of high-density, high-yielding production blocks. While this yield is biologically achievable under intensive precision management, it requires strict alignment of nutrient supply and water delivery to sustain the crop load without inducing stress or alternate bearing in subsequent seasons.

Soil Health & Nutrient Remediation

#

Soil pH Analysis

The soil pH of 6.5 on loam soil is within the ideal agronomic range (6.0 ? 7.0) for *Prunus dulcis*. At this pH:

* Nutrient availability for macronutrients (N, P, K) and secondary nutrients (Ca, Mg, S) is maximized.

* Micronutrient fixation (such as Zinc and Iron) is minimized compared to alkaline soils common in parts of the Central Valley.

#

N-P-K & Liebig's Law of the Minimum

Applying Liebig's Law of the Minimum, which dictates that crop yield is limited by the scarcest essential

AI AgriYield Predictor: Final Advisory Report

resource?reveals potential nutritional bottlenecks for a yield target of 2,350.50 kg/acre:

1. Nitrogen (N - 180 units): Almonds require roughly 30?35 kg of actual N per 1,000 kg of kernel produced, plus maintenance N for tree biomass. For a ~2,350 kg yield, total N demand approaches 220?250 kg/acre. Therefore, 180 units of N is the primary limiting nutrient.

* *Fertilizer Strategy:* Because Urea is the primary source, avoid single heavy applications to prevent ammonia volatilization and nitrate leaching in high-drainage loam.

2. Phosphorus (P - 50 units): Adequate for root dynamics and early flower bud differentiation. Almond P removal is relatively low compared to N and K; 50 units is sufficient.

3. Potassium (K - 150 units): Almond kernels and hulls are heavy K sinks (removing ~40?45 kg K₂O per 1,000 kg yield). To maintain yield and tree vigor, K availability must be closely matched during the kernel filling stage.

Climate Stress Mitigation

#

Water Demand vs. Rainfall

* Rainfall (600 mm / ~23.6 inches): While 600 mm is a favorable natural baseline for Mediterranean climates, mature high-yielding almond orchards in California demand between 900 mm and 1,200 mm (36?48 inches) of evapotranspiration (\$ET_c\$) annually.

* Mitigation: The system operates at a seasonal water deficit of 300?600 mm. Precision irrigation (micro-sprinklers or subsurface drip) scheduled against daily \$ET_c\$ and real-time soil moisture sensors is essential. Regulated Deficit Irrigation (RDI) should be applied strategically during hull split to suppress *Rhizopus* hull rot without reducing kernel weight.

#

Thermal Management

* Temperature (25°C Average): An average temperature of 25°C is highly conducive to photosynthesis and nut development. However, California summer heatwaves frequently exceed 35°C?40°C, causing stomatal closure and photo-inhibition.

* Mitigation: Maintain mid-row cover crops or residue management to reduce orchard floor radiant heat. In extreme thermal spikes, deploy short cooling irrigation sets or foliar kaolin clay sprays to buffer canopy temperatures.

Recommended Field Actions

AI AgriYield Predictor: Final Advisory Report

1. Implement Split-Fertigation for Nitrogen:

* Divide the total Nitrogen budget into 3-4 fertigation applications following the tree demand curve: 20% at leaf-out/fruit set, 30% during rapid nut expansion, 30% during kernel fill, and 20% post-harvest. Increase baseline N from 180 to ~220 units to support the high yield target.

2. Supplemental Potassium Application:

* Fertigate with water-soluble Potassium (e.g., Potassium Thiosulfate or Potassium Nitrate) during the kernel filling stage (May-July) to support the high crop load and prevent spur dieback.

3. Bridge the Irrigation Deficit via ET_c Scheduling:

* Develop a weekly irrigation schedule supplying the additional ~400-500 mm of required crop water using pressure bombs (mid-day stem water potential) to maintain tree water status between -9 to -12 bars during vegetative growth.

4. July Nutrient Tissue Testing:

* Collect critical leaf samples in July to verify that tissue Nitrogen levels remain within the optimal range (2.2%-2.5%) and Potassium remains above 1.4%, making immediate micro-adjustments if deficiencies appear.