

PATIENT: Akoto, Frank
DOB: 52 M
DOCTOR:
ACCN#: 240710005

DRAWN: 7/10/2024 15:15
RECEIVED: 7/10/2024 15:15
PRINTED: 7/10/2024 16:49 **FINAL**

PANEL/TEST	RESULT	REFERENCE RANGE	UNITS
------------	--------	-----------------	-------

BUE,CR

SODIUM	139	136-145	mmol/L
POTASSIUM	4.3	3.5-5.1	mmol/L
CHLORIDE	106	98-107	mmol/L
UREA	3.1	2.1-7.1	mmol/L
CREATININE	66 L	80-115	umol/L
eGFR	122.2		mL/min

Stage of CKD

EGFR RESULTS INTERPRETATION

STAGE 1: 90+ :Normal kidney function but urine findings or structural abnormalities or genetic trait may point to kidney disease

STAGE 2: 60 – 89 :Mildly reduced kidney function, and other findings (as stage 1) point to kidney disease

STAGE 3a: 45 – 59 :Mild to moderate reduced kidney function

STAGE 3b: 30 – 44 :Moderate to severe reduced kidney function

STAGE 4: 15 – 29 :Severely reduced kidney function

STAGE 5: <15 :Very severe, or end-stage kidney failure

LIPID STUDIES

CHOLESTEROL	4.6	3.5-5.2	mmol/L
HDL CHOLESTEROL	1.2	0.73-1.63	mmol/L
CHOL/HDL RATIO	3.8	<4.0	
LDL (Calc.)	3.18		mmol/L
Desirable [≤ 3.4]			
Borderline [3.4-4.1]			
High Risk [> 4.1]			
VLDL (Calc.)	0.22	0.05-0.77	mmol/L
TRIGLYCERIDES	0.49 L	0.71-3.54	mmol/L

PATIENT: Akoto, Frank
DOB: 52 M
DOCTOR:
ACCN#: 240710005

DRAWN: 7/10/2024 15:15
RECEIVED: 7/10/2024 15:15
PRINTED: 7/10/2024 16:49 **FINAL**

PANEL/TEST	RESULT	REFERENCE RANGE	UNITS
LIVER FUNCTION TEST			
TOTAL BILIRUBIN	13.0	1.7-21.0	umol/L
DIRECT BILIRUBIN	2.1	0.0-5.0	mg/dL
INDIRECT BIL	10.9	0.0-19.0	mg/dL
AST (SGOT)	18.5	0.0-40.0	IU/L
ALT (SGPT)	12.2	0.0-41.0	IU/L
ALP	39 L	40-130	IU/L
GGT	23.9	<55	IU/L
GLOBULIN	35	20-45	g/dL
TOTAL PROTEIN	72	66-88	g/L
ALBUMIN	37.5	35-52	g/L
A/G RATIO	1.1	1.1-2.5	Ratio

TUMOR MARKERS			
PSA, TOTAL	2.7	0.0-4.0	ng/mL

Comment:

Normal PSA Levels: 0 - 4.0 ng/mL
Slightly Elevated PSA: 4.1 - 10.0 ng/mL
Moderately Elevated PSA: 10.1 - 20.0 ng/mL
Highly Elevated PSA: 20+ ng/mL

PSA test can detect high levels of PSA that may indicate the presence of prostate cancer. However, many other conditions, such as an enlarged or inflamed prostate, also can increase PSA levels. Results must be interpreted in the context of the patient's clinical presentation and other diagnostic tests.

Approved by: Dr. Kwadwo Mfodwo