



ΕΘΝΙΚΟ  
ΜΕΤΣΟΒΙΟ  
ΠΟΛΥΤΕΧΝΕΙΟ

*National Technical University of Athens*

*School of Applied Mathematical and Physical Sciences*

---

# Study of a non-Autonomous Hamiltonian with Canonical Perturbation Theory

---

Θωμόπουλος Σπυρίδων, ge19042

*Complex Dynamics of Hamiltonian Systems & Applications*

March 2, 2023

---

## **Abstract**

This project was done as part of the class "Complex Dynamics of Hamiltonian Systems & Applications" and it is about the study of a nearly non-integrable Hamiltonian using Canonical Perturbation Theory.

# Contents

|          |  |          |
|----------|--|----------|
| <b>1</b> | <b>Theoretical Introduction</b>              | <b>4</b> |
|          | Canonical Transformation . . . . .           | 5        |
|          | Time Dependent Perturbation Theory . . . . . | 6        |
|          | Final Remarks . . . . .                      | 8        |
| <b>2</b> | <b>Non-autonomous Hamiltonian</b>            | <b>9</b> |

# 1 Theoretical Introduction

Canonical Perturbation Theory (CPT) is the theory which helps us understand how perturbations affect the dynamics of a system which is described from an integrable Hamiltonian. So, if we have a certain Hamiltonian which, after a Canonical Transformation, can be broken into a fully integrable part and a non-integrable part, we can apply CPT and predict how the latter changes the dynamics of the initial system. **MORE ...**

Let's assume that we have an autonomous Hamiltonian  $H'_0(\mathbf{q}, \mathbf{p})$  which fully describes a system via the Hamilton's equations of motion

$$\dot{q}_i = \frac{\partial H'_0(\mathbf{q}, \mathbf{p})}{\partial p_i} \quad (1.1a)$$

$$\dot{p}_i = - \frac{\partial H'_0(\mathbf{q}, \mathbf{p})}{\partial q_i}, i = 1, \dots, n \quad (1.1b)$$

If we can apply a Canonical Transformation on it which maintains the Hamiltonian character of the system, then we call it a Canonical Transformation. Let's further assume that there are as many constants of motion as degrees of freedom,  $2n$ . This means that exists a Canonical Transformation,  $(\mathbf{q}, \mathbf{p}) \rightarrow (\mathbf{J}, \boldsymbol{\theta})$ , which makes the new generalized position ignorable. That is, the transformed Hamiltonian is not dependent on it, thus  $H_0 = H_0(\mathbf{J})$ . These new Canonical Variables are called *Action-Angle Variables*. The system can now be written as

$$\dot{\theta}_i = \frac{\partial H_0(\mathbf{J})}{\partial J_i} \quad (1.2a)$$

$$\dot{J}_i = - \frac{\partial H_0(\mathbf{J})}{\partial \theta_i}, i = 1, \dots, n \quad (1.2b)$$

From the second equation we can easily obtain that the *Action* is constant for all degrees of freedom,  $J_i = \alpha_i$ . **From the first, we can deduce the the derivative of the Angle is constant**, thus the solution of the system is

$$\theta_i = \omega_{0,i} \cdot t + \theta_0 \quad (1.3a)$$

$$J_i = \alpha_i \quad (1.3b)$$

where  $\omega_{0,i} = \partial(H_0(\mathbf{J})/\partial J_i$  Now, we can see that the new variable  $\boldsymbol{\theta}$  is named *Angle* on purpose since, from equation (1a), it can be interpreted as a periodic phase with different frequency for each degree of freedom  $\boldsymbol{\omega} = (\omega_{0,1}, \dots, \omega_{0,n})$ . If we transform back to the initial variables  $(\mathbf{J}, \boldsymbol{\theta}) \rightarrow (\mathbf{q}, \mathbf{p})$  we have the solution of the initial system.

However convenient it seems, the above case is, firstly extremely rare and secondly extremely sensitive. Say for example that the Hamiltonian models a real mechanical system. That system is naturally prone to imperfections which will destroy its integrable character and as a result we will not be able to apply the above method of Canonical Transformation to Action-Angle variables in order to find the solution. Or we may want to increase our system's energy, so we have to impose an external periodic stimulation. All these doesn't mean that we are unable to study the new *perturbed* system.

The new system is now described by a Hamiltonian  $H'(\mathbf{q}, \mathbf{p}, t)$ . Due to the fact that it deviates only a little from the initial one, it can be written as

$$H'(\mathbf{q}, \mathbf{p}, t) = H'_0(\mathbf{q}, \mathbf{p}) + \epsilon H'_1(\mathbf{q}, \mathbf{p}, t) \quad (1.4)$$

Where,  $H'_0$  is our integrable Hamiltonian and  $H'_1$  is the non-integrable perturbation. So, if we apply the same Canonical Transformation as before, the new Hamiltonian,  $H$ , can be written as

$$H(\mathbf{J}, \boldsymbol{\theta}, t) = H_0(\mathbf{J}) + \epsilon H_1(\mathbf{J}, \boldsymbol{\theta}, t) \quad (1.5)$$

Our goal now is to transform the new Hamiltonian in order to push the  $\boldsymbol{\theta}, t$  dependence to higher order  $\epsilon^2$ . In that way, we will have an approximately integrable Hamiltonian to the first order and we will be able to solve it.

## Canonical Transformation

The aforementioned transformation,  $(\mathbf{J}, \boldsymbol{\theta}) \rightarrow (\bar{\mathbf{J}}, \bar{\boldsymbol{\theta}})$  should as well maintain the Hamiltonian structure of our system, so that the new variables  $(\bar{\mathbf{J}}, \bar{\boldsymbol{\theta}})$  are Canonical Variables.

The new Hamiltonian is  $\bar{H} = \bar{H}(\bar{\mathbf{J}}, \bar{\boldsymbol{\theta}})$ . Both the new and the old Hamiltonians should obey the Hamilton's variational principle

$$\left\{ \begin{array}{l} \delta \int_{t_1}^{t_2} \left( J_i \dot{\theta}_i - H(\mathbf{J}, \boldsymbol{\theta}, t) \right) dt = 0 \\ \delta \int_{t_1}^{t_2} \left( \bar{J}_i \dot{\bar{\theta}}_i - \bar{H}(\bar{\mathbf{J}}, \bar{\boldsymbol{\theta}}, t) \right) dt = 0 \end{array} \right\} \Rightarrow$$

$$J_i \dot{\theta}_i - H(\mathbf{J}, \boldsymbol{\theta}, t) = \bar{J}_i \dot{\bar{\theta}}_i - \bar{H}(\bar{\mathbf{J}}, \bar{\boldsymbol{\theta}}, t) + \frac{dF(\mathbf{J}, \boldsymbol{\theta}, \bar{\mathbf{J}}, \bar{\boldsymbol{\theta}}, t)}{dt} \quad (1.6)$$

The F function is called as the *Generating Function* of the transformation. That means that it uniquely defines the transformation. Say that we want it to be dependent only on the old Angles and the new Actions,  $F(\boldsymbol{\theta}, \bar{\mathbf{J}}, t)$ . Our goal is to solve (1.6) for the other two variables,  $\bar{\boldsymbol{\theta}}, \mathbf{J}$ . We have

$$J_i \dot{\theta}_i - H(\mathbf{J}, \boldsymbol{\theta}, t) = \bar{J}_i \dot{\bar{\theta}}_i - \bar{H}(\bar{\mathbf{J}}, \bar{\boldsymbol{\theta}}, t) + \dot{\theta}_i \frac{\partial F}{\partial \theta_i} + \dot{\bar{J}}_i \frac{\partial F}{\partial \bar{J}_i} + \frac{\partial F}{\partial t}$$

We see that there is no way to get rid of the unwanted term (1st on RHS). For that reason, we have to add one arbitrary term to the generating function which will cancel with the  $J_i \dot{\theta}_i$ . Our generating function will be

$$F(\mathbf{J}, \boldsymbol{\theta}, \bar{\mathbf{J}}, \bar{\boldsymbol{\theta}}, t) = -\bar{J}_i \bar{\theta}_i + S(\boldsymbol{\theta}, \bar{\mathbf{J}}, t) \quad (1.7)$$

Now, equation (1.6) can be written as

$$J_i \dot{\theta}_i - H(\mathbf{J}, \boldsymbol{\theta}, t) = \cancel{\bar{J}_i \dot{\bar{\theta}}_i} - \bar{H}(\bar{\mathbf{J}}, \bar{\boldsymbol{\theta}}, t) + \dot{\theta}_i \frac{\partial S}{\partial \theta_i} + \dot{\bar{J}}_i \frac{\partial S}{\partial \bar{J}_i} + \frac{\partial S}{\partial t} - \dot{\bar{J}}_i \bar{\theta}_i - \cancel{\bar{J}_i \dot{\bar{\theta}}_i} \Rightarrow$$

$$\textcolor{red}{J_i \dot{\theta}_i} - H(\mathbf{J}, \boldsymbol{\theta}, t) = -\bar{H}(\bar{\mathbf{J}}, \bar{\boldsymbol{\theta}}, t) + \textcolor{red}{\dot{\theta}_i \frac{\partial S}{\partial \theta_i}} + \textcolor{green}{\dot{\bar{J}}_i \frac{\partial S}{\partial \bar{J}_i}} + \frac{\partial S}{\partial t} - \textcolor{green}{\dot{\bar{J}}_i \bar{\theta}_i} \Rightarrow$$
(1.8)

So we obtain

$$J_i = \partial S / \partial \theta_i \quad (1.9a)$$

$$\bar{\theta}_i = \partial S / \partial \bar{J}_i \quad (1.9b)$$

$$\bar{H} = H + \partial S / \partial t \quad (1.9c)$$

So now we know how each variable can be obtained by the generating function  $S = S(\boldsymbol{\theta}, \bar{\mathbf{J}}, t)$ . It is clear that S uniquely defines the transformation and if it is equal to  $S = \theta_i \bar{J}_i$ , it represents the *identity transformation*, since it leaves both  $\mathbf{J}$  and  $\boldsymbol{\theta}$  unchanged.

## Time Dependent Perturbation Theory

We want to study a system described by (1.5). An integrable system  $H_0$  plus an non-integrable one  $H_1$  which acts as a perturbation on the first. More specifically, what we want to do is to move the  $\bar{\theta}$  dependence of  $\bar{H}_1$ , to higher order utilizing a Canonical Transformation  $(\mathbf{J}, \boldsymbol{\theta}) \rightarrow (\bar{\mathbf{J}}, \bar{\boldsymbol{\theta}})$ .

Since the non-integrable part is a perturbation, we can deduce that the generating function of the wanted trasformation will be nearly identical, with a non-identical first order part:

$$S(\boldsymbol{\theta}, \bar{\mathbf{J}}, t) = \bar{J}_i \theta_i + \epsilon S_1(\boldsymbol{\theta}, \bar{\mathbf{J}}, t) \quad (1.10)$$

From (1.9) we have

$$J_i = \bar{J}_i + \epsilon \frac{\partial S_1}{\partial \theta_i} \quad (1.11a)$$

$$\bar{\theta}_i = \theta_i + \epsilon \frac{\partial S_1}{\partial \bar{J}_i} \quad (1.11b)$$

$$\bar{H}(\bar{\mathbf{J}}, \bar{\boldsymbol{\theta}}, t) = H(\mathbf{J}, \boldsymbol{\theta}, t) + \epsilon \frac{\partial S_1}{\partial t} \quad (1.11c)$$

So we see that for the zero-th order approximation, the new and the old variables are the same. As for the Hamiltonian, we substitute (1.11a), (1.11b) into (1.11c) and we have

$$\begin{aligned} \bar{H}(\bar{\mathbf{J}}, \bar{\boldsymbol{\theta}}, t) &= H(\mathbf{J}(\bar{\mathbf{J}}, \bar{\boldsymbol{\theta}}, t), \boldsymbol{\theta}(\bar{\mathbf{J}}, \bar{\boldsymbol{\theta}}, t), t) + \epsilon \frac{\partial S_1}{\partial t} \xrightarrow{(1.11a,b)} \\ &= H_0(\mathbf{J}(\bar{\mathbf{J}}, \bar{\boldsymbol{\theta}}, t)) + \epsilon H_1(\mathbf{J}(\bar{\mathbf{J}}, \bar{\boldsymbol{\theta}}, t), \boldsymbol{\theta}(\bar{\mathbf{J}}, \bar{\boldsymbol{\theta}}, t), t) + \epsilon \frac{\partial S_1}{\partial t} \xrightarrow{(1.11a)} \\ &= H_0\left(\bar{\mathbf{J}} + \epsilon \frac{\partial S_1}{\partial \boldsymbol{\theta}}\right) + \epsilon H_1(\mathbf{J}(\bar{\mathbf{J}}, \bar{\boldsymbol{\theta}}, t), \boldsymbol{\theta}(\bar{\mathbf{J}}, \bar{\boldsymbol{\theta}}, t), t) + \epsilon \frac{\partial S_1}{\partial t} \xrightarrow{\text{Taylor, up to } (\epsilon^1)} \\ &= H_0(\bar{\mathbf{J}}) + \epsilon \frac{\partial S_1}{\partial \boldsymbol{\theta}} \frac{\partial H_0(\bar{\mathbf{J}})}{\partial \bar{\mathbf{J}}} + \epsilon H_1(\bar{\mathbf{J}}, \bar{\boldsymbol{\theta}}, t) + \epsilon \frac{\partial S_1}{\partial t} \end{aligned}$$

So, if we separate the above equation into  $0^{th}$  and  $1^{st}$  orders, we have

$$(\epsilon^0) : \quad \bar{H}_0 = \bar{H}_0(\bar{\mathbf{J}}) = H_0(\bar{\mathbf{J}}) \quad (1.12a)$$

$$\begin{aligned} (\epsilon^1) : \quad \bar{H}_1 &= \bar{H}_1(\bar{\mathbf{J}}, \bar{\boldsymbol{\theta}}, t) \Rightarrow \\ &= H_1(\bar{\mathbf{J}}, \bar{\boldsymbol{\theta}}, t) + \frac{\partial S_1}{\partial \boldsymbol{\theta}} \cdot \boldsymbol{\omega}_0(\bar{\mathbf{J}}) + \frac{\partial S_1}{\partial t} \end{aligned} \quad (1.12b)$$

Where

$$\boldsymbol{\omega}_0(\bar{\mathbf{J}}) = \frac{\partial H_0(\bar{\mathbf{J}})}{\partial \bar{\mathbf{J}}} \quad (1.13)$$

Now we have to find which  $S_1$  suits us to continue our study, since it defines uniquely the transformation. First of all, if we identify  $\boldsymbol{\theta}$  as a phase, then  $H_1$  has to be a periodic function with respect to  $\boldsymbol{\theta}$  with period  $T_{\theta_i} = 2\pi$ . We also want time periodicity, with period  $T_t = 2\pi/\Omega$ . Since both  $H_1$  and  $\bar{H}_1$  are periodic with respect to both *Angle* and *time*, from equation (1.12b) we have that  $S_1$  should also be a periodic function of *Angle* and *time*

Until now, we have two solid points from which we can continue. Firstly, the periodicity of  $H_1$  and  $S_1$  and secondly, the fact that we know how to solve integrable systems (with *Angle and time* independent Hamiltonian). The above points lead respectively to the Fourier expansion of  $H_1$  and  $S_1$  and the need to make  $\bar{H}_1$  *approximately integrable* (if we could make it exactly integrable we wouldn't need CPT :) ). The latter requires a subtle, yet extremely useful trick which will lead to the choice of  $S_1$  by connecting it with  $H_1$ .

To be more specific, the trick is as simple as separating  $H_1$  into two parts; a mean value with respect to both *Angle and time* plus an oscillating part

$$H_1 = \langle H_1 \rangle_{\bar{\theta}, t} + \{H_1\}_{\bar{\theta}, t} \quad (1.14)$$

Now, if we substitute into (1.12b) we have

$$\begin{aligned} \bar{H}_1 &= H_1(\bar{\mathbf{J}}, \bar{\boldsymbol{\theta}}, t) + \frac{\partial S_1}{\partial \boldsymbol{\theta}} \cdot \boldsymbol{\omega}_0(\bar{\mathbf{J}}) + \frac{\partial S_1}{\partial t} \Rightarrow \\ &= \langle H_1 \rangle_{\bar{\theta}, t}(\bar{\mathbf{J}}) + \{H_1\}_{\bar{\theta}, t}(\bar{\mathbf{J}}, \bar{\boldsymbol{\theta}}, t) + \frac{\partial S_1}{\partial \boldsymbol{\theta}} \cdot \boldsymbol{\omega}_0(\bar{\mathbf{J}}) + \frac{\partial S_1}{\partial t} \end{aligned} \quad (1.15)$$

Here is the crucial point where we will "push" the  $\boldsymbol{\theta} - t$  dependence to higher orders and at the same time we define the Canonical Transformation by defining its gerating function  $S_1$ . Since we can arbitrarily choose  $S_1$ , we choose it in order to satisfy the following relation

$$\{H_1\}_{\bar{\theta}, t}(\bar{\mathbf{J}}, \bar{\boldsymbol{\theta}}, t) + \frac{\partial S_1}{\partial \boldsymbol{\theta}} \cdot \boldsymbol{\omega}_0(\bar{\mathbf{J}}) + \frac{\partial S_1}{\partial t} = 0 \quad (1.16)$$

We finally have that

$$\begin{aligned} \bar{H} &= \bar{H}_0 + \epsilon \bar{H}_1 \Rightarrow \\ &= H_0(\bar{\mathbf{J}}) + \epsilon \langle H_1 \rangle_{\bar{\theta}, t}(\bar{\mathbf{J}}) \end{aligned} \quad (1.17)$$

which is dependent (approximately) only on  $\bar{\mathbf{J}}$  and is an integrable (approximately) Hamiltonian.

There is only one step left to complete the theory. That is to determine  $S_1$  by writing it as a Fourier sum and obtaining the coefficients from (1.16). The Fourier series of  $\{H_1\}_{\bar{\theta}, t}$ <sup>1</sup> and  $S_1$

$$\{H_1\}_{\bar{\theta}, t}(\bar{\mathbf{J}}) = \sum_{\mathbf{l}, m \neq 0}^{\infty} H_{1, \mathbf{l}, m} \cdot e^{i(\mathbf{l} \cdot \bar{\boldsymbol{\theta}} + m \Omega t)} \quad (1.18a)$$

$$S_1 = \sum_{\mathbf{l}, m \neq 0}^{\infty} S_{1, \mathbf{l}, m} \cdot e^{i(\mathbf{l} \cdot \bar{\boldsymbol{\theta}} + m \Omega t)} \quad (1.18b)$$

We substitute (1.18) into (1.16) and we have

$$\begin{aligned} \sum_{\mathbf{l}, m \neq 0}^{\infty} \left( H_{1, \mathbf{l}, m} \cdot e^{i(\mathbf{l} \cdot \bar{\boldsymbol{\theta}} + m \Omega t)} + \boldsymbol{\omega}_0(\bar{\mathbf{J}}) \cdot i\mathbf{l} \cdot S_{1, \mathbf{l}, m} \cdot e^{i(\mathbf{l} \cdot \bar{\boldsymbol{\theta}} + m \Omega t)} + im\Omega S_{1, \mathbf{l}, m} \cdot e^{i(\mathbf{l} \cdot \bar{\boldsymbol{\theta}} + m \Omega t)} \right) &= 0 \Rightarrow \\ \sum_{\mathbf{l}, m \neq 0}^{\infty} \left( H_{1, \mathbf{l}, m} + i \left( \boldsymbol{\omega}_0(\bar{\mathbf{J}}) \cdot \mathbf{l} + m\Omega \right) \cdot S_{1, \mathbf{l}, m} \right) e^{i(\mathbf{l} \cdot \bar{\boldsymbol{\theta}} + m \Omega t)} &= 0 \xrightarrow{\forall \mathbf{l}, m \neq 0} \\ S_{1, \mathbf{l}, m} &= i \frac{H_{1, \mathbf{l}, m}}{\boldsymbol{\omega}_0(\bar{\mathbf{J}}) \cdot \mathbf{l} + m\Omega} \end{aligned} \quad (1.19)$$

where  $H_1$  may act as a perturbation whether  $V_1, V_2 \ll 1$ .

Now, the transformation which "pushes" the *Angle* dependence of  $\bar{H}_1$  is fully defined by equations (1.18b) and (1.19) and our job is just to find the Fourier coefficients of  $H_1$ .

$$S_1 = \sum_{\mathbf{l}, m \neq 0}^{\infty} i \frac{H_{1, \mathbf{l}, m}(\bar{\mathbf{J}})}{\boldsymbol{\omega}_0(\bar{\mathbf{J}}) \cdot \mathbf{l} + m\Omega} \cdot e^{i(\mathbf{l} \cdot \bar{\boldsymbol{\theta}} + m \Omega t)} \quad (1.20)$$

---

<sup>1</sup>The function  $\{H_1\}_{\bar{\theta}, t}$  has no constant part as we have taken out of it as a mean value at (1.14). This means that we will not start the Fourier sum from 0.

## Final Remarks

From the above equation we can see that there may exist certain values of  $\mathbf{l}$  and  $m$ , for which the denominator becomes zero

$$\omega_0(\bar{\mathbf{J}}) \cdot \mathbf{l} + m\Omega = 0 \quad (1.21)$$

This means that the perturbation is no longer a perturbation since it has huge values compared to the integrable part and we have a huge problem.

If we had only the integrable part, then the dynamics of the systems evolve on a  $n$ -dimensional torus with the constant *Action* being the radius and the linearly increasing *Angle* being the phase. There are two cases, which depend on the frequency (time derivative of the *Angle*) of each degree of freedom. If the ratio of each two frequencies is a **rational** number,  $\omega_i/\omega_j \in \mathbb{Q}, \forall i, j$ , then the motion is periodic and the trajectory does not fill the whole "surface" of the torus. On the other hand, if the ratio is **irrational** even for one pair of frequencies,  $\exists i, j : \omega_i/\omega_j \in \mathbb{R}/\mathbb{Q}$ , the motion is not periodic and the trajectory densely fills the surface of the torus.

This behaviour is easily shown with Poincare Maps. To make a Poincare Map we firstly construct a surface on the phase space in which we choose some degrees of freedom to be constant (in our case some *Angles*) and we as we let the system evolve, we see where the trajectory intersects our surface. If we are in the first case of a periodic system, we will see points which will repeat after one period, whereas if we are the second case of irrational ratios, we will see dense regions of points.




Figure 1.1: Poincare Diagram of a 2D periodic system

These were about the integrable system. If we turn on the non-integrable part,



## 2 Non-autonomous Hamiltonian

We will use the theory developed in the above text for systems with  $n$  degrees of freedom, in order to study the following Hamiltonian with one degree of freedom

$$H(J, \theta, t) = \bar{\omega}J + J^k + V_1 \cos(n_1\theta - m_1\Omega t) + V_2 \cos(n_2\theta - m_2\Omega t) \quad (2.1)$$

It is obvious that it can be written as the sum of an integrable and a non-integrable Hamiltonian

$$H_0 = \bar{\omega}J + J^k \quad (2.2a)$$

$$H_1 = V_1 \cos(n_1\theta - m_1\Omega t) + V_2 \cos(n_2\theta - m_2\Omega t) \quad (2.2b)$$

This means that if we "turn off" the non-integrable part by setting  $V_1 = V_2 = 0$ , we can easily find the frequency from the Hamilton's equation of motion for the generalized position

$$\begin{aligned} \omega_0(J) = \dot{\theta} &= \frac{\partial H_0}{\partial J} \Rightarrow \\ &= \bar{\omega} + kJ^{k-1} \end{aligned} \quad (2.3)$$

The non integrable part can be written as a sum of mean value and an oscillating part

$$H_1 = \langle H_1 \rangle_{\theta, t} + \{H_1\}_{\theta, t}$$

But since it is a periodic function with respect to  $\theta, t$ , we have

$$\langle H_1 \rangle_{\theta, t} = 0$$

and since it is already a sum of cosine functions we will not use the Fourier expansion (1.18a) but the following

$$\{H_1\}_{\theta, t} = H_1 = \sum_{i=1,2} H_{1,i} \cos(n_i\theta - m_i\Omega t) \quad (2.4)$$

where

$$H_{1,i} = \begin{cases} V_i & , i = 1, 2 \\ 0 & , \text{otherwise} \end{cases} \quad (2.5)$$

Similarly, we write  $S_1$  as the following sum

$$S_1 = \sum_i S_{1,i} \sin(n_i\theta - m_i\Omega t) \quad (2.6)$$

If we substitute (2.4) and (2.6) into the condition (1.16), we have

$$\begin{aligned} H_1 + \frac{\partial S_1}{\partial \theta} \omega_0(\bar{J}) + \frac{\partial S_1}{\partial t} &= 0 \Rightarrow \\ \sum_i (V_i \cos(n_i\theta - m_i\Omega t) - m_i\Omega S_{1,i} \cos(n_i\theta - m_i\Omega t) - n_i S_{1,i} \cos(n_i\theta - m_i\Omega t) \cdot \omega_0) &= 0 \Rightarrow \\ \sum_i (V_i - m_i\Omega S_{1,i} - n_i S_{1,i} \omega_0) \cos(n_i\theta - m_i\Omega t) &= 0 \Rightarrow \end{aligned}$$

We want it to hold  $\forall i = \{1, 2\}$

$$S_{1,i} = \frac{V_i}{m_i\Omega + n_i(\bar{\omega} + kJ^{k-1})}, i = 1, 2$$

So, for each term of the expansion, we have resonance for the values for which the denominator is zero

$$\begin{aligned} m_i\Omega + n_i(\bar{\omega} + kJ^{k-1}) &= 0 \Rightarrow \\ J_{res,i}^{k-1} &= -\frac{m_i\Omega + n_i\bar{\omega}}{kn_i} \end{aligned} \quad (2.7)$$

---

### QUESTION 3

---

Also, we can compute the approximate (up to first order) constant of motion  $\bar{J}$  from (1.11a)

$$\begin{aligned} \bar{J} &= J - \frac{\partial S_1}{\partial \theta} \Rightarrow \\ &= J - \sum_{i=1,2} S_{1,i} n_i \cos(n_i \theta - m_i \Omega t) \Rightarrow \\ &= J - \frac{V_1 n_1}{m_1 \Omega + n_1(\bar{\omega} + kJ^{k-1})} \cos(n_1 \theta - m_1 \Omega t) - \frac{V_2 n_2}{m_2 \Omega + n_2(\bar{\omega} + kJ^{k-1})} \cos(n_2 \theta - m_2 \Omega t) \end{aligned} \quad (2.8)$$


---

### QUESTION 2

---

If we take either  $V_1 = 0$  or  $V_2 = 0$ , then we can change the position variable with the transformation

$$\hat{\theta} = n_i \theta - m_i \Omega t, \quad i = 1 \text{ or } 2 \quad (2.9)$$

But if we want to use a Canonical Transformation, we can also choose

$$\hat{J} = \alpha J \quad (2.10)$$

where  $\alpha$  is to be determined. In order to demand a Canonical Transformation and by doing so, to find  $\alpha$ , we can use the *symplectic condition*  $\tilde{\mathbf{M}} \mathbf{J} \mathbf{M} = \mathbf{J}$ , where  $\mathbf{M}$  is the Jacobian of the transformation, or the *Poisson Bracket Condition*,  $[\hat{\theta}, \hat{J}] = 1$ , or directly find a generating function.

We will use the *Poisson Bracket Condition* in order to determine  $\alpha$  and then we will find the generating function of the transformation. If we want the transformation to be Canonical, then the following condition must hold

$$\begin{aligned} [\hat{\theta}, \hat{J}] &= 1 \Rightarrow \\ \frac{\partial \hat{J}}{\partial J} \frac{\partial \hat{\theta}}{\partial \theta} - \frac{\partial \hat{J}}{\partial \theta} \frac{\partial \hat{\theta}}{\partial J} &= 1 \xrightarrow{(2.9), (2.10)} \\ \alpha n_i &= 1 \Rightarrow \\ \alpha &= 1/n_i, \quad i = 1 \text{ or } 2 \end{aligned} \quad (2.11)$$

In general, the generating function of the Canonical Transformation can be dependent on any pair of new and old variables. We choose it to be  $F_2 = F_2(\theta, \hat{J})$ . Now, the other two variables, can be expressed with  $F_2$ . For  $\hat{\theta}$

$$\begin{aligned}\hat{\theta} &= \frac{\partial F_2}{\partial \hat{J}} \xrightarrow{(2.9)} \\ n_i \theta - m_i \Omega t &= \frac{\partial F_2}{\partial \hat{J}} \xrightarrow{\theta\text{-indep.}} \\ F_2 &= (n_i \theta - m_i \Omega t) \hat{J} + f(\theta)\end{aligned}\tag{2.12}$$

And for  $J$

$$\begin{aligned}J &= \frac{\partial F_2}{\partial \theta} \xrightarrow{(2.12), (2.10)} \\ n_i \hat{J} &= n_i \hat{J} + f'(\theta) \Rightarrow \\ f(\theta) &= \text{const.}\end{aligned}$$

So, we can ignore  $f(\theta)$  since in our transformation it appears only as a derivative. Finally, the generating function is

$$F_2(\theta, \hat{J}) = (n_i \theta - m_i \Omega t) \hat{J}, \quad i=1 \text{ or } 2\tag{2.13}$$

The new Hamiltonian, according to (1.9c) is

$$\begin{aligned}\hat{H}(\hat{\theta}, \hat{J}) &= H(\hat{\theta}, \hat{J}) + \partial F_2 / \partial t \Rightarrow \\ &= (\bar{\omega} n_i - m_i \Omega) \hat{J} + n_i^k \hat{J}^k + V_i \cos \hat{\theta}\end{aligned}\tag{2.14}$$

And since it depends on  $\hat{\theta}$ , it is not integrable, but again, it can be seen as a sum of an integrable and a non-integrable perturbation. So, from CPT and (1.11a), the first order approximation of the new *Action*, is

$$\hat{\hat{J}} = \hat{J} - \epsilon \frac{\partial S_1}{\partial \hat{\theta}}$$

Where  $S_1$  is the same as the one we calculated (2.6) for the full Hamiltonian (both  $V_1, V_2 \neq 0$ ) without the one sine term, thus

$$\hat{\hat{J}} = \hat{J} - \epsilon \frac{V_i}{m_i \Omega - n_i(\bar{\omega} + k \hat{J}^{k-1})} \cos \hat{\theta}, \quad i=1 \text{ or } 2\tag{2.15}$$

The *resonance Action* can be obtained if we set the above denominator equal to zero

$$\begin{aligned}m_i \Omega - n_i(\bar{\omega} + k \hat{J}_{res}^{k-1}) &= 0 \Rightarrow \\ \hat{J}_{res}^{k-1} &= \frac{m_i \Omega - n_i \bar{\omega}}{n_i k}\end{aligned}\tag{2.16}$$

If we expand our Hamiltonian (2.14) around  $\hat{J}_{res}$ , with  $\Delta \hat{J} = \hat{J} - \hat{J}_{res}$ , we have

$$\begin{aligned}\hat{H} &\simeq \hat{H}(\hat{J}_{res}) + \frac{\partial \hat{H}}{\partial \hat{J}} \Big|_{\hat{J}_{res}} \Delta \hat{J} + \frac{\partial^2 \hat{H}}{\partial \hat{J}^2} \Big|_{\hat{J}_{res}} (\Delta \hat{J})^2 + O(\epsilon^2) \xrightarrow{\frac{\partial \theta}{\partial t}|_{res}=0=\frac{\partial \hat{H}}{\partial \hat{J}}|_{res}} \\ &= \left[ (\bar{\omega} n_i - m_i \Omega) \hat{J}_{res} + n_i^k \hat{J}_{res}^k + V_i \cos \hat{\theta} \right] + \frac{1}{2} k(k-1) \hat{J}_{res} (\Delta \hat{J})^2 \Rightarrow \\ &= V_i \cos \hat{\theta} + \frac{1}{2} k(k-1) \hat{J}_{res} (\Delta \hat{J})^2 + \underbrace{\left[ (\bar{\omega} n_i - m_i \Omega) \hat{J}_{res} + n_i^k \hat{J}_{res}^k \right]}_{\text{constant}}\end{aligned}\tag{2.17}$$

The dynamics of the system won't depend on the constant term of the Hamiltonian, since the equations of motion contain the derivative of the Hamiltonian. So, we can see that the Hamiltonian near the resonance regions is similar to the Hamiltonian of a non-linear pendulum  $H_{nlp}(\theta, J) = A \cos \theta + BJ^2$ .

## QUESTION 4

### Removal of Resonances

When we are near the resonance regions, among others, we have the problem that our new approximate constant of motion  $\bar{J}$  not only is not constant, but it goes to infinity ( we can see this from (2.8) ). So we want to construct a true approximate constant of motion which does not go to infinity. Any function of the "constant"  $\bar{J}$  is going to be a constant, so let's assume the our new constant is  $I(\bar{J})$ . Then we can use its Taylor expansion and (1.11a)

$$\begin{aligned} I(\bar{J}) &= I \left( J - \epsilon \frac{\partial S_1}{\partial \theta} \right) \Rightarrow \\ &= I(J) - \epsilon \frac{\partial S_1}{\partial \theta} \frac{dI}{dJ} \end{aligned} \quad (2.18)$$

We have not yet solved the problem. We see that the term  $\partial S_1 / \partial \theta$  that goes to infity still exists in our relation. But now, since  $I$  is an arbitrary function of  $\bar{J}$ , we can choose it in a way that it cancels that second term at resonances. That is

$$\left. \frac{dI}{dJ} \right|_{resonances} = 0 \quad (2.19)$$

Since the conditions for the resonances are given by (2.7), the condition for eliminating the unwanted term from  $I(\bar{J})$  is

$$\begin{aligned} \frac{dI}{dJ} &= \prod_{i=1,2} C_i \left( m_i \Omega + n_i (\bar{\omega} + k J^{k-1}) \right) \Rightarrow \\ &= C_1 C_2 \left( m_1 \Omega + n_1 (\bar{\omega} + k J^{k-1}) \right) \left( m_2 \Omega + n_2 (\bar{\omega} + k J^{k-1}) \right) \Rightarrow \\ &= C_1 C_2 \left( (m_1 \Omega + n_1 \bar{\omega})(m_2 \Omega + n_2 \bar{\omega}) + [n_1(m_2 \Omega + n_2 \bar{\omega}) + n_2(m_1 \Omega + n_1 \bar{\omega})] J^{k-1} + n_1 n_2 J^{2k-2} \right) \end{aligned} \quad (2.20)$$

So, we have that

$$I(J) = C_1 C_2 \left( (m_1 \Omega + n_1 \bar{\omega})(m_2 \Omega + n_2 \bar{\omega}) J + [n_1(m_2 \Omega + n_2 \bar{\omega}) + n_2(m_1 \Omega + n_1 \bar{\omega})] \frac{J^k}{k} + n_1 n_2 \frac{J^{2k-2}}{2k-1} \right) \quad (2.21)$$

If we combine (2.21) & (2.20) with (2.18) we obtain the new constant of motion which does not go to infinity at the resonances.

## QUESTION 5

We will now study numerically the Hamiltonian (2.1) and we will compare the numerical with the analytical results. In order to do that, we will construct its Poincare Map. In order to do that, we will have to solve numerically the following Hamilton's equations of motion

$$\begin{cases} \dot{\theta} = & \frac{\partial H}{\partial J} \\ \dot{J} = & -\frac{\partial H}{\partial \theta} \end{cases} \Rightarrow$$

$$\begin{cases} \dot{\theta} = & \bar{\omega} + kJ^{k-1} \\ \dot{J} = & V_1 n_1 \sin(n_1 \theta - m_1 \Omega t) + V_2 n_2 \sin(n_2 \theta - m_2 \Omega t) \end{cases}$$