




EFFECTS OF OA IN OLYMPIA OYSTER POPULATIONS

Parental pCO₂ Ambient High




| | <i>Fidalgo Bay</i> | <i>Port Gamble Bay</i> | <i>Skokomish River Delta</i> | <i>Case Inlet</i> |
|------------------|--------------------|------------------------|------------------------------|-------------------|
| Temperature (°C) | 15.4±1.5 | 15.0±1.0 | 16.2±2.7 | 16.8±1.7 |
| DO (mg/L) | 10.6±2.4 | 10.5±1.9 | 10.2±3.9 | 11.2±2.8 |
| Salinity (PSU) | 28.5±3.9 | 31.9±2.0 | 29.6±1.3 | 24.6±1.7 |
| pH | 8.07±0.15 | 7.86±0.17 | 8.01±0.20 | 8.01±0.16 |
| chlorophyll | 227±409 | 225±145 | 572±1536 | 331±613 |




- ENVIRONMENTAL MEMORY?

EFFECTS OF OA IN OLYMPIA OYSTER POPULATIONS

Parental pCO₂ Ambient High



| | <i>Fidalgo Bay</i> | <i>Port Gamble Bay</i> | <i>Skokomish River Delta</i> | <i>Case Inlet</i> |
|------------------|--------------------|------------------------|------------------------------|-------------------|
| Temperature (°C) | 15.4±1.5 | 15.0±1.0 | 16.2±2.7 | 16.8±1.7 |
| DO (mg/L) | 10.6±2.4 | 10.5±1.9 | 10.2±3.9 | 11.2±2.8 |
| Salinity (PSU) | 28.5±3.9 | 31.9±2.0 | 29.6±1.3 | 24.6±1.7 |
| pH | 8.07±0.15 | 7.86±0.17 | 8.01±0.20 | 8.01±0.16 |
| chlorophyll | 227±409 | 225±145 | 572±1536 | 331±613 |



- ENVIRONMENTAL MEMORY?
- SELECTION