

# Lecture 1

## Review of Simple Linear Regression

Reading: ISLR 2021 Chapter 3.1

DSA 8020 Statistical Methods II

Whitney Huang  
Clemson University

Review of Simple Linear Regression
CLEMSON UNIVERSITY
Simple Linear Regression
Parameter Estimation
Residual Analysis
Confidence/Prediction Intervals
Hypothesis Testing

1.1

Notes

---

---

---

---

---

---

---

### Agenda

- 1 Simple Linear Regression
- 2 Parameter Estimation
- 3 Residual Analysis
- 4 Confidence/Prediction Intervals
- 5 Hypothesis Testing

Review of Simple Linear Regression
CLEMSON UNIVERSITY
Simple Linear Regression
Parameter Estimation
Residual Analysis
Confidence/Prediction Intervals
Hypothesis Testing

1.2

Notes

---

---

---

---


---

---

---

### What is Regression Analysis?

**Regression analysis:** A set of statistical procedures for estimating the relationship between **response variable** and **predictor variable(s)**



**Simple linear regression:** The relationship between the response variable and the predictor variable is approximately linear

Review of Simple Linear Regression
CLEMSON UNIVERSITY
Simple Linear Regression
Parameter Estimation
Residual Analysis
Confidence/Prediction Intervals
Hypothesis Testing

1.3

Notes

---

---

---

---

---

---

---

## Simple Linear Regression (SLR)

$y$ : response variable;  $x$ : predictor variable

- In SLR we **assume** there is a **linear relationship** between  $x$  and  $y$ :

$$y = \beta_0 + \beta_1 x + \varepsilon$$

- We need to estimate  $\beta_0$  (**intercept**) and  $\beta_1$  (**slope**) based on observed data  $\{x_i, y_i\}_{i=1}^n$
- We can use the estimated regression equation to
  - make predictions
  - study the relationship between response and predictor
  - control the response
- Yet we need to quantify our **estimation uncertainty** regarding the linear relationship



Notes

---

---


---

---


---

---

**Regression equation:**  $y = \beta_0 + \beta_1 x$



- $\beta_0$ :  $E[y]$  when  $x = 0$
- $\beta_1$ :  $E[\Delta y]$  when  $x$  increases by 1



Notes

---

---

---

---

---

---

## Assumptions about the Random Error $\varepsilon$


In order to estimate  $\beta_0$  and  $\beta_1$ , we make the following assumptions about  $\varepsilon$

- $E[\varepsilon_i] = 0$
- $\text{Var}[\varepsilon_i] = \sigma^2$
- $\text{Cov}[\varepsilon_i, \varepsilon_j] = 0, \quad i \neq j$

Therefore, we have

$$E[y_i] = \beta_0 + \beta_1 x_i, \text{ and } \text{Var}[y_i] = \sigma^2$$

The regression line  $\beta_0 + \beta_1 x$  represents the **conditional mean curve** whereas  $\sigma^2$  measures the magnitude of the **variation** around the regression curve



Notes

---

---

---

---

---

---

Estimation: Method of Least Squares

For given observations  $\{x_i, y_i\}_{i=1}^n$ , choose  $\beta_0$  and  $\beta_1$  to minimize the *sum of squared errors*:

$$\ell(\beta_0, \beta_1) = \sum_{i=1}^n (y_i - (\beta_0 + \beta_1 x_i))^2$$

Solving the above minimization problem requires some knowledge from Calculus (see notes LS\_SLR.pdf)

$$\hat{\beta}_0 = \bar{y} - \hat{\beta}_1 \bar{x} \tag{1}$$

$$\hat{\beta}_1 = \frac{\sum_{i=1}^n (x_i - \bar{x})(y_i - \bar{y})}{\sum_{i=1}^n (x_i - \bar{x})^2} \tag{2}$$

We also need to **estimate**  $\sigma^2$

$$\hat{\sigma}^2 = \frac{\sum_{i=1}^n (y_i - \hat{y}_i)^2}{n - 2}, \tag{3}$$

where

$$\hat{y}_i = \hat{\beta}_0 + \hat{\beta}_1 x_i \tag{4}$$

Review of Simple Linear Regression

CLEMSON UNIVERSITY

Simple Linear Regression

Parameter Estimation

Residual Analysis

Confidence/Prediction Intervals

Hypothesis Testing

1.7

Notes

---

---

---

---

---

---

---

---

Example: Maximum Heart Rate vs. Age

The maximum heart rate `MaxHeartRate` of a person is often said to be related to age `Age` by the equation:

$$\text{MaxHeartRate} = 220 - \text{Age}.$$

Suppose we have 15 people of varying ages are tested for their maximum heart rate (bpm)

- 1 Compute the estimates for the regression coefficients,  $\hat{\beta}_0$  and  $\hat{\beta}_1$ , using Equations (1) and (2)
- 2 Compute the fitted values  $\{\hat{y}_i\}_{i=1}^n$  using Equation (4)
- 3 Compute the estimate for  $\sigma$  by applying the square root of Equation (3)

Review of Simple Linear Regression

CLEMSON UNIVERSITY

Simple Linear Regression

Parameter Estimation

Residual Analysis

Confidence/Prediction Intervals

Hypothesis Testing

1.8

Notes

---

---

---

---

---

---

---

---

Maximum Heart Rate vs. Age

Output from  (R Studio)

```
> fit <- lm(MaxHeartRate ~ Age)
> summary(fit)

Call:
lm(formula = MaxHeartRate ~ Age)

Residuals:
    Min       1Q   Median       3Q      Max
-8.9258 -2.5383  0.3879  3.1867  6.6242

Coefficients:
            Estimate Std. Error t value Pr(>|t|)
(Intercept) 210.04846    2.86694   73.27  < 2e-16 ***
Age         -0.79773     0.06996  -11.40 3.85e-08 ***
---
Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

Residual standard error: 4.578 on 13 degrees of freedom
Multiple R-squared:  0.9091,    Adjusted R-squared:  0.9021
F-statistic: 130 on 1 and 13 DF,  p-value: 3.848e-08
```

Review of Simple Linear Regression

CLEMSON UNIVERSITY

Simple Linear Regression

Parameter Estimation

Residual Analysis

Confidence/Prediction Intervals

Hypothesis Testing

1.9

Notes

---

---

---

---


---

---

---

---

Assessing Linear Regression Fit



**Question:** Is linear relationship between max heart rate and age reasonable?  $\Rightarrow$  [Residual Analysis](#)

Review of Simple Linear Regression

CLEMSON UNIVERSITY

Simple Linear Regression

Parameter Estimation

**Residual Analysis**

Confidence/Prediction Intervals

Hypothesis Testing

1.10

Notes

---

---

---

---

---

---

---

Residuals

- The [residuals](#) are the differences between the observed and fitted values:

$$e_i = y_i - \hat{y}_i,$$

where  $\hat{y}_i = \hat{\beta}_0 + \hat{\beta}_1 x_i$

- Residuals are very useful in assessing the appropriateness of the assumptions on  $\varepsilon_i$ . Recall
  - $E[\varepsilon_i] = 0$
  - $\text{Var}[\varepsilon_i] = \sigma^2$
  - $\text{Cov}[\varepsilon_i, \varepsilon_j] = 0, \quad i \neq j$

Review of Simple Linear Regression

CLEMSON UNIVERSITY

Simple Linear Regression

Parameter Estimation

**Residual Analysis**

Confidence/Prediction Intervals

Hypothesis Testing

1.11

Notes

---

---

---


---

---

---

---

Residuals Against Predictor Plot



Review of Simple Linear Regression

CLEMSON UNIVERSITY

Simple Linear Regression

Parameter Estimation

**Residual Analysis**

Confidence/Prediction Intervals

Hypothesis Testing

1.12

Notes

---

---

---

---

---

---

---

Interpreting Residual Plots




Figure courtesy of Faraway's Linear Models with R (2014, p. 74).

Notes

---

---

---

---


---

---

---

---

Diagnostic Plots in R



Notes

---

---

---

---


---

---

---

---

How (Un)certain We Are?



Can we formally quantify our estimation uncertainty?  
⇒ We need additional (distributional) assumption on  $\epsilon$

Notes

---

---

---

---

---

---

---

---

Normal Error Regression Model

Recall

y\_i = \beta\_0 + \beta\_1 x\_i + \varepsilon\_i

- Further assume \varepsilon\_i \sim N(0, \sigma^2) \Rightarrow y\_i | x\_i \sim N(\beta\_0 + \beta\_1 x\_i, \sigma^2)
- With normality assumption, we can derive the **sampling distribution** of \hat{\beta}\_1 and \hat{\beta}\_0 \Rightarrow

\frac{\hat{\beta}\_1 - \beta\_1}{SE(\hat{\beta}\_1)} \sim t\_{n-2}, \quad \hat{SE}(\hat{\beta}\_1) = \frac{\hat{\sigma}}{\sqrt{\sum\_{i=1}^n (x\_i - \bar{x})^2}}

\frac{\hat{\beta}\_0 - \beta\_0}{SE(\hat{\beta}\_0)} \sim t\_{n-2}, \quad \hat{SE}(\hat{\beta}\_0) = \hat{\sigma} \sqrt{(\frac{1}{n} + \frac{\bar{x}^2}{\sum\_{i=1}^n (x\_i - \bar{x})^2})}

where t\_{n-2} denotes the Student's t distribution with n - 2 degrees of freedom

Review of Simple Linear Regression

CLEMSON UNIVERSITY

Simple Linear Regression

Parameter Estimation

Residual Analysis

Confidence/Prediction Intervals

Hypothesis Testing

1.16

Notes

---

---

---

---

---

---

---

Confidence Intervals for \beta\_0 and \beta\_1

- Recall \frac{\hat{\beta}\_1 - \beta\_1}{SE(\hat{\beta}\_1)} \sim t\_{n-2}, we use this fact to construct a **confidence interval (CI)** for \beta\_1:

[\hat{\beta}\_1 - t\_{\alpha/2, n-2} \hat{SE}(\hat{\beta}\_1), \hat{\beta}\_1 + t\_{\alpha/2, n-2} \hat{SE}(\hat{\beta}\_1)]

where \alpha is the **confidence level** and t\_{\alpha/2, n-2} denotes the 1 - \alpha/2 percentile of a student's t distribution with n - 2 degrees of freedom

- Similarly, we can construct a CI for \beta\_0:

[\hat{\beta}\_0 - t\_{\alpha/2, n-2} \hat{SE}(\hat{\beta}\_0), \hat{\beta}\_0 + t\_{\alpha/2, n-2} \hat{SE}(\hat{\beta}\_0)]

Review of Simple Linear Regression

CLEMSON UNIVERSITY

Simple Linear Regression

Parameter Estimation

Residual Analysis

Confidence/Prediction Intervals

Hypothesis Testing

1.17

Notes

---

---

---

---

---

---

---

Confidence Interval of E(y\_{new})

- We often interested in estimating the **mean** response for an unobserved predictor value, say, x\_{new}. Therefore we would like to construct CI for E[y\_{new}], the corresponding **mean response**
- We need sampling distribution of E(\widehat{y\_{new}}) to form CI:

\frac{E(\widehat{y\_{new}}) - E(y\_{new})}{SE(E(y\_{new}))} \sim t\_{n-2}, \quad \hat{SE}(E(\widehat{y\_{new}})) = \hat{\sigma} \sqrt{(\frac{1}{n} + \frac{(x\_{new} - \bar{x})^2}{\sum\_{i=1}^n (x\_i - \bar{x})^2})}

- CI:

[\hat{y}\_{new} - t\_{\alpha/2, n-2} \hat{SE}(E(\widehat{y\_{new}})), \hat{y}\_{new} + t\_{\alpha/2, n-2} \hat{SE}(E(\widehat{y\_{new}}))]

- **Quiz:** Use this formula to construct CI for \beta\_0

Review of Simple Linear Regression

CLEMSON UNIVERSITY

Simple Linear Regression

Parameter Estimation

Residual Analysis

Confidence/Prediction Intervals

Hypothesis Testing

1.18

Notes

---

---

---

---

---

---

---

Prediction Interval of  $y_{new}$

- Suppose we want to predict the response of a future observation  $y_{new}$  given  $x = x_{new}$
- We need to account for added variability as a new observation does not fall directly on the regression line (i.e.,  $y_{new} = E[y_{new}] + \varepsilon_{new}$ )
- Replace  $\hat{SE}(E(\widehat{y_{new}}))$  by  $\hat{SE}(\hat{y}_{new}) = \hat{\sigma} \sqrt{\left(1 + \frac{1}{n} + \frac{(x_{new} - \bar{x})^2}{\sum_{i=1}^n (x_i - \bar{x})^2}\right)}$  to construct CIs for  $Y_{new}$

Notes

---

---

---


---

---

---

---

Assessing Normality Assumption on  $\varepsilon$



The Q-Q plot is more effective in detecting subtle departures from normality, especially in the tails.

Notes

---

---

---

---

---

---

---

Maximum Heart Rate vs. Age Revisited

The maximum heart rate  $\text{MaxHeartRate}$  ( $\text{HR}_{max}$ ) of a person is often said to be related to age  $\text{Age}$  by the equation:

$$\text{HR}_{max} = 220 - \text{Age}.$$

Suppose we have 15 people of varying ages are tested for their maximum heart rate (bpm)

Age	18	23	25	35	65	54	34	56	72	19	23	42	18	39	37
HR <sub>max</sub>	202	186	187	180	156	169	174	172	153	199	193	174	198	183	178

- Construct the 95% CI for  $\beta_1$
- Compute the estimate for mean  $\text{MaxHeartRate}$  given  $\text{Age} = 40$  and construct the associated 90% CI
- Construct the prediction interval for a new observation given  $\text{Age} = 40$

Notes

---

---

---

---

---

---

---

Maximum Heart Rate vs. Age: Hypothesis Test for Slope

- 1  $H_0 : \beta_1 = 0$  vs.  $H_a : \beta_1 \neq 0$
- 2 Compute the **test statistic**:  
 $t^* = \frac{\hat{\beta}_1 - 0}{SE(\hat{\beta}_1)} = \frac{-0.7977}{0.06996} = -11.40$
- 3 Compute **p-value**:  $P(|t^*| \geq |t_{obs}|) = 3.85 \times 10^{-8}$
- 4 Compare to  $\alpha$  and draw conclusion:

Reject  $H_0$  at  $\alpha = .05$  level, evidence suggests a **negative linear relationship** between MaxHeartRate and Age

Review of Simple Linear Regression

CLEMSON UNIVERSITY

Simple Linear Regression

Parameter Estimation

Residual Analysis

Confidence/Prediction Intervals

Hypothesis Testing

1.22

Notes

---

---

---

---

---

---

---

Maximum Heart Rate vs. Age: Hypothesis Test for Intercept

- 1  $H_0 : \beta_0 = 0$  vs.  $H_a : \beta_0 \neq 0$
- 2 Compute the **test statistic**:  
 $t^* = \frac{\hat{\beta}_0 - 0}{SE(\hat{\beta}_0)} = \frac{210.0485}{2.86694} = 73.27$
- 3 Compute **p-value**:  $P(|t^*| \geq |t_{obs}|) \simeq 0$
- 4 Compare to  $\alpha$  and draw conclusion:

Reject  $H_0$  at  $\alpha = .05$  level, evidence suggests evidence suggests the intercept (the expected MaxHeartRate at age 0) is different from 0

Review of Simple Linear Regression

CLEMSON UNIVERSITY

Simple Linear Regression

Parameter Estimation

Residual Analysis

Confidence/Prediction Intervals

Hypothesis Testing

1.23

Notes

---

---

---

---

---

---

---

Summary

In this lecture, we reviewed

- **Simple Linear Regression**:  
 $y = \beta_0 + \beta_1 x + \varepsilon, \varepsilon \sim N(0, \sigma^2)$
- **Method of Least Squares** for parameter estimation

$$\hat{\beta} = \underset{\beta=(\beta_0, \beta_1)}{\operatorname{argmin}} \sum_{i=1}^n (y_i - (\beta_0 + \beta_1 x_i))^2$$

- **Residual analysis** to check model assumptions
- **Confidence/Prediction Intervals** and **Hypothesis Testing**

Review of Simple Linear Regression

CLEMSON UNIVERSITY

Simple Linear Regression

Parameter Estimation

Residual Analysis

Confidence/Prediction Intervals

Hypothesis Testing

1.24

Notes

---

---

---

---

---

---

---



R Funcations

Fitting linear models

```
object <- lm(formula, data) where the formula
is specified via y ~ x => y is modeled as a linear func-
tion of x
```

```
summary(object);plot(object)
```

```
predict(object, newdata, interval)
```

```
confint(object)
```

Summary of Fits and Diagnostic Plots

Making Predictions and Their Intervals

Confidence Intervals for Model Parameters

Review of Simple  
Linear  
Regression



Simple Linear  
Regression  
Parameter  
Estimation

Residual Analysis  
Confidence/Prediction  
Intervals

Hypothesis Testing

Notes

---

---

---

---

---

---

---

---

Notes

---

---

---

---

---

---

---

---

Notes

---

---

---

---

---

---

---

---