

Scattering from compact objects, Regge poles and the Complex Angular Momentum method

Mohamed Ould El Hadj,^{1,2,*} Tom Stratton,^{2,†} and Sam R. Dolan^{2,‡}

¹*Equipe Physique Théorique, SPE, UMR 6134 du CNRS et de l'Université de Corse,
Université de Corse, Faculté des Sciences, BP 52, F-20250 Corte, France*

²*Consortium for Fundamental Physics, School of Mathematics and Statistics,
University of Sheffield, Hicks Building,
Hounsfield Road, Sheffield S3 7RH, United Kingdom*

(Dated: November 11, 2019)

To be written

CONTENTS

I. Introduction	1
II. The Regge pole spectrum of a compact body	2
A. The model	2
B. Effective potentials	2
C. Boundary conditions	3
D. Partial wave expansions of differential scattering cross sections	3
E. Scattering theory	3
F. CAM representation of the scattering amplitude	4
III. Reconstruction of differential scattering cross sections from Regge pole sums	4
A. Computational methods	5
B. Results and comments	5
IV. WKB ?	5
V. Classification of quasinormal modes	10
Acknowledgments	10
References	11

I. INTRODUCTION

The time-independent scattering of planar waves in the gravitational field of a compact body has been studied in some detail since the 1960s [1–3]. A substantial literature has developed on *black hole scattering*, focussing on the

canonical scenario of a planar wave of frequency ω and spin s [4] impinging upon a black hole of mass M in vacuum [1–3, 5–19]. A dimensionless parameter,

$$M\omega = \pi \frac{r_g}{\lambda}, \quad (1)$$

encapsulates the ratio of the gravitational radius $r_g = 2GM/c^2$ to the wavelength λ (here we adopt geometric units such that $G = c = 1$). The long wavelength ($M\omega \ll 1$), short wavelength ($M\omega \gg 1$) and intermediate regimes have been studied with a combination of perturbative [15, 20–22], semi-classical [10, 23] and numerical methods. The $s = 0$ (scalar) [2, 7, 12, 13, 24], $s = 1/2$ (fermion) [14, 18], $s = 1$ (electromagnetic) [6, 17, 25] and $s = 2$ (gravitational) cases [8, 9, 16] have all been covered.

Time-independent scattering by a compact body with a regular centre, such as a neutron star or white dwarf, has received comparatively less attention. In such a scenario, an electromagnetic wave will not penetrate inside the compact body; on the other hand, a gravitational wave will pass through the body without impediment from the matter distribution. Similarly, a neutron star is expected to be substantially transparent to neutrinos. In these cases, the body scatters the wave indirectly, through its influence on the spacetime curvature. Recent work [26, 27] has explored the *rainbow scattering* phenomenon that arises at short wavelengths, due to a stationary point in the geodesic deflection function associated with a ray that passes somewhat inside the body. In principle, the rainbow angle is a diagnostic of the matter distribution of the body, and thus its nuclear equation of state. [28]

Now describe the application of CAM methods to black holes, and its successes ... The aim of this work is to extend the methods to compact bodies.

Describe QNMs - resonances in the time domain: [29–35] Review of QNMs: [36]. Now describe how these are

* m.ouldelhadj@sheffield.ac.uk

† tstratton1@sheffield.ac.uk

‡ s.dolan@sheffield.ac.uk

connected to Regge poles.

Finally a paragraph on the connections to Mie scattering by a transparent sphere.

[28, 37, 38].

II. THE REGGE POLE SPECTRUM OF A COMPACT BODY

In this section we describe the model for our compact object. Then we recall the partial wave expansions of the differential scattering cross section for plane monochromatic scalar waves impinging upon a compact body. Finally we apply the CAM-approach developed for black-hole scattering in [39] and [38].

A. The model

The gravitating source is assumed to be spherically-symmetric, such that in a coordinate system $\{t, r, \theta, \varphi\}$, the object is described by a diagonal metric $g_{\mu\nu}$ and the line element

$$ds^2 = g_{\mu\nu} dx^\mu dx^\nu = -f(r) dt^2 + h(r)^{-1} dr^2 + r^2 d\sigma_2^2 \quad (2)$$

where $d\sigma_2^2$ denotes the metric on the unit 2-sphere S^2 . In the vacuum exterior of the star (i.e., $r > R$), the radial functions $f(r)$ and $h(r)$ depend only on M , the total mass of body: $f(r) = h(r) = 1 - 2M/r$ by Birkhoff's theorem [40]. In the interior, $f(r)$ and $h(r)$ depend on the matter distribution and equation of state (EoS).

A widely-studied model is that of a polytropic star, with an EoS $p(\rho) = \kappa \rho^{1+1/n}$, where n is the polytropic index (see e.g. [27]). Here we shall consider an incompressible perfect fluid ball of uniform density described by Schwarzschild's interior solution for an incompressible fluid [41], with

$$\rho = \frac{M}{\frac{4}{3}\pi R^3}, \quad (3a)$$

$$p = \rho \frac{\beta(R) - \beta(r)}{\beta(r) - 3\beta(R)}, \quad (3b)$$

$$\beta(r) = \sqrt{3 - 8\pi \rho r^2} \quad (3c)$$

and

$$f(r) = \frac{1}{4} \left(1 - \frac{2Mr^2}{R^3} \right) + \frac{9}{4} \left(1 - \frac{2M}{R} \right) - \frac{3}{2} \sqrt{\left(1 - \frac{2M}{R} \right) \left(1 - \frac{2Mr^2}{R^3} \right)}, \quad (4a)$$

$$h(r) = 1 - \frac{2Mr^2}{R^3}. \quad (4b)$$

The constant-density model can be thought of as representing the $n = 0$ limit of the polytropes. The radial function $h(r)$ is C^0 (i.e. continuous but not differentiable) at the surface of the star $r = R$, and the radial function $f(r)$ is C^1 there (i.e. once but not twice differentiable). In the polytropic cases $n > 0$, the functions are more regular at the surface, but not C^∞ (i.e. not smooth). As we shall see, the breakdown of smoothness leads to consequences for the Regge pole spectrum.

We shall consider a scalar wave $\Phi(x)$ on the compact body spacetime governed by the Klein-Gordon equation

$$\square \Phi \equiv \frac{1}{\sqrt{-g}} \partial_\mu (\sqrt{-g} g^{\mu\nu} \partial_\nu \Phi) = 0 \quad (5)$$

where $g^{\mu\nu}$ is the inverse metric $g = -f(r)h^{-1}(r)r^4 \sin^2 \theta$ and is the metric determinant. Performing a standard separation of variables,

$$\Phi = \frac{1}{r} \sum_{\omega \ell m} \phi_{\omega \ell}(r) Y_{\ell m}(\theta, \phi) e^{-i\omega t}, \quad (6)$$

leads to a radial equation of the form

$$\left[\frac{d^2}{dr_*^2} + \omega^2 - V_\ell(r) \right] \phi_{\omega \ell} = 0, \quad (7)$$

where $V_\ell(r)$ is the effective potential, and r^* denotes the *tortoise coordinate* defined by

$$dr/dr^* = \sqrt{f(r)h(r)}. \quad (8)$$

B. Effective potentials

The effective potential for the scalar field is $V_\ell(r) = V_\ell^{(s=0)}(r)$, where

$$V_\ell^{(s)}(r) \equiv f(r) \left[\frac{\ell(\ell+1)}{r^2} + \frac{(1-s^2)h(r)}{2r} \left(\frac{f'(r)}{f(r)} + \frac{h'(r)}{h(r)} \right) \right] \quad (9)$$

Remarkably, the radial equation for axial gravitational perturbations is identical to Eq. (7) but with the potential replaced by $V_\ell(r) \rightarrow V_\ell^{\text{ax}}(r)$ where [48]

$$V_\ell^{\text{ax}}(r) = V_\ell^{(s=2)} + 8\pi f(r)(p - \rho). \quad (10)$$

Outside the star in the vacuum region ($r > R$), $V_\ell(r)$ reduces to the Regge-Wheeler potential ,

$$V_\ell(r) = \left(1 - \frac{2M}{r}\right) \left(\frac{\ell(\ell+1)}{r^2} + \frac{2M(1-s^2)}{r^3}\right) \quad (11)$$

with $s = 0$ in the scalar-field case, and $s = 2$ in the axial gravitational-wave case. In the exterior, the tortoise coordinates r^* reduces to $r^* = r + 2M \ln[r/(2M) - 1] + \text{const.}$

Effective potentials for the incompressible model are shown in Fig. 1, for two cases: (i) a neutron-star model with $R = 6M$, and (ii) a ultra-compact object with $R = 2.26M$ (i.e. close to the Buchdahl bound). In both cases we observe a discontinuity in $V_\ell(r)$ across the star's surface, due to the C^0 behaviour of $h(r)$. The jump in the potential takes opposite signs in the scalar-field and gravitational-wave cases, with

$$\Delta V^{(s=0)} = +\frac{3Mf(R)}{R^3}, \quad (12a)$$

$$\Delta V^{(ax)} = -\frac{3Mf(R)}{R^3}, \quad (12b)$$

where

$$\Delta V \equiv \lim_{\epsilon \rightarrow 0} \{V(R+\epsilon) - V(R-\epsilon)\}. \quad (13)$$

C. Boundary conditions

It is important to recall that the modes $\phi_{\omega\ell}$ should have a regular behaviour at the origin $r \rightarrow 0$

$$\phi_{\omega\ell}(r) \underset{r \rightarrow 0}{\sim} r^{\ell+1}. \quad (14)$$

The asymptotic behaviour of the modes at spatial infinity $r \rightarrow +\infty$ (i.e., for $r_* \rightarrow +\infty$) is

$$\phi_{\omega\ell}(r) \underset{r_* \rightarrow +\infty}{\sim} A_\ell^{(-)}(\omega) e^{-i\omega r_*} + A_\ell^{(+)}(\omega) e^{+i\omega r_*}. \quad (15)$$

In this last equation, the coefficients $A_\ell^{(-)}(\omega)$ and $A_\ell^{(+)}(\omega)$ are complex amplitudes and we have

$$S_\ell(\omega) = e^{i(\ell+1)\pi} \frac{A_\ell^{(+)}(\omega)}{A_\ell^{(-)}(\omega)}. \quad (16)$$

We can expect then that Regge poles for axial gravitational perturbations are qualitatively the same as for scalar perturbations, and the methods discussed can be easily adapted to deal with either case.




FIG. 1. The effective potential V_ℓ for a quadrupole ($\ell = 2$) perturbation of a compact body of constant density and tenuity $R/M = 6$ (upper) and $R/M = 2.26$ (lower). The scalar field and axial gravitational-wave potentials are indicated as solid/black lines and dotted/red lines, respectively. The horizontal axis is the tortoise coordinate r_* defined in Eq. (8).

D. Partial wave expansions of differential scattering cross sections

E. Scattering theory

We recall that, for the scalar field, the differential scattering cross section is given by (see, e.g., [26] and references therein)

$$\frac{d\sigma}{d\Omega} = |\hat{f}(\omega, \theta)|^2 \quad (17)$$

where

$$\hat{f}(\omega, \theta) = \frac{1}{2i\omega} \sum_{\ell=0}^{\infty} (2\ell+1) [S_\ell(\omega) - 1] P_\ell(\cos \theta) \quad (18)$$

denotes the scattering amplitude. In Eq. (18), the functions $P_\ell(\cos \theta)$ are the Legendre polynomials [42]. We

also recall that the S -matrix elements $S_\ell(\omega)$ appearing in Eq. (18) can be defined from the modes $\phi_{\omega\ell}$ that solve the homogenous radial equation

F. CAM representation of the scattering amplitude

To construct the CAM representation, we follow the steps in section II of the Ref [39] and recall the main results.

By using the Sommerfeld-Watson transformation [43–45] which permits us to write

$$\sum_{\ell=0}^{+\infty} (-1)^\ell F(\ell) = \frac{i}{2} \int_{\mathcal{C}} d\lambda \frac{F(\lambda - 1/2)}{\cos(\pi\lambda)} \quad (19)$$

with a function F without any singularities on the real λ axis, we replace the discrete sum over the ordinary angular momentum ℓ in Eq. (18) by a contour integral in the complex λ plane (i.e., in the complex ℓ plane with $\lambda = \ell + 1/2$). By noting that $P_\ell(\cos \theta) = (-1)^\ell P_\ell(-\cos \theta)$, we obtain

$$\hat{f}(\omega, \theta) = \frac{1}{2\omega} \int_{\mathcal{C}} d\lambda \frac{\lambda}{\cos(\pi\lambda)} \times [S_{\lambda-1/2}(\omega) - 1] P_{\lambda-1/2}(-\cos \theta). \quad (20)$$

In Eqs. (19) and (20), the integration contour encircles counterclockwise the positive real axis of the complex λ plane, i.e., we take $\mathcal{C} =]+\infty + i\epsilon, +i\epsilon[\cup]+i\epsilon, -i\epsilon[\cup [-i\epsilon, +\infty - i\epsilon[$ with $\epsilon \rightarrow 0_+$ (see Fig.1 Ref [39]).

The Legendre function of the first kind $P_{\lambda-1/2}(z)$ denotes the analytic extension of the Legendre polynomials $P_\ell(z)$. It is defined in terms of hypergeometric functions by [42]

$$P_{\lambda-1/2}(z) = F[1/2 - \lambda, 1/2 + \lambda; 1; (1 - z)/2]. \quad (21)$$

In Eq. (20), $S_{\lambda-1/2}(\omega)$ denotes “the” analytic extension of $S_\ell(\omega)$. It is given by [see Eq. (16)]

$$S_{\lambda-1/2}(\omega) = e^{i(\lambda+1/2)\pi} \frac{A_{\lambda-1/2}^{(+)}(\omega)}{A_{\lambda-1/2}^{(-)}(\omega)} \quad (22)$$

where the complex amplitudes $A_{\lambda-1/2}^{(-)}(\omega)$ and $A_{\lambda-1/2}^{(+)}(\omega)$ are defined from the analytic extension of the modes $\phi_{\omega\ell}$, i.e., from the function $\phi_{\omega, \lambda-1/2}$.

It is important to note that the poles of $S_{\lambda-1/2}(\omega)$ in the complex λ plan (i.e., the Regge poles) are defined as the zeros $\lambda_n(\omega)$ with $n = 1, 2, 3, \dots$ of the coefficient $A_{\lambda-1/2}^{(-)}(\omega)$ [see Eq. (22)]

$$A_{\lambda_n(\omega)-1/2}^{(-)}(\omega) = 0. \quad (23)$$

The residue of the matrix $S_{\lambda-1/2}(\omega)$ at the pole $\lambda = \lambda_n(\omega)$ is defined by [see Eq. (22)]

$$r_n(\omega) = e^{i\pi[\lambda_n(\omega)+1/2]} \left[\frac{A_{\lambda-1/2}^{(+)}(\omega)}{\frac{d}{d\lambda} A_{\lambda-1/2}^{(-)}(\omega)} \right]_{\lambda=\lambda_n(\omega)}. \quad (24)$$

These residues play a central role in the complex angular momentum paradigm.

Now, we “deform” the contour \mathcal{C} in Eq. (20) in order to collect, by using Cauchy’s theorem, the Regge poles contributions. This is achieved by following, *mutatis mutandis*, the approach developed in Ref [39] (see more particularly Sec. IIB 3 and Fig. 1). We obtain

$$\hat{f}(\omega, \theta) = \hat{f}^B(\omega, \theta) + \hat{f}^{\text{RP}}(\omega, \theta) \quad (25)$$

where

$$\hat{f}^B(\omega, \theta) = \hat{f}^{\text{B,Re}}(\omega, \theta) + \hat{f}^{\text{B,Im}}(\omega, \theta) \quad (26a)$$

is a background integral contribution with

$$\hat{f}^{\text{B,Re}}(\omega, \theta) = \frac{1}{\pi\omega} \int_{\mathcal{C}_-} d\lambda \lambda S_{\lambda-1/2}(\omega) Q_{\lambda-1/2}(\cos \theta + i0) \quad (26b)$$

and

$$\hat{f}^{\text{B,Im}}(\omega, \theta) = \frac{1}{2\omega} \left(\int_{+i\infty}^0 d\lambda [S_{\lambda-1/2}(\omega) P_{\lambda_n(\omega)-1/2}(-\cos \theta) - S_{-\lambda-1/2}(\omega) e^{i\pi(\lambda+1/2)} P_{\lambda_n(\omega)-1/2}(\cos \theta)] \lambda \right). \quad (26c)$$

The second term in Eq. (25),

$$\hat{f}^{\text{RP}}(\omega, \theta) = -\frac{i\pi}{\omega} \sum_{n=1}^{+\infty} \frac{\lambda_n(\omega) r_n(\omega)}{\cos[\pi\lambda_n(\omega)]} \times P_{\lambda_n(\omega)-1/2}(-\cos \theta), \quad (27)$$

is a sum over the Regge poles lying in the first quadrant of the CAM plane. Of course, Eqs. (25), (26) and (27) provide an exact representation of the scattering amplitude $\hat{f}(\omega, \theta)$ for the scalar field, equivalent to the initial partial wave expansion (18). From this CAM representation, we can extract the contribution $\hat{f}^{\text{RP}}(\omega, \theta)$ given by (27) which, as a sum over Regge poles, is only an approximation of $\hat{f}(\omega, \theta)$, and which provides us with an approximation of the differential scattering cross section (17).

III. RECONSTRUCTION OF DIFFERENTIAL SCATTERING CROSS SECTIONS FROM REGGE POLE SUMS

In this section, we compare the partial wave expansions of the differential scattering cross sections with their

equivalent CAM representations or, more precisely, their Regge pole approximations.

A. Computational methods

To construct the scattering amplitude (18), the background integrals (26b) and (26c) as well as the Regge pole contribution (27), we use, *mutatis mutandis* the computational methods that permitted one of us, in Refs [38, 39] to consider the CAM representation for scattering of the scalar, electromagnetic and gravitational waves by Schwarzschild BH (see also Ref [26]). It is important to remark that, due to the long rang nature of the field propagating on the Schwarzschild BH (outside the compact body), the scattering amplitude (18) and the background integral (26b) suffer a lack of convergence and to overcome this problem, i.e., to accelerate the convergence of this sum and integral, we have used the method described in the Appendix of Ref [39]. We have performed all the numerical calculations by using *Mathematica* [46].

B. Results and comments




FIG. 2. The Regge poles $\lambda_n(\omega)$ for the scalar field. We assume $2M = 1$




FIG. 3. The Regge poles $\lambda_n(\omega)$ for the scalar field. We assume $2M = 1$

IV. WKB ?

The WKB approach developed by Zhang, Wu and Leung [47] to determine the axial w -modes of a variety of stellar models can be adapted in order to obtain analytical approximations for the “broad Regge poles” for a scalar wave on a stellar background. The radial equation for axial gravitational perturbations is identical to Eq. (7) but with the potential replaced by $V_\ell \rightarrow V_\ell^{\text{ax}}$ where [48]

$$V_\ell^{\text{ax}}(r) = f(r) \left[\frac{\ell(\ell+1)}{r^2} - \frac{3h(r)}{2r} \left(\frac{f'(r)}{f(r)} + \frac{h'(r)}{h(r)} \right) + 8\pi(p - \rho) \right] \quad (28)$$

We can expect then that Regge poles for axial gravitational perturbations are qualitatively the same as for scalar perturbations, and the methods discussed can be easily adapted to deal with either case.

Regge poles and quasinormal modes for relativistic stellar models (of which w -modes are a sub-category for gravitational perturbations) both satisfy the same wave




FIG. 4. The scalar cross section of a compact bodies for $2M\omega = 3$ and $R = 6M$, its Regge pole approximation and the background integral contribution.




FIG. 5. The scalar cross section of a compact bodies for $2M\omega = 16$ and $R = 6M$, its Regge pole approximation and the background integral contribution.




FIG. 6. The scalar cross section of a very compact bodies for $2M\omega = 3$ and $R = 2.26M$ and its Regge pole approximation.




FIG. 7. The scalar cross section of a very compact bodies for $2M\omega = 6$ and $R = 2.26M$ and its Regge pole approximation.

TABLE I. The lowest Regge poles $\lambda_n(\omega)$ for the scalar field and the associated residues $r_n(\omega)$. The radius of the compact bodies is $R = 6M$ and we assume $2M = 1$.

n	ω	$\lambda_n^{(S-W)}(\omega)$	$\lambda_n^{(B-R)}(\omega)$	$r_n^{(S-W)}(\omega)$	$r_n^{(B-R)}(\omega)$
1	3	$9.64850 + 2.76784i$	$1.56219 + 2.33072i$	$-12.41483 - 0.10424i$	$-0.184457 + 0.480330i$
	16	$56.00945 + 5.71038i$	$0.62529 + 3.27098i$	$-447.5395 + 25.2912i$	$-0.322061 - 0.088002i$
2	3	$10.71986 + 5.16209i$	$3.81484 + 2.48159i$	$13.8486 + 24.3824i$	$0.290952 + 1.043116i$
	16	$58.442656 + 9.18793i$	$2.64868 + 3.31439i$	$5188.750 - 859.909i$	$-0.381581 - 0.077583i$
3	3	$11.62296 + 7.17454i$	$6.35675 + 2.64104i$	$39.4189 - 12.3554i$	$2.83038 - 0.28686i$
	16	$60.20374 + 12.14965i$	$4.70011 + 3.35821i$	$-29331.71 - 18578.38i$	$-0.456423 - 0.021249i$
4	3	$12.4297 + 8.9960i$	/	$13.2301 - 50.8802i$	/
	16	$61.67700 + 14.84728i$	$6.78093 + 3.40257i$	$-15868.9 + 161199.9i$	$-0.528929 + 0.106794i$
5	3	$13.1734 + 10.6929i$	/	$-33.7366 - 51.7404i$	/
	16	$62.98626 + 17.37165i$	$8.89270 + 3.44762i$	$589920.5 - 79507.8i$	$-0.550038 + 0.330275i$
6	3	$13.8709 + 12.2989i$	/	$-66.4436 - 20.7767i$	/
	16	$64.18605 + 19.76911i$	$11.03720 + 3.49356i$	$-360464. - 1.797518 \times 10^6 i$	$-0.426365 + 0.639191i$
7	3	$14.5322 + 13.8342i$	/	$-73.0825 + 21.9088i$	/
	16	$65.30640 + 22.06743i$	$13.21653 + 3.54058i$	$-4.880638 \times 10^6 + 646112.i$	$-0.038292 + 0.926498i$
8	3	$15.1640 + 15.3122i$	/	$-56.3641 + 59.6187i$	/
	16	$66.36581 + 24.28491i$	$15.43310 + 3.5889i$	$-479098. + 1.1836070 \times 10^7 i$	$0.652285 + 0.920876i$
9	3	$15.7709 + 16.7425i$	/	$-25.0183 + 83.3731i$	/
	16	$67.37659 + 26.43447i$	$17.6898 + 3.6390i$	$2.487209 \times 10^7 + 7.72797 \times 10^6 i$	$1.363464 + 0.248276i$
10	3	$16.3565 + 18.1321i$	/	$11.7631 + 90.6815i$	/
	16	$68.34738 + 28.52564i$	$19.9900 + 3.6910i$	$3.163822 \times 10^7 - 4.265475 \times 10^7 i$	$1.29469 - 1.13096i$

¹ Surface waves

² Broad resonances

TABLE II. The lowest Regge poles $\lambda_n(\omega)$ for the scalar field and the associated residues $r_n(\omega)$. The radius of the compact bodies is $R = 2.26M$ and we assume $2M = 1$.

n	ω	$\lambda_n^{(S-W)}(\omega)$	$\lambda_n^{(B-R)}(\omega)$	$\lambda_n^{(N-R)}(\omega)$	$r_n^{(S-W)}(\omega)$	$r_n^{(B-R)}(\omega)$	$r_n^{(N-R)}(\omega)$
1	3	$5.871590 + 1.553799i$	$1.73455 + 1.64951i$	$6.48474 + 0.68765i$	$-179.7945 + 131.4187i$	$-1.52081 - 2.30968i$	$-2.5672 - 15.3797i$
	6	$12.991923 + 1.754967i$	$1.89664 + 2.13696i$	$13.34118 + 1.13496i$	$4356.193 + 647.790i$	$-0.66176 - 1.31963i$	$-390.218 + 379.906i$
2	3	$5.778805 + 3.228990i$	$3.48084 + 1.45765i$	$7.25606 + 0.24457i$	$428.6893 - 235.0321i$	$16.2123 + 5.2371i$	$-0.272250 - 1.150335i$
	6	$12.705495 + 3.383881i$	$3.74238 + 2.01309i$	$14.18757 + 0.68182i$	$-35075.99 - 9772.94i$	$-2.93679 + 4.83548i$	$-11.3519 + 34.5571i$
3	3	$5.924546 + 4.705899i$	$5.10229 + 1.29099i$	$7.95763 + 0.01764i$	$-404.6185 - 390.8531i$	$70.4849 + 54.1888i$	$-0.0370202 - 0.0048174i$
	6	$12.596259 + 4.982661i$	$5.49829 + 1.89576i$	$14.9017 + 0.2912i$	$82360.19 + 81990.53i$	$6.7872 - 16.9564i$	$0.27028 + 2.27905i$
4	3	$6.144986 + 6.043188i$	/	/	$-471.5443 + 314.3116i$	/	/
	6	$12.614598 + 6.503749i$	$7.17509 + 1.78279i$	$15.5621 + 0.0422i$	$39281.5 - 229393.2i$	$39.6176 + 33.5152i$	$0.1011154 + 0.0020569i$
5	3	$6.398427 + 7.281723i$	/	/	$37.8777 + 546.8945i$	/	/
	6	$12.71646 + 7.95208i$	$8.78112 + 1.67243i$	/	$-356055.5 + 34945.9i$	$2.1175 + 134.5962i$	/
6	3	$6.666837 + 8.447532i$	/	/	$418.7890 + 315.4209i$	/	/
	6	$12.87420 + 9.33552i$	$10.32300 + 1.56317i$	/	$45934.6 + 468157.5i$	$66.944 + 324.598i$	/
7	3	$6.941642 + 9.557619i$	/	/	$499.2703 - 37.6476i$	/	/
	6	$13.06993 + 10.66226i$	$11.80630 + 1.45720i$	/	$558619.4 + 61956.5i$	$833.855 + 78.332i$	/
8	3	$7.218463 + 10.623548i$	/	/	$367.2578 - 307.7533i$	/	/
	6	$13.29184 + 11.93979i$	/	/	$293571.8 - 559756.8i$	/	/
9	3	$7.494953 + 11.653498i$	/	/	$147.3038 - 435.8160i$	/	/
	6	$13.53197 + 13.17461i$	/	/	$-376511.5 - 570254.0i$	/	/
10	3	$7.76982 + 12.65345i$	/	/	$-71.8294 - 437.3469i$	/	/
	6	$13.78485 + 14.37216i$	/	/	$-719306.1 - 20011.7i$	/	/

¹ Surface waves

² Broad resonances

³ Narrow resonances




FIG. 8. Rainbow scattering for compact bodies for $2M\omega = 16$ and $R = 6M$, its Regge pole approximation and different contributions of the sum over Regge poles.




FIG. 9. Rainbow scattering for compact bodies for $2M\omega = 16$ and $R = 2.26M$, its Regge pole approximation and different contributions of the sum over Regge poles.

equation and the same boundary conditions but with different interpretations for the angular momentum index and the frequency. Both types of pole satisfy the regularity condition at the origin (14) and the condition of a purely outgoing wave in the far field

$$\phi_{\omega, \lambda-1/2}^{\text{out}}(r) \underset{r \rightarrow +\infty}{\sim} A_{\lambda-1/2}^{(+)}(\omega) e^{+i\omega r_*}. \quad (29)$$

Thus, the Regge poles are solutions of Eq. (7) for which the Wronskian of $\phi_{\omega, \lambda-1/2}$ and $\phi_{\omega, \lambda-1/2}^{\text{out}}$ vanishes (*i.e.*, $A_{\lambda_n(\omega)-1/2}^{(-)}(\omega) = 0$)

$$W[\phi_{\omega, \lambda-1/2}, \phi_{\omega, \lambda-1/2}^{\text{out}}] = 0 \quad (30)$$

It has been shown that, the asymptotic expressions of $\phi_{\omega, \lambda-1/2}$ and $\phi_{\omega, \lambda-1/2}^{\text{out}}$ can be derived in certain regions, but using the standard WKB approximation. In the high-frequency domain we have (see Refs ...)

$$\phi_{\omega, \lambda-1/2} \underset{\omega \rightarrow \infty}{=} \omega r_* j_{\lambda-1/2}(\omega r_*) \quad 0 \leq r_* \leq R_* \quad (31)$$

where $j_{\lambda-1/2}$ is the spherical Bessel function of the first kind and

$$\phi_{\omega, \lambda-1/2}^{\text{out}} \underset{\omega \rightarrow \infty}{=} \begin{cases} e^{i\omega(r_*-R_*)} + \mathcal{R}e^{-i\omega(r_*-R_*)} & 1/\omega \leq r_* < R_*, \\ (1 + \mathcal{R})e^{i\omega(r_*-R_*)} & R_* \leq r_* < \infty. \end{cases} \quad (32)$$

Here

$$R_* = \int_0^R dr \frac{1}{\sqrt{f(r)h(r)}} \quad (33)$$

and \mathcal{R} is a reflection coefficient with the definition given in Ref. [49]. Because the potential has a direct discontinuity at the surface of the compact body (see. Refs [47, 49] for more details), we have for our model (*i.e.*, compact body with a constant density)

$$\mathcal{R} = a\omega^{-2} \quad (34)$$

with

$$a = -\left(\frac{i}{2}\right)^2 \lim_{\epsilon \rightarrow 0^+} \left[V_{\lambda-1/2}(r) \Big|_{r=R+\epsilon} - V_{\lambda-1/2}(r) \Big|_{r=R-\epsilon} \right] = \frac{3M(R-2M)}{4R^4} \quad (35)$$

Now, by inserting the high frequency approximation for $j_{\lambda-1/2}(\omega r_*)$ into Eq. (31) [50],

$$\phi_{\omega, \lambda-1/2} \underset{\omega \rightarrow \infty}{\approx} -\sin\left(\frac{(\lambda-1/2)\pi}{2} - \omega r_*\right) \quad 0 \leq r_* \leq R_* \quad (36)$$

and Eq. (32) into the condition (30), we obtain

$$e^{i\pi(\lambda-1/2)-2i\omega R_*} = -\mathcal{R}. \quad (37)$$

We then solve Eq. (37) and we obtain

$$\lambda_n \approx \frac{2\omega R_*}{\pi} - \left(2n + \frac{1}{2}\right) + \frac{i}{\pi} \ln\left(\frac{1}{\mathcal{R}}\right) \quad (38)$$

This corresponds to the series of Regge poles with spacing $|\Delta\lambda_n| \approx 2$ with the almost constant imaginary part. Of course, they lie in the first quadrant of the CAM plan with a positive real part

$$\frac{2\omega R_*}{\pi} - \left(2n + \frac{1}{2}\right) > 0. \quad (39)$$

The overtones are labelled by $n = 1, 2, \dots$ and n has an upper limit

$$n \leq \left\lfloor \frac{\omega R_*}{\pi} - \frac{1}{4} \right\rfloor. \quad (40)$$

In other words, there is a finite number of the “broad Regge poles”.

TABLE III. The lowest Regge poles $\lambda_n(\omega)$ for the scalar field versus WBK results given by Eq. (38). The radius of the compact bodies is $R = 6M$ and we assume $2M = 1$.

n	ω	$\lambda_n^{(\text{B-R})}(\omega)$	$\lambda_n^{(\text{B-R, WBK})}(\omega)$
1	3	$1.56219 + 2.33072i$	$1.592793 + 2.189767i$
	16	$0.62529 + 3.27098i$	$0.661564 + 3.255453i$
2	3	$3.81484 + 2.48159i$	$3.592793 + 2.189767i$
	16	$2.64868 + 3.31439i$	$2.661564 + 3.255453i$
3	3	$6.35675 + 2.64104i$	$5.592793 + 2.189767i$
	16	$4.70011 + 3.35821i$	$4.661564 + 3.255453i$
4	3	/	/
	16	$6.78093 + 3.40257i$	$6.661564 + 3.255453i$
5	3	/	/
	16	$8.89270 + 3.44762i$	$8.661564 + 3.255453i$
6	3	/	/
	16	$11.03720 + 3.49356i$	$10.661564 + 3.255453i$
7	3	/	/
	16	$13.21653 + 3.54058i$	$12.661564 + 3.255453i$
8	3	/	/
	16	$15.4331 + 3.5889i$	$14.661564 + 3.255453i$
9	3	/	/
	16	$17.6898 + 3.6390i$	$16.661564 + 3.384517i$
10	3	/	/
	16	$19.9900 + 3.6910i$	$18.661564 + 3.255453i$

V. CLASSIFICATION OF QUASINORMAL MODES

The study of stellar quasinormal modes began with a general relativistic extension of the Newtonian treatment of fluid pulsations of stars. Newly formed neutron stars, the remnants of supernovae collapse, are predicted to pulsate with a large initial energy [51]. It was and still is of interest to study how these pulsations lose energy in the form of gravitational waves. Recently, the gravitational waves from a neutron star merger were detected... In particular the emitted radiation should carry a signature of the modes of oscillation allowed inside the star. These fluid modes can be classified in analogy to their real Newtonian counterparts, with an additional damping time due to emission of GWs. Later, Kokkotas and Schutz showed that an additional family of modes existed [30, 32], dubbed ω -modes. These modes are characterised by a negligible excitation of fluid motion (and in the axial sector no fluid motion). They're highly damped and correspond to excitations of the dynamical perturbed space-time. For an excellent review of (gravitational) quasinormal modes in relativistic stars and black holes see [36].

We consider the massless scalar field which only cou-

ples to the body via gravitation. The scalar modes obey an equation of motion very similar to the axial gravitational wave sector. Accordingly, a similar spectrum for scalar modes can be expected (with no fluid excitation). Whilst these are arguably of less astrophysical interest, they are nonetheless a good model for axial GWs and allow us to make a first step towards a CAM approach for perturbed relativistic stars. We will classify the scalar ω -modes as the axial GW modes have been, we have

1. Curvature modes: These modes exist for all relativistic stars. They are rapidly damped, and the damping is larger for less compact bodies.
2. Trapped modes: These are the scalar analogue of the axial trapped modes considered by Chandrasekhar and Ferrari [31]. They exist for bodies with $R/M < 3$. These models have an effective radial potential with a local minimum somewhere inside the star, and local maximum at the photon sphere $r = 3M$. The trapped modes are essentially the first few curvature modes with energy less than the potential barrier. There are a finite number of them, and the number increases with the depth of the potential well.
3. Interface modes: also known as ω_{II} -modes, these were discovered by Leins *et al.* [33]. They used a generalisation of Leaver's continued fraction method as well as a 'Wronskain method' that allowed them to accurately compute these modes which are characterised by very rapid damping (large imaginary part). This second branch of modes is the most similar to the Schwarzschild black hole quasinormal modes. We have taken their continued fraction method one step further (see section...) in order to be confident with our results.

ACKNOWLEDGMENTS

M. O. E. H. wish to thank Antoine Folacci for various discussions concerning this work. T.S. acknowledges financial support from EPSRC. S.R.D. acknowledges financial support from the European Union's Horizon 2020 research and innovation programme under the H2020-MSCA-RISE-2017 Grant No. FunFiCO-777740, and from the Science and Technology Facilities Council (STFC) under Grant No. ST/P000800/1.

-
- [1] W. W. Hildreth, *The Interaction of Scalar Gravitational Waves with the Schwarzschild Metric.*, Ph.D. thesis (1964).
- [2] Richard A Matzner, “Scattering of massless scalar waves by a schwarzschild “singularity”,” *Journal of Mathematical Physics* **9**, 163–170 (1968).
- [3] CV Vishveshwara, “Scattering of gravitational radiation by a Schwarzschild black-hole,” *Nature* **227**, 936 (1970).
- [4] P. L. Chrzanowski, R. A. Matzner, V. D. Sandberg, and M. P. Ryan, “Zero Mass Plane Waves in Nonzero Gravitational Backgrounds,” *Phys. Rev.* **D14**, 317–326 (1976).
- [5] Bahram Mashhoon, “Scattering of Electromagnetic Radiation from a Black Hole,” *Phys. Rev.* **D7**, 2807–2814 (1973).
- [6] R Fabbri, “Scattering and absorption of electromagnetic waves by a schwarzschild black hole,” *Physical Review D* **12**, 933 (1975).
- [7] Norma G. Sanchez, “Elastic Scattering of Waves by a Black Hole,” *Phys. Rev.* **D18**, 1798 (1978).
- [8] Richard A. Matzner and Michael P. Jr. Ryan, “Scattering of gravitational radiation from vacuum black holes.” *The Astrophysical Journal Supplement Series* **36**, 451–481 (1978).
- [9] F. A. Handler and R. A. Matzner, “Gravitational wave scattering,” *Phys. Rev.* **D22**, 2331–2348 (1980).
- [10] Richard A. Matzner, Cécile DeWitt-Morette, Bruce Nelson, and Tian-Rong Zhang, “Glory scattering by black holes,” *Phys. Rev.* **D31**, 1869 (1985).
- [11] J. A. H. Futterman, F. A. Handler, and R. A. Matzner, *Scattering from black holes* (Cambridge University Press, 2012).
- [12] N. Andersson, “Scattering of massless scalar waves by a Schwarzschild black hole: A Phase integral study,” *Phys. Rev.* **D52**, 1808–1820 (1995).
- [13] Kostas Glampedakis and Nils Andersson, “Scattering of scalar waves by rotating black holes,” *Class. Quant. Grav.* **18**, 1939–1966 (2001), [arXiv:gr-qc/0102100 \[gr-qc\]](#).
- [14] Sam Dolan, Chris Doran, and Anthony Lasenby, “Fermion scattering by a Schwarzschild black hole,” *Phys. Rev.* **D74**, 064005 (2006), [arXiv:gr-qc/0605031 \[gr-qc\]](#).
- [15] Sam R. Dolan, “Scattering of long-wavelength gravitational waves,” *Phys. Rev.* **D77**, 044004 (2008), [arXiv:0710.4252 \[gr-qc\]](#).
- [16] Sam R. Dolan, “Scattering and Absorption of Gravitational Plane Waves by Rotating Black Holes,” *Class. Quant. Grav.* **25**, 235002 (2008), [arXiv:0801.3805 \[gr-qc\]](#).
- [17] Luis C. B. Crispino, Sam R. Dolan, and Ednilton S. Oliveira, “Electromagnetic wave scattering by Schwarzschild black holes,” *Phys. Rev. Lett.* **102**, 231103 (2009), [arXiv:0905.3339 \[gr-qc\]](#).
- [18] Ion I. Cotaescu, Cosmin Crucean, and Ciprian A. Sporea, “Partial wave analysis of the Dirac fermions scattered from Schwarzschild black holes,” *Eur. Phys. J.* **C76**, 102 (2016), [arXiv:1409.7201 \[gr-qc\]](#).
- [19] Alexander Gußmann, “Scattering of massless scalar waves by magnetically charged black holes in Einstein–Yang–Mills–Higgs theory,” *Class. Quant. Grav.* **34**, 065007 (2017), [arXiv:1608.00552 \[hep-th\]](#).
- [20] W. K. De Logi and S. J. Kovacs, “Gravitational Scattering of Zero Rest Mass Plane Waves,” *Phys. Rev.* **D16**, 237–244 (1977).
- [21] E. Guadagnini, “Gravitons scattering from classical matter,” *Class. Quant. Grav.* **25**, 095012 (2008), [arXiv:0803.2855 \[gr-qc\]](#).
- [22] Francesco Sorge, “On the gravitational scattering of gravitational waves,” *Class. Quant. Grav.* **32**, 035007 (2015).
- [23] P. Anninos, C. DeWitt-Morette, R. A. Matzner, P. Yioutas, and T. R. Zhang, “Orbiting cross-sections: Application to black hole scattering,” *Phys. Rev.* **D46**, 4477–4494 (1992).
- [24] Luiz C. S. Leite, Carolina L. Benone, and Luís C. B. Crispino, “On-axis scattering of scalar fields by charged rotating black holes,” *Phys. Lett.* **B795**, 496–501 (2019), [arXiv:1907.04746 \[gr-qc\]](#).
- [25] Luís C. B. Crispino, Sam R. Dolan, Atsushi Higuchi, and Ednilton S. de Oliveira, “Scattering from charged black holes and supergravity,” *Phys. Rev.* **D92**, 084056 (2015), [arXiv:1507.03993 \[gr-qc\]](#).
- [26] Sam R. Dolan and Tom Stratton, “Rainbow scattering in the gravitational field of a compact object,” *Phys. Rev.* **D95**, 124055 (2017), [arXiv:1702.06127 \[gr-qc\]](#).
- [27] Tom Stratton and Sam R. Dolan, “Rainbow scattering of gravitational plane waves by a compact body,” *Phys. Rev.* **D100**, 024007 (2019), [arXiv:1903.00025 \[gr-qc\]](#).
- [28] Yasusada Nambu, Sousuke Noda, and Yuichiro Sakai, “Wave Optics in Spacetimes with Compact Gravitating Object,” (2019), [arXiv:1905.01793 \[gr-qc\]](#).
- [29] Steven L. Detweiler and L. Lindblom, “On the nonradial pulsations of general relativistic stellar models,” *Astrophys. J.* **292**, 12–15 (1985).
- [30] Kostas D. Kokkotas and Bernard F. Schutz, “Normal Modes of a Model Radiating System,” *Gen. Rel. Grav.* **18**, 913 (1986).
- [31] Subrahmanyan Chandrasekhar and Valeria Ferrari, “On the non-radial oscillations of a star iii. a reconsideration of the axial modes,” *Proceedings of the Royal Society of London A: Mathematical, Physical and Engineering Sciences* **434**, 449–457 (1991).
- [32] K. D. Kokkotas and Bernard F. Schutz, “W-modes: A New family of normal modes of pulsating relativistic stars,” *Mon. Not. Roy. Astron. Soc.* **225**, 119 (1992).

- [33] M. Leins, H. P. Nollert, and M. H. Soffel, “Nonradial oscillations of neutron stars: A New branch of strongly damped normal modes,” *Phys. Rev.* **D48**, 3467–3472 (1993).
- [34] Nils Andersson, Yasufumi Kojima, and Kostas D. Kokkotas, “On the oscillation spectra of ultracompact stars: An Extensive survey of gravitational wave modes,” *Astrophys. J.* **462**, 855 (1996), [arXiv:gr-qc/9512048 \[gr-qc\]](#).
- [35] Nils Andersson, “Two simple models for gravitational-wave modes of compact stars,” *General Relativity and Gravitation* **28**, 1433–1445 (1996).
- [36] Kostas D. Kokkotas and Bernd G. Schmidt, “Quasinormal modes of stars and black holes,” *Living Rev. Rel.* **2**, 2 (1999), [arXiv:gr-qc/9909058 \[gr-qc\]](#).
- [37] Luiz C. S. Leite, Sam R. Dolan, and Luís C. B. Crispino, “Absorption of electromagnetic and gravitational waves by Kerr black holes,” *Phys. Lett.* **B774**, 130–134 (2017), [arXiv:1707.01144 \[gr-qc\]](#).
- [38] Antoine Folacci and Mohamed Ould El Hadj, “Regge pole description of scattering of gravitational waves by a Schwarzschild black hole,” *Phys. Rev.* **D100**, 064009 (2019), [arXiv:1906.01441 \[gr-qc\]](#).
- [39] Antoine Folacci and Mohamed Ould El Hadj, “Regge pole description of scattering of scalar and electromagnetic waves by a Schwarzschild black hole,” *Phys. Rev.* **D99**, 104079 (2019), [arXiv:1901.03965 \[gr-qc\]](#).
- [40] Nils Voje Johansen and Finn Ravndal, “On the discovery of Birkhoff’s theorem,” *Gen. Rel. Grav.* **38**, 537–540 (2006), [arXiv:physics/0508163 \[physics\]](#).
- [41] S. L. Shapiro and S. A. Teukolsky, *Black Holes, White Dwarfs, and Neutron Stars: The Physics of Compact Objects* (Wiley, New-York, 1983).
- [42] M. Abramowitz and I. A. Stegun, *Handbook of Mathematical Functions* (Dover, New-York, 1965).
- [43] G. N. Watson, “The diffraction of electric waves by the Earth,” *Proc. R. Soc. London A* **95**, 83 (1918).
- [44] A. Sommerfeld, *Partial Differential Equations of Physics* (Academic Press, New York, 1949).
- [45] R. G. Newton, *Scattering Theory of Waves and Particles*, 2nd ed. (Springer-Verlag, New York, 1982).
- [46] Wolfram Research, Inc., “*Mathematica*, Version 10.0,” (Wolfram Research, Inc., Champaign, IL, 2014).
- [47] Y. J. Zhang, J. Wu, and P. T. Leung, “High-frequency behavior of w-mode pulsations of compact stars,” *Phys. Rev.* **D83**, 064012 (2011), [arXiv:1101.0319 \[gr-qc\]](#).
- [48] Vitor Cardoso, Luís C. B. Crispino, Caio F. B. Macedo, Hirotada Okawa, and Paolo Pani, “Light rings as observational evidence for event horizons: long-lived modes, ergoregions and nonlinear instabilities of ultra-compact objects,” *Phys. Rev. D* **D90**, 044069 (2014), [arXiv:1406.5510 \[gr-qc\]](#).
- [49] M V Berry, “Semiclassically weak reflections above analytic and non-analytic potential barriers,” *Journal of Physics A: Mathematical and General* **15**, 3693–3704 (1982).
- [50] F. W. J. Olver, D. W. Lozier, R. F. Boisvert, and C. W. Clark, *NIST Handbook of Mathematical Functions* (Cambridge University Press, 2010).
- [51] Kip S Thorne and Alfonso Campolattaro, “Non-Radial Pulsation of General-Relativistic Stellar Models. I. Analytic Analysis for $L_\ell = 2$,” *The Astrophysical Journal* **149**, 591 (1967).