


# NOT (SO MANY) LABELS



# IT'S THE TIME TO DISCO

- Leverages existing TOPUP capability to train
- Much faster approach



Published in final edited form as:

*Magn Reson Imaging*. 2019 December ; 64: 62–70. doi:10.1016/j.mri.2019.05.008.

## Synthesized b0 for diffusion distortion correction (Synb0-DisCo)

Kurt G Schilling<sup>1</sup>, Justin Blaber<sup>3</sup>, Yuankai Huo<sup>2</sup>, Allen Newton<sup>1,5</sup>, Colin Hansen<sup>2</sup>, Vishwesh Nath<sup>3</sup>, Andrea T. Shafer<sup>4</sup>, Owen Williams<sup>4</sup>, Susan M. Resnick<sup>4</sup>, Baxter Rogers<sup>1</sup>, Adam W Anderson<sup>1</sup>, Bennett A Landman<sup>1,2,3</sup>

<sup>1</sup>Vanderbilt University Institute of Imaging Science, Vanderbilt University, Nashville, TN

<sup>2</sup>Department of Electrical Engineering, Vanderbilt University, Nashville, TN

<sup>3</sup>Electrical Engineering & Computer Science, Vanderbilt University, Nashville, TN

<sup>4</sup>Laboratory of Behavioral Neuroscience, National Institute on Aging, National Institutes of Health, Baltimore, MD

<sup>5</sup>Department of Radiology and Radiological Sciences, Vanderbilt University Medical Center, Nashville, TN

## Abstract

Diffusion magnetic resonance images typically suffer from spatial distortions due to susceptibility induced off-resonance fields, which may affect the geometric fidelity of the reconstructed volume and cause mismatches with anatomical images. State-of-the art susceptibility correction (for example, FSL's TOPUP algorithm) typically requires data acquired twice with reverse phase