
A
TM 0.5
FG5
FG8
FG3
FG4
FG7
FG1
FG2
FG6
fes reductase ccg
dna glycosylase muty
dna glycosylase nth
fbic fo synthase
ferredoxin 2fe2s
aconitase
ipm isomerase leuc
fadh dienoyl coa
glutamate synthase
aldehyde fd oxidoreductase
dihydroxyacid dehydratase ilvd
ferredoxin 4fe4s
ferredoxin reductase fprb
cobg precorrin
nitrite sulfite reductase
cdgsh neet
nadh dehydrogenase
phnj cp lyase
suf biogenesis
isph lytb
quinolinate synthase nada
fhuf siderophore reductase
radical sam miab
ispg hmbpp synthase
thic
radical sam lipa
radical sam rlmn
radical sam biob
radical sam
radical sam hemn
radical sam moaa
radical sam pqqe
sufu assembly
whib
rieske qcra
nifu assembly
serine dehydratase
sdh frd
duf779 fes candidate
sera dehydrogenase
rieske nitrite reductase
rieske steroid oxygenase
molybdoenzyme
rieske
TM-score
1.0
0.8
0.6
0.4
0.2
0.0
B
all 44 families pLDDT ≥ 76
pTM 0.80
cdgsh neet
ferredoxin 2fe2s
nifu assembly
pTM
Mean per-model pLDDT
C
Wilcoxon signed-rank, p < 0.001
19 families in 8 fold groups
Mean TM-score to other families
between fold groups
within fold group
FG1 (n=5)
FG2 (n=2)
FG3 (n=2)
FG4 (n=2)
FG5 (n=2)
FG6 (n=2)
FG7 (n=2)
FG8 (n=2)