


Review

Experimental and in silico approaches to study CES substrate selectivity

Sergio R. Ribone ¹* and Mario A. Quevedo ¹

¹ Universidad Nacional de Córdoba. Facultad de Ciencias Químicas. Departamento de Ciencias Farmacéuticas. Consejo Nacional de Investigaciones Científicas y Técnicas (CONICET), Unidad de Investigación y Desarrollo en Tecnología Farmacéutica (UNITEFA), Córdoba (X5000HUA), Argentina.
* Correspondence: ; Tel.: (optional; include country code; if there are multiple corresponding authors, add author initials) +xx-xxxx-xxx-xxxx (F.L.)

Esta muy bien el abstract, solo lo siento un poco cortados en oraciones breves al inicio, con lo que hilaria un poco mas la conexion de algunas ideas.

Abstract

hCES

Human carboxylesterases (CES) are enzymes that play a central role in the metabolism and biotransformation of diverse substances. The two most relevant isoforms, CES1 and CES2, catalyze the hydrolysis of numerous approved drugs and prodrugs. Understanding CES isoform substrate specificity is crucial for multiple research areas. Various experimental and computational methodologies have been developed to quantify CES kinetic parameters (k_{cat} and K_M) for different substrates. This review will focus in the recent advancements in these methodologies to study substrate selectivity between CES1 and CES2. Experimental measurements commonly use recombinant CES or human tissue microsomes. Chromatographic methods, such as HPLC and LC/MS, are widely employed because they allow separation of substrates from hydrolysis products. Computational approaches are typically divided into two categories: molecular docking, molecular dynamic simulation and free-energy of binding to study CES–substrate affinity, and hybrid QM/MM simulations to evaluate catalytic turnover rates. Both approaches have demonstrated high accuracy and often agree with experimental results. The advantages of integrating experimental and computational techniques are evident in several studies that elucidates the structural and mechanistic basis of CES substrate selectivity. Given the biological relevance of CES-mediated catalysis, this review aims to provide a concise resource for continued exploration of CES isoform specificity and its implications for drug and prodrug design.

Que tal si explicitamos?: la prediccion de metabolismo de nuevas drogas y el potencial de desarrollo de inhibidores¹

Keywords: Carboxylesterases; CES1; CES2; substrate; selectivity; HPLC; LC/MS; Docking; MD simulation; QM/MM

Received:
Revised:
Accepted:
Published:

Citation: Ribone, S.R., Quevedo, M.A. Experimental and in silico approaches to study CES substrate selectivity. *J. Xenobiot.* **2025**, *1*, 0. <https://doi.org/>
Copyright: © 2025 by the authors. Submitted to *J. Xenobiot.* for possible open access publication under the terms and conditions of the Creative Commons Attribution (CC BY) license (<https://creativecommons.org/licenses/by/4.0/>).

1. Introduction

Human Carboxylesterases (CES, EC 3.1.1.1) are serine hydrolase enzymes responsible for the metabolism and biotransformation of diverse endogenous and exogenous substances containing esters, thioesters, amides, carbonates and carbamates moieties in their structures[1–3]. Based on amino acid sequence homology, human CES are classified into five isoforms (CES1–CES5). Among these, CES1 and CES2 are the most clinically relevant isoforms[4–6].
CES1 and CES2 share 47% of protein sequence identity but exhibit distinct substrate specificities and tissue distributions. Specifically, CES1 is primarily expressed in the liver, whereas CES2 is predominantly found in the intestines[1,3,4,7]. It is widely accepted in the literature that the substrate binding and hydrolysis selectivity of each isoform is primarily determined by the size of the acyl and alkyl moieties of the respective substrate

molecular structures[4,6–8], ^{with} ~~In this context~~, CES1 preferentially catalyzes ^{ing} the hydrolysis of substrates with smaller alkyl than acyl groups, as seen with drugs and prodrugs like clopidogrel, oseltamivir and meperidine[2,8,9]. Conversely, CES2 tends to hydrolyze compounds with ^{larger alkyl than acyl groups} ~~smaller acyl than alkyl groups~~, examples of which include halperidol, procaine and flutamide[1,3,8].

Conectaría estas dos ideas porque me parece que son parte del mismo objetivo de discusión

Despite this general trend, several exceptions to substrate selectivity have been reported. Notably, drugs and prodrugs like irinotecan, propanil, oxybutinin and procaine have been shown to be metabolized by both CES isoforms with similar efficiency [8,10]. Another noteworthy example is the differential CES-mediated biotransformation rates of pyrethroid derivatives, where *cis* and *trans* isomers of permethrin exhibit distinct hydrolysis pattern by CES1 and CES2, despite having identical acyl and alkyl group sizes[11].

Me parece que hay que expandir un poquito esta sección, con algunos ejemplos más y detallando si es que los ácidos y alquiles son considerados "grandes" o "chicos" y cuál es la CES que la hidroliza. Tiene que quedar la sensación de que la "general rule" no alcanza para pensar en drug design que contemple algún tipo de predicción del metabolismo.

Given the critical role of CES-mediated catalysis in various physiological processes, understanding the structural determinants of substrate specificity among CES isoforms is crucial for several areas of research. One such area is the design of prodrugs with optimized biopharmaceutical profiles. This approach aims to avoid undesirable early biotransformation of the prodrug or to enable site-specific bioactivation targeting a particular CES isoform[2,12]. In addition, the development of specific inhibitors for CES1 and CES2 offers promising therapeutic potential ^{to} ~~for~~ managing metabolic diseases. For instance, the CES1 inhibitor GR148672X has been explored as a candidate for treating hypertriglyceridemia, obesity and atherosclerosis[13].

Le sumaría "la importancia de considerar interacciones entre fármacos respecto del metabolismo por CES. Hay un ejemplo de un ensayo clínico (NCT04603391) que hacen con ese objetivo."

^{In} ~~Over~~ recent years, the selective imaging of *in vivo* enzyme activity has emerged as a powerful method for the study of biological systems, due to its ability to provide real-time, noninvasive monitoring within living organisms[5,14–17]. In this context, the design of fluorescent probes that specifically target CES1 and CES2 offers a promising strategy for visualizing their activity in complex biological systems.

Hay varios ensayos clínicos ya terminados o reclutando que son interesantes para discutir respecto de la relevancia de caracterizar el perfil de sustrato CES:

- NCT03051282
- NCT01708369
- NCT02192359: Administran células madre que sobreexpresan CES para activar irinotecan y tumores.
- NCT02147535: estudian el gene coding expression de CES1 y su consecuencia sobre el metilfenidato (incluir este sustrato que está siendo muy estudiado).

Enzymology has long been a foundational field for studying enzyme structure and function, as well as advancing our understanding of biological phenomena such as intermediary metabolism, molecular biology, and cellular signaling and regulation[18]. Early enzymology focused primarily on experimental techniques aimed at analyzing the catalytic properties and molecular specificity of enzymes[18]. The first one studies the thermodynamic and kinetic of the enzymatic reaction, measuring free-energy of reaction and activation. The second studies the specificity of different molecules for the enzymes, not only limited to the substrates, but also any other molecule that satisfy the specificity criteria for the enzyme, as is the case for potential inhibitors[18].

Due to the complexity of enzymes and the challenges associated with studying biomolecular reactions, many questions and mechanisms remain unclear. Computational enzymology, defined as the study of enzymes and their reaction mechanisms by molecular modeling and simulation [19], has the unique potential to investigate the dynamic behavior and reactions of biomolecules at atomic resolution. This approach can address unresolved issues by complementing and interpreting findings from experimental enzymology[19–21].

^{Starting} ~~Since the beginning~~ in 1976, with the pioneering work of Warshel and Levitt (Nobel laureates in 2013), computational enzymology has rapidly evolved over the past two decades. This progress has been driven by close collaboration between experimental and computational enzymologists, enhancing our ability to explain and interpret experimental data[19–21].

Building on the symbiotic relationship between experimental and computational enzymology, this review will focus on the recent advancements in methodologies devel-

oped to study substrate selectivity between the two most important isoforms of human carboxylesterase: CES1 and CES2. Usualmente los reviews hacen una breve descripcion de las secciones vienen antes de empezar, explicitando su importancia

2. Part I: Experimental enzymology

2.1. Kinetic parameters

Reescribir con mas claridad

During the study of the enzymatic properties of different substrates, there are important kinetic parameters determine by the Michaelis-Menten fundamental equation of enzyme kinetics[18,22]. The first parameter is the **dissociation constant**, also known as Michaelis constant (K_M), which reflects the affinity between the enzyme and the corresponding substrate. A lower ~~the~~ K_M value indicates stronger binding within the enzyme-substrate complex [2,18,23,24]. The second key kinetic parameter is the **turnover number**, also referred as catalytic constant (k_{cat}), ~~This value~~ ^{determined by the} ~~represents the number of catalytic cycles the enzyme can complete per unit of time when it is fully saturated with substrate. A higher k_{cat} value signifies greater substrate turnover and more efficient metabolism by the enzyme [2,18,23,24].~~ ^{catalytic turnover ?}

The ratio between these two kinetic parameters (k_{cat} / K_M) is known as **specificity constant**, as it reflects the ability of an enzyme to discriminate between ~~structurally similar~~ ^{different} substrates. A higher k_{cat} / K_M ratio indicates that the enzyme shows high affinity (low K_M value) and high catalytic rate for the substrate. Because of this, the specificity constant is often used as a measure of **catalytic efficiency**, indicating if the ligand is a good or poor substrate for the enzyme[18,24].

To obtain these kinetic parameters to quantify the substrate selectivity for a specific enzyme, experimental enzymology is used to ~~measure the progress of an enzyme-catalyzed reaction. Like any other chemical reaction, the progress can be monitored either by measuring the formation of the product, or the consumption of the substrate. Reliable detection methods for product formation or substrate depletion are essential for a successful enzyme assay[18].~~ ^{characterize} ~~the progress~~ ^{enzyme mediated substrate hydrolysis...} Capaz aqui introduciria la Table 1 como compendio general que luego se van detallando

Esto seria un ejemplo del resumen de secciones que pondria arriba

2.2. Enzyme sources

This section outlines the different CES enzymatic sources used in various analytical methodologies for calculating kinetic parameters. The two primary sources are *ex vivo* tissues, including purified human tissues and human microsomes, ~~and pure recombinant enzymes (Table 1).~~ ^{as well as}

Table 1. Enzymatic sources and analytical methods used for the exploration of CES kinetic parameters on different substrates.

Substrates	Ex vivo tissue		Recombinant enzyme	Analytical method	Reference
	Human tissue	Human miocrosomes			
Irinotecan	Liver	-	-	HPLC-FL	[25,26]
Methylphenidate	Liver	-	CES1/CES2	LC/MS	[27]
Oseltamivir	-	HLM/HIM	-	HPLC-UV	[8,28]
Temocapril	-	HLM/HIM	CES1/CES2	HPLC-UV	[8,29]
Aspirin	-	HLM/HIM	CES1/CES2	HPLC-UV	[29,30]
Clopidogrel	-	HLM/HIM	CES1/CES2	HPLC-UV	[8,30]
Flutamide	-	HLM/HIM	CES1/CES2	HPLC-UV	[31,32]
Pyrethroids	-	-	CES1/CES2	FL	[23,33]
Fluorescein diacetate	-	HLM/HIM	CES1/CES2	FL	[34]
Plasugrel	-	-	CES1/CES2	LC/MS	[35]
Heroin	-	-	CES1/CES2	HPLC-UV	[36]
Cocaine	-	-	CES1/CES2	HPLC-UV	[36,37]
Oxybutynin	-	HLM	CES1/CES2	LC/MS	[38]
Prilocaine	-	HLM	CES1/CES2	HPLC-UV	[39]
Lidocaine	-	HLM	CES1/CES2	HPLC-UV	[39]
Clofibrate	-	-	CES1/CES2	HPLC-UV	[8]
Fenofibrate	-	HLM/HIM	CES1/CES2	HPLC-UV	[8,40]
Imidapril	-	-	CES1/CES2	HPLC-UV	[8]
Enalapril	-	HLM	CES1/CES2	LC/MS	[41]
Sacubitril	Liver	-	CES1/CES2	LC/MS	[42]
Anordrin	-	HLM/HIM	CES1/CES2	LC/MS	[43]

HLM: Human liver microsomes. HIM: human intestine microsomes. FL: Fluorescence

2.2.1. Ex vivo tissue

Since the early 2000’s, the enzymatic studies of CES1 and CES2 substrate hydrolysis have relied on purified isoforms sourced from human liver. In several studies, human liver tissue is processed by homogenization and centrifugation, followed by separation of CES1 and CES2 isoforms using various chromatographic columns[25–27]. Using this method, it was determined that the produg irinotecan, along with other metabolites, is primarily bioactivated by human CES2 isoform [25,26].

The second source of CES is human tissue microsomes, which are small vesicles derived from fragmented cell membranes, mainly endoplasmic reticulum. These human microsomes can be obtained by differential centrifugation of the corresponding tissue or purchased from different biological supply companies. Human liver microsomes (HLM) are commonly used as enzyme source for measuring metabolic stability, as they contain key metabolizing enzymes, including CES. As mentioned in the introduction, CES1 is predominantly expressed in the liver, making HLM a valuable source for studies of CES1 substrate selectivity. Following similar tissue distribution patterns human intestines microsomes (HIM) have been used as a source for CES2 in selectivity assays. Using this approach, studies have shown that the antiviral produgs oseltamivir and temocapril are preferentially activated by CES1 (HLM) over CES2 (HIM)[28,29].

2.2.2. Recombinant enzyme

Shortly after the use of human tissue as enzyme source, it became evident that a more purified version of both CES isoforms were necessary to performed accurate selectivity enzymatic experiments with different substrates. Morton and Potter developed a method for cloning and expressing CES using baculovirus to infect *Spodoptera frugiperda* insect cells [44]. This technique enabled the production of recombinant CES1 and CES2, which were

Usualmente esta bueno
incluir una minima
descripcion del hallazgo
mas alla de lo que hicieron

then used to measure the ~~enzymatic properties~~ ^{no son "enzymatic properties"} of several substrates (Table 1). The hydrolysis of pyrethroids by human CES1 and CES2 have been studied using this recombinant enzyme source [23,33]. Additionally, the hydrolysis specificity of CES1 and CES2 for drugs of abuse, such as heroin and cocaine [36], as well as other drugs and prodrugs, have also been studied using recombinant enzymes (Table 1) [8,38,39].

As recombinant CES enzymes became commercially available from multiple suppliers, research groups were able to continue exploring the metabolism selectivity of CES isoforms across different substrates. For example, studies on the angiotensin-converting enzyme inhibitors enalapril and ramipril showed that both drugs ~~were~~ ^{are} selectively hydrolyzed by CES1 [41]. Recombinant CES enzymes have also been used to determine the predominant role of CES isoforms in the human bioactivation of prodrugs such as sacubitril [42] and anordrin [43]. Additionally, recombinant CES enzyme have facilitated pre-clinical evaluations of the bioactivation rates for structurally designed prodrugs, including atorvastatin [45,46], indomethacin [47,48] and haloperidol [49].

2.3. Reported analytical methods

As mentioned earlier, obtaining accurate kinetic parameters requires a reliable analytical method to quantify the progress of the enzyme-catalyzed reaction. In this section, the different analytical methods used for the determination of the described kinetic parameters for different substrates and both CES isoforms will be described.

2.3.1. Spectrophotometry

Spectrophotometric methods have been widely used to quantitatively determine total CES activity, by measuring the absorbance of *p*-nitrophenol, produced through the hydrolysis of *p*-nitrophenyl acetate (pNPA), at 405 nm (Figure 1) [36,50,51]. This approach was applied to the functional characterization of recombinant human CES expressed in *E. coli* as an alternative method for obtaining the human enzyme [51].




Figure 1. Hydrolysis of *p*-nitrophenyl acetate (pNPA) by CES to produced *p*-nitrophenol.

Esta info en el cuadro me
parece muy bien y cuando se
cita algo estaria bueno un par
de lineas con este tipo de main
findings

The absorbance properties of *p*-nitrophenol ~~have~~ ^{has} also been used to evaluate the kinetic parameters of various *p*-nitrophenyl esters derivatives with CES1 and CES2 [36]. The authors observed a correlation between the affinity constant (K_M) and the calculated water/octanol partition coefficients (clogP) values, concluding that the affinity of the substrates for both CES isoforms is directly related to their lipophilicity properties [36]. In addition, kinetic data for naphthyl esters derivatives were obtained by measuring the formation of naphthol at 230nm [50].

This method is simple and rapid, but it has several disadvantages. One key issue is the potential ~~overlapping~~ interference between substrates and products, which complicates experiments in complex biological systems. Additionally, the method requires ~~a larger~~ ^{higher amounts..} amount of enzyme, as it is performed in a UV cuvette with a total volume of 1 ml [51].

•El metodo de cuantificacion por UV, no se puede aplicar a well plates tambien?

Habria que describir en que consiste el metodo.

2.3.2. Fluorescence

In contrast to the previously described analytical method, the fluorescent probe-based approach is not only simple but also highly selective and sensitive. These probes quantify CES activity by detecting changes in fluorescence intensity. Known as "off-on" fluorescent probes, they initially exhibited little or no fluorescence, but the hydrolyzed products release strong fluorescence in presence of CES.

This methodology has been used to determine the kinetic parameters of pyrethroids-like substrates containing 6-methoxy-2-naphthaldehyde. The fluorescence of this moiety is measured with an excitation wavelength of 330 nm and an emission wavelength of 465 nm[23]. In this study, it was observed that the stereoisomeric centers of these derivatives presented a differential impact on hydrolysis by the two CES isoforms. Specifically, the presence of an (*R*)-enantiomer carbon adjacent to the ester carbonyl carbon resulted in a greater preference for CES2 hydrolysis than CES1[23]. These findings highlight the importance of the three-dimensional disposition of the substrate groups within the catalytic site of the enzyme for CES hydrolysis selectivity.

In another study, fluorescein diacetate was used as a substrate to assess CES1 and CES2 selectivity. The hydrolysis of fluorescein diacetate by the CES enzymes releases fluorescein, which ~~was quantify~~ ^{is quantified} at an excitation wavelength of 483 nm and emission at 525 nm. The results indicated that fluorescein diacetate is a selective CES2 substrate (Figure 2)[34].

De nuevo remarcar que para cada ejemplo de aplicacion citado se deberia avanzar en alguna explicacion del "finding" como se hace aca.





Figure 2. hydrolysis of fluorescent probe substrates. (a) Fluorescein diacetate. (b) BODIPY ester.

The high fluorescence quantum yield and photochemical stability of BODIPY dyes make them excellent candidates for this analytical methodology. As a result, a BODIPY ester was designed as a specific substrate for CES1. The acid product formed after ~~CES~~ ^{CES1} hydrolysis was used to measured the kinetic parameters at an excitation wavelength of 505 nm and emission at 560 nm (Figure 2)[52].

Tener en cuenta que la cromatografía como tal es una técnica de separación, a la cual se le puede acoplar tanto UV como fluorescencia, con lo cual hay que buscarle el sentido aquí.

2.3.3. Chromatography

Diverse chromatographic methods have been widely used to determine the kinetic parameters of numerous substrates (e.g., drugs, prodrugs, ~~pyrethroids~~ ^{its}) due to their high sensitivity, accuracy, rapid analysis and reproducibility. One additional advantage is the ability to separate substrate from products for precise quantification. The two most commonly used techniques are the high performance liquid chromatography (HPLC) and liquid chromatography-tandem mass spectrometry (LC/MS).

In the case of HPLC, an ~~additional~~ ^{no sería "additional", por la LC no es de detección} detection technique is required to quantify the different ligands after chromatographic separation, with the two previously described methodologies being the most frequently used. The spectrophotometric methodology usually ~~implement~~ ^{couple} an UV-detector (HPLC-UV), as most ligands exhibit absorbance in the ultraviolet spectrum. This methodology has been widely used to study the CES kinetic parameters of various therapeutic drugs and prodrugs. ^{cit}

To investigate substrate specificity among CES isoforms, a study was conducted on 13 compounds, including clopidogrel, clofibrate, oseltamivir, mycophenolate mofetil, procaine and temocapril, among others (Table 1), using HPLC with specific conditions for each metabolite (e.g., mobile phase, column and UV wavelength)[8]. This research group also applied the HPLC-UV method to ~~analyzed~~ ^{study} the kinetics of other drugs, such as flutamide[31,32], prilocaine and lidocaine[39]. Based on the structural characteristics of these compounds, the authors proposed the already discussed general substrate selectivity pattern for CES: CES1 preferentially hydrolyzes ligands with smaller alkyl than acyl moiety (e.g., clofibrate, lidocaine, temocapril), while CES2 favors those with larger alkyl than acyl groups (e.g., flutamide, procaine).

The metabolism of the abuse drugs cocaine and heroine by CES1 and CES2 was also studied using this methodology[36]. For cocaine, hydrolysis was monitored by quantifying the formation of benzoylecgonine and benzoic acid at 235nm. After incubation with both CES isoforms, the results showed that cocaine was exclusively metabolized by CES2, producing benzoic acid and ecgonine methyl ester. The second potential metabolic pathway, forming benzoylecgonine and methanol, was not detected[36]. ^{Si fue detectada con CES1?}

For heroine, hydrolysis was assessed by monitoring the formation of 6-acetylmorphine, also at 235nm. Both CES isoforms were found to hydrolyze ~~heroin~~ ^{heroin}, with CES1 exhibiting a higher catalytic efficiency (k_{cat}/K_M) compared to CES2[36]. This result represents an exception to the general CES1 substrate specificity, as heroin has a larger size alkyl group than acyl moiety in its structure.

A different approach was presented in several studies by Takahashi et al., in which all the investigated substrates were structurally diverse indomethacin-derived prodrugs. The formation of indomethacin, monitored at 254 nm, was used as the analytical marker to determine the kinetic parameters[47,48,53]. By synthesizing prodrugs bearing a variety of alkyl moieties, the authors were able to investigate how different structural features influence the hydrolysis behavior of both CES isoforms. Specifically, they examined: 1) the effect of the steric hindrance on the carbon adjacent to the carbonyl group, 2) the influence of electron density around the carbonyl group and 3) the chiral recognition ability between CES1 and CES2. ^{Lo juntaría porque la idea es enunciar el uso de la técnica y una breve descripción de los resultados}

Overall, the results indicated that the indomethacin prodrugs were mainly hydrolyzed by CES1, because the drug structure represents the acyl group, which in all the cases was larger than the corresponding alkyl group. However, prodrugs with aryl-containing alkyl moieties exhibited reduced or even no selectivity between CES isoforms, demonstrating that steric hindrance near the ester carbonyl carbon plays a crucial role in determining metabolic selectivity.

Como sentido a incluir la cromatografía, me parece que estaría bueno centrar en este punto de la LC/MS que efectivamente tiene más sensibilidad y especificidad que los otros métodos de detección. Entonces quizás esta sección la cambiaría a LC-MS/MS

Porque sería un "different approach"? Estamos discutiendo la cromatografía más que la estrategia de diseño de análogos.

This research group also conducted similar studies by synthesizing prodrugs derivatives of haloperidol[49] and atorvastatin [46], using the absorbance properties of these drugs as detection methods for the HPLC technique.

On the other hand, LC/MS does not require an additional method for the quantification of the desired kinetic parameters, as both substrate and product are detected directly by the mass spectrometer following their liquid chromatographic separation. This analytical technique has primarily been used to investigate the metabolic pathway of a diverse set of prodrugs.

Interesting results were observed in the hydrolysis study of capecitabine, a carbamate prodrug of 5-fluorouracil used in the treatment of colorectal cancer[54]. The study showed that capecitabine is hydrolyzed equally by both CES1 and CES2, suggesting that substrates containing carbamate groups can be metabolized by either isoform, independent of the size of their acyl or alkyl moieties. Other prodrugs studied include plasugrel, which exhibited selectivity for CES2[35], sacubitril, selectively activated by CES1[42], and anordrin, which showed similar hydrolysis parameters with both CES isoforms[43].

In a more recent study, the hydrolysis characteristics of cocaine by CES1 were re-examined using LC/MS methodology[37]. This study revealed that cocaine is hydrolyzed by CES1, producing benzoylecgonine and methanol, which contrast with earlier findings using HPLC-UV where this hydrolysis product was not detected (Figure 3)[36]. These results suggest that LC/MS provides a more accurate analytical approach for studying cocaine hydrolysis compared to HPLC-UV.

reescríbelo en inglés




Figure 3. Different obtained cocaine metabolic pathway from CES1 and CES2 hydrolysis experimental studies with HPLC-UV and LC/MS analytical methodologies.

This section highlights the importance of experimental enzymology in studying CES substrate selectivity. The gathered information shows that, despite the general structural rules established for substrate selectivity between CES1 and CES2, several exceptions and underlying structural properties remain unclear. Given that CES binding and hydrolysis involve subtle intermolecular interactions, computational enzymology emerges as a crucial tool for analyzing CES substrate selectivity at the atomistic level.

A partir de aca me parece que ya lo pondria en la seccion de "Computational enzymology", ya que al fin y al cabo la bases de datos son la fuente de datasets para el modelado teorico.

2.4. Experimental CES kinetic parameter databases

In the era of bioinformatics, providing reliable kinetic parameter data and effective data management systems is essential to support researchers in retrieving enzymatic information from the vast amount of biological data published annually, such as those related to CES enzymes. In this context, online databases have become invaluable tools for granting researchers access to this information. This section will focus on two of the the most well-known and widely used online databases ~~containing~~ ^{are} for obtaining human CES enzymatic parameters for different substrates.

2.4.1. BRENDA database

The BRAunschweig ENzyme Database (BRENDA) is the oldest collection of enzyme-related data ^{compiled} from the scientific literature, established in 1987 at the German National Research Centre for Biotechnology in Braunschweig [55]. Today, the BRENDA website (www.brenda-enzymes.org/) is accessed by over 100,000 users each month. The site is very intuitive, allowing users to search by entering a text, such as enzyme name, ligand name, EC class, inhibitors, etc., or through structured-based queries by drawing substrate/products or ligand substructures[55].

The search for CES (EC 3.1.1.1) ^{enzymatic} information on BRENDA, yielded 2079 substrates/products and 67 natural ^{substances} ~~products from the database~~. ^{Among this data,...} From these ligands, 849 K_M values, 550 turnover numbers (k_{cat}) and 232 k_{cat} / K_M values (catalytic efficiency) can be retrieved from diverse bibliographic sources. In addition, the reaction diagrams and references associated with these substrates ^{are} ~~were~~ provided.

However, two main disadvantage were noted: 1) it is not possible to filtrate information based on the organism origin of the enzyme, and 2) the kinetic information can not easily be separate by CES isoforms. Despite this limitations, BRENDA remains a valuable starting point for searchif kinetic parameters related to CES1 and CES2 substrate hydrolysis.

2.4.2. SABIO-RK database

SABIO-RK is a manually curated database containing enzymatic biochemical reactions and their kinetic parameters (sabiork.h-its.org/). ~~The authors have established that the database is a valuable resource for both experimental and computational enzymology researchers[56].~~ ^{constituting a ...} Data in SABIO-RK are primarily extracted manually from the literature and stored in a structured and standardized format. The database includes essential data to describe the characteristics of biochemical reactions, the corresponding biological source, kinetic properties and experimental conditions[56].

The regular search for CES name in SABIO-RK website returned 519 entries, which is a ^{less} ~~smaller~~ amount of information compared to BRENDA. The SABIO-RK website offers an advance search feature that allows filtering by various conditions, such as "organisms: *Homo sapiens*", which narrows the results to 168 entries related to human CES data. Additionally, it is possible to filter the results to include only data from recombinant CES enzyme (82 entries). However, like BRENDA database, SABIO-RK also has the limitation of not separating kinetic parameters by CES isoforms. In conclusion, while SABIO-RK contains less information than BRENDA, it offers better organization and more intuitive filtering options, making it easier to retrieve relevant data.

3. Part II: Computational enzymology

3.1. Enzyme sources: structures

At the beginning of any computational molecular modeling substrate-enzyme complex study, the information regarding the structure of both, the enzyme and the substrate, is ~~needed~~. ^{needed} The structures of the ligands are relatively easy retrieved from different biological

No me convece el titulo, sobre todo porque queda como repetitivo con la seccion experimental. Talvez utilizaria algo del estilo: "Availability of CES1 and CES2 structural templates" o algo del estilo.

En realidad no es "necesaria", sino que es en algun punto "deseada". Mas que nada porque entiendo que luego incluiras algo respecto a prediccion de CES2 y docking molecular?

La estucutura de los sustratos no es critica de estar disponible segun mi opinion.

databases, such as those mentioned in the previous section. On the other hand, obtaining the structures of the macromolecules (enzymes) can be more challenging. This is because enzyme structures are often distributed across different sources and databases. In this section, the ~~reported sources~~^{availability of} for obtaining the three-dimensional structures of CES1 and CES2 enzymes will be ~~displayed~~^{discussed}.

3.1.1. CES1 crystal structures

Since 2003, different crystallographic structures of CES1 have been reported, as summarized in Table 2. The first published structures involved CES1 complexes with ligands from several categories, including metabolites of abuse drugs like cocaine and heroin (homatropine and naloxone methiodide, respectively)[57], therapeutic drugs (tacrine, tamoxifen and mevastatin)[58,59], and endogenous substrates (cholate, taurocholate and coenzyme A) [60] (Table 2). Notably, the crystal structure of CES1-tamoxifen complex revealed that tacrine interacts in the catalytic binding site of CES1 in four different binding modes[58]. These findings highlights the promiscuous nature of CES1, suggesting that its ability to hydrolyzed a variety of substrates is driven by its capacity to interact with these ligands in multiple conformations[58].

Table 2. Information related to CES1 reported crystallographic structures.

Substrates	Classification	Resolution (Å)	PDB code	Reference
Homatropine	M.L.	2.80	1MX5	[57]
Naloxone methiodide	M.L.	2.90	1MX9	[57]
Tacrine	Drug	2.40	1MX1	[58]
Tamoxifen	Drug	3.20	1YA4	[59]
Mevastatin	Drug	3.00	1YA8	[59]
Ethylacetate	M.L.	3.00	1YAH	[59]
Benzil	M.L.	3.20	1YAJ	[59]
Cholate/Palmitate	E.S.	3.00	2DQY	[60]
CoenzymeA	E.S.	2.00	2H7C	[60]
CoenzymeA/Palmitate	E.S.	2.80	2DQZ	[60]
Taurocholate	E.S.	3.20	2DR0	[60]
Soman	N.A.	2.70	2HRQ	[61]
Tabun	N.A.	2.70	2HRR	[61]
Cyclosarin	N.A.	3.10	3K9B	[62]
-	-	2.20	4AB1	[63]
-	-	1.86	5A7F	[64]
-	-	2.67	8EOR	[65]
F-3	C.I.	1.83	9KWL	[66]
F-4	C.I.	1.89	9KWM	[66]

M.L.: Metabolite ligand. E.S.: Endogenous substrate. N.A.: Nerve agent. C.I.: covalent inhibitor.

Different types of covalent ligands have also been studied through crystallographic structures. The first group to be investigated included the organophosphorus nerve agents soman, tabun and cyclosarin [61,62]. More recently, attention shifted to a second family focused on the covalent binding mechanism of serine-selective electrophilic warheads. Two crystal structures were ^{obtained} generated showing CES1 covalently bound to 2,2,2-trifluoroacetophenone derivatives at the catalytic serine (Table 2)[66]. Additionally, there are crystallographic structures of CES1 in the absence of substrates[63–65]. One of these structures exhibited the highest resolution of any reported CES1 crystal structure to date (1.86 Å).

For conducting molecular modeling studies focused on substrate selectivity between CES isoforms, the optimal approach is to use the CES1 structure with the highest possible resolution and complexed with non-covalent ligands or in a substrate-free state. On the

Poner entre parentesis el
codigo PDB.

other hand, it is not advisable to use the crystal structures of CES1 bound to covalent ligands. This is because the catalytic residues, along with other residues in the catalytic binding site, are in conformations suitable for binding covalent inhibitors, rather than for accommodating non-covalent substrates.

Lo mismo que antes con el titulo: dejar la explicacion de los Hom. Mod. para el parrafo.

3.1.2. CES2 homology modeling structures

The crystallographic structure of CES2 remains unresolved to this day. As a result, homology modeling is necessary to obtain the three-dimensional structure of this CES isoform in order to model the binding mode of substrates in the catalytic site of the enzyme. Different research publications have reported distinct strategies for generating the homology model structure of CES2. One of the most common methodologies involves using tools provided by the Swiss Institute of Bioinformatics server (www.expasy.org/). In the early publications, the process of generating the CES2 model followed a two-steps procedure: first, the amino acid sequence of human CES2 was retrieved from the Swiss-Prot database, and then submitted to the Swiss-Model[67] server for fully automated protein structure homology modeling[68–70]. Over time, this methodology became more automated, allowing users to directly download pre-existing CES2 models created by other researchers via the Swiss-Model server[24,71,72].

The second approach involves generating the CES2 structure using homology modeling software. Several studies have reported the use of the open-source *Modeler* software[73] to generate the homology model of CES2 and perform subsequent molecular modeling studies on this isoform[74,75].

podriamos referir a AlphaFold en general, y destacar AlphaFold3 como herramienta reciente

In recent years, *AlphaFold2*, a powerful neural network-based model methodology has been developed for predicting the three-dimensional structures of proteins[76]. Since its beginning in 2021, this machine learning approach has generated more than 200 million protein structures, all freely available for download from the AlphaFold database (alphafold.ebi.ac.uk). To the best of our knowledge, despite its accessibility and remarkable accuracy, an AlphaFold-predicted human CES2 structure has not yet been utilized for the molecular modeling of this enzyme with any substrate.

3.2. Molecular modeling methods

To analyze the results from experimental enzymology studies of the CES-substrate complexes, a series of molecular modeling methodologies can be employed. These computational techniques have been widely used to investigate, at an atomistic level, the kinetic parameters of different substrates for both CES isoforms. This section is organized into two subsections: 1) molecular modeling approaches (molecular docking, molecular dynamics and free-energy of interaction analysis) used to study the affinity constant (K_M) and 2) hybrid QM/MM simulations employed to examine the catalytic constant (k_{cat}).

3.2.1. Molecular modeling methods to analyze affinity constant - K_M

Typically, the study of the recognition and binding of CES substrates...

Every molecular modeling campaign begins with a molecular docking procedure, aimed to identify substrate conformations that yield the lowest-energy interactions with residues in the CES catalytic binding site. In the subsequent stage, the CES-substrate complex is subjected to molecular dynamics (MD) simulation to characterize its dynamic behavior and stability in an explicit aqueous environment and at physiological temperature. Finally, a free-energy interaction analysis is performed based on the MD simulation data obtained for the CES-substrate complex. A small subset of studies incorporate all three of these molecular modeling methods in the investigation of CES-substrate complexes. Although a few studies incorporate all three of these molecular modeling approaches when investigating CES-substrate interactions, most rely solely on molecular docking to identify the optimal substrate conformation within the CES catalytic binding site.

"may be biased to conformations favoring covalent inhibitors rather than modeling the intrinsic substrate recognition associated to the apo-protein"

Existe alguna explicacion de porque no se pudo obtener el cristal de CES2? Una busqueda rapida indica que es debido a los sitios de glicosilacion. Tambien veo que recientemente se reporto cristal de CES2 de raton: DOI:10.3390/ijms232113101 Estaria bueno decir algo al respecto.

AlphFold1 aparecion en 2018. Capaz convenga haver review de AlphFold mas que de AlphaFold3.

Siento que seria mas claro dividir los molecular modeling methods en "Modeling of substrate:enzyme recognition" y "Modeling of substrate:enzyme reactivity"

Estos trabajos son del 2010, y una búsqueda rápida muestra estudios anteriores de docking sobre CES. Mas bien enfocaría en los reviews o trabajos mas representativos y recientes de docking sobre CES.

Early...
The first molecular docking studies investigating the selectivity of CES1 and CES2 were conducted by the group of Vistolli et al. [68,77]. In these works, 40 known CES substrates were subjected to molecular docking protocols using two crystallographic structures of CES1 (PDB codes: 1MX9 and 1YAJ) and a homology modeling structure of CES2. Different scoring functions were calculated and correlated with the experimentally reported K_M values to develop predictive models of substrate affinity toward both CES isoforms. In both cases, the results revealed a strong correlation between K_M and the calculated lipophilic interaction scores, highlighting the central role of hydrophobic interactions, primarily attributable to the abundance of apolar residues within the catalytic binding site [68,77].

A combination of molecular docking and MD simulations have been employed to assist in the design of selective inhibitors of human CES2. These studies were focused on the design, synthesis and structure-activity relationship of glycyrrhetic acid and benzofuranone derivatives (Figure 4) [69,78]. In the first work, molecular docking was performed using the most active and selective glycyrrhetic acid derivative against both CES isoforms to elucidate its 1000-fold preference for CES2 over CES1 [69]. The results indicated that this derivative displayed a greater number of hydrogen bond interactions with residues in the CES2 catalytic binding site residues than in CES1. In addition, the conformation adopted by the inhibitor in CES1 complex was positioned farther from the catalytic serine residue than in the CES2 complex [69]. In the second study, a similar molecular docking analysis was performed for the most active and selective benzofuranone derivative against both isoforms. The results were consistent: the inhibitor exhibited more favorable interactions and a higher number binding site contacts with CES2 residues compared to CES1 [78]. Furthermore, MD simulation and free-energy decomposition analyses of the CES2-inhibitor complex revealed that the complex remain stable over 50 ns of simulation and that the hydrophobic interactions played a major role in the binding of the inhibitor to CES2 [78]. As shown in Figure 4, both selective CES2 inhibitors share the presence of a carboxylic acid group in their molecular structures.

Nunca de hablo de este residuo, lo que evidencia que es necesario en etapas tempranas revisar el mecanismos catalítico de las CES.

Dado que se destaca este punto, me pregunto que implica la presencia de grupos ácidos carboxílicos para la selectividad?




Figure 4. Molecular structures of CES2 selective inhibitors. (a) Glycyrrhetic acid derivative. (b) Benzofuranone derivative.

In a more recent study [24], molecular docking, MD simulation and free-energy interaction decomposition analyses were performed on two families of previously studied substrate families: five *p*-nitrophenyl ester derivatives [36] and two pyrethroid stereoisomers [23], using both CES1 and CES2. Consistent with earlier findings, hydrophobic interactions were found to contribute substantially to the CES-ligand free-energy of interaction, correlating well with experimentally determined affinity constants (K_M) [24]. An additional key ob-

En realidad me imagino que para anticipar si una molécula es buen o mal sustrato de CES, se necesita verificar si la afinidad es alta y original, una baja K_M , pero además que el modo de unión que se asocia a esa alta afinidad sea catalíticamente productiva, es decir que implique el posicionamiento del grupo a hidrolizar en las cercanías de la serina catalítica y ocupando el oxianion hole.

Esto es lo que pienso es más útil en la introducción para que quede claro el mecanismo catalítico.


servation was that the ^{higher} increase binding cavity volume of CES2, relative to CES1, enabled one pyrethroid stereoisomer to adopt a distinct interaction pattern and thereby maintain a higher affinity (lower K_M) for CES2 compared ^{compared to the other} with its corresponding stereoisomer. Overall, the substrates analyzed in this work formed the most favorable interaction patterns with residues in the respective CES catalytic binding sites, resulting in the highest attainable affinity for each isoform[24].

3.2.2. Hybrid QM/MM simulation to analyzed catalytic constant - k_{cat} Lo sacaría del título y explicaría en el texto


The study of enzyme-ligand catalytic constant through computational chemistry requires specialized methodologies to analyzed the free-energy of activation associated with the enzyme-catalyzed hydrolytic reaction. The most commonly employed approach for this purpose is hybrid QM/MM simulation. In this method, the ligand and the catalytic residues directly involved in the hydrolysis reaction are modeled using quantum mechanics (QM), while the remaining enzyme residues are described with a molecular mechanics (MM) force field.

The general hydrolysis mechanism of esterases, such as CES, is well established and involves two consecutive reaction steps. First, the acylation step, where the carbonyl carbon of the substrate is attacked by the hydroxyl moiety from the catalytic serine, leading to the formation of an acylated serine intermediate and the release of the alkyl group of the substrate. Second, the deacylation reaction, where the carbonyl carbon of the acylated serine is attacked by the oxygen of a water molecule, yielding the carboxylic acid portion of the substrate and regenerating the free serine residue, thereby allowing a new catalytic cycle to begin (Figure 5)[3,24,37].

1) Acylation



2) Deacylation



En el texto, donde se mencionan estas etapas, habría que citar la Figura.

Figure 5. Acylation and deacylation reaction pathways for CES1 and CES2 catalyzed hydrolysis of the ester ligands. *p*-nitrophenyl ester derivatives were used as example. Reactant state (RS), first transition state (TS1), intermediate (INT), second transition state (TS2) and product state (PS).

There ^{are} relatively few reported studies employing hybrid QM/MM simulation for substrates hydrolyzed by human CES, likely because of the greater complexity of this methodology compared with the molecular modeling methodologies described in the previous section. Given the importance of human CES in the metabolism of abuse drugs, several

works have investigated the hydrolysis mechanism of cocaine by CES1 and CES2. The first study reported the hydrolysis mechanism of cocaine by CES1 using hybrid QM/MM simulation alongside the experimental determination of the corresponding kinetic parameters (K_M and k_{cat}) [37]. The QM region was parameterized with the semi-empirical SCC-DFTB method, while the MM region with CHARMM27 force field. The reaction was modeled using an umbrella sampling approach, where the reaction coordinate (RC) was defined as a linear combination of two covalent bond distances: one forming and one breaking. The simulations indicated that CES1 catalyzes the hydrolysis of cocaine to benzoylecgonine and methanol (Figure 1) through a single-step acylation reaction followed by a single-step deacylation stage, each showing a distinct transition state (TS). The combined free-energy profile of both reaction steps revealed that the acylation TS is the rate-limiting reaction, with a free energy barrier of 20.1 kcal/mol. This result is in close agreement with the experimental free-energy barrier, derived from the catalytic constant, which was 21.5 kcal/mol, demonstrating the high accuracy of the hybrid QM/MM methodology employed in this study [37].

A second article reported a study similar to the previous one, exploring the hydrolysis of cocaine by CES2 to produce ecgonine methyl ester and benzoic acid (Figure 1). In this study, the QM region was parameterized using the semi-empirical PM6 method, and the MM area was modeled with ~~an Amber18~~ ^{the ff99SB} force field (ff99SB). An umbrella sampling approach was again employed to follow the two-steps hydrolysis reaction, in which two covalent bond distances, one bond forming and one bond breaking, were used as reaction coordinates. The resulting 2D-PMF profiles for the full catalytic cycle showed that each reaction stage proceeds through a tetrahedral intermediate structure and involves two transition states. Among these, TS₄ (associated with the formation of benzoic acid) was identified as the rate-limiting step of cocaine hydrolysis by the CES2, displaying an activation free-energy barrier of 19.5 kcal/mol. This lower energetic barrier respect to CES1 is consistent with previous experimental findings that reported a higher turnover number for CES2 than for CES1.

In the previous section, a study involving two families of substrates, *p*-nitrophenyl esters and pyrethroid stereoisomers, in complex with CES1 and CES2 was described. In this work, two *p*-nitrophenyl ester derivatives and two pyrethroid stereoisomers were subjected to hybrid QM/MM simulation to analyze their hydrolysis by both CES isoforms [24]. The QM/MM simulation protocol followed was similar to the earlier study of cocaine with CES1 [37]; the QM region was parameterized using the semi-empirical SCC-DFTB3 method, while the MM region was modeled with Amber20 force field (ff14SB). As in the previous example, the study employed an umbrella sampling approach; however, in this case, four covalent bond distances, two bonds forming and two bonds breaking, were combined linearly using a linear combination of distances (LCOD) to define the reaction coordinate (RC).

Across all modeled reactions, substrate hydrolysis by both CES isoforms proceeded through a ^{concerted?} single-step acylation stage followed by ~~a single-step~~ ^{another concerted} deacylation stage, each involving one transition state (TS). Analysis of the results showed that the rate-limiting step for each substrate corresponded to the TS associated with the highest molecular steric hindrance encountered during the course of the hydrolysis, acylation or deacylation stage [24]. The authors concluded that the CES selectivity is not solely determined by the molecular size of the alkyl or acyl groups of the substrates, but instead arises from a more complex scenario governed by the initial conformation of the ligand within the CES binding site [24].

Overall, all studies discussed in this section showed a good correlation between the experimental catalytic constant (k_{cat}) and the calculated free-energy of activation from the

rate-limiting TS reaction stage, demonstrating the reliability of the hybrid QM/MM methodology. This was observed despite the differences in the proposed hydrolysis mechanisms, which involved either two or four transition-state structures.

No existe la "Part II"


4. Part III: Combination of experimental and computational enzymology

This final section of the article is devoted to studies that combined experimental and computational methodologies to investigate the hydrolysis properties of various substrates by both CES isoforms.

In the first study, the hydrolysis of fenofibrate by CES1 and CES2 was examined to identify the primary enzyme responsible for its metabolism in humans[40]. The kinetic parameters were determined by measuring the formation rate of fenofibric acid (the hydrolytic metabolite of fenofibrate) through HPLC-UV in the presence of human microsoms and recombinant CES1 and CES2. These results exhibited that the affinity constant was slightly higher for CES1, but the major difference was observed in the catalytic constant, with CES1 exhibiting a substantially faster reaction rate than CES2[40].

The authors performed a molecular docking study of fenofibrate within the binding site of both CES isoforms. The docking results showed that fenofibrate adopted a more favorable binding pose and received a better docking score into the catalytic binding site of CES1 than CES2, consistent with the experimental affinity results. Hybrid QM/MM simulations were not performed in this publication[40]. This study provides useful insight into the factors underlying the pronounced differences in fenofibrate hydrolysis between the two CES isoforms.

The second study, reported the design and CES selectivity of a family of four fluorescent substrates derived from a naphthalimide scaffold [17]. Enzymatic assay was monitored by measuring the emission intensity of the hydrolysis product at 520 nm (excitation 450 nm) after incubation with CES1 and CES2. Only the two derivatives containing amide or carbamate moieties (NIC-1 and NIC-2, figure 6) underwent specific hydrolysis in the presence of CES2. In contrast, the other two carbamates in which the nitrogen was fully substituted with carbon atoms (NIC-3 and NIC-4, figure 6), exhibited inhibitory effects of both CES isoforms.



Este punto me parece muy interesante. Como interpretan los resultados los autores para convertir un buen sustrato en un buen inhibidor? Sospecho que los sustratos tienen buena K_M y buena k_{cat} , mientras que los inhibidores tienen buena K_M y mala k_{cat} ?. Entonces son inhibidores competitivos?

Figure 6. Structure of the fluorescent substrates derivatives of naphthalimide (NIC)[17].

Los sustratos per se son fluorescentes?

To investigate the shift in activity from CES substrate to inhibitor, the binding properties of NIC-4 toward CES1 were analyzed using molecular modeling methods with CES1 and compared with the known substrate *p*-nitrophenyl acetate (pNPA)[17]. Molecular docking followed by MD simulations, revealed that both NIC-4 and pNPA displayed simi-

Aca esto explica lo que me pregunté antes. Es un buen caso para destacar la importancia de comprender a nivel atomístico los eventos asociados al funcionamiento de las CES, tanto para la predicción de substratos como para el diseño de inhibidores.

lar interaction patterns with the residues of the CES1 binding site. In addition, a steered MD simulation was performed in which the structures and the binding site residues were parameterized with SCC-DFTB method, while the remaining part of the enzyme with Amber16 ff14SB force field. The reaction coordinate was defined as the distance between the catalytic serine oxygen and the carbonyl carbon of the substrate, representing the nucleophilic attack in the hydrolysis reaction. During the simulation of pNPA hydrolysis by CES1, the proton transfer from the catalytic serine to the nitrogen atom of the catalytic histidine occurred spontaneously, following the normal acylation mechanism with an estimated energetic barrier of 20 kcal/mol. In contrast, during the simulation of NIC-4 with CES1, this proton transfer did not occur, because the methyl group positioned near the potential nucleophilic center of NIC-4 (Figure 6) sterically blocked the transfer pathway, forcing the serine hydroxyl group to orient away from the histidine imidazole. These molecular modeling results enable the authors to explained the mechanism of CES inhibition induced by NIC-4[17].

5. Conclusions

The objective of this article was to highlight the importance of combining experimental and computational techniques to elucidate substrate selectivity between the two major human CES isoforms: CES1 and CES2. The first section showed that chromatographic methodologies (HPLC and LC/MS) are widely used by different research groups for the experimental determination of CES kinetic parameters (K_M and k_{cat}) for a variety of substrates. In some cases, this methodology is complemented by spectrophotometric techniques, such as HPLC-UV, to measure enzymatic activity. The corresponding kinetic data and experimental methodologies can be found in several freely accessible enzymatic databases, including BRENDA and SABIO-RK.

The second section summarized the computational strategies employed to model substrate affinity, through molecular docking, MD simulation and free-energy of binding analysis, and catalytic turnover numbers, through hybrid QM/MM simulation, for the studied CES-substrate complexes. These computational methodologies demonstrated high accuracy and provided results consistent with experimentally reported data.

The advantages of integrating experimental and computational approaches became clear in the examples discussed in the final section, where the combined use of both methodologies enabled the elucidation of structural factors underlying substrate selectivity toward specific CES isoforms.

This collaborative strategy should be more widely adopted to accelerate advances in understanding CES substrate selectivity for future compounds. Given the biological relevance of CES-mediated catalysis, further investigation into the structural determinants governing isoform specific substrate recognition remains a significant scientific challenge, offering valuable insights for the rational design of drugs and prodrugs with optimized biopharmaceutical properties. It is hoped that this article will serve as a useful resource to support continued exploration in this field.

Author Contributions: For research articles with several authors, a short paragraph specifying their individual contributions must be provided. The following statements should be used “Conceptualization, X.X. and Y.Y.; methodology, X.X.; software, X.X.; validation, X.X., Y.Y. and Z.Z.; formal analysis, X.X.; investigation, X.X.; resources, X.X.; data curation, X.X.; writing—original draft preparation, X.X.; writing—review and editing, X.X.; visualization, X.X.; supervision, X.X.; project administration, X.X.; funding acquisition, Y.Y. All authors have read and agreed to the published version of the manuscript.”, please turn to the [CRediT taxonomy](#) for the term explanation. Authorship must be limited to those who have contributed substantially to the work reported.

Funding: Please add: “This research received no external funding” or “This research was funded by NAME OF FUNDER grant number XXX.” and and “The APC was funded by XXX”. Check carefully that the details given are accurate and use the standard spelling of funding agency names at <https://search.crossref.org/funding>, any errors may affect your future funding.

Institutional Review Board Statement: In this section, you should add the Institutional Review Board Statement and approval number, if relevant to your study. You might choose to exclude this statement if the study did not require ethical approval. Please note that the Editorial Office might ask you for further information. Please add “The study was conducted in accordance with the Declaration of Helsinki, and approved by the Institutional Review Board (or Ethics Committee) of NAME OF INSTITUTE (protocol code XXX and date of approval).” for studies involving humans. OR “The animal study protocol was approved by the Institutional Review Board (or Ethics Committee) of NAME OF INSTITUTE (protocol code XXX and date of approval).” for studies involving animals. OR “Ethical review and approval were waived for this study due to REASON (please provide a detailed justification).” OR “Not applicable” for studies not involving humans or animals.

Informed Consent Statement: Any research article describing a study involving humans should contain this statement. Please add “Informed consent was obtained from all subjects involved in the study.” OR “Patient consent was waived due to REASON (please provide a detailed justification).” OR “Not applicable” for studies not involving humans. You might also choose to exclude this statement if the study did not involve humans.

Written informed consent for publication must be obtained from participating patients who can be identified (including by the patients themselves). Please state “Written informed consent has been obtained from the patient(s) to publish this paper” if applicable.

Data Availability Statement: We encourage all authors of articles published in MDPI journals to share their research data. In this section, please provide details regarding where data supporting reported results can be found, including links to publicly archived datasets analyzed or generated during the study. Where no new data were created, or where data is unavailable due to privacy or ethical restrictions, a statement is still required. Suggested Data Availability Statements are available in section “MDPI Research Data Policies” at <https://www.mdpi.com/ethics>.

Acknowledgments: In this section you can acknowledge any support given which is not covered by the author contribution or funding sections. This may include administrative and technical support, or donations in kind (e.g., materials used for experiments). Where GenAI has been used for purposes such as generating text, data, or graphics, or for study design, data collection, analysis, or interpretation of data, please add “During the preparation of this manuscript/study, the author(s) used [tool name, version information] for the purposes of [description of use]. The authors have reviewed and edited the output and take full responsibility for the content of this publication.”

Conflicts of Interest: Declare conflicts of interest or state “The authors declare no conflicts of interest.” Authors must identify and declare any personal circumstances or interest that may be perceived as inappropriately influencing the representation or interpretation of reported research results. Any role of the funders in the design of the study; in the collection, analyses or interpretation of data; in the writing of the manuscript; or in the decision to publish the results must be declared in this section. If there is no role, please state “The funders had no role in the design of the study; in the collection, analyses, or interpretation of data; in the writing of the manuscript; or in the decision to publish the results”.

Abbreviations

The following abbreviations are used in this manuscript:

MDPI	Multidisciplinary Digital Publishing Institute
DOAJ	Directory of open access journals
TLA	Three letter acronym
LD	Linear dichroism

References

- Wang, D.; Zou, L.; Jin, Q.; Hou, J.; Ge, G.; Yang, L. Human carboxylesterases: a comprehensive review. *Acta Pharmaceutica Sinica B* **2018**, *8*, 699–712. <https://doi.org/https://doi.org/10.1016/j.apsb.2018.05.005>.
- Fukami, T.; Yokoi, T. The emerging role of human esterases. *Drug metabolism and pharmacokinetics* **2012**, *27*, 466–477.
- Hosokawa, M. Structure and catalytic properties of carboxylesterase isozymes involved in metabolic activation of prodrugs. *Molecules* **2008**, *13*, 412–431.
- Casey Laizure, S.; Herring, V.; Hu, Z.; Witbrodt, K.; Parker, R.B. The role of human carboxylesterases in drug metabolism: have we overlooked their importance? *Pharmacotherapy: The Journal of Human Pharmacology and Drug Therapy* **2013**, *33*, 210–222.
- Dai, J.; Hou, Y.; Wu, J.; Shen, B. A minireview of recent reported carboxylesterase fluorescent probes: Design and biological applications. *ChemistrySelect* **2020**, *5*, 11185–11196.
- Xu, Y.; Zhang, C.; He, W.; Liu, D. Regulations of xenobiotics and endobiotics on carboxylesterases: a comprehensive review. *European journal of drug metabolism and pharmacokinetics* **2016**, *41*, 321–330.
- Di, L. The impact of carboxylesterases in drug metabolism and pharmacokinetics. *Current drug metabolism* **2019**, *20*, 91–102.
- Fukami, T.; Kariya, M.; Kurokawa, T.; Iida, A.; Nakajima, M. Comparison of substrate specificity among human arylacetamide deacetylase and carboxylesterases. *European Journal of Pharmaceutical Sciences* **2015**, *78*, 47–53. <https://doi.org/10.1016/j.ejps.2015.07.006>.
- Zhang, C.; Xu, Y.; Zhong, Q.; Li, X.; Gao, P.; Feng, C.; Chu, Q.; Chen, Y.; Liu, D. In Vitro Evaluation of the Inhibitory Potential of Pharmaceutical Excipients on Human Carboxylesterase 1A and 2. *PLOS ONE* **2014**, *9*, 1–8. <https://doi.org/10.1371/journal.pone.0093819>.
- Honda, S.; Fukami, T.; Tsujiguchi, T.; Zhang, Y.; Nakano, M.; Nakajima, M. Hydrolase activities of cynomolgus monkey liver microsomes and recombinant CES1, CES2, and AADAC. *European Journal of Pharmaceutical Sciences* **2021**, *161*, 105807.
- Yang, D.; Wang, X.; Chen, Y.t.; Deng, R.; Yan, B. Pyrethroid insecticides: isoform-dependent hydrolysis, induction of cytochrome P450 3A4 and evidence on the involvement of the pregnane X receptor. *Toxicology and applied pharmacology* **2009**, *237*, 49–58.
- Pratt, S.E.; Durland-Busby, S.; Shepard, R.L.; Heinz-Taheny, K.; Iversen, P.W.; Dantzig, A.H. Human carboxylesterase-2 hydrolyzes the prodrug of gemcitabine (LY2334737) and confers prodrug sensitivity to cancer cells. *Clinical Cancer Research* **2013**, *19*, 1159–1168.
- Bachovchin, D.A.; Cravatt, B.F. The pharmacological landscape and therapeutic potential of serine hydrolases. *Nature reviews Drug discovery* **2012**, *11*, 52–68.
- Zhang, M.m.; Li, P.; Hai, F.; Jia, Y. Determination of carboxylesterase 2 by fluorescence probe to guide pancreatic adenocarcinoma profiling. *Chemical Physics Letters* **2021**, *785*, 139143.
- Elkhanoufi, S.; Stefania, R.; Alberti, D.; Baroni, S.; Aime, S.; Geninatti Crich, S. Highly sensitive “off/on” EPR probes to monitor enzymatic activity. *Chemistry—A European Journal* **2022**, *28*, e202104563.
- Dai, J.; Zhao, Y.; Hou, Y.; Zhong, G.; Gao, R.; Wu, J.; Shen, B.; Zhang, X. Detection of carboxylesterase 1 and carbamates with a novel fluorescent protein chromophore based probe. *Dyes and Pigments* **2021**, *192*, 109444. <https://doi.org/https://doi.org/10.1016/j.dyepig.2021.109444>.
- Jia, Y.; Wang, J.; Li, P.; Ma, X.; Han, K. Directionally Modified Fluorophores for Super-Resolution Imaging of Target Enzymes: A Case Study with Carboxylesterases. *Journal of Medicinal Chemistry* **2021**, *64*, 16177–16186. <https://doi.org/10.1021/acs.jmedchem.1c01469>.
- Punekar, N.S. *Enzymes: catalysis, kinetics and mechanisms*; Vol. 10, Springer, 2025.
- Van Der Kamp, M.W.; Mulholland, A.J. Combined quantum mechanics/molecular mechanics (QM/MM) methods in computational enzymology. *Biochemistry* **2013**, *52*, 2708–2728.
- Lonsdale, R.; Ranaghan, K.E.; Mulholland, A.J. Computational enzymology. *Chemical communications* **2010**, *46*, 2354–2372.
- Lodola, A.; Mulholland, A.J. Computational enzymology. In *Biomolecular Simulations: Methods and Protocols*; Springer, 2012; pp. 67–89.
- Michaelis, L.; Menten, M.L.; et al. Die kinetik der invertinwirkung. *Biochem. z* **1913**, *49*, 352.
- Nishi, K.; Huang, H.; Kamita, S.G.; Kim, I.H.; Morisseau, C.; Hammock, B.D. Characterization of pyrethroid hydrolysis by the human liver carboxylesterases hCE-1 and hCE-2. *Archives of biochemistry and biophysics* **2006**, *445*, 115–123.
- Ribone, S.R.; Estrin, D.A.; Quevedo, M.A. Exploring human carboxylesterases 1 and 2 selectivity of two families of substrates at an atomistic level. *Biochimica et Biophysica Acta (BBA)-Proteins and Proteomics* **2025**, *1873*, 141069.
- Humerickhouse, R.; Lohrbach, K.; Li, L.; Bosron, W.F.; Dolan, M.E. Characterization of CPT-11 hydrolysis by human liver carboxylesterase isoforms hCE-1 and hCE-2. *Cancer research* **2000**, *60*, 1189–1192.
- Sanghani, S.P.; Quinney, S.K.; Fredenburg, T.B.; Davis, W.I.; Murry, D.J.; Bosron, W.F. HYDROLYSIS OF IRINOTECAN AND ITS OXIDATIVE METABOLITES, 7-ETHYL-10-[4-N-(5-AMINOPENTANOIC ACID)-1-PIPERIDINO] CARBOXYLOXY-CAMPTOTHECIN AND 7-ETHYL-10-[4-(1-PIPERIDINO)-1-AMINO]-CARBOXYLOXYCAMPTOTHECIN, BY HUMAN CAR-

- BOXYLESTERASES CES1A1, CES2, AND A NEWLY EXPRESSED CARBOXYLESTERASE ISOENZYME, CES3. *Drug Metabolism and Disposition* **2004**, 32, 505–511. <https://doi.org/https://doi.org/10.1124/dmd.32.5.505>.
27. Sun, Z.; Murry, D.J.; Sanghani, S.P.; Davis, W.I.; Kedishvili, N.Y.; Zou, Q.; Hurley, T.D.; Bosron, W.F. Methylphenidate Is Stereoselectively Hydrolyzed by Human Carboxylesterase CES1A1. *The Journal of Pharmacology and Experimental Therapeutics* **2004**, 310, 469–476. <https://doi.org/https://doi.org/10.1124/jpet.104.067116>.
 28. Shi, D.; Yang, J.; Yang, D.; LeCluyse, E.L.; Black, C.; You, L.; Akhlaghi, F.; Yan, B. Anti-Influenza Prodrug Oseltamivir Is Activated by Carboxylesterase Human Carboxylesterase 1, and the Activation Is Inhibited by Antiplatelet Agent Clopidogrel. *The Journal of Pharmacology and Experimental Therapeutics* **2006**, 319, 1477–1484. <https://doi.org/https://doi.org/10.1124/jpet.106.111807>.
 29. Imai, T.; Taketani, M.; Shii, M.; Hosokawa, M.; Chiba, K. Substrate specificity of carboxylesterase isozymes and their contribution to hydrolase activity in human liver and small intestine. *Drug metabolism and disposition* **2006**, 34, 1734–1741.
 30. Tang, M.; Mukundan, M.; Yang, J.; Charpentier, N.; LeCluyse, E.L.; Black, C.; Yang, D.; Shi, D.; Yan, B. Antiplatelet Agents Aspirin and Clopidogrel Are Hydrolyzed by Distinct Carboxylesterases, and Clopidogrel Is Transesterified in the Presence of Ethyl Alcohol. *The Journal of Pharmacology and Experimental Therapeutics* **2006**, 319, 1467–1476. <https://doi.org/https://doi.org/10.1124/jpet.106.110577>.
 31. Watanabe, A.; Fukami, T.; Nakajima, M.; Takamiya, M.; Aoki, Y.; Yokoi, T. Human arylacetamide deacetylase is a principal enzyme in flutamide hydrolysis. *Drug metabolism and disposition* **2009**, 37, 1513–1520.
 32. Kobayashi, Y.; Fukami, T.; Shimizu, M.; Nakajima, M.; Yokoi, T. Contributions of Arylacetamide Deacetylase and Carboxylesterase 2 to Flutamide Hydrolysis in Human Liver. *Drug Metabolism and Disposition* **2012**, 40, 1080–1084. <https://doi.org/10.1124/dmd.112.044537>.
 33. Ross, M.K.; Borazjani, A.; Edwards, C.C.; Potter, P.M. Hydrolytic metabolism of pyrethroids by human and other mammalian carboxylesterases. *Biochemical pharmacology* **2006**, 71, 657–669.
 34. Wang, J.; Williams, E.T.; Bourgea, J.; Wong, Y.N.; Patten, C.J. Characterization of recombinant human carboxylesterases: fluorescein diacetate as a probe substrate for human carboxylesterase 2. *Drug metabolism and disposition* **2011**, 39, 1329–1333.
 35. Williams, E.T.; Jones, K.O.; Ponsler, G.D.; Lowery, S.M.; Perkins, E.J.; Wrighton, S.A.; Ruterbories, K.J.; Kazui, M.; Farid, N.A. The biotransformation of prasugrel, a new thienopyridine prodrug, by the human carboxylesterases 1 and 2. *Drug Metabolism and Disposition* **2008**, 36, 1227–1232.
 36. Hatfield, M.; Tsurkan, L.; Hyatt, J.; Yu, X.; Edwards, C.; Hicks, L.; Wadkins, R.; Potter, P. Biochemical and molecular analysis of carboxylesterase-mediated hydrolysis of cocaine and heroin. *British journal of pharmacology* **2010**, 160, 1916–1928.
 37. Yao, J.; Chen, X.; Zheng, F.; Zhan, C.G. Catalytic reaction mechanism for drug metabolism in human carboxylesterase-1: cocaine hydrolysis pathway. *Molecular pharmaceutics* **2018**, 15, 3871–3880.
 38. Sato, Y.; Miyashita, A.; Iwatsubo, T.; Usui, T. Conclusive identification of the oxybutynin-hydrolyzing enzyme in human liver. *Drug Metabolism and Disposition* **2012**, 40, 902–906.
 39. Higuchi, R.; Fukami, T.; Nakajima, M.; Yokoi, T. Prilocaine- and Lidocaine-Induced Methemoglobinemia Is Caused by Human Carboxylesterase-, CYP2E1-, and CYP3A4-Mediated Metabolic Activation. *Drug Metabolism and Disposition* **2013**, 41, 1220–1230. <https://doi.org/10.1124/dmd.113.051714>.
 40. Li, H.X.; Sun, M.R.; Zhang, Y.; Song, L.L.; Zhang, F.; Song, Y.Q.; Hou, X.D.; Ge, G.B. Human Carboxylesterase 1A Plays a Predominant Role in Hydrolysis of the Anti-Dyslipidemia Agent Fenofibrate in Humans. *Drug Metabolism and Disposition* **2023**, 51, 1490–1498. <https://doi.org/https://doi.org/10.1124/dmd.123.001365>.
 41. Thomsen, R.; Rasmussen, H.B.; Linnet, K. In Vitro Drug Metabolism by Human Carboxylesterase 1: Focus on Angiotensin-Converting Enzyme Inhibitors. *Drug Metabolism and Disposition* **2014**, 42, 126–133. <https://doi.org/https://doi.org/10.1124/dmd.113.053512>.
 42. Shi, J.; Wang, X.; Nguyen, J.; Wu, A.H.; Bleske, B.E.; Zhu, H.J. Sacubitril Is Selectively Activated by Carboxylesterase 1 (CES1) in the Liver and the Activation Is Affected by CES1 Genetic Variation. *Drug Metabolism and Disposition* **2016**, 44, 554–559. <https://doi.org/https://doi.org/10.1124/dmd.115.068536>.
 43. Jiang, J.; Chen, X.; Zhong, D. Predominant contributions of carboxylesterase 1 and 2 in hydrolysis of anordrin in humans. *Xenobiotica* **2018**, 48, 533–540.
 44. Morton, C.L.; Potter, P.M. Comparison of Escherichia coli, Saccharomyces cerevisiae, Pichia pastoris, Spodoptera frugiperda, and COS7 cells for recombinant gene expression: Application to a rabbit liver carboxylesterase. *Molecular biotechnology* **2000**, 16, 193–202.
 45. Mizoi, K.; Takahashi, M.; Haba, M.; Hosokawa, M. Synthesis and evaluation of atorvastatin esters as prodrugs metabolically activated by human carboxylesterases. *Bioorganic & Medicinal Chemistry Letters* **2016**, 26, 921–923.
 46. Takahashi, M.; Sakai, S.; Takahashi, K.; Hosokawa, M. Species Differences in Carboxylesterases Among Humans, Cynomolgus Monkeys, and Mice in the Hydrolysis of Atorvastatin Derivatives. *Biopharmaceutics & Drug Disposition* **2025**, 46, 49–57.

47. Takahashi, M.; Ogawa, T.; Kashiwagi, H.; Fukushima, F.; Yoshitsugu, M.; Haba, M.; Hosokawa, M. Chemical synthesis of an indomethacin ester prodrug and its metabolic activation by human carboxylesterase 1. *Bioorganic & medicinal chemistry letters* **2018**, *28*, 997–1000. 742–744
48. Takahashi, M.; Takani, D.; Haba, M.; Hosokawa, M. Investigation of the chiral recognition ability of human carboxylesterase 1 using indomethacin esters. *Chirality* **2020**, *32*, 73–80. 745–746
49. Takahashi, M.; Uehara, T.; Nonaka, M.; Minagawa, Y.; Yamazaki, R.; Haba, M.; Hosokawa, M. Synthesis and evaluation of haloperidol ester prodrugs metabolically activated by human carboxylesterase. *European Journal of Pharmaceutical Sciences* **2019**, *132*, 125–131. 747–749
50. Wadkins, R.M.; Morton, C.L.; Weeks, J.K.; Oliver, L.; Wierdl, M.; Danks, M.K.; Potter, P.M. Structural constraints affect the metabolism of 7-ethyl-10-[4-(1-piperidino)-1-piperidino] carbonyloxycamptothecin (CPT-11) by carboxylesterases. *Molecular pharmacology* **2001**, *60*, 355–362. 750–752
51. Boonyuen, U.; Promnares, K.; Junkree, S.; Day, N.P.; Imwong, M. Efficient in vitro refolding and functional characterization of recombinant human liver carboxylesterase (CES1) expressed in *E. coli*. *Protein Expression and Purification* **2015**, *107*, 68–75. 753–754
52. Ding, L.; Tian, Z.; Hou, J.; Dou, T.; Jin, Q.; Wang, D.; Zou, L.; Zhu, Y.; Song, Y.; Cui, J.; et al. Sensing carboxylesterase 1 in living systems by a practical and isoform-specific fluorescent probe. *Chinese Chemical Letters* **2019**, *30*, 558–562. 755–756
53. Takahashi, M.; Hirota, I.; Nakano, T.; Kotani, T.; Takani, D.; Shiratori, K.; Choi, Y.; Haba, M.; Hosokawa, M. Effects of steric hindrance and electron density of ester prodrugs on controlling the metabolic activation by human carboxylesterase. *Drug Metabolism and Pharmacokinetics* **2021**, *38*, 100391. <https://doi.org/10.1016/j.dmpk.2021.100391>. 757–759
54. Quinney, S.; Sanghani, S.; Davis, W.; Hurley, T.; Sun, Z.; Murry, D.; Bosron, W. Hydrolysis of Capecitabine to 5-Deoxy-5-fluorocytidine by Human Carboxylesterases and Inhibition by Loperamide. *The Journal of Pharmacology and Experimental Therapeutics* **2005**, *313*, 1011–1016. <https://doi.org/10.1124/jpet.104.081265>. 760–762
55. Schomburg, I.; Jeske, L.; Ulbrich, M.; Placzek, S.; Chang, A.; Schomburg, D. The BRENDA enzyme information system—From a database to an expert system. *Journal of biotechnology* **2017**, *261*, 194–206. 763–764
56. Wittig, U.; Kania, R.; Bittkowski, M.; Wetsch, E.; Shi, L.; Jong, L.; Golebiewski, M.; Rey, M.; Weidemann, A.; Rojas, I.; et al. Data extraction for the reaction kinetics database SABIO-RK. *Perspectives in Science* **2014**, *1*, 33–40. 765–766
57. Bencharit, S.; Morton, C.L.; Xue, Y.; Potter, P.M.; Redinbo, M.R. Structural basis of heroin and cocaine metabolism by a promiscuous human drug-processing enzyme. *Nature Structural & Molecular Biology* **2003**, *10*, 349–356. 767–768
58. Bencharit, S.; Morton, C.L.; Hyatt, J.L.; Kuhn, P.; Danks, M.K.; Potter, P.M.; Redinbo, M.R. Crystal structure of human carboxylesterase 1 complexed with the Alzheimer's drug tacrine: from binding promiscuity to selective inhibition. *Chemistry & Biology* **2003**, *10*, 341–349. 769–771
59. Fleming, C.D.; Bencharit, S.; Edwards, C.C.; Hyatt, J.L.; Tsurkan, L.; Bai, F.; Fraga, C.; Morton, C.L.; Howard-Williams, E.L.; Potter, P.M.; et al. Structural insights into drug processing by human carboxylesterase 1: tamoxifen, mevastatin, and inhibition by benzil. *Journal of molecular biology* **2005**, *352*, 165–177. 772–774
60. Bencharit, S.; Edwards, C.C.; Morton, C.L.; Howard-Williams, E.L.; Kuhn, P.; Potter, P.M.; Redinbo, M.R. Multisite promiscuity in the processing of endogenous substrates by human carboxylesterase 1. *Journal of molecular biology* **2006**, *363*, 201–214. 775–776
61. Fleming, C.D.; Edwards, C.C.; Kirby, S.D.; Maxwell, D.M.; Potter, P.M.; Cerasoli, D.M.; Redinbo, M.R. Crystal structures of human carboxylesterase 1 in covalent complexes with the chemical warfare agents soman and tabun. *Biochemistry* **2007**, *46*, 5063–5071. 777–778
62. Hemmert, A.C.; Otto, T.C.; Wierdl, M.; Edwards, C.C.; Fleming, C.D.; MacDonald, M.; Cashman, J.R.; Potter, P.M.; Cerasoli, D.M.; Redinbo, M.R. Human carboxylesterase 1 stereoselectively binds the nerve agent cyclosarin and spontaneously hydrolyzes the nerve agent sarin. *Molecular pharmacology* **2010**, *77*, 508–516. 779–781
63. Greenblatt, H.M.; Otto, T.C.; Kirkpatrick, M.G.; Kovaleva, E.; Brown, S.; Buchman, G.; Cerasoli, D.M.; Sussman, J.L. Structure of recombinant human carboxylesterase 1 isolated from whole cabbage looper larvae. *Structural Biology and Crystallization Communications* **2012**, *68*, 269–272. 782–784
64. Arena de Souza, V.; Scott, D.J.; Nettleship, J.E.; Rahman, N.; Charlton, M.H.; Walsh, M.A.; Owens, R.J. Comparison of the structure and activity of glycosylated and aglycosylated human carboxylesterase 1. *PLoS One* **2015**, *10*, e0143919. 785–786
65. Su, C.C.; Lyu, M.; Zhang, Z.; Miyagi, M.; Huang, W.; Taylor, D.J.; Yu, E.W. High-resolution structural-omics of human liver enzymes. *Cell reports* **2023**, *42*. 787–788
66. Gai, C.; Zhang, Y.; Zhang, S.; Hu, X.; Song, Y.Q.; Zhuang, X.; Chai, X.; Zou, Y.; Ge, G.B.; Zhao, Q. The study of halogen effect on the reactivity of the serine-targeting covalent warheads. *Frontiers in Chemistry* **2024**, *12*, 1504453. 789–790
67. Waterhouse, A.; Bertoni, M.; Bienert, S.; Studer, G.; Tauriello, G.; Gumienny, R.; Heer, F.T.; de Beer, T.A.P.; Rempfer, C.; Bordoli, L.; et al. SWISS-MODEL: homology modelling of protein structures and complexes. *Nucleic acids research* **2018**, *46*, W296–W303. 791–792
68. Vistoli, G.; Pedretti, A.; Mazzolari, A.; Testa, B. Homology modeling and metabolism prediction of human carboxylesterase-2 using docking analyses by GriDock: a parallelized tool based on AutoDock 4.0. *Journal of computer-aided molecular design* **2010**, *24*, 771–787. 793–795

69. Zou, L.W.; Li, Y.G.; Wang, P.; Zhou, K.; Hou, J.; Jin, Q.; Hao, D.C.; Ge, G.B.; Yang, L. Design, synthesis, and structure-activity relationship study of glycyrrhetic acid derivatives as potent and selective inhibitors against human carboxylesterase 2. *European Journal of Medicinal Chemistry* **2016**, *112*, 280–288. <https://doi.org/10.1016/j.ejmech.2016.02.020>.
70. Song, Y.Q.; Guan, X.Q.; Weng, Z.M.; Wang, Y.Q.; Chen, J.; Jin, Q.; Fang, S.Q.; Fan, B.; Cao, Y.F.; Hou, J.; et al. Discovery of a highly specific and efficacious inhibitor of human carboxylesterase 2 by large-scale screening. *International Journal of Biological Macromolecules* **2019**, *137*, 261–269. <https://doi.org/10.1016/j.ijbiomac.2019.06.235>.
71. Qian, X.K.; Zhang, J.; Song, P.F.; Zhao, Y.S.; Ma, H.Y.; Jin, Q.; Wang, D.D.; Guan, X.Q.; Li, S.Y.; Bao, X.; et al. Discovery of pyrazolones as novel carboxylesterase 2 inhibitors that potently inhibit the adipogenesis in cells. *Bioorganic & Medicinal Chemistry* **2021**, *40*, 116187.
72. Lv, C.S.; Wang, B.; Zheng, Y.Z.; Wang, Y.H.; Zhang, J.; Jiang, Z.J.; Wang, J.G.; Zou, L.W. Discovery of cardiovascular drugs as effective inhibitors of human carboxylesterase. *International Journal of Biological Macromolecules* **2025**, p. 143967.
73. Webb, B.; Sali, A. Protein structure modeling with MODELLER. In *Functional genomics: Methods and protocols*; Springer, 2017; pp. 39–54.
74. Wang, Y.L.; Dong, P.P.; Liang, J.H.; Li, N.; Sun, C.P.; Tian, X.G.; Huo, X.K.; Zhang, B.J.; Ma, X.C.; Lv, C.Z. Phytochemical constituents from *Uncaria rhynchophylla* in human carboxylesterase 2 inhibition: kinetics and interaction mechanism merged with docking simulations. *Phytomedicine* **2018**, *51*, 120–127.
75. Choudhary, S.; Silakari, O. hCES1 and hCES2 mediated activation of epalrestat-antioxidant mutual prodrugs: Unwinding the hydrolytic mechanism using in silico approaches. *Journal of Molecular Graphics and Modelling* **2019**, *91*, 148–163.
76. Jumper, J.; Evans, R.; Pritzel, A.; Green, T.; Figurnov, M.; Ronneberger, O.; Tunyasuvunakool, K.; Bates, R.; Žídek, A.; Potapenko, A.; et al. Highly accurate protein structure prediction with AlphaFold. *nature* **2021**, *596*, 583–589.
77. Vistoli, G.; Pedretti, A.; Mazzolari, A.; Testa, B. In silico prediction of human carboxylesterase-1 (hCES1) metabolism combining docking analyses and MD simulations. *Bioorganic & medicinal chemistry* **2010**, *18*, 320–329.
78. Yang, Z.; Cao, Z.; Wang, W.; Chen, Y.; Huang, W.; Jiao, S.; Chen, S.; Chen, L.; Liu, Y.; Mao, J.; et al. Design, synthesis, and biological evaluation studies of novel carboxylesterase 2 inhibitors for the treatment of irinotecan-induced delayed diarrhea. *Bioorganic Chemistry* **2023**, *138*, 106625.

Disclaimer/Publisher's Note: The statements, opinions and data contained in all publications are solely those of the individual author(s) and contributor(s) and not of MDPI and/or the editor(s). MDPI and/or the editor(s) disclaim responsibility for any injury to people or property resulting from any ideas, methods, instructions or products referred to in the content.